Other Stable Particle Searches

OMITTED FROM SUMMARY TABLE
MINIREVIEW GOES HERE

Concentration of Heavy (Charge +1) Stable Particles in Matter

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<4\times10^{-17}$	95	[1] YAMAGATA	93	SPEC	Deep sea water, m=5–1600m_p
$<6\times10^{-15}$	95	[2] VERKERK	92	SPEC	Water, m= 10^5 to 3×10^7 GeV
$<7\times10^{-15}$	95	[2] VERKERK	92	SPEC	Water, m= 10^4, 6×10^7 GeV
$<9\times10^{-15}$	95	[2] VERKERK	92	SPEC	Water, m= 10^8 GeV
$<3\times10^{-23}$	90	[3] HEMMICK	90	SPEC	Water, $m=1000m_p$
$<2\times10^{-21}$	90	[3] HEMMICK	90	SPEC	Water, $m=5000m_p$
$<3\times10^{-20}$	90	[3] HEMMICK	90	SPEC	Water, $m=10000m_p$
$<1.\times10^{-29}$		SMITH	82B	SPEC	Water, m=30–400m_p
$<2.\times10^{-28}$		SMITH	82B	SPEC	Water, m=12–1000m_p
$<1.\times10^{-14}$		SMITH	82B	SPEC	Water, $m>1000\ m_p$
$<(0.2–1.)\times10^{-21}$		SMITH	79	SPEC	Water, m=6–350 m_p

[1] YAMAGATA 93 used deep sea water at 4000 m since the concentration is enhanced in deep sea due to gravity.

[2] VERKERK 92 looked for heavy isotopes in sea water and put a bound on concentration of stable charged massive particle in sea water. The above bound can be translated into into a bound on charged dark matter particle (5×10^6 GeV), assuming the local density, ρ=0.3 GeV/cm^3, and the mean velocity $\langle v\rangle$=300 km/s.

[3] See HEMMICK 90 Fig. 7 for other masses 100–10000 m_p.

Concentration of Heavy (Charge −1) Stable Particles

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<4\times10^{-20}$	90	[4] HEMMICK	90	SPEC	C, $M=100m_p$
$<8\times10^{-20}$	90	[4] HEMMICK	90	SPEC	C, $M=1000m_p$
$<2\times10^{-16}$	90	[4] HEMMICK	90	SPEC	C, $M=10000m_p$
$<6\times10^{-13}$	90	[4] HEMMICK	90	SPEC	Li, $M=1000m_p$
$<1\times10^{-11}$	90	[4] HEMMICK	90	SPEC	Be, $M=1000m_p$
$<6\times10^{-14}$	90	[4] HEMMICK	90	SPEC	B, $M=1000m_p$
$<4\times10^{-17}$	90	[4] HEMMICK	90	SPEC	O, $M=1000m_p$
$<4\times10^{-15}$	90	[4] HEMMICK	90	SPEC	F, $M=1000m_p$
$<1.5\times10^{-13}$/nucleon	68	[5] NORMAN	89	SPEC	^{206}Pb X^-
$<1.2\times10^{-12}$/nucleon	68	[5] NORMAN	87	SPEC	^{56,58}Fe X^-

[4] See HEMMICK 90 Fig. 7 for other masses 100–10000 m_p.

[5] Bound valid up to $m_{X^-}\sim$ 100 TeV.

Light (between μ and e Masses) Particle MASS

VALUE (m_e)	EVTS	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
none 110–180	0	[6] VIERTEL	78	CNTR	$\tau>2.\times10^{-5}$ s
none 2–13	0	[7] BLAGOV	75	CNTR	Spinor, $\tau>2\times10^{-10}$ s
none 2–10.6	0	[7] BLAGOV	75	CNTR	Scalar, $\tau>2\times10^{-10}$ s
none 5–175	0	COWARD	63	CNTR	Spinor, $\tau>22\times10^{-10}$
none 5–175	0	COWARD	63	CNTR	Scalar, $\tau>68\times10^{-10}$
none 6–25	0	BELOUSOV	60	CNTR	Spinor, $\tau>1\times10^{-8}$
none 2–25	0	GORBUNOV	60	CC	Spinor, $\tau>1\times10^{-9}$

[6] VIERTEL 78 searches for $\mu^+\rightarrow X^+\nu$. Finds BR $<8.5\times10^{-6}$ in mass range given above (CL = 90%). Best limit BR $<5.\times10^{-7}$ (CL = 90%) is found at mass = 80 MeV.

[7] BLAGOV 75 bounds on lifetime depend on mass and improve as mass decreases. At 2 GeV the experiment is sensitive to $\tau > 3 \times 10^{-11}$ s for spinor, $\tau > 5 \times 10^{-11}$ s for scalar.

Highly Ionizing Particle Flux

VALUE ($m^{-2}yr^{-1}$)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.4	95	0	KINOSHITA	81B PLAS	Z/β 30–100

Production of New Penetrating Non-ν Like States in Beam Dump

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
	[8] LOSECCO	81 CALO	28 GeV protons

[8] No excess neutral-current events leads to σ(production) $\times$ σ(interaction)$\times$acceptance $< 2.26 \times 10^{-71}$ cm^4/nucleon^2 (CL = 90%) for light neutrals. Acceptance depends on models (0.1 to 4. $\times 10^{-4}$).

Branching Fraction of Z^0 to a Pair of Stable Charged Heavy Fermions

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<1 \times 10^{-3}$	95	AKRAWY	90O OPAL	m = 29–40 GeV

Heavy Particle Production Cross Section in $e^+ e^-$

Ratio to $\sigma(e^+ e^- \to \mu^+ \mu^-)$. See also entries in Free Quark Search and Magnetic Monopole Searches.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<2 \times 10^{-3}$	90		[9] BUSKULIC	93C ALEP	Q=1, m=32–72 GeV
$<(10^{-2}$–1)	95		[10] ADACHI	90C TOPZ	Q = 1, m = 1–16, 18–27 GeV
$<7 \times 10^{-2}$	90		[11] ADACHI	90E TOPZ	Q = 1, m = 5–25 GeV
$<1.6 \times 10^{-2}$	95	0	[12] KINOSHITA	82 PLAS	Q=3–180, m <14.5 GeV
$<5.0 \times 10^{-2}$	90	0	[13] BARTEL	80 JADE	Q=(3,4,5)/3 2–12 GeV

[9] BUSKULIC 93C is a CERN-LEP experiment with $W_{cm} = m_Z$. The limit is for a pair or single production of heavy particles with unusual ionization loss in TPC. See their Fig. 5 and Table 1.

[10] ADACHI 90C is a KEK-TRISTAN experiment with W_{cm} = 52–60 GeV. The limit is for pair production of a scalar or spin-1/2 particle. See Figs. 3 and 4.

[11] ADACHI 90E is KEK-TRISTAN experiment with W_{cm} = 52–61.4 GeV. The above limit is for inclusive production cross section normalized to $\sigma(e^+ e^- \to \mu^+ \mu^-)\cdot\beta(3-\beta^2)/2$, where $\beta = (1 - 4m^2/W_{cm}^2)^{1/2}$. See the paper for the assumption about the production mechanism.

[12] KINOSHITA 82 is SLAC PEP experiment at W_{cm} = 29 GeV using lexan and ^{39}Cr plastic sheets sensitive to highly ionizing particles.

[13] BARTEL 80 is DESY-PETRA experiment with W_{cm} = 27–35 GeV. Above limit is for inclusive pair production and ranges between 1. $\times 10^{-1}$ and 1. $\times 10^{-2}$ depending on mass and production momentum distributions. (See their figures 9, 10, 11).

Heavy Particle Production Cross Section in $p\overline{p}$

Limits are for a particle decaying to two hadronic jets.

Units(pb)	CL%	Mass(GeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<2603	95	200	[14] ABE	93G CDF	1.8 TeV $p\overline{p} \to$ 2jets
< 44	95	400	[14] ABE	93G CDF	1.8 TeV $p\overline{p} \to$ 2jets
< 7	95	600	[14] ABE	93G CDF	1.8 TeV $p\overline{p} \to$ 2jets

[14] ABE 93G gives cross section times branching ratio into light (d, u, s, c, b) quarks for Γ = 0.02 M. Their Table II gives limits for M = 200–900 GeV and Γ = (0.02–0.2) M.

Heavy Particle Production Cross Section

VALUE (nb)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					

<0.05	95		[15] ABE	92J	CDF	m=50–200 GeV
<30–130			[16] CARROLL	78	SPEC	m=2–2.5 GeV
<100		0	[17] LEIPUNER	73	CNTR	m=3–11 GeV

[15] ABE 92J look for pair production of unit-charged particles which leave detector before decaying. Limit shown here is for m=50 GeV. See their Fig. 5 for different charges and stronger limits for higher mass.

[16] CARROLL 78 look for neutral, $S = -2$ dihyperon resonance in $pp \rightarrow 2K^+X$. Cross section varies within above limits over mass range and p_{lab} = 5.1–5.9 GeV/c.

[17] LEIPUNER 73 is an NAL 300 GeV p experiment. Would have detected particles with lifetime greater than 200 ns.

Heavy Particle Production Cross Section

VALUE (cm^2/N)	CL%	EVTS	DOCUMENT ID		TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •							
$<(4–0.3)\times10^{-31}$	95		[18] AKESSON	91	CNTR	0	m = 0–5 GeV
$<2.5\times10^{-35}$		0	[19] GUSTAFSON	76	CNTR	0	$\tau > 10^{-7}$ s

[18] AKESSON 91 limit is from weakly interacting neutral long-lived particles produced in pN reaction at 450 GeV/c performed at CERN SPS. Bourquin-Gaillard formula is used as the production model. The above limit is for $\tau > 10^{-7}$ s. For $\tau > 10^{-9}$ s, $\sigma < 10^{-30}$ cm^{-2}/nucleon is obtained.

[19] GUSTAFSON 76 is a 300 GeV FNAL experiment looking for heavy (m >2 GeV) long-lived neutral hadrons in the M4 neutral beam. The above typical value is for m = 3 GeV and assumes an interaction cross section of 1 mb. Values as a function of mass and interaction cross section are given in figure 2.

Heavy Particle Production Differential Cross Section

VALUE ($cm^2 sr^{-1} GeV^{-1}$)	CL%	EVTS	DOCUMENT ID		TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •							
$<2.6\times10^{-36}$	90	0	[20] BALDIN	76	CNTR	–	Q= 1, m=2.1–9.4 GeV
$<2.2\times10^{-33}$	90	0	[21] ALBROW	75	SPEC	±	Q= ±1, m=4–15 GeV
$<1.1\times10^{-33}$	90	0	[21] ALBROW	75	SPEC	±	Q= ±2, m=6–27 GeV
$<8.\times10^{-35}$	90	0	[22] JOVANOV...	75	CNTR	±	m=15–26 GeV
$<1.5\times10^{-34}$	90	0	[22] JOVANOV...	75	CNTR	±	Q= ±2, m=3–10 GeV
$<6.\times10^{-35}$	90	0	[22] JOVANOV...	75	CNTR	±	Q= ±2, m=10–26 GeV
$<1.\times10^{-31}$	90	0	[23] APPEL	74	CNTR	±	m=3.2–7.2 GeV
$<5.8\times10^{-34}$	90	0	[24] ALPER	73	SPEC	±	m=1.5–24 GeV
$<1.2\times10^{-35}$	90	0	[25] ANTIPOV	71B	CNTR	–	Q=–, m=2.2–2.8
$<2.4\times10^{-35}$	90	0	[26] ANTIPOV	71C	CNTR	–	Q=–, m=1.2–1.7, 2.1–4
$<2.4\times10^{-35}$	90	0	BINON	69	CNTR	–	Q=–, m=1–1.8 GeV
$<1.5\times10^{-36}$		0	[27] DORFAN	65	CNTR		Be target m=3–7 GeV
$<3.0\times10^{-36}$		0	[27] DORFAN	65	CNTR		Fe target m=3–7 GeV

[20] BALDIN 76 is a 70 GeV Serpukhov experiment. Value is per Al nucleus at $\theta = 0$. For other charges in range -0.5 to -3.0, CL = 90% limit is $(2.6\times10^{-36})/|(\text{charge})|$ for mass range (2.1–9.4 GeV) $\times$ |(charge)|. Assumes stable particle interacting with matter as do antiprotons.

[21] ALBROW 75 is a CERN ISR experiment with E_{cm} = 53 GeV. θ = 40 mr. See figure 5 for mass ranges up to 35 GeV.

[22] JOVANOVICH 75 is a CERN ISR 26+26 and 15+15 GeV pp experiment. Figure 4 covers ranges Q = 1/3 to 2 and m = 3 to 26 GeV. Value is per GeV momentum.

[23] APPEL 74 is NAL 300 GeV pW experiment. Studies forward production of heavy (up to 24 GeV) charged particles with momenta 24–200 GeV (−charge) and 40–150 GeV (+charge). Above typical value is for 75 GeV and is per GeV momentum per nucleon.

[24] ALPER 73 is CERN ISR 26+26 GeV pp experiment. p >0.9 GeV, 0.2 < β <0.65.

[25] ANTIPOV 71B is from same 70 GeV p experiment as ANTIPOV 71C and BINON 69.

[26] ANTIPOV 71C limit inferred from flux ratio. 70 GeV p experiment.

[27] DORFAN 65 is a 30 GeV/c p experiment at BNL. Units are per GeV momentum per nucleus.

Long-Lived Heavy Particle Invariant Cross Section

VALUE ($cm^2/GeV^2/N$)	CL%	EVTS	DOCUMENT ID		TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •							
$< 5 \times 10^{-35}$–7×10^{-33}	90	0	[28] BERNSTEIN	88	CNTR		
$< 5 \times 10^{-37}$–7×10^{-35}	90	0	[28] BERNSTEIN	88	CNTR		
$<2.5 \times 10^{-36}$	90	0	[29] THRON	85	CNTR	−	Q= 1, m=4–12 GeV
$<1. \times 10^{-35}$	90	1	[29] THRON	85	CNTR	+	Q= 1, m=4–12 GeV
$<6. \times 10^{-33}$	90	0	[30] ARMITAGE	79	SPEC		m=1.87 GeV
$<1.5 \times 10^{-33}$	90	0	[30] ARMITAGE	79	SPEC		m=1.5–3.0 GeV
		0	[31] BOZZOLI	79	CNTR	±	Q = (2/3, 1, 4/3, 2)
$<1.1 \times 10^{-37}$	90	0	[32] CUTTS	78	CNTR		m=4–10 GeV
$<3.0 \times 10^{-37}$	90	0	[33] VIDAL	78	CNTR		m=4.5–6 GeV

[28] BERNSTEIN 88 limits apply at $x = 0.2$ and $p_T = 0$. Mass and lifetime dependence of limits are shown in the regions: m = 1.5–7.5 GeV and $\tau = 10^{-8}$–2×10^{-6} s. First number is for hadrons; second is for weakly interacting particles.

[29] THRON 85 is FNAL 400 GeV proton experiment. Mass determined from measured velocity and momentum. Limits are for $\tau > 3 \times 10^{-9}$ s.

[30] ARMITAGE 79 is CERN-ISR experiment at $E_{cm} = 53$ GeV. Value is for $x = 0.1$ and $p_T = 0.15$. Observed particles at $m = 1.87$ GeV are found all consistent with being antideuterons.

[31] BOZZOLI 79 is CERN-SPS 200 GeV pN experiment. Looks for particle with τ larger than 10^{-8} s. See their figure 11–18 for production cross-section upper limits vs mass.

[32] CUTTS 78 is pBe experiment at FNAL sensitive to particles of $\tau > 5 \times 10^{-8}$ s. Value is for $-0.3 < x < 0$ and $p_T = 0.175$.

[33] VIDAL 78 is FNAL 400 GeV proton experiment. Value is for $x = 0$ and $p_T = 0$. Puts lifetime limit of $< 5 \times 10^{-8}$ s on particle in this mass range.

Long-Lived Heavy Particle Production (σ(Heavy Particle) / $\sigma(\pi)$)

VALUE	EVTS	DOCUMENT ID		TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$<10^{-8}$		[34] NAKAMURA	89	SPEC	±	Q= (−5/3,±2)
	0	[35] BUSSIERE	80	CNTR	±	Q= (2/3,1,4/3,2)

[34] NAKAMURA 89 is KEK experiment with 12 GeV protons on Pt target. The limit applies for mass $\lesssim$ 1.6 GeV and lifetime $\gtrsim 10^{-7}$ s.

[35] BUSSIERE 80 is CERN-SPS experiment with 200–240 GeV protons on Be and Al target. See their figures 6 and 7 for cross-section ratio vs mass.

Production and Capture of Long-Lived Massive Particles

VALUE (10^{-36} cm^2)	EVTS	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<20 to 800	0	[36] ALEKSEEV	76	ELEC	τ=5 ms to 1 day
<200 to 2000	0	[36] ALEKSEEV	76B	ELEC	τ=100 ms to 1 day
<1.4 to 9	0	[37] FRANKEL	75	CNTR	τ=50 ms to 10 hours
<0.1 to 9	0	[38] FRANKEL	74	CNTR	τ=1 to 1000 hours

[36] ALEKSEEV 76 and ALEKSEEV 76B are 61–70 GeV p Serpukhov experiment. Cross section is per Pb nucleus.

[37] FRANKEL 75 is extension of FRANKEL 74.

[38] FRANKEL 74 looks for particles produced in thick Al targets by 300–400 GeV/c protons.

Heavy Particle Flux in Cosmic Rays

VALUE ($cm^{-2}sr^{-1}s^{-1}$)	CL%	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •						

< 1.8	$\times 10^{-12}$	90		[39] ASTONE	93	CNTR		$m \geq 1.5 \times 10^{-13}$gram
< 1.1	$\times 10^{-14}$	90		[40] AHLEN	92	MCRO		$10^{-10} < m <$ 0.1 gram
~ 6	$\times 10^{-9}$		2	[41] SAITO	90			$Q \simeq 14$, $m \simeq 370 m_p$
< 1.4	$\times 10^{-12}$	90	0	[42] MINCER	85	CALO		$m \geq 1$ TeV
< 3.2	$\times 10^{-11}$	90	0	[43] NAKAMURA	85	CNTR		$m > 1.5 \times 10^{-13}$gram
				[44] SAKUYAMA	83B	PLAS		$m \sim 1$ TeV
< 1.7	$\times 10^{-11}$	99	0	[45] BHAT	82	CC		
$< 1.$	$\times 10^{-9}$	90	0	[46] MARINI	82	CNTR	$\pm$	$Q = 1$, $m \sim 4.5 m_p$
< 3.5	$\times 10^{-11}$	90	0	[47] ULLMAN	81	CNTR		Planck-mass 10^{19}GeV
$< 7.$	$\times 10^{-11}$	90	0	[47] ULLMAN	81	CNTR		$m = 1. \times 10^{-16}$ GeV or less
2.	$\times 10^{-9}$		3	[48] YOCK	81	SPRK	$\pm$	$Q = 1$, $m \sim 4.5 m_p$
			3	[48] YOCK	81	SPRK		Fractionally charged
3.0	$\times 10^{-9}$		3	[49] YOCK	80	SPRK		$m \sim 4.5\ m_p$
(4 ± 1)	$\times 10^{-11}$		3	GOODMAN	79	ELEC		$m \geq 5$ GeV
< 1.3	$\times 10^{-9}$	90		[50] BHAT	78	CNTR	$\pm$	$m > 1$ GeV
< 1.0	$\times 10^{-9}$		0	BRIATORE	76	ELEC		
$< 7.$	$\times 10^{-10}$	90	0	YOCK	75	ELEC	$\pm$	$Q > 7e$ or $< -7e$
$> 6.$	$\times 10^{-9}$		5	[51] YOCK	74	CNTR		$m > 6$ GeV
< 3.0	$\times 10^{-8}$		0	DARDO	72	CNTR		
< 1.5	$\times 10^{-9}$		0	TONWAR	72	CNTR		$m > 10$ GeV
< 3.0	$\times 10^{-10}$		0	BJORNBOE	68	CNTR		$m > 5$ GeV
< 5.0	$\times 10^{-11}$	90	0	JONES	67	ELEC		m=5–15 GeV

[39] ASTONE 93 searched for quark matter ("nuclearites") in the velocity/c range $= 10^{-3}$–1. Their Table 1 gives a compilation of searches for nuclearites.

[40] AHLEN 92 searched for quark matter ("nuclearites"). The bound applies to velocity/c $< 2.5 \times 10^{-3}$. See their Fig. 3 for other velocity/c and heavier mass range.

[41] SAITO 90 candidates carry about 450 MeV/nucleon. Cannot be accounted for by conventional backgrounds. Consistent with strange quark matter hypothesis.

[42] MINCER 85 is high statistics study of calorimeter signals delayed by 20–200 ns. Calibration with AGS beam shows they can be accounted for by rare fluctuations in signals from low-energy hadrons in the shower. Claim that previous delayed signals including BJORNBOE 68, DARDO 72, BHAT 82, SAKUYAMA 83B below may be due to this fake effect.

[43] NAKAMURA 85 at KEK searched for quark-matter. These might be lumps of strange quark matter with roughly equal numbers of u, d, s quarks. These lumps or nuclearites were assumed to have velocity/c of 10^{-4}–10^{-3}.

[44] SAKUYAMA 83B analyzed 6000 extended air shower events. Increase of delayed particles and change of lateral distribution above 10^{17} eV may indicate production of very heavy parent at top of atmosphere.

[45] BHAT 82 observed 12 events with delay $> 2. \times 10^{-8}$ s and with more than 40 particles. 1 eV has good hadron shower. However all events are delayed in only one of two detectors in cloud chamber, and could not be due to strongly interacting massive particle.

[46] MARINI 82 applied PEP-counter for TOF. Above limit is for velocity = 0.54 of light. Limit is inconsistent with YOCK 80 YOCK 81 events if isotropic dependence on zenith angle is assumed.

[47] ULLMAN 81 is sensitive for heavy slow singly charge particle reaching earth with vertical velocity 100–350 km/s.

[48] YOCK 81 saw another 3 events with $Q = \pm 1$ and m about $4.5 m_p$ as well as 2 events with $m > 5.3 m_p$, $Q = \pm 0.75 \pm 0.05$ and $m > 2.8 m_p$, $Q = \pm 0.70 \pm 0.05$ and 1 event with $m = (9.3 \pm 3.) m_p$, $Q = \pm 0.89 \pm 0.06$ as possible heavy candidates.

[49] YOCK 80 events are with charge exactly or approximately equal to unity.

[50] BHAT 78 is at Kolar gold fields. Limit is for $\tau > 10^{-6}$ s.

[51] YOCK 74 events could be tritons.

Long-Lived Particle Search at Hadron Collisions

Limits are for cross section times branching ratio.

VALUE (pb/nucleon)	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<2	90	0	[52] BADIER	86	BDMP	$\tau = (0.05$–$1.) \times 10^{-8}$s

[52] BADIER 86 looked for long-lived particles at 300 GeV π^- beam dump. The limit applies for nonstrongly interaction neutral or charged particles with mass >2 GeV. The limit applies for particle modes, $\mu^+\pi^-$, $\mu^+\mu^-$, $\pi^+\pi^-$X, $\pi^+\pi^-\pi^\pm$ etc. See their figure 5 for the contours of limits in the mass-τ plane for each mode.

Long-Lived Heavy Particle Cross Section

VALUE (pb/sr)	CL%	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<34	95	[53] RAM	94	SPEC	1015<$m_{X^{++}}$<1085 MeV
<75	95	[53] RAM	94	SPEC	920<$m_{X^{++}}$<1025 MeV

[53] RAM 94 search for a long-lived doubly-charged fermion X^{++} with mass between m_N and m_N+m_π and baryon number +1 in the reaction $pp \to X^{++}n$. No candidate is found. The limit is for the cross section at 15° scattering angle at 460 MeV incident energy and applies for $\tau(X^{++}) \gg 0.1\,\mu$s.

REFERENCES FOR Other Stable Particle Searches

RAM	94	PR D49 3120	+Abegg, Ashery, Frekers, Helmer+ (TELA, TRIU)
ABE	93G	PRL 71 2542	+Albrow, Akimoto, Amidei, Anway-Wiese+ (CDF Collab.)
ASTONE	93	PR D47 4770	+Bassan, Bonifazi, Coccia+ (ROMA, ROMAI, CATA, FRAS)
BUSKULIC	93C	PL B303 198	+Decamp, Goy, Lees, Minard+ (ALEPH Collab.)
YAMAGATA	93	PR D47 1231	+Takamori, Utsunomiya (KONAN)
ABE	92J	PR D46 R1889	+Amidei, Anway-Weiss+ (CDF Collab.)
AHLEN	92	PRL 69 1860	+Ambrosio, Antolini, Auriemma, Baker+ (MACRO Collab.)
VERKERK	92	PRL 68 1116	+Grynberg, Pichard, Spiro, Zylberajch+ (ENSP, SACL, PAST)
AKESSON	91	ZPHY C52 219	+Almehed, Angelis, Atherton, Aubry+ (HELIOS Collab.)
ADACHI	90C	PL B244 352	+Aihara, Doser, Enomoto+ (TOPAZ Collab.)
ADACHI	90E	PL B249 336	+Anazawa, Doser, Enomoto, Fujii+ (TOPAZ Collab.)
AKRAWY	90O	PL B252 290	+Alexander, Allison, Allport, Anderson+ (OPAL Collab.)
HEMMICK	90	PR D41 2074	+Elmore+ (ROCH, MICH, OHIO, RAL, LANL, STON)
SAITO	90	PRL 65 2094	+Hatano, Fukada, Oda (ICRR, KOBE)
NAKAMURA	89	PR D39 1261	+Kobayashi, Konaka, Imai, Masaike+ (KYOT, TMTC)
NORMAN	89	PR D39 2499	+Chadwick, Lesko, Larimer, Hoffman (LBL)
BERNSTEIN	88	PR D37 3103	+Shea, Winstein, Cousins, Greenhalgh+ (STAN, WISC)
NORMAN	87	PRL 58 1403	+Gazes, Bennett (LBL)
BADIER	86	ZPHY C31 21	+Bemporad, Boucrot, Callot+ (NA3 Collab.)
MINCER	85	PR D32 541	+Freudenreich, Goodman+ (UMD, GMAS, NSF)
NAKAMURA	85	PL 161B 417	+Horie, Takahashi, Tanimori (KEK, INUS)
THRON	85	PR D31 451	+Cardello, Cooper, Teig+ (YALE, FNAL, IOWA)
SAKUYAMA	83B	LNC 37 17	+Nuzuki (MEIS)
Also	83	LNC 36 389	Sakuyama, Watanabe (MEIS)
Also	83D	NC 78A 147	Sakuyama, Watanabe (MEIS)
Also	83C	NC 6C 371	Sakuyama, Watanabe (MEIS)
BHAT	82	PR D25 2820	+Gupta, Murthy, Sreekantan+ (TATA)
KINOSHITA	82	PRL 48 77	+Price, Fryberger (UCB, SLAC)
MARINI	82	PR D26 1777	+Peruzzi, Piccolo+ (FRAS, LBL, NWES, STAN, HAWA)
SMITH	82B	NP B206 333	+Bennett, Homer, Lewin, Walford, Smith (RAL)
KINOSHITA	81B	PR D24 1707	+Price (UCB)
LOSECCO	81	PL 102B 209	+Sulak, Galik, Horstkotte+ (MICH, PENN, BNL)
ULLMAN	81	PRL 47 289	(LEHM, BNL)
YOCK	81	PR D23 1207	(AUCK)
BARTEL	80	ZPHY C6 295	+Canzler, Lords, Drumm+ (JADE Collab.)
BUSSIERE	80	NP B174 1	+Giacomelli, Lesquoy+ (BGNA, SACL, LAPP)
YOCK	80	PR D22 61	(AUCK)
ARMITAGE	79	NP B150 87	+Benz, Bobbink+ (CERN, DARE, FOM, MCHS, UTRE)
BOZZOLI	79	NP B159 363	+Bussiere, Giacomelli+ (BGNA, LAPP, SACL, CERN)
GOODMAN	79	PR D19 2572	+Ellsworth, Ito, Macfall, Siohan+ (UMD)
SMITH	79	NP B149 525	+Bennett (RHEL)
BHAT	78	Pramana 10 115	+Murthy (TATA)
CARROLL	78	PRL 41 777	+Chiang, Johnson, Kycia, Ki+ (BNL, PRIN)
CUTTS	78	PRL 41 363	+Dulude+ (BROW, FNAL, ILL, BARI, MIT, WARS)
VIDAL	78	PL 77B 344	+Herb, Lederman+ (COLU, FNAL, STON, UCB)
VIERTEL	78	LNC 22 235	+Hahn, Schacher (BERN)
ALEKSEEV	76	SJNP 22 531	+Zaitsev, Kalinina, Kruglov+ (JINR)
		Translated from YAF 22 1021.	
ALEKSEEV	76B	SJNP 23 633	+Zaitsev, Kalinina, Kruglov+ (JINR)
		Translated from YAF 23 1190.	
BALDIN	76	SJNP 22 264	+Vertogradov, Vishnevsky, Grishkevich+ (JINR)
		Translated from YAF 22 512.	
BRIATORE	76	NC 31A 553	+Dardo, Piazzoli, Mannocchi+ (LCGT, FRAS, FREIB)
GUSTAFSON	76	PRL 37 474	+Ayre, Jones, Longo, Murthy (MICH)
ALBROW	75	NP B97 189	+Barber+ (CERN, DARE, FOM, LANC, MCHS, UTRE)
BLAGOV	75	SJNP 21 158	+Komar, Murashova, Syreishchikova+ (LEBD)
		Translated from YAF 21 300.	

FRANKEL	75	PR D12 2561	+Frati, Resvanis, Yang, Nezrick	(PENN, FNAL)
JOVANOV...	75	PL 56B 105	Jovanovich+	(MANI, AACH, CERN, GENO, HARV+)
YOCK	75	NP B86 216		(AUCK, SLAC)
APPEL	74	PRL 32 428	+Bourquin, Gaines, Lederman+	(COLU, FNAL)
FRANKEL	74	PR D9 1932	+Frati, Resvanis, Yang, Nezrick	(PENN, FNAL)
YOCK	74	NP B76 175		(AUCK)
ALPER	73	PL 46B 265	+	(CERN, LIVP, LUND, BOHR, RHEL, STOH, BERG+)
LEIPUNER	73	PRL 31 1226	+Larsen, Sessoms, Smith, Williams+	(BNL, YALE)
DARDO	72	NC 9A 319	+Navarra, Penengo, Sitte	(TORI)
TONWAR	72	JPA 5 569	+Naranan, Sreekantan	(TATA)
ANTIPOV	71B	NP B31 235	+Denisov, Donskov, Gorin, Kachanov+	(SERP)
ANTIPOV	71C	PL 34B 164	+Denisov, Donskov, Gorin, Kachanov+	(SERP)
BINON	69	PL 30B 510	+Duteil, Kachanov, Khromov, Kutyin+	(SERP)
BJORNBOE	68	NC B53 241	+Damgard, Hansen+	(BOHR, TATA, BERN, BERG)
JONES	67	PR 164 1584		(MICH, WISC, LBL, UCLA, MINN, COSU, COLO+)
DORFAN	65	PRL 14 999	+Eades, Lederman, Lee, Ting	(COLU)
COWARD	63	PR 131 1782	+Gittelman, Lynch, Ritson	(STAN)
BELOUSOV	60	JETP 11 1143	+Rusakov, Tamm, Cerenkov	(LEBD)
		Translated from ZETF 38 1589.		
GORBUNOV	60	JETP 11 51	+Spiridonov, Cerenkov	(LEBD)
		Translated from ZETF 38 69.		