

$f_2(1950)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

$f_2(1950)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1960±30	¹ BARBERIS	97B	OMEG	450 $pp \rightarrow$ $pp2(\pi^+\pi^-)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1918±12	ANTINORI	95	OMEG	300,450 $pp \rightarrow$ $pp2(\pi^+\pi^-)$
~ 1996	HASAN	94	RVUE	$\bar{p}p \rightarrow \pi\pi$
~ 1990	² OAKDEN	94	RVUE	0.36–1.55 $\bar{p}p \rightarrow$ $\pi\pi$
1950±15	³ ASTON	91	LASS 0	11 $K^-p \rightarrow$ $\Lambda K \bar{K} \pi\pi$

¹ Possibly two states.

² From solution B of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$. See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

³ Cannot determine spin to be 2.

$f_2(1950)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
460±40	⁴ BARBERIS	97B	OMEG	450 $pp \rightarrow$ $pp2(\pi^+\pi^-)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
390±60	ANTINORI	95	OMEG	300,450 $pp \rightarrow$ $pp2(\pi^+\pi^-)$
~ 134	HASAN	94	RVUE	$\bar{p}p \rightarrow \pi\pi$
~ 100	⁵ OAKDEN	94	RVUE	0.36–1.55 $\bar{p}p \rightarrow$ $\pi\pi$
250±50	⁶ ASTON	91	LASS 0	11 $K^-p \rightarrow$ $\Lambda K \bar{K} \pi\pi$

⁴ Possibly two states.

⁵ From solution B of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$. See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

⁶ Cannot determine spin to be 2.

$f_2(1950)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K^*(892)\bar{K}^*(892)$	seen
Γ_2 $\pi^+\pi^-$	seen
Γ_3 $\pi^+\pi^-\pi^+\pi^-$	possibly seen
Γ_4 $a_2(1320)\pi$	
Γ_5 $f_2(1270)\pi\pi$	

$f_2(1950)$ BRANCHING RATIOS

$\Gamma(K^*(892)\bar{K}^*(892))/\Gamma_{\text{total}}$	Γ_1/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u> <u>COMMENT</u>
seen	ASTON 91 LASS 0 11 $K^-p \rightarrow \Lambda K \bar{K} \pi \pi$

$\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}}$	Γ_4/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
possibly seen	BARBERIS 97B OMEG 450 $pp \rightarrow pp2(\pi^+\pi^-)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$f_2(1950)$ REFERENCES

BARBERIS	97B	PL B413 217	D. Barberis+	(WA102 Collab.)
KLOET	96	PR D53 6120	+Myhrer	(RUTG, NORD)
ANTINORI	95	PL B353 589	+Barberis, Bayes+	(ATHU, BARI, BIRM, CERN, JINR) JP
HASAN	94	PL B334 215	+Bugg	(LOQM)
OAKDEN	94	NPA 574 731	+Pennington	(DURH)
ASTON	91	NP B21 5 (suppl)	+Awaji+	(LASS Collab.)

OTHER RELATED PAPERS

ALBRECHT	88N	PL B212 528	+	(ARGUS Collab.)
ALBRECHT	87Q	PL B198 255	+Binder+	(ARGUS Collab.)
ARMSTRONG	87C	ZPHY C34 33	+Bloodworth+	(CERN, BIRM, BARI, ATHU, CURIN+)