

# STRANGE MESONS

## ( $S = \pm 1, C = B = 0$ )

$$K^+ = u\bar{s}, K^0 = d\bar{s}, \bar{K}^0 = \bar{d}s, K^- = \bar{u}s, \text{ similarly for } K^{*'}s$$

 **$K^\pm$** 

$$I(J^P) = \frac{1}{2}(0^-)$$

$$\text{Mass } m = 493.677 \pm 0.016 \text{ MeV } [u] \quad (S = 2.8)$$

$$\text{Mean life } \tau = (1.2386 \pm 0.0024) \times 10^{-8} \text{ s} \quad (S = 2.0)$$

$$c\tau = 3.713 \text{ m}$$

**Slope parameter  $g$  [v]**

(See Particle Listings for quadratic coefficients)

$$K^+ \rightarrow \pi^+ \pi^+ \pi^- = -0.2154 \pm 0.0035 \quad (S = 1.4)$$

$$K^- \rightarrow \pi^- \pi^- \pi^+ = -0.217 \pm 0.007 \quad (S = 2.5)$$

$$K^\pm \rightarrow \pi^\pm \pi^0 \pi^0 = 0.594 \pm 0.019 \quad (S = 1.3)$$

 **$K^\pm$  decay form factors [a,w]**

$$K_{e3}^+ \quad \lambda_+ = 0.0286 \pm 0.0022$$

$$K_{\mu 3}^+ \quad \lambda_+ = 0.032 \pm 0.008 \quad (S = 1.6)$$

$$K_{\mu 3}^+ \quad \lambda_0 = 0.006 \pm 0.007 \quad (S = 1.6)$$

$$K_{e3}^+ \quad |f_S/f_+| = 0.084 \pm 0.023 \quad (S = 1.2)$$

$$K_{e3}^+ \quad |f_T/f_+| = 0.38 \pm 0.11 \quad (S = 1.1)$$

$$K_{\mu 3}^+ \quad |f_T/f_+| = 0.02 \pm 0.12$$

$$K^+ \rightarrow e^+ \nu_e \gamma \quad |F_A + F_V| = 0.148 \pm 0.010$$

$$K^+ \rightarrow \mu^+ \nu_\mu \gamma \quad |F_A + F_V| < 0.23, \text{ CL} = 90\%$$

$$K^+ \rightarrow e^+ \nu_e \gamma \quad |F_A - F_V| < 0.49$$

$$K^+ \rightarrow \mu^+ \nu_\mu \gamma \quad |F_A - F_V| = -2.2 \text{ to } 0.3$$

 $K^-$  modes are charge conjugates of the modes below.

<b><math>K^+</math> DECAY MODES</b>	Fraction (, i/, )	Scale factor/ Confidence level	$p$ (MeV/c)
$\mu^+ \nu_\mu$	(63.51±0.18) %	S=1.3	236
$e^+ \nu_e$	( 1.55±0.07) × 10 <sup>-5</sup>		247
$\pi^+ \pi^0$	(21.16±0.14) %	S=1.1	205
$\pi^+ \pi^+ \pi^-$	( 5.59±0.05) %	S=1.8	125
$\pi^+ \pi^0 \pi^0$	( 1.73±0.04) %	S=1.2	133
$\pi^0 \mu^+ \nu_\mu$	( 3.18±0.08) %	S=1.5	215

 Called  $K_{\mu 3}^+$ .

$\pi^0 e^+ \nu_e$		( 4.82±0.06 ) %	S=1.3	228
Called $K_{e3}^+$ .				
$\pi^0 \pi^0 e^+ \nu_e$		( 2.1 ±0.4 ) × 10 <sup>-5</sup>		206
$\pi^+ \pi^- e^+ \nu_e$		( 3.91±0.17 ) × 10 <sup>-5</sup>		203
$\pi^+ \pi^- \mu^+ \nu_\mu$		( 1.4 ±0.9 ) × 10 <sup>-5</sup>		151
$\pi^0 \pi^0 \pi^0 e^+ \nu_e$		< 3.5 × 10 <sup>-6</sup>	CL=90%	135
$\pi^+ \gamma \gamma$	[x]	( 1.10±0.32 ) × 10 <sup>-6</sup>		227
$\pi^+ 3\gamma$	[x]	< 1.0 × 10 <sup>-4</sup>	CL=90%	227
$\mu^+ \nu_\mu \nu \bar{\nu}$		< 6.0 × 10 <sup>-6</sup>	CL=90%	236
$e^+ \nu_e \nu \bar{\nu}$		< 6 × 10 <sup>-5</sup>	CL=90%	247
$\mu^+ \nu_\mu e^+ e^-$		( 1.3 ±0.4 ) × 10 <sup>-7</sup>		236
$e^+ \nu_e e^+ e^-$		( 3.0 <sup>+3.0</sup> <sub>-1.5</sub> ) × 10 <sup>-8</sup>		247
$\mu^+ \nu_\mu \mu^+ \mu^-$		< 4.1 × 10 <sup>-7</sup>	CL=90%	185
$\mu^+ \nu_\mu \gamma$	[x,y]	( 5.50±0.28 ) × 10 <sup>-3</sup>		236
$\pi^+ \pi^0 \gamma$	[x,y]	( 2.75±0.15 ) × 10 <sup>-4</sup>		205
$\pi^+ \pi^0 \gamma(\text{DE})$	[x,z]	( 1.8 ±0.4 ) × 10 <sup>-5</sup>		205
$\pi^+ \pi^+ \pi^- \gamma$	[x,y]	( 1.04±0.31 ) × 10 <sup>-4</sup>		125
$\pi^+ \pi^0 \pi^0 \gamma$	[x,y]	( 7.5 <sup>+5.5</sup> <sub>-3.0</sub> ) × 10 <sup>-6</sup>		133
$\pi^0 \mu^+ \nu_\mu \gamma$	[x,y]	< 6.1 × 10 <sup>-5</sup>	CL=90%	215
$\pi^0 e^+ \nu_e \gamma$	[x,y]	( 2.62±0.20 ) × 10 <sup>-4</sup>		228
$\pi^0 e^+ \nu_e \gamma(\text{SD})$	[aa]	< 5.3 × 10 <sup>-5</sup>	CL=90%	228
$\pi^0 \pi^0 e^+ \nu_e \gamma$		< 5 × 10 <sup>-6</sup>	CL=90%	206

**Lepton Family number (LF), Lepton number (L),  $\Delta S = \Delta Q$  (SQ)  
violating modes, or  $\Delta S = 1$  weak neutral current (S1) modes**

$\pi^+ \pi^+ e^- \bar{\nu}_e$	SQ	< 1.2 × 10 <sup>-8</sup>	CL=90%	203
$\pi^+ \pi^+ \mu^- \bar{\nu}_\mu$	SQ	< 3.0 × 10 <sup>-6</sup>	CL=95%	151
$\pi^+ e^+ e^-$	S1	( 2.74±0.23 ) × 10 <sup>-7</sup>		227
$\pi^+ \mu^+ \mu^-$	S1	( 5.0 ±1.0 ) × 10 <sup>-8</sup>		172
$\pi^+ \nu \bar{\nu}$	S1	( 4.2 <sup>+9.7</sup> <sub>-3.5</sub> ) × 10 <sup>-10</sup>		227
$\mu^- \nu e^+ e^+$	LF	< 2.0 × 10 <sup>-8</sup>	CL=90%	236
$\mu^+ \nu_e$	LF	[d] < 4 × 10 <sup>-3</sup>	CL=90%	236
$\pi^+ \mu^+ e^-$	LF	< 2.1 × 10 <sup>-10</sup>	CL=90%	214
$\pi^+ \mu^- e^+$	LF	< 7 × 10 <sup>-9</sup>	CL=90%	214
$\pi^- \mu^+ e^+$	L	< 7 × 10 <sup>-9</sup>	CL=90%	214
$\pi^- e^+ e^+$	L	< 1.0 × 10 <sup>-8</sup>	CL=90%	227
$\pi^- \mu^+ \mu^+$	L	[d] < 1.5 × 10 <sup>-4</sup>	CL=90%	172
$\mu^+ \bar{\nu}_e$	L	[d] < 3.3 × 10 <sup>-3</sup>	CL=90%	236
$\pi^0 e^+ \bar{\nu}_e$	L	< 3 × 10 <sup>-3</sup>	CL=90%	228

**$K^0$** 

$$I(J^P) = \frac{1}{2}(0^-)$$

 50%  $K_S$ , 50%  $K_L$ 

 Mass  $m = 497.672 \pm 0.031$  MeV

 $m_{K^0} - m_{K^\pm} = 3.995 \pm 0.034$  MeV ( $S = 1.1$ )

 $|m_{K^0} - m_{\bar{K}^0}| / m_{\text{average}} < 10^{-18}$  [bb]

 **$K_S^0$** 

$$I(J^P) = \frac{1}{2}(0^-)$$

 Mean life  $\tau = (0.8934 \pm 0.0008) \times 10^{-10}$  s

 $c\tau = 2.6783$  cm

**CP-violation parameters** [cc]

 $\text{Im}(\eta_{+-0}) = -0.002 \pm 0.009$ 
 $\text{Im}(\eta_{000})^2 < 0.1$ , CL = 90%

<b><math>K_S^0</math> DECAY MODES</b>	Fraction (, ; /, )	Scale factor/ Confidence level	$p$ (MeV/c)
$\pi^+ \pi^-$	$(68.61 \pm 0.28) \%$	S=1.2	206
$\pi^0 \pi^0$	$(31.39 \pm 0.28) \%$	S=1.2	209
$\pi^+ \pi^- \gamma$	[y, dd] $(1.78 \pm 0.05) \times 10^{-3}$		206
$\gamma \gamma$	$(2.4 \pm 0.9) \times 10^{-6}$		249
$\pi^+ \pi^- \pi^0$	$(3.4 \pm_{-0.9}^{+1.1}) \times 10^{-7}$		133
$3\pi^0$	$< 3.7 \times 10^{-5}$	CL=90%	139
$\pi^\pm e^\mp \nu$	[ee] $(6.70 \pm 0.07) \times 10^{-4}$	S=1.1	229
$\pi^\pm \mu^\mp \nu$	[ee] $(4.69 \pm 0.06) \times 10^{-4}$	S=1.1	216

 **$\Delta S = 1$  weak neutral current (S1) modes**

$\mu^+ \mu^-$	S1	$< 3.2 \times 10^{-7}$	CL=90%	225
$e^+ e^-$	S1	$< 1.4 \times 10^{-7}$	CL=90%	249
$\pi^0 e^+ e^-$	S1	$< 1.1 \times 10^{-6}$	CL=90%	231

**$K_L^0$** 

$$I(J^P) = \frac{1}{2}(0^-)$$

$$m_{K_L} - m_{K_S} = (0.5301 \pm 0.0014) \times 10^{10} \hbar s^{-1}$$

$$= (3.489 \pm 0.009) \times 10^{-12} \text{ MeV}$$

$$\text{Mean life } \tau = (5.17 \pm 0.04) \times 10^{-8} \text{ s} \quad (S = 1.1)$$

$$c\tau = 15.51 \text{ m}$$

**Slope parameter  $g$**  <sup>[v]</sup>

(See Particle Listings for quadratic coefficients)

$$K_L^0 \rightarrow \pi^+ \pi^- \pi^0 = 0.670 \pm 0.014 \quad (S = 1.6)$$

 **$K_L$  decay form factors** <sup>[w]</sup>

$$K_{e3}^0 \quad \lambda_+ = 0.0300 \pm 0.0016 \quad (S = 1.2)$$

$$K_{\mu 3}^0 \quad \lambda_+ = 0.034 \pm 0.005 \quad (S = 2.3)$$

$$K_{\mu 3}^0 \quad \lambda_0 = 0.025 \pm 0.006 \quad (S = 2.3)$$

$$K_{e3}^0 \quad |f_S/f_+| < 0.04, \text{ CL} = 68\%$$

$$K_{e3}^0 \quad |f_T/f_+| < 0.23, \text{ CL} = 68\%$$

$$K_{\mu 3}^0 \quad |f_T/f_+| = 0.12 \pm 0.12$$

$$K_L \rightarrow e^+ e^- \gamma: \quad \alpha_{K^*} = -0.28 \pm 0.08$$

**CP-violation parameters** <sup>[cc]</sup>

$$\delta = (0.327 \pm 0.012)\%$$

$$|\eta_{00}| = (2.275 \pm 0.019) \times 10^{-3} \quad (S = 1.1)$$

$$|\eta_{+-}| = (2.285 \pm 0.019) \times 10^{-3}$$

$$|\eta_{00}/\eta_{+-}| = 0.9956 \pm 0.0023 \text{ [ff]} \quad (S = 1.8)$$

$$\epsilon'/\epsilon = (1.5 \pm 0.8) \times 10^{-3} \text{ [ff]} \quad (S = 1.8)$$

$$\phi_{+-} = (43.5 \pm 0.6)^\circ$$

$$\phi_{00} = (43.4 \pm 1.0)^\circ$$

$$\phi_{00} - \phi_{+-} = (-0.1 \pm 0.8)^\circ$$

$$j \text{ for } K_L^0 \rightarrow \pi^+ \pi^- \pi^0 = 0.0011 \pm 0.0008$$

$$|\eta_{+-\gamma}| = (2.35 \pm 0.07) \times 10^{-3}$$

$$\phi_{+-\gamma} = (44 \pm 4)^\circ$$

$$|\epsilon'_{+-\gamma}|/\epsilon < 0.3, \text{ CL} = 90\%$$

**$\Delta S = -\Delta Q$  in  $K_{l3}^0$  decay**

$$\text{Re } x = 0.006 \pm 0.018 \quad (S = 1.3)$$

$$\text{Im } x = -0.003 \pm 0.026 \quad (S = 1.2)$$

 **$CPT$ -violation parameters**

$$\text{Re } \Delta = 0.018 \pm 0.020$$

$$\text{Im } \Delta = 0.02 \pm 0.04$$

$K_L^0$ DECAY MODES	Fraction ( , j/, )	Scale factor/ Confidence level	$p$ (MeV/c)
$3\pi^0$	(21.12 $\pm$ 0.27 ) %	S=1.1	139
$\pi^+ \pi^- \pi^0$	(12.56 $\pm$ 0.20 ) %	S=1.7	133
$\pi^\pm \mu^\mp \nu$ Called $K_{\mu 3}^0$ .	[gg] (27.17 $\pm$ 0.25 ) %	S=1.1	216
$\pi^\pm e^\mp \nu_e$ Called $K_{e 3}^0$ .	[gg] (38.78 $\pm$ 0.27 ) %	S=1.1	229
$2\gamma$	( 5.92 $\pm$ 0.15 ) $\times 10^{-4}$		249
$3\gamma$	< 2.4 $\times 10^{-7}$	CL=90%	249
$\pi^0 2\gamma$	[hh] ( 1.70 $\pm$ 0.28 ) $\times 10^{-6}$		231
$\pi^0 \pi^\pm e^\mp \nu$	[gg] ( 5.18 $\pm$ 0.29 ) $\times 10^{-5}$		207
$(\pi \mu \text{atom})\nu$	( 1.06 $\pm$ 0.11 ) $\times 10^{-7}$		–
$\pi^\pm e^\mp \nu_e \gamma$	[y,gg,hh] ( 3.62 $^{+0.26}_{-0.21}$ ) $\times 10^{-3}$		229
$\pi^+ \pi^- \gamma$	[y,hh] ( 4.61 $\pm$ 0.14 ) $\times 10^{-5}$		206
$\pi^0 \pi^0 \gamma$	< 5.6 $\times 10^{-6}$		209

**Charge conjugation  $\times$  Parity ( $CP$ ,  $CPV$ ) or Lepton Family number ( $LF$ )  
violating modes, or  $\Delta S = 1$  weak neutral current ( $S1$ ) modes**

$\pi^+ \pi^-$	$CPV$	( 2.067 $\pm$ 0.035 ) $\times 10^{-3}$	S=1.1	206
$\pi^0 \pi^0$	$CPV$	( 9.36 $\pm$ 0.20 ) $\times 10^{-4}$		209
$\mu^+ \mu^-$	$S1$	( 7.2 $\pm$ 0.5 ) $\times 10^{-9}$	S=1.4	225
$\mu^+ \mu^- \gamma$	$S1$	( 3.25 $\pm$ 0.28 ) $\times 10^{-7}$		225
$e^+ e^-$	$S1$	< 4.1 $\times 10^{-11}$	CL=90%	249
$e^+ e^- \gamma$	$S1$	( 9.1 $\pm$ 0.5 ) $\times 10^{-6}$		249
$e^+ e^- \gamma \gamma$	$S1$ [hh]	( 6.5 $\pm$ 1.2 ) $\times 10^{-7}$		249
$\pi^+ \pi^- e^+ e^-$	$S1$ [hh]	< 4.6 $\times 10^{-7}$	CL=90%	206
$\mu^+ \mu^- e^+ e^-$	$S1$	( 2.9 $^{+6.7}_{-2.4}$ ) $\times 10^{-9}$		225
$e^+ e^- e^+ e^-$	$S1$	( 4.1 $\pm$ 0.8 ) $\times 10^{-8}$	S=1.2	249
$\pi^0 \mu^+ \mu^-$	$CP, S1$ [ii]	< 5.1 $\times 10^{-9}$	CL=90%	177
$\pi^0 e^+ e^-$	$CP, S1$ [ii]	< 4.3 $\times 10^{-9}$	CL=90%	231
$\pi^0 \nu \bar{\nu}$	$CP, S1$ [jj]	< 5.8 $\times 10^{-5}$	CL=90%	231
$e^\pm \mu^\mp$	$LF$ [gg]	< 3.3 $\times 10^{-11}$	CL=90%	238
$e^\pm e^\pm \mu^\mp \mu^\mp$	$LF$ [gg]	< 6.1 $\times 10^{-9}$	CL=90%	–

**$K^*(892)$** 

$$I(J^P) = \frac{1}{2}(1^-)$$

 $K^*(892)^\pm$  mass  $m = 891.66 \pm 0.26$  MeV $K^*(892)^0$  mass  $m = 896.10 \pm 0.28$  MeV ( $S = 1.4$ ) $K^*(892)^\pm$  full width  $\Gamma = 50.8 \pm 0.9$  MeV $K^*(892)^0$  full width  $\Gamma = 50.5 \pm 0.6$  MeV ( $S = 1.1$ )

<b><math>K^*(892)</math> DECAY MODES</b>	Fraction ( $i, j, \dots$ )	Confidence level	$p$ (MeV/c)
$K\pi$	$\sim 100$	%	291
$K^0\gamma$	$(2.30 \pm 0.20) \times 10^{-3}$		310
$K^\pm\gamma$	$(9.9 \pm 0.9) \times 10^{-4}$		309
$K\pi\pi$	$< 7$	$\times 10^{-4}$ 95%	224

 **$K_1(1270)$** 

$$I(J^P) = \frac{1}{2}(1^+)$$

Mass  $m = 1273 \pm 7$  MeV [ $m$ ]Full width  $\Gamma = 90 \pm 20$  MeV [ $m$ ]

<b><math>K_1(1270)</math> DECAY MODES</b>	Fraction ( $i, j, \dots$ )	$p$ (MeV/c)
$K\rho$	$(42 \pm 6) \%$	76
$K_0^*(1430)\pi$	$(28 \pm 4) \%$	—
$K^*(892)\pi$	$(16 \pm 5) \%$	301
$K\omega$	$(11.0 \pm 2.0) \%$	—
$Kf_0(1370)$	$(3.0 \pm 2.0) \%$	—

 **$K_1(1400)$** 

$$I(J^P) = \frac{1}{2}(1^+)$$

Mass  $m = 1402 \pm 7$  MeVFull width  $\Gamma = 174 \pm 13$  MeV ( $S = 1.6$ )

<b><math>K_1(1400)</math> DECAY MODES</b>	Fraction ( $i, j, \dots$ )	$p$ (MeV/c)
$K^*(892)\pi$	$(94 \pm 6) \%$	401
$K\rho$	$(3.0 \pm 3.0) \%$	298
$Kf_0(1370)$	$(2.0 \pm 2.0) \%$	—
$K\omega$	$(1.0 \pm 1.0) \%$	285
$K_0^*(1430)\pi$	not seen	—

**$K^*(1410)$** 

$$I(J^P) = \frac{1}{2}(1^-)$$

Mass  $m = 1414 \pm 15$  MeV (S = 1.3)Full width  $\Gamma = 232 \pm 21$  MeV (S = 1.1)

<b><math>K^*(1410)</math> DECAY MODES</b>	Fraction (, $i$ /, )	Confidence level	$p$ (MeV/c)
$K^*(892)\pi$	> 40 %	95%	408
$K\pi$	( $6.6 \pm 1.3$ ) %		611
$K\rho$	< 7 %	95%	309

 **$K_0^*(1430)$  <sup>[kk]</sup>**

$$I(J^P) = \frac{1}{2}(0^+)$$

Mass  $m = 1429 \pm 6$  MeVFull width  $\Gamma = 287 \pm 23$  MeV

<b><math>K_0^*(1430)</math> DECAY MODES</b>	Fraction (, $i$ /, )	$p$ (MeV/c)
$K\pi$	( $93 \pm 10$ ) %	621

 **$K_2^*(1430)$** 

$$I(J^P) = \frac{1}{2}(2^+)$$

 $K_2^*(1430)^\pm$  mass  $m = 1425.6 \pm 1.5$  MeV (S = 1.1) $K_2^*(1430)^0$  mass  $m = 1432.4 \pm 1.3$  MeV $K_2^*(1430)^\pm$  full width  $\Gamma = 98.5 \pm 2.7$  MeV (S = 1.1) $K_2^*(1430)^0$  full width  $\Gamma = 109 \pm 5$  MeV (S = 1.9)

<b><math>K_2^*(1430)</math> DECAY MODES</b>	Fraction (, $i$ /, )	Scale factor/ Confidence level	$p$ (MeV/c)
$K\pi$	( $49.9 \pm 1.2$ ) %		622
$K^*(892)\pi$	( $24.7 \pm 1.5$ ) %		423
$K^*(892)\pi\pi$	( $13.4 \pm 2.2$ ) %		375
$K\rho$	( $8.7 \pm 0.8$ ) %	S=1.2	331
$K\omega$	( $2.9 \pm 0.8$ ) %		319
$K^+\gamma$	( $2.4 \pm 0.5$ ) $\times 10^{-3}$	S=1.1	627
$K\eta$	( $1.5^{+3.4}_{-1.0}$ ) $\times 10^{-3}$	S=1.3	492
$K\omega\pi$	< 7.2 $\times 10^{-4}$	CL=95%	110
$K^0\gamma$	< 9 $\times 10^{-4}$	CL=90%	631

**$K^*(1680)$** 

$$I(J^P) = \frac{1}{2}(1^-)$$

Mass  $m = 1717 \pm 27$  MeV (S = 1.4)Full width  $\Gamma = 322 \pm 110$  MeV (S = 4.2)

<b><math>K^*(1680)</math> DECAY MODES</b>	Fraction (, $i/j$ , )	$p$ (MeV/c)
$K\pi$	$(38.7 \pm 2.5)$ %	779
$K\rho$	$(31.4^{+4.7}_{-2.1})$ %	571
$K^*(892)\pi$	$(29.9^{+2.2}_{-4.7})$ %	615

 **$K_2(1770)$  <sup>[11]</sup>**

$$I(J^P) = \frac{1}{2}(2^-)$$

Mass  $m = 1773 \pm 8$  MeVFull width  $\Gamma = 186 \pm 14$  MeV

<b><math>K_2(1770)</math> DECAY MODES</b>	Fraction (, $i/j$ , )	$p$ (MeV/c)
$K\pi\pi$		—
$K_2^*(1430)\pi$	dominant	287
$K^*(892)\pi$	seen	653
$Kf_2(1270)$	seen	—
$K\phi$	seen	441
$K\omega$	seen	608

 **$K_3^*(1780)$** 

$$I(J^P) = \frac{1}{2}(3^-)$$

Mass  $m = 1776 \pm 7$  MeV (S = 1.1)Full width  $\Gamma = 159 \pm 21$  MeV (S = 1.3)

<b><math>K_3^*(1780)</math> DECAY MODES</b>	Fraction (, $i/j$ , )	Confidence level	$p$ (MeV/c)
$K\rho$	$(31 \pm 9)$ %		612
$K^*(892)\pi$	$(20 \pm 5)$ %		651
$K\pi$	$(18.8 \pm 1.0)$ %		810
$K\eta$	$(30 \pm 13)$ %		715
$K_2^*(1430)\pi$	< 16 %	95%	284



**$K_2(1820)$   $[mm]$** 

$$I(J^P) = \frac{1}{2}(2^-)$$

 Mass  $m = 1816 \pm 13$  MeV

 Full width  $\Gamma = 276 \pm 35$  MeV

<b><math>K_2(1820)</math> DECAY MODES</b>	Fraction (, $j$ /, )	$p$ (MeV/c)
$K_2^*(1430)\pi$	seen	325
$K^*(892)\pi$	seen	680
$K f_2(1270)$	seen	186
$K\omega$	seen	638

 **$K_4^*(2045)$** 

$$I(J^P) = \frac{1}{2}(4^+)$$

 Mass  $m = 2045 \pm 9$  MeV ( $S = 1.1$ )

 Full width  $\Gamma = 198 \pm 30$  MeV

<b><math>K_4^*(2045)</math> DECAY MODES</b>	Fraction (, $j$ /, )	$p$ (MeV/c)
$K\pi$	$(9.9 \pm 1.2) \%$	958
$K^*(892)\pi\pi$	$(9 \pm 5) \%$	800
$K^*(892)\pi\pi\pi$	$(7 \pm 5) \%$	764
$\rho K\pi$	$(5.7 \pm 3.2) \%$	742
$\omega K\pi$	$(5.0 \pm 3.0) \%$	736
$\phi K\pi$	$(2.8 \pm 1.4) \%$	591
$\phi K^*(892)$	$(1.4 \pm 0.7) \%$	363