

N(1700) D_{13}

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-) \text{ Status: } ***$$

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters **111B** (1982).

The various partial-wave analyses do not agree very well.

N(1700) BREIT-WIGNER MASS

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|-----------------------|------|--|
| 1650 to 1750 (\approx 1700) OUR ESTIMATE | | | |
| 1737 \pm 44 | MANLEY | 92 | IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$ |
| 1675 \pm 25 | CUTKOSKY | 80 | IPWA $\pi N \rightarrow \pi N$ |
| 1731 \pm 15 | HOEHLER | 79 | IPWA $\pi N \rightarrow \pi N$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 1791 \pm 46 | BATINIC | 95 | DPWA $\pi N \rightarrow N\pi, N\eta$ |
| 1709 | CRAWFORD | 80 | DPWA $\gamma N \rightarrow \pi N$ |
| 1650 | SAXON | 80 | DPWA $\pi^- p \rightarrow \Lambda K^0$ |
| 1690 to 1710 | BAKER | 78 | DPWA $\pi^- p \rightarrow \Lambda K^0$ |
| 1719 | BARBOUR | 78 | DPWA $\gamma N \rightarrow \pi N$ |
| 1670 \pm 10 | ¹ BAKER | 77 | IPWA $\pi^- p \rightarrow \Lambda K^0$ |
| 1690 | ¹ BAKER | 77 | DPWA $\pi^- p \rightarrow \Lambda K^0$ |
| 1660 | ² LONGACRE | 77 | IPWA $\pi N \rightarrow N\pi\pi$ |
| 1710 | ³ LONGACRE | 75 | IPWA $\pi N \rightarrow N\pi\pi$ |

N(1700) BREIT-WIGNER WIDTH

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|-----------------------|------|--|
| 50 to 150 (\approx 100) OUR ESTIMATE | | | |
| 250 \pm 220 | MANLEY | 92 | IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$ |
| 90 \pm 40 | CUTKOSKY | 80 | IPWA $\pi N \rightarrow \pi N$ |
| 110 \pm 30 | HOEHLER | 79 | IPWA $\pi N \rightarrow \pi N$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 215 \pm 60 | BATINIC | 95 | DPWA $\pi N \rightarrow N\pi, N\eta$ |
| 166 | CRAWFORD | 80 | DPWA $\gamma N \rightarrow \pi N$ |
| 70 | SAXON | 80 | DPWA $\pi^- p \rightarrow \Lambda K^0$ |
| 70 to 100 | BAKER | 78 | DPWA $\pi^- p \rightarrow \Lambda K^0$ |
| 126 | BARBOUR | 78 | DPWA $\gamma N \rightarrow \pi N$ |
| 90 \pm 25 | ¹ BAKER | 77 | IPWA $\pi^- p \rightarrow \Lambda K^0$ |
| 100 | ¹ BAKER | 77 | DPWA $\pi^- p \rightarrow \Lambda K^0$ |
| 600 | ² LONGACRE | 77 | IPWA $\pi N \rightarrow N\pi\pi$ |
| 300 | ³ LONGACRE | 75 | IPWA $\pi N \rightarrow N\pi\pi$ |

N(1700) POLE POSITION

REAL PART

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|-----------------------|------|--|
| 1630 to 1730 (\approx 1680) OUR ESTIMATE | | | |
| 1700 | ⁴ HOEHLER | 93 | SPED $\pi N \rightarrow \pi N$ |
| 1660 \pm 30 | CUTKOSKY | 80 | IPWA $\pi N \rightarrow \pi N$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| not seen | ARNDT | 91 | DPWA $\pi N \rightarrow \pi N$ Soln SM90 |
| 1710 or 1678 | ⁵ LONGACRE | 78 | IPWA $\pi N \rightarrow N\pi\pi$ |
| 1616 or 1613 | ² LONGACRE | 77 | IPWA $\pi N \rightarrow N\pi\pi$ |

– 2×IMAGINARY PART

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|-----------------------|------|--|
| 50 to 150 (\approx 100) OUR ESTIMATE | | | |
| 120 | ⁴ HOEHLER | 93 | SPED $\pi N \rightarrow \pi N$ |
| 90 \pm 40 | CUTKOSKY | 80 | IPWA $\pi N \rightarrow \pi N$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| not seen | ARNDT | 91 | DPWA $\pi N \rightarrow \pi N$ Soln SM90 |
| 607 or 567 | ⁵ LONGACRE | 78 | IPWA $\pi N \rightarrow N\pi\pi$ |
| 577 or 575 | ² LONGACRE | 77 | IPWA $\pi N \rightarrow N\pi\pi$ |

N(1700) ELASTIC POLE RESIDUE

MODULUS $|r|$

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|-------------|-------------|------|--------------------------------|
| 5 | HOEHLER | 93 | SPED $\pi N \rightarrow \pi N$ |
| 6 \pm 3 | CUTKOSKY | 80 | IPWA $\pi N \rightarrow \pi N$ |

PHASE θ

| VALUE ($^\circ$) | DOCUMENT ID | TECN | COMMENT |
|--------------------|-------------|------|--------------------------------|
| 0 \pm 50 | CUTKOSKY | 80 | IPWA $\pi N \rightarrow \pi N$ |

N(1700) DECAY MODES

The following branching fractions are our estimates, not fits or averages.

| Mode | Fraction (Γ_i/Γ) |
|---------------------------------------|--------------------------------|
| Γ_1 $N\pi$ | 5–15 % |
| Γ_2 $N\eta$ | |
| Γ_3 ΛK | <3 % |
| Γ_4 ΣK | |
| Γ_5 $N\pi\pi$ | 85–95 % |
| Γ_6 $\Delta\pi$ | |
| Γ_7 $\Delta(1232)\pi$, S-wave | |
| Γ_8 $\Delta(1232)\pi$, D-wave | |
| Γ_9 $N\rho$ | <35 % |

| | | |
|---------------|-----------------------------------|---------------|
| Γ_{10} | $N\rho, S=1/2, D\text{-wave}$ | |
| Γ_{11} | $N\rho, S=3/2, S\text{-wave}$ | |
| Γ_{12} | $N\rho, S=3/2, D\text{-wave}$ | |
| Γ_{13} | $N(\pi\pi)_{S\text{-wave}}^{I=0}$ | |
| Γ_{14} | $p\gamma$ | 0.01–0.05 % |
| Γ_{15} | $p\gamma, \text{helicity}=1/2$ | 0.0–0.024 % |
| Γ_{16} | $p\gamma, \text{helicity}=3/2$ | 0.002–0.026 % |
| Γ_{17} | $n\gamma$ | 0.01–0.13 % |
| Γ_{18} | $n\gamma, \text{helicity}=1/2$ | 0.0–0.09 % |
| Γ_{19} | $n\gamma, \text{helicity}=3/2$ | 0.01–0.05 % |

$N(1700)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$ Γ_1/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|--|
| 0.05 to 0.15 OUR ESTIMATE | | | |
| 0.01 ± 0.02 | MANLEY | 92 | IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$ |
| 0.11 ± 0.05 | CUTKOSKY | 80 | IPWA $\pi N \rightarrow \pi N$ |
| 0.08 ± 0.03 | HOEHLER | 79 | IPWA $\pi N \rightarrow \pi N$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 0.04 ± 0.05 | BATINIC | 95 | DPWA $\pi N \rightarrow N\pi, N\eta$ |

$\Gamma(N\eta)/\Gamma_{\text{total}}$ Γ_2/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|--------------------------------------|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 0.10 ± 0.06 | BATINIC | 95 | DPWA $\pi N \rightarrow N\pi, N\eta$ |

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1700) \rightarrow \Lambda K$ $(\Gamma_1\Gamma_3)^{1/2}/\Gamma$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|--|
| –0.06 to +0.04 OUR ESTIMATE | | | |
| –0.012 | BELL | 83 | DPWA $\pi^- p \rightarrow \Lambda K^0$ |
| –0.012 | SAXON | 80 | DPWA $\pi^- p \rightarrow \Lambda K^0$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| –0.04 | ⁶ BAKER | 78 | DPWA See SAXON 80 |
| –0.03 ± 0.004 | ¹ BAKER | 77 | IPWA $\pi^- p \rightarrow \Lambda K^0$ |
| –0.03 | ¹ BAKER | 77 | DPWA $\pi^- p \rightarrow \Lambda K^0$ |
| +0.026 ± 0.019 | DEVENISH | 74B | Fixed- t dispersion rel. |

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1700) \rightarrow \Sigma K$ $(\Gamma_1\Gamma_4)^{1/2}/\Gamma$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|-----------------------------------|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| not seen | LIVANOS | 80 | DPWA $\pi p \rightarrow \Sigma K$ |
| <0.017 | ⁷ DEANS | 75 | DPWA $\pi N \rightarrow \Sigma K$ |

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the $\Delta(1620) S_{31}$ coupling to $\Delta(1232)\pi$.

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1700) \rightarrow \Delta(1232)\pi$, S-wave $(\Gamma_1 \Gamma_7)^{1/2} / \Gamma$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------------------|-----------------------|------|--|
| 0.00 to ±0.08 OUR ESTIMATE | | | |
| +0.02±0.03 | MANLEY | 92 | IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$ |
| 0.00 | ² LONGACRE | 77 | IPWA $\pi N \rightarrow N\pi\pi$ |
| -0.16 | ³ LONGACRE | 75 | IPWA $\pi N \rightarrow N\pi\pi$ |

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1700) \rightarrow \Delta(1232)\pi$, D-wave $(\Gamma_1 \Gamma_8)^{1/2} / \Gamma$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------------------|-----------------------|------|--|
| ±0.04 to ±0.20 OUR ESTIMATE | | | |
| +0.10±0.09 | MANLEY | 92 | IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$ |
| -0.12 | ² LONGACRE | 77 | IPWA $\pi N \rightarrow N\pi\pi$ |
| +0.14 | ³ LONGACRE | 75 | IPWA $\pi N \rightarrow N\pi\pi$ |

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1700) \rightarrow N\rho$, S=3/2, S-wave $(\Gamma_1 \Gamma_{11})^{1/2} / \Gamma$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------------------|-----------------------|------|--|
| ±0.01 to ±0.13 OUR ESTIMATE | | | |
| -0.04±0.06 | MANLEY | 92 | IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$ |
| -0.07 | ² LONGACRE | 77 | IPWA $\pi N \rightarrow N\pi\pi$ |
| +0.07 | ³ LONGACRE | 75 | IPWA $\pi N \rightarrow N\pi\pi$ |

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1700) \rightarrow N(\pi\pi)_{S=0}^{I=0}$, S-wave $(\Gamma_1 \Gamma_{13})^{1/2} / \Gamma$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------------------|-----------------------|------|--|
| ±0.02 to ±0.28 OUR ESTIMATE | | | |
| +0.02±0.02 | MANLEY | 92 | IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$ |
| 0.00 | ² LONGACRE | 77 | IPWA $\pi N \rightarrow N\pi\pi$ |
| +0.2 | ³ LONGACRE | 75 | IPWA $\pi N \rightarrow N\pi\pi$ |

N(1700) PHOTON DECAY AMPLITUDES

$N(1700) \rightarrow p\gamma$, helicity-1/2 amplitude $A_{1/2}$

| VALUE (GeV ^{-1/2}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|---|
| -0.018±0.013 OUR ESTIMATE | | | |
| -0.016±0.014 | CRAWFORD | 83 | IPWA $\gamma N \rightarrow \pi N$ |
| -0.002±0.013 | AWAJI | 81 | DPWA $\gamma N \rightarrow \pi N$ |
| -0.028±0.007 | ARAI | 80 | DPWA $\gamma N \rightarrow \pi N$ (fit 1) |
| -0.029±0.006 | ARAI | 80 | DPWA $\gamma N \rightarrow \pi N$ (fit 2) |
| -0.024±0.019 | CRAWFORD | 80 | DPWA $\gamma N \rightarrow \pi N$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| -0.033±0.021 | BARBOUR | 78 | DPWA $\gamma N \rightarrow \pi N$ |
| -0.014±0.025 | FELLER | 76 | DPWA $\gamma N \rightarrow \pi N$ |

$N(1700) \rightarrow p\gamma$, helicity-3/2 amplitude $A_{3/2}$

| <u>VALUE (GeV^{-1/2})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|---|
| -0.002±0.024 OUR ESTIMATE | | | |
| -0.009±0.012 | CRAWFORD | 83 | IPWA $\gamma N \rightarrow \pi N$ |
| 0.029±0.014 | AWAJI | 81 | DPWA $\gamma N \rightarrow \pi N$ |
| -0.002±0.005 | ARAI | 80 | DPWA $\gamma N \rightarrow \pi N$ (fit 1) |
| 0.014±0.005 | ARAI | 80 | DPWA $\gamma N \rightarrow \pi N$ (fit 2) |
| -0.017±0.014 | CRAWFORD | 80 | DPWA $\gamma N \rightarrow \pi N$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| -0.014±0.025 | BARBOUR | 78 | DPWA $\gamma N \rightarrow \pi N$ |
| 0.0 ±0.014 | FELLER | 76 | DPWA $\gamma N \rightarrow \pi N$ |

$N(1700) \rightarrow n\gamma$, helicity-1/2 amplitude $A_{1/2}$

| <u>VALUE (GeV^{-1/2})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|---|
| 0.000±0.050 OUR ESTIMATE | | | |
| 0.006±0.024 | AWAJI | 81 | DPWA $\gamma N \rightarrow \pi N$ |
| -0.002±0.013 | FUJII | 81 | DPWA $\gamma N \rightarrow \pi N$ |
| -0.052±0.030 | ARAI | 80 | DPWA $\gamma N \rightarrow \pi N$ (fit 1) |
| -0.055±0.030 | ARAI | 80 | DPWA $\gamma N \rightarrow \pi N$ (fit 2) |
| 0.052±0.035 | CRAWFORD | 80 | DPWA $\gamma N \rightarrow \pi N$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| +0.050±0.042 | BARBOUR | 78 | DPWA $\gamma N \rightarrow \pi N$ |

$N(1700) \rightarrow n\gamma$, helicity-3/2 amplitude $A_{3/2}$

| <u>VALUE (GeV^{-1/2})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|---|
| -0.003±0.044 OUR ESTIMATE | | | |
| -0.033±0.017 | AWAJI | 81 | DPWA $\gamma N \rightarrow \pi N$ |
| 0.018±0.018 | FUJII | 81 | DPWA $\gamma N \rightarrow \pi N$ |
| -0.037±0.036 | ARAI | 80 | DPWA $\gamma N \rightarrow \pi N$ (fit 1) |
| -0.035±0.024 | ARAI | 80 | DPWA $\gamma N \rightarrow \pi N$ (fit 2) |
| 0.041±0.030 | CRAWFORD | 80 | DPWA $\gamma N \rightarrow \pi N$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| +0.035±0.030 | BARBOUR | 78 | DPWA $\gamma N \rightarrow \pi N$ |

$N(1700) \quad \gamma p \rightarrow \Lambda K^+$ AMPLITUDES

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $p\gamma \rightarrow N(1700) \rightarrow \Lambda K^+$ (E_{2-} amplitude)

| <u>VALUE (units 10⁻³)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> |
|---|--------------------|-------------|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | |
| 4.09 | TANABE | 89 DPWA |

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $p\gamma \rightarrow N(1700) \rightarrow \Lambda K^+$ (M_{2-} amplitude)

| <u>VALUE (units 10⁻³)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> |
|---|--------------------|-------------|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | |
| -7.09 | TANABE | 89 DPWA |

$p\gamma \rightarrow N(1700) \rightarrow \Lambda K^+$ phase angle θ (E_{2-} amplitude)

| | | |
|------------------------|--------------------|-------------|
| <u>VALUE (degrees)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> |
|------------------------|--------------------|-------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|--------|--------|----|------|
| - 35.9 | TANABE | 89 | DPWA |
|--------|--------|----|------|

N(1700) FOOTNOTES

- ¹ The two BAKER 77 entries are from an IPWA using the Barrelet-zero method and from a conventional energy-dependent analysis.
- ² LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- ³ From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- ⁴ See HOEHLER 93 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.
- ⁵ LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.
- ⁶ The overall phase of BAKER 78 couplings has been changed to agree with previous conventions.
- ⁷ The range given is from the four best solutions.

N(1700) REFERENCESFor early references, see Physics Letters **111B** 70 (1982).

| | | | | |
|----------|-----|------------------------|--|--------------------|
| BATINIC | 95 | PR C51 2310 | +Slaus, Svarc, Nefkens | (BOSK, UCLA) |
| Also | 98 | PR C57 1004 (erratum) | M. Batinic+ | |
| HOEHLER | 93 | πN Newsletter 9 1 | | (KARL) |
| MANLEY | 92 | PR D45 4002 | +Saleski | (KENT) IJP |
| Also | 84 | PR D30 904 | Manley, Arndt, Goradia, Teplitz | (VPI) |
| ARNDT | 91 | PR D43 2131 | +Li, Roper, Workman, Ford | (VPI, TELE) IJP |
| TANABE | 89 | PR C39 741 | +Kohno, Bennhold | (MANZ) |
| Also | 89 | NC 102A 193 | Kohno, Tanabe, Bennhold | (MANZ) |
| BELL | 83 | NP B222 389 | +Blissett, Broome, Daley, Hart, Lintern+ | (RL) IJP |
| CRAWFORD | 83 | NP B211 1 | +Morton | (GLAS) |
| PDG | 82 | PL 111B | Roos, Porter, Aguilar-Benitez+ | (HELS, CIT, CERN) |
| AWAJI | 81 | Bonn Conf. 352 | +Kajikawa | (NAGO) |
| Also | 82 | NP B197 365 | Fujii, Hayashii, Iwata, Kajikawa+ | (NAGO) |
| FUJII | 81 | NP B187 53 | +Hayashii, Iwata, Kajikawa+ | (NAGO, OSAK) |
| ARAI | 80 | Toronto Conf. 93 | | (INUS) |
| Also | 82 | NP B194 251 | Arai, Fujii | (INUS) |
| CRAWFORD | 80 | Toronto Conf. 107 | | (GLAS) |
| CUTKOSKY | 80 | Toronto Conf. 19 | +Forsyth, Babcock, Kelly, Hendrick | (CMU, LBL) IJP |
| Also | 79 | PR D20 2839 | Cutkosky, Forsyth, Hendrick, Kelly | (CMU, LBL) IJP |
| LIVANOS | 80 | Toronto Conf. 35 | +Baton, Coutures, Kochowski, Neveu | (SACL) IJP |
| SAXON | 80 | NP B162 522 | +Baker, Bell, Blissett, Bloodworth+ | (RHEL, BRIS) IJP |
| HOEHLER | 79 | PDAT 12-1 | +Kaiser, Koch, Pietarinen | (KARLT) IJP |
| Also | 80 | Toronto Conf. 3 | Koch | (KARLT) IJP |
| BAKER | 78 | NP B141 29 | +Blissett, Bloodworth, Broome+ | (RL, CAVE) IJP |
| BARBOUR | 78 | NP B141 253 | +Crawford, Parsons | (GLAS) |
| LONGACRE | 78 | PR D17 1795 | +Lasinski, Rosenfeld, Smadja+ | (LBL, SLAC) |
| BAKER | 77 | NP B126 365 | +Blissett, Bloodworth, Broome, Hart+ | (RHEL) IJP |
| LONGACRE | 77 | NP B122 493 | +Dolbeau | (SACL) IJP |
| Also | 76 | NP B108 365 | Dolbeau, Triantis, Neveu, Cadiet | (SACL) IJP |
| FELLER | 76 | NP B104 219 | +Fukushima, Horikawa, Kajikawa+ | (NAGO, OSAK) IJP |
| DEANS | 75 | NP B96 90 | +Mitchell, Montgomery+ | (SFLA, ALAH) IJP |
| LONGACRE | 75 | PL 55B 415 | +Rosenfeld, Lasinski, Smadja+ | (LBL, SLAC) IJP |
| DEVENISH | 74B | NP B81 330 | +Froggatt, Martin | (DESY, NORD, LOUC) |