

$f_2(1565)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

Seen in antinucleon-nucleon annihilation at rest. See also minireview under non- $q\bar{q}$ candidates. (See the index for the page number.)
Needs confirmation.

$f_2(1565)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1544±17 OUR AVERAGE	Error	includes scale factor of 1.8.	See the ideogram below.
1550±10±20	AMELIN	00 VES	37 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
1575±18	BERTIN	98 OBLX	0.05-0.405 $\bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$
1507±15	¹ BERTIN	97C OBLX	0.0 $\bar{p} p \rightarrow \pi^+ \pi^- \pi^0$
1565±20	MAY	90 ASTE	0.0 $\bar{p} p \rightarrow \pi^+ \pi^- \pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1598±11± 9	BAKER	99B SPEC	0 $\bar{p} p \rightarrow \omega \omega \pi^0$
1534±20	² ABELE	96C RVUE	Compilation
~ 1552	³ AMSLER	95D CBAR	0.0 $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta \eta, \pi^0 \pi^0 \eta$
1598±72	BALOSHIN	95 SPEC	40 $\pi^- C \rightarrow K_S^0 K_S^0 X$
1566 ⁺⁸⁰ ₋₅₀	⁴ ANISOVICH	94 CBAR	0.0 $\bar{p} p \rightarrow 3\pi^0, \eta \eta \pi^0$
1502± 9	ADAMO	93 OBLX	$\bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$
1488±10	⁵ ARMSTRONG	93C E760	$\bar{p} p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$
1508±10	⁵ ARMSTRONG	93D E760	$\bar{p} p \rightarrow 3\pi^0 \rightarrow 6\gamma$
1525±10	⁵ ARMSTRONG	93D E760	$\bar{p} p \rightarrow \eta \pi^0 \pi^0 \rightarrow 6\gamma$
~ 1504	⁶ WEIDENAUER	93 ASTE	0.0 $\bar{p} N \rightarrow 3\pi^- 2\pi^+$
1540±15	⁵ ADAMO	92 OBLX	$\bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$
1515±10	⁷ AKER	91 CBAR	0.0 $\bar{p} p \rightarrow 3\pi^0$
1477± 5	BRIDGES	86C DBC	0.0 $\bar{p} N \rightarrow 3\pi^- 2\pi^+$

¹ T-matrix pole.

² T-matrix pole, large coupling to $\rho\rho$ and $\omega\omega$, could be $f_2(1640)$.


³ Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.

⁴ From a simultaneous analysis of the annihilations $\bar{p} p \rightarrow 3\pi^0, \pi^0 \eta \eta$ including AKER 91 data.

⁵ J^P not determined, could be partly $f_0(1500)$.

⁶ J^P not determined.

⁷ Superseded by AMSLER 95B,



$f_2(1565)$ mass (MeV)

$f_2(1565)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
131 ± 14 OUR AVERAGE			
130 ± 20 ± 40	AMELIN	00 VES	37 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
119 ± 24	BERTIN	98 OBLX	0.05–0.405 $\bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$
130 ± 20	⁸ BERTIN	97C OBLX	0.0 $\bar{p} p \rightarrow \pi^+ \pi^- \pi^0$
170 ± 40	MAY	90 ASTE	0.0 $\bar{p} p \rightarrow \pi^+ \pi^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
180 ± 60	⁹ ABELE	96C RVUE	Compilation
~ 142	¹⁰ AMSLER	95D CBAR	0.0 $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta \eta, \pi^0 \pi^0 \eta$
263 ± 101	BALOSHIN	95 SPEC	40 $\pi^- C \rightarrow K_S^0 K_S^0 X$
166 ⁺ ₋₂₀ 80	¹¹ ANISOVICH	94 CBAR	0.0 $\bar{p} p \rightarrow 3\pi^0, \eta \eta \pi^0$
130 ± 10	¹² ADAMO	93 OBLX	$\bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$
148 ± 27	¹³ ARMSTRONG	93C E760	$\bar{p} p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$
103 ± 15	¹³ ARMSTRONG	93D E760	$\bar{p} p \rightarrow 3\pi^0 \rightarrow 6\gamma$
111 ± 10	¹³ ARMSTRONG	93D E760	$\bar{p} p \rightarrow \eta \pi^0 \pi^0 \rightarrow 6\gamma$
~ 206	¹⁴ WEIDENAUER	93 ASTE	0.0 $\bar{p} N \rightarrow 3\pi^- 2\pi^+$
132 ± 37	¹³ ADAMO	92 OBLX	$\bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$
120 ± 10	¹⁵ AKER	91 CBAR	0.0 $\bar{p} p \rightarrow 3\pi^0$
116 ± 9	BRIDGES	86C DBC	0.0 $\bar{p} N \rightarrow 3\pi^- 2\pi^+$

⁸ T-matrix pole.

⁹ T-matrix pole, large coupling to $\rho\rho$ and $\omega\omega$, could be $f_2(1640)$.

¹⁰ Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.

¹¹ From a simultaneous analysis of the annihilations $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta\eta$ including AKER 91 data.

¹² Supersedes ADAMO 92.

¹³ J^P not determined, could be partly $f_0(1500)$.

¹⁴ J^P not determined.

¹⁵ Superseded by AMSLER 95B,

$f_2(1565)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi$	seen
Γ_2 $\pi^+\pi^-$	seen
Γ_3 $\pi^0\pi^0$	seen
Γ_4 $\rho^0\rho^0$	seen
Γ_5 $2\pi^+2\pi^-$	seen
Γ_6 $\eta\eta$	seen
Γ_7 $a_2(1320)\pi$	not seen
Γ_8 $\omega\omega$	seen

$f_2(1565)$ BRANCHING RATIOS

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen BAKER 99B SPEC 0 $\bar{p}p \rightarrow \omega\omega\pi^0$

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen BERTIN 98 OBLX 0.05–0.405 $\bar{p}p \rightarrow$

not seen ¹⁶ ANISOVICH 94B RVUE $\bar{p}p \rightarrow \pi^+\pi^+\pi^-\pi^0$

seen MAY 89 ASTE $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$

¹⁶ ANISOVICH 94B is from a reanalysis of MAY 90.

$\Gamma(\pi^+\pi^-)/\Gamma(\rho^0\rho^0)$ Γ_2/Γ_4

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.042 ± 0.013 BRIDGES 86B DBC $\bar{p}N \rightarrow 3\pi^-2\pi^+$

$\Gamma(\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE DOCUMENT ID TECN COMMENT

seen AMSLER 95B CBAR 0.0 $\bar{p}p \rightarrow 3\pi^0$

