b-baryon ADMIXTURE (Λ_b , Ξ_b , Σ_b , Ω_b)

b-baryon ADMIXTURE MEAN LIFE

Each measurement of the b-baryon mean life is an average over an admixture of various b baryons which decay weakly. Different techniques emphasize different admixtures of produced particles, which could result in a different b-baryon mean life. More b-baryon flavor specific channels are not included in the measurement.

"OUR EVALUATION" is an average of the data listed below performed by the LEP B Lifetimes Working Group as described in our review "Production and Decay of bflavored Hadrons" in the B^{\pm} Section of these Listings. The averaging procedure takes into account correlations between the measurements and asymmetric lifetime errors.

$VALUE (10^{-12} \text{ s})$	EVTS	DOCUMENT ID	TECN	COMMENT		
1.208±0.051 OUR EVALUATION						
$1.16 \pm 0.20 \pm 0.08$		$^{ m 1}$ ABREU	99w DLPH	$e^+e^- \rightarrow Z$		
$1.19 \ \pm 0.14 \ \pm 0.07$		² ABREU		$e^+e^- ightarrow Z$		
$1.20\ \pm0.08\ \pm0.06$		³ BARATE	98D ALEP	$e^+e^- o Z$		
$1.10 \ ^{+0.19}_{-0.17} \ \pm 0.09$		ABREU	96D DLPH	Excess $\Lambda\mu^-$ impact parameters		
$1.16 \pm 0.11 \pm 0.06$		AKERS	96 OPAL	Excess $\Lambda \ell^-$, decay lengths and impact parameters		
• • • We do not use	the following	g data for averages	s, fits, limits,	•		
$1.14 \pm 0.08 \pm 0.04$		⁴ ABREU	99w DLPH	$e^+e^- ightarrow Z$		
$1.46 \begin{array}{l} +0.22 \\ -0.21 \end{array} \begin{array}{l} +0.07 \\ -0.09 \end{array}$		ABREU	96D DLPH	Repl. by ABREU 99W		
$1.27 \ ^{+0.35}_{-0.29} \ \pm 0.09$		ABREU	95s DLPH	Repl. by ABREU 99W		
$1.05 \ ^{+0.12}_{-0.11} \ \pm 0.09$	290	BUSKULIC	95L ALEP	Repl. by BARATE 98D		
$1.04 \ ^{+0.48}_{-0.38} \ \pm 0.10$	11	⁵ ABREU	93F DLPH	Excess $\Lambda\mu^-$, decay lengths		
$1.05 \ ^{+0.23}_{-0.20} \ \pm 0.08$	157	⁶ AKERS	93 OPAL	Excess $\Lambda \ell^-$, decay lengths		
$1.12 \begin{array}{c} +0.32 \\ -0.29 \end{array} \pm 0.16$	101	⁷ BUSKULIC	92ı ALEP	Excess $\Lambda \ell^-$, impact parameters		

¹ Measured using $\Lambda \ell^-$ decay length.

Created: 12/18/2000 15:07

² Measured using $p\ell^-$ decay length.

³ Measured using the excess of $\Lambda \ell^-$, lepton impact parameter.

⁴This ABREU 99W result is the combined result of the $\Lambda \ell^-$, $p\ell^-$, and excess $\Lambda \mu^$ impact parameter measurements.

⁵ ABREU 93F superseded by ABREU 96D.

⁶ AKERS 93 superseded by AKERS 96.

⁷ BUSKULIC 921 superseded by BUSKULIC 95L.

b-baryon ADMIXTURE $(\Lambda_b, \Xi_b, \Sigma_b, \Omega_b)$

These branching fractions are actually an average over weakly decaying b-baryons weighted by their production rates in Z decay (or high-energy $p\overline{p}$), branching ratios, and detection efficiencies. They scale with the LEP b-baryon production fraction B($b \rightarrow b$ -baryon) and are evaluated for our value B($b \rightarrow b$ -baryon) = (11.6 \pm 2.0)%.

The branching fractions B(b-baryon $\to \Lambda \ell^- \overline{\nu}_\ell$ anything) and B($\Lambda_b^0 \to \Lambda_c^+ \ell^- \overline{\nu}_\ell$ anything) are not pure measurements because the underlying measured products of these with B($b \to b$ -baryon) were used to determine B($b \to b$ -baryon), as described in the note "Production and Decay of b-Flavored Hadrons."

	Mode	Fraction (Γ_i/Γ)	
Γ_1	$p\mu^-\overline{ u}$ anything	(4.2 + 1.8) %	
Γ_2	$oldsymbol{ ho} \ell \overline{ u}_\ell$ anything	($4.1\pm~1.0$) %	
Γ_3	<i>p</i> anything	(51 ± 17) %	
Γ_4	$arLambda \ell^- \overline{ u}_\ell$ anything	(2.7± 0.8) %	
Γ_5	$arLambda\ell^+ u_\ell$ anything		
Γ_6	arLambdaanything		
Γ_7	$arLambda_{m{c}}^+ \ell^- \overline{ u}_\ell$ anything		
Γ ₈	$\Lambda/\overline{\Lambda}$ anything	(28 ± 7)%	
_	$ar{\Xi}^-\ell^-ar{ u}_\ell$ anything	$(4.8 \pm 1.3) \times 10^{-3}$	

b-baryon ADMIXTURE (Λ_b , Ξ_b , Σ_b , Ω_b) BRANCHING RATIOS

$\Gamma(p\mu^-\overline{\nu}anything)/\Gamma_{total}$ Γ_1/Γ_{NALUE} EVTS DOCUMENT ID TECN COMMENT $0.042^{+0.016}_{-0.013}\pm0.007$ 125 8 ABREU 958 DLPH $e^+e^- \rightarrow Z$

⁸ ABREU 95S reports [B(b-baryon $\to p\mu^-\overline{\nu}$ anything) \times B($\overline{b} \to b$ -baryon)] = 0.0049 \pm 0.0011 $^+$ 0.0015. We divide by our best value B($\overline{b} \to b$ -baryon) = (11.6 \pm 2.0) \times 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\ell\overline{ u}_\ell$ anything)/ $\Gamma_{ ext{total}}$				Γ_2/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
0.041±0.007±0.007	9 BARATE	98V ALEP	$e^+e^- \rightarrow Z$	

⁹ BARATE 98V reports [B(b-baryon $\to p\ell\overline{\nu}_\ell$ anything) \times B($\overline{b}\to b$ -baryon)] = (4.72 \pm 0.66 \pm 0.44) \times 10⁻³. We divide by our best value B($\overline{b}\to b$ -baryon) = (11.6 \pm 2.0) \times 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(ho\ell\overline{ u}_\ell$ anything)/ $\Gamma(ho$ anything)	Γ_2/Γ_3
(perfangumg)/1 (pangumg)	12/13

VALUEDOCUMENT IDTECNCOMMENT $\mathbf{0.080 \pm 0.012 \pm 0.014}$ BARATE98V ALEP $e^+e^- \rightarrow Z$

HTTP://PDG.LBL.GOV Page 2 Created: 12/18/2000 15:07

$\Gamma(\Lambda \ell^- \overline{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}$

 Γ_{4}/Γ

The values and averages in this section serve only to show what values result if one assumes our $B(b \to b\text{-baryon})$. They cannot be thought of as measurements since the underlying product branching fractions were also used to determinine $B(b \to b\text{-baryon})$ as described in the note on "Production and Decay of b-Flavored Hadrons."

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.027±0.008 OUR AVER	RAGE				
$0.028\!\pm\!0.004\!\pm\!0.005$		¹⁰ BARATE	98D	ALEP	$e^+e^- \rightarrow Z$
$0.025\!\pm\!0.003\!\pm\!0.004$		¹¹ AKERS	96	OPAL	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$
$0.026 \pm 0.006 \pm 0.004$	262	¹² ABREU	95 S	DLPH	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$
$0.053 \!\pm\! 0.010 \!\pm\! 0.009$	290	¹³ BUSKULIC	95L	ALEP	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$
• • • We do not use the	followin	ng data for averages	, fits,	limits,	etc. • • •
seen	157	¹⁴ AKERS	93	OPAL	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$
$0.060 \pm 0.018 \pm 0.010$	101	¹⁵ BUSKULIC	921	ALEP	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$
¹⁰ BARATE 98D reports [B(b -baryon $\to \Lambda \ell^- \overline{\nu}_\ell$ anything) \times B($\overline{b} \to b$ -baryon)] = 0.00326 \pm 0.00016 \pm 0.00039. We divide by our best value B($\overline{b} \to b$ -baryon) = (11.6 \pm 2.0) \times 10 ⁻² . Our first error is their experiment's error and our second error					
is the systematic error from using our best value. Measured using the excess of $\Lambda\ell^-$, lepton impact parameter.					
		on $ ightarrow$ $arLambda \ell^- \overline{ u}_\ell$ anyth	ing) >	× В(<u>Б</u> –	$_{ ightarrow}$ <i>b</i> -baryon)] = 0.00291 \pm

- ¹¹ AKERS 96 reports [B(*b*-baryon $\to \Lambda \ell^- \overline{\nu}_\ell$ anything) \times B($\overline{b} \to b$ -baryon)] = 0.00291 \pm 0.00023 \pm 0.00025. We divide by our best value B($\overline{b} \to b$ -baryon) = (11.6 \pm 2.0) \times 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ¹² ABREU 95S reports [B(b-baryon $\to \Lambda \ell^- \overline{\nu}_\ell$ anything) \times B($\overline{b} \to b$ -baryon)] = 0.0030 \pm 0.0006 \pm 0.0004. We divide by our best value B($\overline{b} \to b$ -baryon) = (11.6 \pm 2.0) \times 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ¹³ BUSKULIC 95L reports [B(b-baryon $\to \Lambda \ell^- \overline{\nu}_\ell$ anything) \times B($\overline{b} \to b$ -baryon)] = 0.0061 \pm 0.0006 \pm 0.0010. We divide by our best value B($\overline{b} \to b$ -baryon) = (11.6 \pm 2.0) \times 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ¹⁴ AKERS 93 superseded by AKERS 96.
- ¹⁵ BUSKULIC 92I reports $[B(b\text{-baryon} \to \Lambda \ell^- \overline{\nu}_\ell \text{ anything}) \times B(\overline{b} \to b\text{-baryon})] = 0.0070 \pm 0.0010 \pm 0.0018$. We divide by our best value $B(\overline{b} \to b\text{-baryon}) = (11.6 \pm 2.0) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Superseded by BUSKULIC 95L.

$\Gamma(\Lambda \ell^+ \nu_{\ell} \text{ anything}) / \Gamma(\Lambda \text{ anything})$

 Γ_5/Γ_6

Created: 12/18/2000 15:07

(*) 0)/ ()	υ,			<i>3</i> / 0
VALUE	DOCUMENT ID	TECN	COMMENT	
$0.080 \pm 0.012 \pm 0.008$	ABBIENDI	99L OPAL	$e^+e^- ightarrow Z$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $0.070\pm0.012\pm0.007$ ACKERSTAFF 97N OPAL Repl. by ABBIENDI 99L

$\Gamma(\Lambda/\overline{\Lambda})$ anything $\Gamma(\Lambda/\overline{\Lambda})$				Γ_8/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
0.28±0.07 OUR AVERAGE				
$0.30 \pm 0.04 \pm 0.05$	¹⁶ ABBIENDI	99L OPAL	$e^+e^- ightarrow Z$	
$0.19^{\color{red}+0.11}_{-0.07}\!\pm\!0.03$	¹⁷ ABREU	95C DLPH	$e^+e^- ightarrow Z$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $0.34 \pm 0.05 \pm 0.06$ 18 ACKERSTAFF 97N OPAL Repl. by ABBIENDI 99L

- ¹⁶ ABBIENDI 99L reports [B(*b*-baryon $\rightarrow \Lambda/\overline{\Lambda}$ anything) \times B($\overline{b} \rightarrow b$ -baryon)] = 0.035 \pm 0.0032 \pm 0.0035. We divide by our best value B($\overline{b} \rightarrow b$ -baryon) = (11.6 \pm 2.0) \times 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.
- 17 ABREU 95C reports $0.28^{+0.17}_{-0.12}$ for B(\$\overline{b}\$ \rightarrow \$b\$-baryon) = 0.08 \pm 0.02. We rescale to our best value B(\$\overline{b}\$ \rightarrow \$b\$-baryon) = (11.6 \pm 2.0) \times 10 $^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ¹⁸ ACKERSTAFF 97N reports [B(b-baryon $\to \Lambda/\overline{\Lambda}$ anything) \times B($\overline{b} \to b$ -baryon)] = 0.0393 \pm 0.0046 \pm 0.0037. We divide by our best value B($\overline{b} \to b$ -baryon) = (11.6 \pm 2.0) \times 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Xi^-\ell^-\overline{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}$

 Γ_0/Γ

VALUE	DOCUMENT ID	TI	ECN	COMMENT
0.0048±0.0013 OUR AVERAGE				
$0.0047\!\pm\!0.0012\!\pm\!0.0008$	¹⁹ BUSKULIC	96T A	LEP	Excess $\Xi^-\ell^-$ over $\Xi^-\ell^+$
$0.0051 \!\pm\! 0.0020 \!\pm\! 0.0009$	²⁰ ABREU			Excess $\Xi^-\ell^-$ over $\Xi^-\ell^+$

- 19 BUSKULIC 96T reports [B(b-baryon $\to \Xi^-\ell^-\overline{\nu}_\ell$ anything) \times B(\$\overline{b}\$ \$\to\$ b-baryon)] = 0.00054 \pm 0.00011 \pm 0.00008. We divide by our best value B(\$\overline{b}\$ \$\to\$ b-baryon) = (11.6 \pm 2.0) \times 10 $^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ²⁰ ABREU 95V reports $[B(b\text{-baryon} \to \overline{\Xi}^-\ell^-\overline{\nu}_\ell \text{ anything}) \times B(\overline{b} \to b\text{-baryon})] = 0.00059 \pm 0.00021 \pm 0.0001$. We divide by our best value $B(\overline{b} \to b\text{-baryon}) = (11.6 \pm 2.0) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

b-baryon ADMIXTURE (Λ_b , Ξ_b , Σ_b , Ω_b) REFERENCES

Created: 12/18/2000 15:07