			NODE=S091
$B_c^{\pm}$	$I(J^P) = 0(0^-)$ I, J, P need con	firmation.	
Quantum numbers sho	wn are quark-model predictions.		NODE=S091
	$B_c^{\pm}$ MASS		NODE=S091205
<ul> <li>VALUE (GeV)</li> <li>6.4 ±0.39±0.13</li> <li>● ● We do not use the followin</li> <li>6.32±0.06</li> </ul>	DOCUMENT ID     TECN     CC       1     ABE     98M CDF     p7       g data for averages, fits, limits, etc     2     ACKERSTAFF     980 OPAL     e <sup>-1</sup>	$\frac{DMMENT}{\overline{p} \ 1.8 \ \text{TeV}}$ $\therefore \bullet \bullet \bullet$ $+ e^{-} \rightarrow Z$	NODE=S091M
<sup>1</sup> ABE 98M observed $20.4^{+6.2}_{-5.5}$ > 4.8 standard deviations. Th	events in the $B_c^+ \rightarrow J/\psi(1s) \ell \nu$ the mass value is estimated from $m(t)$	$\ell$ with a significance of $(J/\psi(1S) \ell)$ .	NODE=S091M;LINKAGE=A
an estimated background of 0	$A_{\rm c}^{\rm c}$ and date events in the $B_{\rm c} \rightarrow J/$ .63 $\pm$ 0.20 events.	$\psi(13)\pi^+$ channel with	NODE=S091M;LINKAGE=D
	$B_c^{\pm}$ MEAN LIFE		NODE=S091210
VALUE (10 <sup>-12</sup> s)	DOCUMENT ID TECN CO	OMMENT	NODE=S091T
$0.46^{+0.18}_{-0.16}{\pm}0.03$	<sup>3</sup> АВЕ 98м CDF р7	p 1.8 TeV	
<sup>3</sup> The lifetime is measured from	the $J/\psi(1S)\ell$ decay vertices.	I	NODE=S091T;LINKAGE=A
$B_c^+$ DEC	CAY MODES $\times B(\overline{b} \rightarrow B_c)$		NODE=S091215;NODE=S091
$B_c^-$ modes are charge co	onjugates of the modes below.		NODE=S091
Mode	Fraction $(\Gamma_i/\Gamma)$	Confidence level	
The following quanities $\Gamma_i/\Gamma \times B(\overline{b} \to B_c)$ .	are not pure branching ratios; rath	er the fraction	CLUMP=A;NODE=S091
$\Gamma_1 = J/\psi(1S)\ell^+ u_\ell$ anything	$(5.2^{+2.4}_{-2.1}) \times 1$	<sub>10</sub> -5	DESIG=1;OUR EVAL
$ \begin{array}{ll} \Gamma_2 & J/\psi(1S)\pi^+ \\ \Gamma_3 & J/\psi(1S)\pi^+\pi^+\pi^- \\ \Gamma_4 & J/\psi(1S)a_1(1260) \\ \Gamma_5 & D^*(2010)^+\overline{D}{}^0 \end{array} $	$< 8.2 \times 1 \\ < 5.7 \times 1 \\ < 1.2 \times 1 \\ < 6.2 \times 1 $	$\begin{array}{cccc} 10^{-5} & 90\% \\ 10^{-4} & 90\% \\ 10^{-3} & 90\% \\ 10^{-3} & 90\% \end{array}$	DESIG=2;OUR EVAL DESIG=3;OUR EVAL DESIG=4;OUR EVAL DESIG=5;OUR EVAL
B_c^+	BRANCHING RATIOS		NODE=S091225
$\frac{\Gamma(J/\psi(1S)\ell^+\nu_\ell \text{ anything})}{\frac{VALUE}{2L}}$	$\begin{array}{c} \mathbf{b} \\ \mathbf{b} \\ \underline{b} \\ \underline{b} \\ \underline{b} \\ \underline{b} \\ \underline{b} \\ \underline{c} \\ $	Γ <sub>1</sub> /Γ × Β	NODE=S091R1 NODE=S091R1
$(5.2^{+2.4}_{-2.1}) \times 10^{-5}$	<sup>4</sup> ABE 98M CDF <i>p</i>	₱ 1.8 TeV	
• • • We do not use the followin $< 1.6 \times 10^{-4}  90$ $< 1.9 \times 10^{-4}  90$ $< 1.2 \times 10^{-4}  90$	g data for averages, fits, limits, etc <sup>5</sup> ACKERSTAFF 980 OPAL e <sup>-</sup> <sup>6</sup> ABREU 97E DLPH e <sup>-</sup> <sup>7</sup> BARATE 97H ALEP e <sup>-</sup>	$\begin{array}{ccc} & \bullet \bullet \bullet \bullet \\ & + e^- \to Z \\ & + e^- \to Z \\ & + e^- \to Z \end{array}$	
<sup>4</sup> ABE 98M result is derived from $[\sigma(B^+)  imes B(B^+  o J/\psi(1S))]$ by using PDG 98 values of B(	m the measurement of $[\sigma(B_c) \times B(R_c)] = 0.132 \stackrel{+0.041}{_{-}0.037} (\text{stat}) \pm 0.03$ $b \rightarrow B^+)$ and $B(B^+ \rightarrow J/\psi(15))$	$\begin{array}{l} B_{c} \rightarrow J/\psi(1S) \ell \nu_{\ell})] \ / \\ h1(sys) {+0.032 \atop -0.020} (lifteime) \\ \delta ) \ K^{+}). \end{array}$	NODE=S091R1;LINKAGE=C
<sup>5</sup> ACKERSTAFF 980 reports E $6.95 \times 10^{-5}$ at 90%CL. We	NODE=S091R1;LINKAGE=D		
$^{\circ}$ ABREU 97E value listed is for $ au_{B_c} = 1.4$ ps.	an assumed $\tau_{B_c}=$ 0.4 ps and imp	roves to $1.6  imes 10^{-4}$ for	NODE=S091R1;LINKAGE=A
<sup>7</sup> BARATE 97H reports $B(Z \rightarrow at 90\%$ CL. We rescale to our candidate event is found, co which gives $m_{B_c} = 5.96^{+0.2}_{-0.1}$	$B_c X)/B(Z \rightarrow q q) \cdot B(B_c \rightarrow J/\psi)$ PDG 96 values of $B(Z \rightarrow b \overline{b})$ . A matrix matrix of $B(Z \rightarrow b \overline{b})$ is a second seco	$\begin{array}{l} (1S)\ell\nu_\ell) < 5.2 \times 10^{-5} \\ {\rm B}_c^+ \to ~J/\psi(1S)\mu^+\nu_\mu \\ {\rm und \ sources} \ 2 \times 10^{-3}, \end{array}$	NODE=S091R1;LINKAGE=B

Page 1

7/6/2000 09:36

			7/6/2000 09:36	Page 2
$\Gamma(J/\psi(1S)\pi^+)/VALUE$	$\Gamma_{\text{total}} \times B(\overline{b} \to B_c)$	<b>Г<sub>2</sub>/Г × В</b>	NODE=S091R2 NODE=S091R2	
<8.2 × 10 <sup>-5</sup>	90 8 BARATE 97H ALEP $e^+$	$e^- \rightarrow Z$		
• • • We do not us	e the following data for averages, fits, limits, etc.	• • •		
$< 2.4 \times 10^{-4}$	90 <sup>9</sup> ACKERSTAFF 980 OPAL $e^+$	$e^- \rightarrow Z$		
$< 3.4 \times 10^{-4}$	90 $10$ ABREU 97E DLPH $e^+$	$e^- \rightarrow Z$		
$< 2.0 \times 10^{-5}$	95 <sup>11</sup> ABE 96R CDF $p\overline{p}$	1.8 TeV		
<sup>8</sup> BARATE 97H re at 90%CL. We	ports $B(Z \rightarrow B_C X)/B(Z \rightarrow q q) \cdot B(B_C \rightarrow J/\psi$ rescale to our PDG 96 values of $B(Z \rightarrow b\overline{b})$ .	$(1S)\pi) < 3.6  imes 10^{-5}$	NODE=S091R2;LINKAGI	E=B
$^9$ ACKERSTAFF $1.06  imes 10^{-4}$ at	980 reports $B(Z \rightarrow B_c X)/B(Z \rightarrow q q) \times B(B_c Q)$ 90%CL. We rescale to our PDG 98 values of $B(Z)$	$f_{2} \rightarrow J/\psi(1S)\pi^{+}) < S_{2} \rightarrow b\overline{b}.$	NODE=S091R2;LINKAG	E=D
$^{10}$ ABREU 97E val $ au_{B_c}=$ 1.4 ps.	ue listed is for an assumed $\tau_{B_c} = 0.4 \text{ ps and improved}$	oves to $2.7 \times 10^{-4}$ for	NODE=S091R2;LINKAGI	E=A
<sup>11</sup> ABE 96R report $J/\psi(1S)K^+$ < 0.17 ps< $\tau_{B_c}$ < and B(B <sup>+</sup> $\rightarrow$	s B( $b \rightarrow B_c X$ )/B( $b \rightarrow B^+ X$ )·B( $B_c^+ \rightarrow J/$ ( 0.053 at 95%CL for $\tau_{B_c} = 0.8$ ps. It changes 1.6 ps. We rescale to our PDG 96 values of B( $b \rightarrow U/\psi(15) K^+$ ) = 0.00101 + 0.00014	$(\psi(1S) \pi^+)/B(B^+ \rightarrow from 0.15 to 0.04 for B^+) = 0.378 \pm 0.022$	NODE=S091R2;LINKAGI	E=C
$\Gamma(J/\psi(1S)\pi^+\pi^-)$ VALUE	$^{+}\pi^{-})/\Gamma_{\text{total}} \times B(\overline{b} \to B_{c})$ $\stackrel{CL\%}{=} 2000000000000000000000000000000000000$	Г <sub>3</sub> /Г × В	NODE=S091R3 NODE=S091R3	
<5.7 × 10 <sup>=4</sup>	90 <sup>12</sup> ABREU 97E DLPH $e^+$	$e^- \rightarrow Z$		
<sup>12</sup> ABREU 97E val	ue listed is independent of 0.4 ps< $\tau_{\mbox{\it B}_{\rm C}} <$ 1.4 ps.		NODE=S091R3;LINKAGI	E=A
$\frac{\Gamma(J/\psi(1S)a_1(12))}{(J^2+V^2)} = \frac{V^2}{(J^2+V^2)^2}$	$60))/\Gamma_{\text{total}} \times B(\overline{b} \to B_c)$ $\qquad \qquad $	$\Gamma_4/\Gamma \times B$ <u>MMENT</u>	NODE=S091R4 NODE=S091R4	
$^{13}$ ACKERSTAFF 9 $< 5.29 \times 10^{-4}$	at 90%CL. We rescale to our PDG 98 values of E		NODE=S091R4;LINKAG	E=D
$\frac{\Gamma(D^*(2010)^+ \overline{D}^0)}{\frac{VALUE}{2}}$	$\frac{)/\Gamma_{\text{total}} \times B(\overline{b} \to B_c)}{\frac{CL\%}{200}}$	Γ <sub>5</sub> /Γ × Β	NODE=S091R5 NODE=S091R5	
<6.2 × 10 <sup>-3</sup>	90 <sup>14</sup> BARATE 98Q ALEP $e^+$	$e^- \rightarrow Z$		
<sup>14</sup> BARATE 98Q r 90%CL. We reso	eports $B(Z \rightarrow B_{c}X) \times B(B_{c} \rightarrow D^{*}(2010)^{+}\overline{L})$ cale to our PDG 98 values of $B(Z \rightarrow b\overline{b})$ .	$(\overline{b}^0) < 1.9  imes 10^{-3}$ at $($	NODE=S091R5;LINKAG	E=A

 $B_c^{\pm}$  REFERENCES

ABE Also ACKERSTAFF	98M 98R 98O	PRL 81 2432 PR D58 112004 PL B420 157 EPL C4 387	F. Abe <i>et al.</i> F. Abe <i>et al.</i> K. Ackerstaff <i>et al.</i>	(CDF Collab.) (CDF Collab.) (OPAL Collab.) (ALEPH Collab.)
PDG ABREU	98Q 98 97E	EPJ C4 387 EPJ C3 1 PL B398 207	C. Caso <i>et al.</i> P. Abreu <i>et al.</i>	(DELPHI Collab.)
BARATE ABE PDG	97H 96R 96	PL B402 213 PRL 77 5176 PR D54 1	R. Barate <i>et al.</i> F. Abe <i>et al.</i>	`(ALEPH Collab.) (CDF Collab.)

NODE=S091

REFID=46120
REFID=46488
REFID=46029
REFID=46146
REFID=45838
REFID=45322
REFID=45480
REFID=45159
REFID=44495