

N(1440) P₁₁ $I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$ Status: ***

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters **111B** (1982).

N(1440) BREIT-WIGNER MASS

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|--------------------------|------|-------------------------------------|
| 1430 to 1470 (≈ 1440) OUR ESTIMATE | | | |
| 1462 \pm 10 | MANLEY 92 | IPWA | $\pi N \rightarrow \pi N & N\pi\pi$ |
| 1440 \pm 30 | CUTKOSKY 80 | IPWA | $\pi N \rightarrow \pi N$ |
| 1410 \pm 12 | HOEHLER 79 | IPWA | $\pi N \rightarrow \pi N$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 1479 \pm 80 | VRANA 00 | DPWA | Multichannel |
| 1463 \pm 7 | ARNDT 96 | IPWA | $\gamma N \rightarrow \pi N$ |
| 1467 | ARNDT 95 | DPWA | $\pi N \rightarrow N\pi$ |
| 1421 \pm 18 | BATINIC 95 | DPWA | $\pi N \rightarrow N\pi, N\eta$ |
| 1465 | LI 93 | IPWA | $\gamma N \rightarrow \pi N$ |
| 1471 | CUTKOSKY 90 | IPWA | $\pi N \rightarrow \pi N$ |
| 1411 | CRAWFORD 80 | DPWA | $\gamma N \rightarrow \pi N$ |
| 1472 | ¹ BAKER 79 | DPWA | $\pi^- p \rightarrow n\eta$ |
| 1417 | BARBOUR 78 | DPWA | $\gamma N \rightarrow \pi N$ |
| 1460 | BERENDS 77 | IPWA | $\gamma N \rightarrow \pi N$ |
| 1380 | ² LONGACRE 77 | IPWA | $\pi N \rightarrow N\pi\pi$ |
| 1390 | ³ LONGACRE 75 | IPWA | $\pi N \rightarrow N\pi\pi$ |

N(1440) BREIT-WIGNER WIDTH

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|--------------------------|------|-------------------------------------|
| 250 to 450 (≈ 350) OUR ESTIMATE | | | |
| 391 \pm 34 | MANLEY 92 | IPWA | $\pi N \rightarrow \pi N & N\pi\pi$ |
| 545 \pm 170 | CUTKOSKY 90 | IPWA | $\pi N \rightarrow \pi N$ |
| 340 \pm 70 | CUTKOSKY 80 | IPWA | $\pi N \rightarrow \pi N$ |
| 135 \pm 10 | HOEHLER 79 | IPWA | $\pi N \rightarrow \pi N$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 490 \pm 120 | VRANA 00 | DPWA | Multichannel |
| 360 \pm 20 | ARNDT 96 | IPWA | $\gamma N \rightarrow \pi N$ |
| 440 | ARNDT 95 | DPWA | $\pi N \rightarrow N\pi$ |
| 250 \pm 63 | BATINIC 95 | DPWA | $\pi N \rightarrow N\pi, N\eta$ |
| 315 | LI 93 | IPWA | $\gamma N \rightarrow \pi N$ |
| 334 | CRAWFORD 80 | DPWA | $\gamma N \rightarrow \pi N$ |
| 113 | ¹ BAKER 79 | DPWA | $\pi^- p \rightarrow n\eta$ |
| 331 | BARBOUR 78 | DPWA | $\gamma N \rightarrow \pi N$ |
| 279 | BERENDS 77 | IPWA | $\gamma N \rightarrow \pi N$ |
| 200 | ² LONGACRE 77 | IPWA | $\pi N \rightarrow N\pi\pi$ |
| 200 | ³ LONGACRE 75 | IPWA | $\pi N \rightarrow N\pi\pi$ |

N(1440) POLE POSITION**REAL PART**

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|-----------------------|------|--|
| 1345 to 1385 (\approx 1365) OUR ESTIMATE | | | |
| 1346 | ⁴ ARNDT | 95 | DPWA $\pi N \rightarrow N\pi$ |
| 1385 | ⁵ HOEHLER | 93 | SPED $\pi N \rightarrow \pi N$ |
| 1370 | CUTKOSKY | 90 | IPWA $\pi N \rightarrow \pi N$ |
| 1375±30 | CUTKOSKY | 80 | IPWA $\pi N \rightarrow \pi N$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 1383 | VRANA | 00 | DPWA Multichannel |
| 1360 | ⁶ ARNDT | 91 | DPWA $\pi N \rightarrow \pi N$ Soln SM90 |
| 1381 or 1379 | ⁷ LONGACRE | 78 | IPWA $\pi N \rightarrow N\pi\pi$ |
| 1360 or 1333 | ² LONGACRE | 77 | IPWA $\pi N \rightarrow N\pi\pi$ |

-2×IMAGINARY PART

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|-----------------------|------|--|
| 160 to 260 (\approx 210) OUR ESTIMATE | | | |
| 176 | ⁴ ARNDT | 95 | DPWA $\pi N \rightarrow N\pi$ |
| 164 | ⁵ HOEHLER | 93 | SPED $\pi N \rightarrow \pi N$ |
| 228 | CUTKOSKY | 90 | IPWA $\pi N \rightarrow \pi N$ |
| 180±40 | CUTKOSKY | 80 | IPWA $\pi N \rightarrow \pi N$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 316 | VRANA | 00 | DPWA Multichannel |
| 252 | ⁶ ARNDT | 91 | DPWA $\pi N \rightarrow \pi N$ Soln SM90 |
| 209 or 210 | ⁷ LONGACRE | 78 | IPWA $\pi N \rightarrow N\pi\pi$ |
| 167 or 234 | ² LONGACRE | 77 | IPWA $\pi N \rightarrow N\pi\pi$ |

N(1440) ELASTIC POLE RESIDUE**MODULUS |r|**

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|--------------------|------|--|
| 42 | ⁴ ARNDT | 95 | DPWA $\pi N \rightarrow N\pi$ |
| 40 | HOEHLER | 93 | SPED $\pi N \rightarrow \pi N$ |
| 74 | CUTKOSKY | 90 | IPWA $\pi N \rightarrow \pi N$ |
| 52±5 | CUTKOSKY | 80 | IPWA $\pi N \rightarrow \pi N$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 109 | ⁶ ARNDT | 91 | DPWA $\pi N \rightarrow \pi N$ Soln SM90 |

PHASE θ

| VALUE (°) | DOCUMENT ID | TECN | COMMENT |
|---|--------------------|------|--|
| -101 | ⁴ ARNDT | 95 | DPWA $\pi N \rightarrow N\pi$ |
| -84 | CUTKOSKY | 90 | IPWA $\pi N \rightarrow \pi N$ |
| -100±35 | CUTKOSKY | 80 | IPWA $\pi N \rightarrow \pi N$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -93 | ⁶ ARNDT | 91 | DPWA $\pi N \rightarrow \pi N$ Soln SM90 |

N(1440) DECAY MODES

The following branching fractions are our estimates, not fits or averages.

| Mode | Fraction (Γ_i/Γ) |
|---|--------------------------------|
| $\Gamma_1 N\pi$ | 60–70 % |
| $\Gamma_2 N\eta$ | |
| $\Gamma_3 N\pi\pi$ | 30–40 % |
| $\Gamma_4 \Delta\pi$ | 20–30 % |
| $\Gamma_5 \Delta(1232)\pi, P\text{-wave}$ | |
| $\Gamma_6 N\rho$ | <8 % |
| $\Gamma_7 N\rho, S=1/2, P\text{-wave}$ | |
| $\Gamma_8 N\rho, S=3/2, P\text{-wave}$ | |
| $\Gamma_9 N(\pi\pi)^{I=0}_{S\text{-wave}}$ | 5–10 % |
| $\Gamma_{10} p\gamma$ | 0.035–0.048 % |
| $\Gamma_{11} p\gamma, \text{ helicity}=1/2$ | 0.035–0.048 % |
| $\Gamma_{12} n\gamma$ | 0.009–0.032 % |
| $\Gamma_{13} n\gamma, \text{ helicity}=1/2$ | 0.009–0.032 % |

N(1440) BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$

0.6 to 0.7 OUR ESTIMATE

| VALUE | DOCUMENT ID | TECN | COMMENT | Γ_1/Γ |
|--|-------------|------|-------------------------------------|-------------------|
| 0.69 ± 0.03 | MANLEY 92 | IPWA | $\pi N \rightarrow \pi N & N\pi\pi$ | |
| 0.68 ± 0.04 | CUTKOSKY 80 | IPWA | $\pi N \rightarrow \pi N$ | |
| 0.51 ± 0.05 | HOEHLER 79 | IPWA | $\pi N \rightarrow \pi N$ | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.72 ± 0.05 | VRANA 00 | DPWA | Multichannel | |
| 0.68 | ARNDT 95 | DPWA | $\pi N \rightarrow N\pi$ | |
| 0.56 ± 0.08 | BATINIC 95 | DPWA | $\pi N \rightarrow N\pi, N\eta$ | |

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1440) \rightarrow N\eta$

0.6 to 0.7 OUR ESTIMATE

| VALUE | DOCUMENT ID | TECN | COMMENT | $(\Gamma_1\Gamma_2)^{1/2}/\Gamma$ |
|--|--------------------------|------|-----------------------------|-----------------------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| seen | ¹ BAKER 79 | DPWA | $\pi^- p \rightarrow n\eta$ | |
| +0.328 | ⁸ FELTESSE 75 | DPWA | 1488–1745 MeV | |

$\Gamma(N\eta)/\Gamma_{\text{total}}$

0.6 to 0.7 OUR ESTIMATE

| VALUE | DOCUMENT ID | TECN | COMMENT | Γ_2/Γ |
|-----------------|-------------|------|--------------|-------------------|
| 0.00 ± 0.01 | VRANA 00 | DPWA | Multichannel | |

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase

ambiguity is resolved by choosing a negative sign for the $\Delta(1620)$ S_{31} coupling to $\Delta(1232)\pi$.

$$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}} \text{ in } N\pi \rightarrow N(1440) \rightarrow \Delta(1232)\pi, P\text{-wave} \quad (\Gamma_1 \Gamma_5)^{1/2} / \Gamma$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

+0.37 to +0.41 OUR ESTIMATE

| | | | | |
|--------------|--------------|----|------|-------------------------------------|
| +0.39 ± 0.02 | MANLEY | 92 | IPWA | $\pi N \rightarrow \pi N & N\pi\pi$ |
| +0.41 | 2,9 LONGACRE | 77 | IPWA | $\pi N \rightarrow N\pi\pi$ |
| +0.37 | 3 LONGACRE | 75 | IPWA | $\pi N \rightarrow N\pi\pi$ |

$$\Gamma(\Delta(1232)\pi, P\text{-wave}) / \Gamma_{\text{total}} \quad \Gamma_5 / \Gamma$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

0.16 ± 0.01

| DOCUMENT ID | TECN | COMMENT |
|-------------|------|-------------------|
| VRANA | 00 | DPWA Multichannel |

$$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}} \text{ in } N\pi \rightarrow N(1440) \rightarrow N\rho, S=1/2, P\text{-wave} \quad (\Gamma_1 \Gamma_7)^{1/2} / \Gamma$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

±0.07 to ±0.25 OUR ESTIMATE

| | | | | |
|-------|--------------|----|------|-----------------------------|
| -0.11 | 2,9 LONGACRE | 77 | IPWA | $\pi N \rightarrow N\pi\pi$ |
| +0.23 | 3 LONGACRE | 75 | IPWA | $\pi N \rightarrow N\pi\pi$ |

$$\Gamma(N\rho, S=1/2, P\text{-wave}) / \Gamma_{\text{total}} \quad \Gamma_7 / \Gamma$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

0.00 ± 0.01

| DOCUMENT ID | TECN | COMMENT |
|-------------|------|-------------------|
| VRANA | 00 | DPWA Multichannel |

$$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}} \text{ in } N\pi \rightarrow N(1440) \rightarrow N\rho, S=3/2, P\text{-wave} \quad (\Gamma_1 \Gamma_8)^{1/2} / \Gamma$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

+0.18

| DOCUMENT ID | TECN | COMMENT |
|--------------|------|----------------------------------|
| 2,9 LONGACRE | 77 | IPWA $\pi N \rightarrow N\pi\pi$ |

$$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}} \text{ in } N\pi \rightarrow N(1440) \rightarrow N(\pi\pi)_{S\text{-wave}}^{I=0} \quad (\Gamma_1 \Gamma_9)^{1/2} / \Gamma$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

±0.17 to ±0.25 OUR ESTIMATE

| | | | | |
|--------------|--------------|----|------|-------------------------------------|
| +0.24 ± 0.03 | MANLEY | 92 | IPWA | $\pi N \rightarrow \pi N & N\pi\pi$ |
| -0.18 | 2,9 LONGACRE | 77 | IPWA | $\pi N \rightarrow N\pi\pi$ |
| -0.23 | 3 LONGACRE | 75 | IPWA | $\pi N \rightarrow N\pi\pi$ |

$$\Gamma(N(\pi\pi)_{S\text{-wave}}^{I=0}) / \Gamma_{\text{total}} \quad \Gamma_9 / \Gamma$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

0.12 ± 0.01

| DOCUMENT ID | TECN | COMMENT |
|-------------|------|-------------------|
| VRANA | 00 | DPWA Multichannel |

N(1440) PHOTON DECAY AMPLITUDES

$$N(1440) \rightarrow p\gamma, \text{ helicity-1/2 amplitude } A_{1/2}$$

$$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}} \text{ in } N(1440) \rightarrow p\gamma, \text{ helicity-1/2 amplitude } A_{1/2}$$

| VALUE (GeV ^{-1/2}) | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-------------|------|---------|
|------------------------------|-------------|------|---------|

-0.065 ± 0.004 OUR ESTIMATE

| | | | | |
|----------------|----------|----|------|--------------------------------------|
| -0.063 ± 0.005 | ARNDT | 96 | IPWA | $\gamma N \rightarrow \pi N$ |
| -0.069 ± 0.018 | CRAWFORD | 83 | IPWA | $\gamma N \rightarrow \pi N$ |
| -0.063 ± 0.008 | AWAJI | 81 | DPWA | $\gamma N \rightarrow \pi N$ |
| -0.069 ± 0.004 | ARAI | 80 | DPWA | $\gamma N \rightarrow \pi N$ (fit 1) |

| | | | | |
|--|--------------|----|------|--------------------------------------|
| -0.066 ± 0.004 | ARAI | 80 | DPWA | $\gamma N \rightarrow \pi N$ (fit 2) |
| -0.079 ± 0.009 | BRATASHEV... | 80 | DPWA | $\gamma N \rightarrow \pi N$ |
| -0.068 ± 0.015 | CRAWFORD | 80 | DPWA | $\gamma N \rightarrow \pi N$ |
| -0.0584 ± 0.0148 | ISHII | 80 | DPWA | Compton scattering |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| -0.085 ± 0.003 | LI | 93 | IPWA | $\gamma N \rightarrow \pi N$ |
| -0.129 | 10 WADA | 84 | DPWA | Compton scattering |
| -0.075 ± 0.015 | BARBOUR | 78 | DPWA | $\gamma N \rightarrow \pi N$ |
| -0.125 | 11 NOELLE | 78 | | $\gamma N \rightarrow \pi N$ |
| -0.076 | BERENDS | 77 | IPWA | $\gamma N \rightarrow \pi N$ |
| -0.087 ± 0.006 | FELLER | 76 | DPWA | $\gamma N \rightarrow \pi N$ |

$N(1440) \rightarrow n\gamma$, helicity-1/2 amplitude $A_{1/2}$

| VALUE (GeV $^{-1/2}$) | DOCUMENT ID | TECN | COMMENT |
|--|-------------|------|---|
| +0.040±0.010 OUR ESTIMATE | | | |
| 0.045±0.015 | ARNDT | 96 | IPWA $\gamma N \rightarrow \pi N$ |
| 0.037±0.010 | AWAJI | 81 | DPWA $\gamma N \rightarrow \pi N$ |
| 0.030±0.003 | FUJII | 81 | DPWA $\gamma N \rightarrow \pi N$ |
| 0.023±0.009 | ARAI | 80 | DPWA $\gamma N \rightarrow \pi N$ (fit 1) |
| 0.019±0.012 | ARAI | 80 | DPWA $\gamma N \rightarrow \pi N$ (fit 2) |
| 0.056±0.015 | CRAWFORD | 80 | DPWA $\gamma N \rightarrow \pi N$ |
| -0.029±0.035 | TAKEDA | 80 | DPWA $\gamma N \rightarrow \pi N$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.085±0.006 | LI | 93 | IPWA $\gamma N \rightarrow \pi N$ |
| +0.059±0.016 | BARBOUR | 78 | DPWA $\gamma N \rightarrow \pi N$ |
| 0.062 | 11 NOELLE | 78 | $\gamma N \rightarrow \pi N$ |

$N(1440)$ FOOTNOTES

¹ BAKER 79 finds a coupling of the $N(1440)$ to the $N\eta$ channel near (but slightly below) threshold.

² LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.

³ From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.

⁴ ARNDT 95 also finds a second-sheet pole with real part = 1383 MeV, -2×imaginary part = 210 MeV, and residue with modulus 92 MeV and phase = -54°.

⁵ See HOEHLER 93 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.

⁶ ARNDT 91 (Soln SM90) also finds a second-sheet pole with real part = 1413 MeV, -2 × imaginary part = 256 MeV, and residue = (78-153*i*) MeV.

⁷ LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.

⁸ An alternative which cannot be distinguished from this is to have a P_{13} resonance with $M = 1530$ MeV, $\Gamma = 79$ MeV, and elasticity = +0.271.

⁹ LONGACRE 77 considers this coupling to be well determined.

¹⁰ WADA 84 is inconsistent with other analyses; see the Note on N and Δ Resonances.

¹¹ Converted to our conventions using $M = 1486$ MeV, $\Gamma = 613$ MeV from NOELLE 78.

N(1440) REFERENCES

For early references, see Physics Letters **111B** 70 (1982).

| | | | | |
|--------------|----|------------------------|---|-------------------|
| VRANA | 00 | PRPL 328 181 | T.P. Vrana, S.A. Dytman,, T.-S.H. Lee | (PITT+) |
| ARNDT | 96 | PR C53 430 | R.A. Arndt, I.I. Strakovsky, R.L. Workman | (VPI) |
| ARNDT | 95 | PR C52 2120 | R.A. Arndt <i>et al.</i> | (VPI, BRCO) |
| BATINIC | 95 | PR C51 2310 | M. Batinic <i>et al.</i> | (BOSK, UCLA) |
| Also | 98 | PR C57 1004 (erratum) | M. Batinic <i>et al.</i> | |
| HOEHLER | 93 | πN Newsletter 9 1 | G. Hohler | (KARL) |
| LI | 93 | PR C47 2759 | Z.J. Li <i>et al.</i> | (VPI) |
| MANLEY | 92 | PR D45 4002 | D.M. Manley, E.M. Saleski | (KENT) IJP |
| Also | 84 | PR D30 904 | D.M. Manley <i>et al.</i> | (VPI) |
| ARNDT | 91 | PR D43 2131 | R.A. Arndt <i>et al.</i> | (VPI, TELE) IJP |
| CUTKOSKY | 90 | PR D42 235 | R.E. Cutkosky, S. Wang | (CMU) |
| WADA | 84 | NP B247 313 | Y. Wada <i>et al.</i> | (INUS) |
| CRAWFORD | 83 | NP B211 1 | R.L. Crawford, W.T. Morton | (GLAS) |
| PDG | 82 | PL 111B | M. Roos <i>et al.</i> | (HELS, CIT, CERN) |
| AWAJI | 81 | Bonn Conf. 352 | N. Awaji, R. Kajikawa | (NAGO) |
| Also | 82 | NP B197 365 | K. Fujii <i>et al.</i> | (NAGO) |
| FUJII | 81 | NP B187 53 | K. Fujii <i>et al.</i> | (NAGO, OSAK) |
| ARAI | 80 | Toronto Conf. 93 | I. Arai | (INUS) |
| Also | 82 | NP B194 251 | I. Arai, H. Fujii | (INUS) |
| BRATASHEV... | 80 | NP B166 525 | A.S. Bratashevsky <i>et al.</i> | (KFTI) |
| CRAWFORD | 80 | Toronto Conf. 107 | R.L. Crawford | (GLAS) |
| CUTKOSKY | 80 | Toronto Conf. 19 | R.E. Cutkosky <i>et al.</i> | (CMU, LBL) IJP |
| Also | 79 | PR D20 2839 | R.E. Cutkosky <i>et al.</i> | (CMU, LBL) IJP |
| ISHII | 80 | NP B165 189 | T. Ishii <i>et al.</i> | (KYOT, INUS) |
| TAKEDA | 80 | NP B168 17 | H. Takeda <i>et al.</i> | (TOKY, INUS) |
| BAKER | 79 | NP B156 93 | R.D. Baker <i>et al.</i> | (RHEL) IJP |
| HOEHLER | 79 | PDAT 12-1 | G. Hohler <i>et al.</i> | (KARLT) IJP |
| Also | 80 | Toronto Conf. 3 | R. Koch | (KARLT) IJP |
| BARBOUR | 78 | NP B141 253 | I.M. Barbour, R.L. Crawford, N.H. Parsons | (GLAS) |
| LONGACRE | 78 | PR D17 1795 | R.S. Longacre <i>et al.</i> | (LBL, SLAC) |
| NOELLE | 78 | PTP 60 778 | P. Noelle | (NAGO) |
| BERENDS | 77 | NP B136 317 | F.A. Berends, A. Donnachie | (LEID, MCHS) IJP |
| LONGACRE | 77 | NP B122 493 | R.S. Longacre, J. Dolbeau | (SACL) IJP |
| Also | 76 | NP B108 365 | J. Dolbeau <i>et al.</i> | (SACL) IJP |
| FELLER | 76 | NP B104 219 | P. Feller <i>et al.</i> | (NAGO, OSAK) IJP |
| FELTESSE | 75 | NP B93 242 | J. Feltesse <i>et al.</i> | (SACL) IJP |
| LONGACRE | 75 | PL 55B 415 | R.S. Longacre <i>et al.</i> | (LBL, SLAC) IJP |