38. CROSS-SECTION FORMULAE FOR SPECIFIC PROCESSES

Revised October 2001 by R.N. Cahn (LBNL).

38.1. Leptoproduction

See section on Structure Functions (Sec. 14 of this *Review*).

38.2. e^+e^- annihilation

For pointlike, spin-1/2 fermions, the differential cross section in the c.m. for $e^+e^- \to f\overline{f}$ via single photon annihilation is (θ is the angle between the incident electron and the produced fermion; $N_c = 1$ if f is a lepton and $N_c = 3$ if f is a quark).

$$\frac{d\sigma}{d\Omega} = N_c \frac{\alpha^2}{4s} \beta\big[1 + \cos^2\theta + (1-\beta^2)\sin^2\theta\big] Q_f^2 \ , \tag{38.1}$$

where β is the velocity of the final state fermion in the c.m. and Q_f is the charge of the fermion in units of the proton charge. For $\beta \to 1$,

$$\sigma = N_c \frac{4\pi\alpha^2}{3s} Q_f^2 = N_c \frac{86.8 Q_f^2 \ nb}{s} \ . \tag{38.2}$$

where s is in GeV^2 units.

At higher energies, the Z^0 (mass M_Z and width Γ_Z) must be included. If the mass of a fermion f is much less than the mass of the Z^0, then the differential cross section for $e^+e^- \to f\overline{f}$ is

$$\begin{aligned} \frac{d\sigma}{d\Omega} = \ & N_c \frac{\alpha^2}{4s} \Big\{ (1+\cos^2\theta) \Big[Q_f^2 - 2\chi_1 v_e v_f Q_f + \chi_2 (a_e^2 + v_e^2)(a_f^2 + v_f^2) \Big] \\ & + 2\cos\theta \left[-2\chi_1 a_e a_f Q_f + 4\chi_2 a_e a_f v_e v_f \right] \Big\} \end{aligned} \tag{38.3}$$

where

$$\begin{aligned} \chi_1 = \ & \frac{1}{16\sin^2\theta_W \cos^2\theta_W} \frac{s(s-M_Z^2)}{(s-M_Z^2)^2 + M_Z^2\Gamma_Z^2} \ , \\ \chi_2 = \ & \frac{1}{256\sin^4\theta_W \cos^4\theta_W} \frac{s^2}{(s-M_Z^2)^2 + M_Z^2\Gamma_Z^2} \ , \\ a_e = \ & -1 \ , \\ v_e = \ & -1 + 4\sin^2\theta_W \ , \\ a_f = \ & 2T_{3f} \ , \\ v_f = \ & 2T_{3f} - 4Q_f \sin^2\theta_W \ , \end{aligned} \tag{38.4}$$

where $T_{3f} = 1/2$ for u, c and neutrinos, while $T_{3f} = -1/2$ for d, s, b, and negatively charged leptons.

At LEP II it may be possible to produce the orthodox Higgs boson, H, (see the mini-review on Higgs bosons) in the reaction $e^+e^- \to HZ^0$, which proceeds dominantly through a virtual Z^0. The Standard Model prediction for the cross section [3] is

$$\sigma(e^+e^- \to HZ^0) = \frac{\pi\alpha^2}{24} \cdot \frac{2K}{\sqrt{s}} \cdot \frac{K^2 + 3M_Z^2}{(s - M_Z^2)^2} \cdot \frac{1 - 4\sin^2\theta_W + 8\sin^4\theta_W}{\sin^4\theta_W \cos^4\theta_W} \ . \tag{38.5}$$

where K is the c.m. momentum of the produced H or Z^0. Near the production threshold, this formula needs to be corrected for the finite width of the Z^0.

38.3. Two-photon process at e^+e^- colliders

When an e^+ and an e^- collide with energies E_1 and E_2, they emit dn_1 and dn_2 virtual photons with energies ω_1 and ω_2 and 4-momenta q_1 and q_2. In the equivalent photon approximation, the cross section for $e^+e^- \to e^+e^-X$ is related to the cross section for $\gamma\gamma \to X$ by (Ref. 1)

$$d\sigma_{e^+e^-\to e^+e^-X}(s) = dn_1\, dn_2\, d\sigma_{\gamma\gamma\to X}(W^2) \tag{38.6}$$

where $s = 4E_1E_2$, $W^2 = 4\omega_1\omega_2$ and

$$dn_i = \frac{\alpha}{\pi}\left[1 - \frac{\omega_i}{E_i} + \frac{\omega_i^2}{2E_i^2} - \frac{m_e^2\omega_i^2}{(-q_i^2)E_i^2}\right] \frac{d\omega_i}{\omega_i} \frac{d(-q_i^2)}{(-q_i^2)} \ . \tag{38.7}$$

After integration (including that over q_i^2 in the region $m_e^2\omega_i^2/E_i(E_i - \omega_i) \le -q_i^2 \le (-q^2)_{\max}$), the cross section is

$$\begin{aligned} \sigma_{e^+e^-\to e^+e^-X}(s) =& \frac{\alpha^2}{\pi^2} \int_{z_{th}}^1 \frac{dz}{z} \left[f(z) \left(\ln \frac{(-q^2)_{\max}}{m_e^2 z} - 1 \right)^2 \right. \\ & \left. - \tfrac{1}{3} \left(\ln \frac{1}{z} \right)^3 \right] \sigma_{\gamma\gamma\to X}(zs) \ ; \\ f(z) =& \left(1 + \tfrac{1}{2}z\right)^2 \ln \frac{1}{z} - \tfrac{1}{2}(1 - z)(3 + z) \ ; \\ z =& \frac{W^2}{s} \ . \end{aligned} \tag{38.8}$$

The quantity $(-q^2)_{\max}$ depends on properties of the produced system X, in particular, $(-q^2)_{\max} \sim m_\rho^2$ for hadron production ($X = h$) and $(-q^2)_{\max} \sim W^2$ for lepton pair production ($X = \ell^+\ell^-$, $\ell = e, \mu, \tau$).

For production of a resonance of mass m_R and spin $J \ne 1$

$$\begin{aligned} \sigma_{e^+e^-\to e^+e^-R}(s) =& (2J + 1) \frac{8\alpha^2\Gamma_{R\to\gamma\gamma}}{m_R^3} \\ & \times \left[f(m_R^2/s) \left(\ln \frac{s m_V^2}{m_e^2 m_R^2} - 1 \right)^2 - \frac{1}{3} \left(\ln \frac{s}{m_R^2} \right)^3 \right] \end{aligned} \tag{38.9}$$

where m_V is the mass that enters into the form factor of the $\gamma\gamma \to R$ transition: $m_V \sim m_\rho$ for $R = \pi^0, \eta, f_2(1270), \ldots$, $m_V \sim m_R$ for $R = c\bar{c}$ or $b\bar{b}$ resonances.

38.4. Inclusive hadronic reactions

One-particle inclusive cross sections $Ed^3\sigma/d^3p$ for the production of a particle of momentum p are conveniently expressed in terms of rapidity (see above) and the momentum p_T transverse to the beam direction (defined in the center-of-mass frame)

$$E\frac{d^3\sigma}{d^3p} = \frac{d^3\sigma}{d\phi\, dy\, p_T\, dp_T} \ . \tag{38.10}$$

In the case of processes where p_T is large or the mass of the produced particle is large (here large means greater than 10 GeV), the parton model can be used to calculate the rate. Symbolically

$$\sigma_{\text{hadronic}} = \sum_{ij} \int f_i(x_1,\, Q^2)\; f_j(x_2,\, Q^2)\; dx_1\; dx_2\; \widehat{\sigma}_{\text{partonic}} \ , \tag{38.11}$$

where $f_i(x,\, Q^2)$ is the parton distribution introduced above and Q is a typical momentum transfer in the partonic process and $\widehat{\sigma}$ is the partonic cross section. Some examples will help to clarify. The production of a W^+ in pp reactions at rapidity y in the center-of-mass frame is given by

$$\begin{aligned}\frac{d\sigma}{dy} = \frac{G_F\, \pi\sqrt{2}}{3} & \\ \times\, \tau \Big[\cos^2\theta_c & \Big(u(x_1\ ,\ M_W^2)\, \overline{d}\,(x_2, M_W^2) \\ & \quad + \ u(x_2\ ,\ M_W^2)\, \overline{d}\,(x_1, M_W^2) \Big) \\ + \ \sin^2\theta_c & \Big(u(x_1\ ,\ M_W^2)\, \overline{s}\,(x_2\ ,\ M_W^2) \\ & \quad + \ s(x_2, M_W^2)\, \overline{u}\,(x_1, M_W^2) \Big) \Big] \ ,\end{aligned} \tag{38.12}$$

where $x_1 = \sqrt{\tau}\, e^y$, $x_2 = \sqrt{\tau}\, e^{-y}$, and $\tau = M_W^2/s$. Similarly the production of a jet in pp (or $p\overline{p}$) collisions is given by

$$\begin{aligned}\frac{d^3\sigma}{d^2p_T\; dy} = \sum_{ij} \int & f_i(x_1\ ,\ p_T^2)\; f_j(x_2, p_T^2) \\ \times & \left[\widehat{s}\; \frac{d\widehat{\sigma}}{d\,\widehat{t}} \right]_{ij} dx_1\; dx_2\; \delta(\,\widehat{s} + \,\widehat{t}\, + \widehat{u}) \ ,\end{aligned} \tag{38.13}$$

where the summation is over quarks, gluons, and antiquarks. Here

$$s = (p_1 + p_2)^2 \ , \tag{38.14}$$

$$t = (p_1 - p_{\text{jet}})^2 \ , \tag{38.15}$$

$$u = (p_2 - p_{\text{jet}})^2 \ , \tag{38.16}$$

p_1 and p_2 are the momenta of the incoming p and p (or $\overline{p}$) and $\widehat{s}$, $\widehat{t}$, and $\widehat{u}$ are s, t, and u with $p_1 \to x_1 p_1$ and $p_2 \to x_2 p_2$. The partonic cross section $\widehat{s}[(d\widehat{\sigma})/(d\widehat{t})]$ can be found in Ref. 2. Example: for the process $gg \to q\overline{q}$,

$$\widehat{s}\,\frac{d\sigma}{dt} = 3\alpha_s^2\,\frac{(\widehat{t}^{\,2}+\widehat{u}^2)}{8\widehat{s}}\,\left[\frac{4}{9\,\widehat{t}\,\widehat{u}} - \frac{1}{\widehat{s}^{\,2}}\right] . \tag{38.17}$$

The prediction of Eq. (38.13) is compared to data from the UA1 and UA2 collaborations in Fig. 39.1 in the Plots of Cross Sections and Related Quantities section of this *Review.*

The associated production of a Higgs boson and a gauge boson is analogous to the process $e^+e^- \to HZ^0$ in Sec. 38.2. The required parton-level cross sections [4], averaged over initial quark colors, are

$$\begin{aligned}
\sigma(q_i\overline{q}_j \to W^{\pm}H) &= \frac{\pi\alpha^2|V_{ij}|^2}{36\sin^4\theta_W}\cdot\frac{2K}{\sqrt{s}}\cdot\frac{K^2+3M_W^2}{(s-M_W^2)^2} \\
\sigma(q\overline{q} \to Z^0H) &= \frac{\pi\alpha^2(a_q^2+v_q^2)}{144\sin^4\theta_W\cos^4\theta_W}\cdot\frac{2K}{\sqrt{s}}\cdot\frac{K^2+3M_Z^2}{(s-M_Z^2)^2} .
\end{aligned}$$

Here V_{ij} is the appropriate element of the Kobayashi-Maskawa matrix and K is the c.m. momentum of the produced H. The axial and vector couplings are defined as in Sec. 38.2.

38.5. One-particle inclusive distributions

In order to describe one-particle inclusive production in e^+e^- annihilation or deep inelastic scattering, it is convenient to introduce a fragmentation function $D_i^h\,(z,Q^2)$ where $D_i^h\,(z,Q^2)$ is the number of hadrons of type h and momentum between zp and $(z+dz)p$ produced in the fragmentation of a parton of type i. The Q^2 evolution is predicted by QCD and is similar to that of the parton distribution functions [see section on Quantum Chromodynamics (Sec. 9 of this *Review*)]. The $D_i^h(z,Q^2)$ are normalized so that

$$\sum_h \int zD_i^h\,(z,Q^2)dz = 1 . \tag{38.18}$$

If the contributions of the Z boson and three-jet events are neglected, the cross section for producing a hadron h in e^+e^- annihilation is given by

$$\frac{1}{\sigma_{\text{had}}}\frac{d\sigma}{dz} = \frac{\sum_i e_i^2\,D_i^h\,(z,Q^2)}{\sum_i e_i^2} , \tag{38.19}$$

where e_i is the charge of quark-type i, σ_{had} is the total hadronic cross section, and the momentum of the hadron is $zE_{\text{cm}}/2$.

In the case of deep inelastic muon scattering, the cross section for producing a hadron of energy E_h is given by

$$\frac{1}{\sigma_{\rm tot}}\,\frac{d\sigma}{dz} = \frac{\sum_i e_i^2\; q_i(x,Q^2)\; D_i^h(z,Q^2)}{\sum_i e_i^2\; q_i(x,Q^2)}\ , \tag{38.20}$$

where $E_h = \nu z$. (For the kinematics of deep inelastic scattering, see Sec. 37.4.2 of the Kinematics section of this *Review.*) The fragmentation functions for light and heavy quarks have a different z dependence; the former peak near $z = 0$. They are illustrated in Figs. 15.5a and 15.5b in the section on "Fragmentation Functions in e^+e^- Annihilation" (Sec. 15 of this *Review*).

References:

1. V.M. Budnev, I.F. Ginzburg, G.V. Meledin, and V.G. Serbo,Phys. Reports **15C**, 181 (1975);

   See also S. Brodsky, T. Kinoshita, and H. Terazawa, Phys. Rev. **D4**, 1532 (1971).
2. G.F. Owens, F. Reya, and M. Glück, Phys. Rev. **D18**, 1501 (1978).
3. B.W. Lee, C. Quigg, and B. Thacker, Phys. Rev. **D16**, 1519 (1977).
4. E. Eichten, I. Hinchliffe, K. Lane, and C. Quigg, Rev. Mod. Phys. **56**, 579 (1984).