

Average Hadron Multiplicities in Hadronic  $e^+e^-$  Annihilation Events

**Table 39.1:** Average hadron multiplicities per hadronic  $e^+e^-$  annihilation event at  $\sqrt{s} = 10$ , 29–35, 91, and 130–200 GeV. The rates given include decay products from resonances with  $c\tau < 10$  cm, and include the corresponding anti-particle state. Correlations of the systematic uncertainties were considered for the calculation of the averages. (Updated July 2001 by O. Biebel.)

Particle	$\sqrt{s} = 10$ GeV	$\sqrt{s} = 29\text{--}35$ GeV	$\sqrt{s} = 91$ GeV	$\sqrt{s} = 130\text{--}200$ GeV
<b>Pseudoscalar mesons:</b>				
$\pi^+$	$6.6 \pm 0.2$	$10.3 \pm 0.4$	$16.99 \pm 0.27$	$21.24 \pm 0.39$
$\pi^0$	$3.2 \pm 0.3$	$5.83 \pm 0.28$	$9.42 \pm 0.32$	
$K^+$	$0.90 \pm 0.04$	$1.48 \pm 0.09$	$2.242 \pm 0.063$	$2.81 \pm 0.19$
$K^0$	$0.91 \pm 0.05$	$1.48 \pm 0.07$	$2.049 \pm 0.026$	$2.10 \pm 0.12$
$\eta$	$0.20 \pm 0.04$	$0.61 \pm 0.07$	$0.946 \pm 0.075$	
$\eta'(958)$	$0.03 \pm 0.01$	$0.26 \pm 0.10$	$0.152 \pm 0.020$	
$D^+$	$0.16 \pm 0.03$	$0.17 \pm 0.03$	$0.175 \pm 0.016$	
$D^0$	$0.37 \pm 0.06$	$0.45 \pm 0.07$	$0.454 \pm 0.030$	
$D_s^+$	$0.13 \pm 0.02$	$0.45 \pm 0.20^{(a)}$	$0.131 \pm 0.021$	
$B^+, B_d^0$	—	—	$0.165 \pm 0.026^{(b)}$	
$B_s^0$	—	—	$0.057 \pm 0.013^{(b)}$	
<b>Scalar mesons:</b>				
$f_0(980)$	$0.024 \pm 0.006$	$0.05 \pm 0.02^{(c)}$	$0.146 \pm 0.012$	
$a_0(980)^\pm$	—	—	$0.27 \pm 0.11^{(d)}$	
<b>Vector mesons:</b>				
$\rho(770)^0$	$0.35 \pm 0.04$	$0.81 \pm 0.08$	$1.231 \pm 0.098$	
$\rho(770)^\pm$	—	—	$2.40 \pm 0.43^{(d)}$	
$\omega(782)$	$0.30 \pm 0.08$	—	$1.08 \pm 0.12$	
$K^*(892)^+$	$0.27 \pm 0.03$	$0.64 \pm 0.05$	$0.715 \pm 0.059$	
$K^*(892)^0$	$0.29 \pm 0.03$	$0.56 \pm 0.06$	$0.738 \pm 0.024$	
$\phi(1020)$	$0.044 \pm 0.003$	$0.085 \pm 0.011$	$0.0963 \pm 0.0032$	
$D^*(2010)^+$	$0.22 \pm 0.04$	$0.43 \pm 0.07$	$0.1973 \pm 0.0057^{(e)}$	
$D^*(2007)^0$	$0.23 \pm 0.06$	$0.27 \pm 0.11$	—	
$D_s^*(2112)^+$	—	—	$0.101 \pm 0.048^{(f)}$	
$B^*(g)$	—	—	$0.288 \pm 0.026$	
$J/\psi(1S)$	—	—	$0.0052 \pm 0.0004^{(h)}$	
$\psi(2S)$	—	—	$0.0023 \pm 0.0004^{(h)}$	
$\Upsilon(1S)$	—	—	$0.00014 \pm 0.00007^{(h)}$	
<b>Pseudovector mesons:</b>				
$\chi_{c1}(3510)$	—	—	$0.0041 \pm 0.0011^{(h)}$	
<b>Tensor mesons:</b>				
$f_2(1270)$	$0.09 \pm 0.02$	$0.14 \pm 0.04$	$0.166 \pm 0.020$	
$f_2'(1525)$	—	—	$0.012 \pm 0.006$	
$K_2^*(1430)^+$	—	$0.09 \pm 0.03$	—	
$K_2^*(1430)^0$	—	$0.12 \pm 0.06$	$0.084 \pm 0.022^{(h)}$	
$B^{**}(i)$	—	—	$0.118 \pm 0.024$	
<b>Baryons:</b>				
$p$	$0.253 \pm 0.016$	$0.640 \pm 0.050$	$1.048 \pm 0.045$	$1.41 \pm 0.18$
$\Lambda$	$0.080 \pm 0.007$	$0.205 \pm 0.010$	$0.3915 \pm 0.0065$	$0.39 \pm 0.03$
$\Sigma^0$	$0.023 \pm 0.008$	—	$0.076 \pm 0.011$	
$\Sigma^-$	—	—	$0.081 \pm 0.010$	
$\Sigma^+$	—	—	$0.107 \pm 0.011$	
$\Sigma^\pm$	—	—	$0.174 \pm 0.009$	
$\Xi^-$	$0.0059 \pm 0.0007$	$0.0176 \pm 0.0027$	$0.0258 \pm 0.0010$	
$\Delta(1232)^{++}$	$0.040 \pm 0.010$	—	$0.085 \pm 0.014$	
$\Sigma(1385)^-$	$0.006 \pm 0.002$	$0.017 \pm 0.004$	$0.0240 \pm 0.0017$	
$\Sigma(1385)^+$	$0.005 \pm 0.001$	$0.017 \pm 0.004$	$0.0239 \pm 0.0015$	
$\Sigma(1385)^\pm$	$0.0106 \pm 0.0020$	$0.033 \pm 0.008$	$0.0462 \pm 0.0028$	
$\Xi(1530)^0$	$0.0015 \pm 0.0006$	—	$0.0055 \pm 0.0005$	
$\Omega^-$	$0.0007 \pm 0.0004$	$0.014 \pm 0.007$	$0.0016 \pm 0.0003$	
$\Lambda_c^+$	$0.100 \pm 0.030^{(j)}$	$0.110 \pm 0.050$	$0.078 \pm 0.017$	
$\Lambda_b^0$	—	—	$0.031 \pm 0.016$	
$\Sigma_c^{++}, \Sigma_c^0$	$0.014 \pm 0.007$	—	—	
$\Lambda(1520)$	$0.008 \pm 0.002$	—	$0.0222 \pm 0.0027$	

## Notes for Table 39.1:

- (a)  $B(D_s \rightarrow \eta\pi, \eta'\pi)$  was used (RPP1994).
- (b) The Standard Model  $B(Z \rightarrow b\bar{b}) = 0.217$  was used.
- (c)  $x_p = p/p_{\text{beam}} > 0.1$  only.
- (d) Both charge states.
- (e)  $B(D_s^*(2010)^+ \rightarrow D^0\pi^+) \times B(D^0 \rightarrow K^-\pi^+)$  has been used (RPP2000).
- (f)  $B(D_s^* \rightarrow D_s^+\gamma)$ ,  $B(D_s^+ \rightarrow \phi\pi^+)$ ,  $B(\phi \rightarrow K^+K^-)$  have been used (RPP1998).
- (g) Any charge state (*i.e.*,  $B_d^*$ ,  $B_u^*$ , or  $B_s^*$ ).
- (h)  $B(Z \rightarrow \text{hadrons}) = 0.699$  was used (RPP1994).
- (i) Any charge state (*i.e.*,  $B_d^{**}$ ,  $B_u^{**}$ , or  $B_s^{**}$ ).
- (j) The value was derived from the cross section of  $A_c^+ \rightarrow p\pi K$ , assuming the branching fraction to be  $(3.2 \pm 0.7)\%$  (RPP1992).

## References for Table 39.1:

- RPP1992**: Phys. Rev. **D45** (1992) and references therein.
- RPP1994**: Phys. Rev. **D50**, 1173 (1994) and references therein.
- RPP1996**: Phys. Rev. **D54**, 1 (1996) and references therein.
- RPP1998**: Eur. Phys. J. **C3**, 1 (1998) and references therein.
- RPP2000**: Eur. Phys. J. **C15**, 1 (2000) and references therein.
- R. Marshall, Rep. Prog. Phys. **52**, 1329 (1989)
- A. De Angelis, J. Phys. **G19**, 1233 (1993) and references therein.
- ALEPH**: D. Buskulic *et al.*: Phys. Lett. **B295**, 396 (1992); Z. Phys. **C64**, 361 (1994); **C69**, 15 (1996); **C69**, 379 (1996); **C73**, 409 (1997); and  
R. Barate *et al.*: Z. Phys. **C74**, 451 (1997); Phys. Reports **294**, 1 (1998); Eur. Phys. J. **C5**, 205 (1998); **C16**, 597 (2000); **C16**, 613 (2000).
- ARGUS**: H. Albrecht *et al.*: Phys. Lett. **230B**, 169 (1989); Z. Phys. **C44**, 547 (1989); **C46**, 15 (1990); **C54**, 1 (1992); **C58**, 199 (1993); **C61**, 1 (1994); Phys. Rep. **276**, 223 (1996).
- CELLO**: H.J. Behrend *et al.*: Z. Phys. **C46**, 397 (1990); **C47**, 1 (1990).
- CLEO**: D. Bortoletto *et al.*, Phys. Rev. **D37**, 1719 (1988).
- Crystal Ball**: Ch. Bieler *et al.*, Z. Phys. **C49**, 225 (1991).
- DELPHI**: P. Abreu *et al.*: Z. Phys. **C57**, 181 (1993); **C59**, 533 (1993); **C61**, 40 (1994); **C65**, 587 (1995); **C67**, 543 (1995); **C68**, 353 (1995); **C73**, 61 (1996); Nucl. Phys. **B444**, 3 (1995); Phys. Lett. **B341**, 109 (1994); **B345**, 598 (1995); **B361**, 207 (1995); **B372**, 172 (1996); **B379**, 309 (1996); **B416**, 233 (1998); **B449**, 364 (1999); **B475**, 429 (2000); Eur. Phys. J. **C6**, 19 (1999); **C5**, 585 (1998); **C18**, 203 (2000).
- HRS**: S. Abachi *et al.*, Phys. Rev. Lett. **57**, 1990 (1986); and  
M. Derrick *et al.*, Phys. Rev. **D35**, 2639 (1987).
- L3**: M. Acciarri *et al.*: Phys. Lett. **B328**, 223 (1994); **B345**, 589 (1995); **B371**, 126 (1996); **B371**, 137 (1996); **B393**, 465 (1997); **B404**, 390 (1997); **B407**, 351 (1997); **B407**, 389 (1997), erratum *ibid.* **B427**, 409 (1998); **B453**, 94 (1999); **B479**, 79 (2000).
- MARK II**: H. Schellman *et al.*, Phys. Rev. **D31**, 3013 (1985); and  
G. Wormser *et al.*, Phys. Rev. Lett. **61**, 1057 (1988).
- JADE**: W. Bartel *et al.*, Z. Phys. **C20**, 187 (1983); and  
D.D. Pietzl *et al.*, Z. Phys. **C46**, 1 (1990).
- OPAL**: R. Akers *et al.*: Z. Phys. **C63**, 181 (1994); **C66**, 555 (1995); **C67**, 389 (1995); **C68**, 1 (1995); and  
G. Alexander *et al.*: Phys. Lett. **B358**, 162 (1995); Z. Phys. **C70**, 197 (1996); **C72**, 1 (1996); **C72**, 191 (1996); **C73**, 569 (1997); **C73**, 587 (1997); Phys. Lett. **B370**, 185 (1996); and  
K. Ackerstaff *et al.*: Z. Phys. **C75**, 192 (1997); Phys. Lett. **B412**, 210 (1997); Eur. Phys. J. **C1**, 439 (1998); **C4**, 19 (1998); **C5**, 1 (1998); **C5**, 411 (1998); and  
G. Abbiendi *et al.*: Eur. Phys. J. **C16**, 185 (2000); **C16**, 185 (2000).
- PLUTO**: Ch. Berger *et al.*, Phys. Lett. **104B**, 79 (1981).
- SLD**: K. Abe, Phys. Rev. **D59**, 052001 (1999).
- TASSO**: H. Aihara *et al.*, Z. Phys. **C27**, 27 (1985).
- TPC**: H. Aihara *et al.*, Phys. Rev. Lett. **53**, 2378 (1984).