

p $I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$ Status: *****p MASS (atomic mass units u)**

The mass is known much more precisely in u (atomic mass units) than in MeV. The conversion from u to MeV, $1\text{ u} = 931.494013 \pm 0.000037\text{ MeV}/c^2$ (MOHR 99, the 1998 CODATA value), involves the relatively poorly known electronic charge.

VALUE (u)	DOCUMENT ID	TECN	COMMENT
1.00727646688±0.0000000013	MOHR 99	RVUE	1998 CODATA value
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.007276470 ± 0.000000012	COHEN 87	RVUE	1986 CODATA value

p MASS (MeV)

The mass is known much more precisely in u (atomic mass units) than in MeV; see the previous data block. The conversion from u to MeV, $1\text{ u} = 931.494013 \pm 0.000037\text{ MeV}/c^2$ (MOHR 99, the 1998 CODATA value), involves the relatively poorly known electronic charge.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
938.271998±0.000038	MOHR 99	RVUE	1998 CODATA value
• • • We do not use the following data for averages, fits, limits, etc. • • •			
938.27231 ± 0.00028	COHEN 87	RVUE	1986 CODATA value
938.2796 ± 0.0027	COHEN 73	RVUE	1973 CODATA value

 $|m_p - m_{\bar{p}}|/m_p$

A test of *CPT* invariance. Note that the \bar{p}/p charge-to-mass ratio, given below, is much better determined.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6 \times 10^{-8}$	90	¹ HORI 01	SPEC	$\bar{p}e^-$ He atom
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<5 \times 10^{-7}$		² TORII 99	SPEC	$\bar{p}e^-$ He atom
¹ HORI 01 uses the more-precisely-known constraint on the \bar{p} charge-to-mass ratio of GABRIELSE 99 (see below) to get this result. This is not independent of the HORI 01 value for $ q_p + q_{\bar{p}} /e$, below.				
² TORII 99 uses the more-precisely-known constraint on the \bar{p} charge-to-mass ratio of GABRIELSE 95 (see below) to get this result. This is not independent of the TORII 99 value for $ q_p + q_{\bar{p}} /e$, below.				

\bar{p}/p CHARGE-TO-MASS RATIO, $| \frac{q_{\bar{p}}}{m_{\bar{p}}} | / (\frac{q_p}{m_p})$

A test of *CPT* invariance. Listed here are measurements involving the *inertial* masses. For a discussion of what may be inferred about the ratio of \bar{p} and p *gravitational* masses, see ERICSON 90; they obtain an upper bound of 10^{-6} – 10^{-7} for violation of the equivalence principle for \bar{p} 's.

VALUE	DOCUMENT ID	TECN	COMMENT
$0.99999999991 \pm 0.0000000009$	GABRIELSE 99	TRAP	Penning trap
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.0000000015 ± 0.0000000011	³ GABRIELSE 95	TRAP	Penning trap
1.000000023 ± 0.000000042	⁴ GABRIELSE 90	TRAP	Penning trap
³ Equation (2) of GABRIELSE 95 should read $M(\bar{p})/M(p) = 0.999\ 999\ 9985$ (11) (G. Gabrielse, private communication).			
⁴ GABRIELSE 90 also measures $m_{\bar{p}}/m_{e^-} = 1836.152660 \pm 0.000083$ and $m_p/m_{e^-} = 1836.152680 \pm 0.000088$. Both are completely consistent with the 1986 CODATA (COHEN 87) value for m_p/m_{e^-} of 1836.152701 ± 0.000037 .			

$$(| \frac{q_{\bar{p}}}{m_{\bar{p}}} | - \frac{q_p}{m_p}) / \frac{q_p}{m_p}$$

A test of *CPT* invariance. Taken from the \bar{p}/p charge-to-mass ratio, above.

VALUE	DOCUMENT ID
$(-9 \pm 9) \times 10^{-11}$ OUR EVALUATION	

$$| q_p + q_{\bar{p}} | / e$$

A test of *CPT* invariance. Note that the \bar{p}/p charge-to-mass ratio, given above, is much better determined. See also a similar test involving the electron.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6 \times 10^{-8}$	90	⁵ HORI 01	SPEC	$\bar{p}e^-$ -He atom
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<5 \times 10^{-7}$		⁶ TORII 99	SPEC	$\bar{p}e^-$ -He atom
$<2 \times 10^{-5}$		⁷ HUGHES 92	RVUE	
⁵ HORI 01 uses the more-precisely-known constraint on the \bar{p} charge-to-mass ratio of GABRIELSE 99 (see above) to get this result. This is not independent of the HORI 01 value for $ m_p - m_{\bar{p}} / m_p$, above.				
⁶ TORII 99 uses the more-precisely-known constraint on the \bar{p} charge-to-mass ratio of GABRIELSE 95 (see above) to get this result. This is not independent of the TORII 99 value for $ m_p - m_{\bar{p}} / m_p$, above.				
⁷ HUGHES 92 uses recent measurements of Rydberg-energy and cyclotron-frequency ratios.				

$|q_p + q_e|/e$

See DYLLA 73 for a summary of experiments on the neutrality of matter.

See also “*n* CHARGE” in the neutron Listings.

VALUE	DOCUMENT ID	COMMENT	
$<1.0 \times 10^{-21}$	8 DYLLA	73	Neutrality of SF ₆
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$<3.2 \times 10^{-20}$	9 SENGUPTA	00	binary pulsar
$<0.8 \times 10^{-21}$	MARINELLI	84	Magnetic levitation
8 Assumes that $q_n = q_p + q_e$.			
9 SENGUPTA 00 uses the difference between the observed rate of rotational energy loss by the binary pulsar PSR B1913+16 and the rate predicted by general relativity to set this limit. See the paper for assumptions.			

p MAGNETIC MOMENT

See the “Note on Baryon Magnetic Moments” in the Λ Listings.

VALUE (μ_N)	DOCUMENT ID	TECN	COMMENT
2.792847337 ± 0.000000029	MOHR	99	RVUE 1998 CODATA value
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.792847386 ± 0.000000063	COHEN	87	RVUE 1986 CODATA value
2.7928456 ± 0.0000011	COHEN	73	RVUE 1973 CODATA value

\bar{p} MAGNETIC MOMENT

A few early results have been omitted.

VALUE (μ_N)	DOCUMENT ID	TECN	COMMENT
-2.800 ± 0.008 OUR AVERAGE			
-2.8005 ± 0.0090	KREISSL	88	CNTR \bar{p} ²⁰⁸ Pb 11 → 10 X-ray
-2.817 ± 0.048	ROBERTS	78	CNTR
-2.791 ± 0.021	HU	75	CNTR Exotic atoms

$(\mu_p + \mu_{\bar{p}}) / \mu_p$

A test of *CPT* invariance. Calculated from the p and \bar{p} magnetic moments, above.

VALUE	DOCUMENT ID
(-2.6 ± 2.9) × 10⁻³ OUR EVALUATION	

p ELECTRIC DIPOLE MOMENT

A nonzero value is forbidden by both T invariance and P invariance.

VALUE (10^{-23} ecm)	EVTS	DOCUMENT ID	TECN	COMMENT
$- 3.7 \pm 6.3$		CHO	89	NMR TI F molecules
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 400	DZUBA	85	THEO	Uses ^{129}Xe moment
130 \pm 200	¹⁰ WILKENING	84		
900 \pm 1400	¹¹ WILKENING	84		
700 \pm 900	1G	HARRISON	69	MBR Molecular beam

¹⁰ This WILKENING 84 value includes a finite-size effect and a magnetic effect.

¹¹ This WILKENING 84 value is more cautious than the other and excludes the finite-size effect, which relies on uncertain nuclear integrals.

p ELECTRIC POLARIZABILITY $\bar{\alpha}_p$

In the 2004 Review, we will probably switch to listing measurements of $\bar{\alpha}_p + \bar{\beta}_p$ and $\bar{\alpha}_p - \bar{\beta}_p$ instead of $\bar{\alpha}_p$ and $\bar{\beta}_p$ separately.

VALUE (10^{-4} fm 3)	DOCUMENT ID	TECN	COMMENT
12.0 \pm 0.7 OUR AVERAGE			
$11.82 \pm 0.98^{+0.52}_{-0.98}$	¹² BLANPIED	01 LEGS	$p(\vec{\gamma},\gamma)$, $p(\vec{\gamma},\pi^0)$, $p(\vec{\gamma},\pi^+)$
$11.9 \pm 0.5 \pm 1.3$	¹³ OL莫斯DEL...	01 CNTR	γp Compton scattering
$12.1 \pm 0.8 \pm 0.5$	¹⁴ MACGIBBON	95 RVUE	global average
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$11.7 \pm 0.8 \pm 0.7$	¹⁵ BARANOV	01 RVUE	Global average
$12.5 \pm 0.6 \pm 0.9$	MACGIBBON	95 CNTR	γp Compton scattering
$9.8 \pm 0.4 \pm 1.1$	HALLIN	93 CNTR	γp Compton scattering
$10.62^{+1.25+1.07}_{-1.19-1.03}$	ZIEGER	92 CNTR	γp Compton scattering
$10.9 \pm 2.2 \pm 1.3$	¹⁶ FEDERSPIEL	91 CNTR	γp Compton scattering

¹² BLANPIED 01 gives $\bar{\alpha}_p + \bar{\beta}_p$ and $\bar{\alpha}_p - \bar{\beta}_p$. The separate $\bar{\alpha}_p$ and $\bar{\beta}_p$ are provided to us by A. Sandorfi. The first error above is statistics plus systematics; the second is from the model.

¹³ This OLMSDELEON 01 result uses the TAPS data alone, and does not use the (re-evaluated) sum-rule constraint that $\bar{\alpha} + \bar{\beta} = (13.8 \pm 0.4) \times 10^{-4}$ fm 3 . See the paper for a discussion.

¹⁴ MACGIBBON 95 combine the results of ZIEGER 92, FEDERSPIEL 91, and their own experiment to get a “global average” in which model errors and systematic errors are treated in a consistent way. See MACGIBBON 95 for a discussion.

¹⁵ BARANOV 01 combines the results of 10 experiments from 1958 through 1995 to get a global average that takes into account both systematic and model errors and does not use the theoretical constraint on the sum $\alpha_p + \beta_p$.

¹⁶ FEDERSPIEL 91 obtains for the (static) electric polarizability α_p , defined in terms of the induced electric dipole moment by $\mathbf{D} = 4\pi\epsilon_0\alpha_p\mathbf{E}$, the value $(7.0 \pm 2.2 \pm 1.3) \times 10^{-4}$ fm 3 .

p MAGNETIC POLARIZABILITY $\bar{\beta}_p$

The electric and magnetic polarizabilities are subject to a dispersion sum-rule constraint $\bar{\alpha} + \bar{\beta} = (14.2 \pm 0.5) \times 10^{-4} \text{ fm}^3$. Errors here are anticorrelated with those on $\bar{\alpha}_p$ due to this constraint.

In the 2004 *Review*, we will probably switch to listing measurements of $\bar{\alpha}_p + \bar{\beta}_p$ and $\bar{\alpha}_p - \bar{\beta}_p$ instead of $\bar{\alpha}_p$ and $\bar{\beta}_p$ separately.

VALUE (10^{-4} fm^3)	DOCUMENT ID	TECN	COMMENT
1.6 ± 0.6 OUR AVERAGE			
$1.43 \pm 0.98^{+0.52}_{-0.98}$	17 BLANPIED 01 LEGS	$p(\vec{\gamma},\gamma)$, $p(\vec{\gamma},\pi^0)$, $p(\vec{\gamma},\pi^+)$	
$1.2 \pm 0.7 \pm 0.5$	18 OL莫斯DEL... 01 CNTR	γp Compton scattering	
$2.1 \pm 0.8 \pm 0.5$	19 MACGIBBON 95 RVUE	global average	
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$2.3 \pm 0.9 \pm 0.7$	20 BARANOV 01 RVUE	Global average	
$1.7 \pm 0.6 \pm 0.9$	MACGIBBON 95 CNTR	γp Compton scattering	
$4.4 \pm 0.4 \pm 1.1$	HALLIN 93 CNTR	γp Compton scattering	
$3.58^{+1.19}_{-1.25}{}^{+1.03}_{-1.07}$	ZIEGER 92 CNTR	γp Compton scattering	
$3.3 \pm 2.2 \pm 1.3$	FEDERSPIEL 91 CNTR	γp Compton scattering	
17 BLANPIED 01 gives $\bar{\alpha}_p + \bar{\beta}_p$ and $\bar{\alpha}_p - \bar{\beta}_p$. The separate $\bar{\alpha}_p$ and $\bar{\beta}_p$ are provided to us by A. Sandorfi. The first error above is statistics plus systematics; the second is from the model.			
18 This OL莫斯ДЕЛЕОН 01 result uses the TAPS data alone, and does not use the (re-evaluated) sum-rule constraint that $\bar{\alpha} + \bar{\beta} = (13.8 \pm 0.4) \times 10^{-4} \text{ fm}^3$. See the paper for a discussion.			
19 MACGIBBON 95 combine the results of ZIEGER 92, FEDERSPIEL 91, and their own experiment to get a “global average” in which model errors and systematic errors are treated in a consistent way. See MACGIBBON 95 for a discussion.			
20 BARANOV 01 combines the results of 10 experiments from 1958 through 1995 to get a global average that takes into account both systematic and model errors and does not use the theoretical constraint on the sum $\alpha_p + \beta_p$.			

p CHARGE RADIUS

VALUE (fm)	DOCUMENT ID	COMMENT
0.870 ± 0.008 OUR AVERAGE		
$0.830 \pm 0.040 \pm 0.040$	21 ESCHRICH 01 $e p \rightarrow e p$	
0.883 ± 0.014	MELNIKOV 00 1S Lamb Shift in H	
0.865 ± 0.020	MCCORD 91 $e p \rightarrow e p$	
0.862 ± 0.012	SIMON 80 $e p \rightarrow e p$	
0.880 ± 0.030	BORKOWSKI 74 $e p \rightarrow e p$	
• • • We do not use the following data for averages, fits, limits, etc. • • •		
0.880 ± 0.015	ROSENFELDR.00 $e p + \text{Coul. corrections}$	
0.847 ± 0.008	MERGELL 96 $e p + \text{disp. relations}$	
0.877 ± 0.024	WONG 94 reanalysis of Mainz $e p$ data	
0.810 ± 0.020	AKIMOV 72 $e p \rightarrow e p$	
0.800 ± 0.025	FREREJACQ... 66 $e p \rightarrow e p$ (CH_2 tgt.)	
0.805 ± 0.011	HAND 63 $e p \rightarrow e p$	
21 ESCHRICH 01 actually gives $\langle r^2 \rangle = (0.69 \pm 0.06 \pm 0.06) \text{ fm}^2$.		

p MEAN LIFE

A test of baryon conservation. See the “*p* Partial Mean Lives” section below for limits that depend on decay modes. *p* = proton, *n* = bound neutron.

LIMIT (years)	PARTICLE	CL%	DOCUMENT ID	TECN	COMMENT
>1.6 × 10²⁵	<i>p, n</i>	22,23	EVANS	77	
>4 × 10 ²³	<i>p</i>	95	TRETYAK	01	$d \rightarrow n + ?$
>1.9 × 10 ²⁴	<i>p</i>	90	24 BERNABEI	00B	DAMA
>3 × 10 ²³	<i>p</i>		23 DIX	70	CNTR
>3 × 10 ²³	<i>p, n</i>		23,25 FLEROV	58	

22 Mean lifetime of nucleons in ^{130}Te nuclei.

23 Converted to mean life by dividing half-life by $\ln(2) = 0.693$.

24 BERNABEI 00B looks for the decay of a $^{128}_{53}\text{I}$ nucleus following the disappearance of a proton in the otherwise-stable $^{129}_{54}\text{Xe}$ nucleus. The *p* decay is to neutrinos or to “nothing,” and thus doesn’t conserve charge as well as baryon number.

25 Mean lifetime of nucleons in ^{232}Th nuclei.

\bar{p} MEAN LIFE

Of the two astrophysical limits here, that of GEER 00D involves considerably more refinements in its modeling. The other limits come from direct observations of stored antiprotons. See also “ \bar{p} Partial Mean Lives” after “*p* Partial Mean Lives,” below, for exclusive-mode limits. The best (life-time/branching fraction) limit there is 7×10^5 years, for $\bar{p} \rightarrow e^- \gamma$. We advance only the exclusive-mode limits to our Summary Tables.

LIMIT (years)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
>8 × 10 ⁵	90		26 GEER	00D	\bar{p}/p ratio, cosmic rays
>0.28			GABRIELSE	90	TRAP Penning trap
>0.08	90	1	BELL	79	CNTR Storage ring
>1 × 10 ⁷			GOLDEN	79	SPEC \bar{p}/p ratio, cosmic rays
>3.7 × 10 ⁻³			BREGMAN	78	CNTR Storage ring

26 GEER 00D uses agreement between a model of galactic \bar{p} production and propagation and the observed \bar{p}/p cosmic-ray spectrum to set this limit.

***p* DECAY MODES**

See the "Note on Nucleon Decay" in our 1994 edition (Phys. Rev. **D50**, 1673) for a short review.

The "partial mean life" limits tabulated here are the limits on τ/B_i , where τ is the total mean life and B_i is the branching fraction for the mode in question. For N decays, p and n indicate proton and neutron partial lifetimes.

Mode	Partial mean life (10^{30} years)	Confidence level
Antilepton + meson		
$\tau_1 \quad N \rightarrow e^+ \pi^-$	> 158 (n), > 1600 (p)	90%
$\tau_2 \quad N \rightarrow \mu^+ \pi^-$	> 100 (n), > 473 (p)	90%
$\tau_3 \quad N \rightarrow \nu \pi$	> 112 (n), > 25 (p)	90%
$\tau_4 \quad p \rightarrow e^+ \eta$	> 313	90%
$\tau_5 \quad p \rightarrow \mu^+ \eta$	> 126	90%
$\tau_6 \quad n \rightarrow \nu \eta$	> 158	90%
$\tau_7 \quad N \rightarrow e^+ \rho^-$	> 217 (n), > 75 (p)	90%
$\tau_8 \quad N \rightarrow \mu^+ \rho^-$	> 228 (n), > 110 (p)	90%
$\tau_9 \quad N \rightarrow \nu \rho$	> 19 (n), > 162 (p)	90%
$\tau_{10} \quad p \rightarrow e^+ \omega$	> 107	90%
$\tau_{11} \quad p \rightarrow \mu^+ \omega$	> 117	90%
$\tau_{12} \quad n \rightarrow \nu \omega$	> 108	90%
$\tau_{13} \quad N \rightarrow e^+ K^-$	> 17 (n), > 150 (p)	90%
$\tau_{14} \quad p \rightarrow e^+ K_S^0$	> 120	90%
$\tau_{15} \quad p \rightarrow e^+ K_L^0$	> 51	90%
$\tau_{16} \quad N \rightarrow \mu^+ K^-$	> 26 (n), > 120 (p)	90%
$\tau_{17} \quad p \rightarrow \mu^+ K_S^0$	> 150	90%
$\tau_{18} \quad p \rightarrow \mu^+ K_L^0$	> 83	90%
$\tau_{19} \quad N \rightarrow \nu K$	> 86 (n), > 670 (p)	90%
$\tau_{20} \quad n \rightarrow \nu K_S^0$	> 51	90%
$\tau_{21} \quad p \rightarrow e^+ K^*(892)^0$	> 84	90%
$\tau_{22} \quad N \rightarrow \nu K^*(892)$	> 78 (n), > 51 (p)	90%
Antilepton + mesons		
$\tau_{23} \quad p \rightarrow e^+ \pi^+ \pi^-$	> 82	90%
$\tau_{24} \quad p \rightarrow e^+ \pi^0 \pi^0$	> 147	90%
$\tau_{25} \quad n \rightarrow e^+ \pi^- \pi^0$	> 52	90%
$\tau_{26} \quad p \rightarrow \mu^+ \pi^+ \pi^-$	> 133	90%
$\tau_{27} \quad p \rightarrow \mu^+ \pi^0 \pi^0$	> 101	90%
$\tau_{28} \quad n \rightarrow \mu^+ \pi^- \pi^0$	> 74	90%
$\tau_{29} \quad n \rightarrow e^+ K^0 \pi^-$	> 18	90%

Lepton + meson

τ_{30}	$n \rightarrow e^- \pi^+$	> 65	90%
τ_{31}	$n \rightarrow \mu^- \pi^+$	> 49	90%
τ_{32}	$n \rightarrow e^- \rho^+$	> 62	90%
τ_{33}	$n \rightarrow \mu^- \rho^+$	> 7	90%
τ_{34}	$n \rightarrow e^- K^+$	> 32	90%
τ_{35}	$n \rightarrow \mu^- K^+$	> 57	90%

Lepton + mesons

τ_{36}	$p \rightarrow e^- \pi^+ \pi^+$	> 30	90%
τ_{37}	$n \rightarrow e^- \pi^+ \pi^0$	> 29	90%
τ_{38}	$p \rightarrow \mu^- \pi^+ \pi^+$	> 17	90%
τ_{39}	$n \rightarrow \mu^- \pi^+ \pi^0$	> 34	90%
τ_{40}	$p \rightarrow e^- \pi^+ K^+$	> 75	90%
τ_{41}	$p \rightarrow \mu^- \pi^+ K^+$	> 245	90%

Antilepton + photon(s)

τ_{42}	$p \rightarrow e^+ \gamma$	> 670	90%
τ_{43}	$p \rightarrow \mu^+ \gamma$	> 478	90%
τ_{44}	$n \rightarrow \nu \gamma$	> 28	90%
τ_{45}	$p \rightarrow e^+ \gamma \gamma$	> 100	90%
τ_{46}	$n \rightarrow \nu \gamma \gamma$	> 219	90%

Three (or more) leptons

τ_{47}	$p \rightarrow e^+ e^+ e^-$	> 793	90%
τ_{48}	$p \rightarrow e^+ \mu^+ \mu^-$	> 359	90%
τ_{49}	$p \rightarrow e^+ \nu \nu$	> 17	90%
τ_{50}	$n \rightarrow e^+ e^- \nu$	> 257	90%
τ_{51}	$n \rightarrow \mu^+ e^- \nu$	> 83	90%
τ_{52}	$n \rightarrow \mu^+ \mu^- \nu$	> 79	90%
τ_{53}	$p \rightarrow \mu^+ e^+ e^-$	> 529	90%
τ_{54}	$p \rightarrow \mu^+ \mu^+ \mu^-$	> 675	90%
τ_{55}	$p \rightarrow \mu^+ \nu \nu$	> 21	90%
τ_{56}	$p \rightarrow e^- \mu^+ \mu^+$	> 6	90%
τ_{57}	$n \rightarrow 3\nu$	> 0.0005	90%
τ_{58}	$n \rightarrow 5\nu$	—	—

Inclusive modes

τ_{59}	$N \rightarrow e^+ \text{anything}$	> 0.6 (n, p)	90%
τ_{60}	$N \rightarrow \mu^+ \text{anything}$	> 12 (n, p)	90%
τ_{61}	$N \rightarrow \nu \text{anything}$	—	—
τ_{62}	$N \rightarrow e^+ \pi^0 \text{anything}$	> 0.6 (n, p)	90%
τ_{63}	$N \rightarrow 2 \text{ bodies, } \nu\text{-free}$	—	—

$\Delta B = 2$ dinucleon modes

The following are lifetime limits per iron nucleus.

τ_{64}	$p p \rightarrow \pi^+ \pi^+$	> 0.7	90%
τ_{65}	$p n \rightarrow \pi^+ \pi^0$	> 2	90%
τ_{66}	$n n \rightarrow \pi^+ \pi^-$	> 0.7	90%
τ_{67}	$n n \rightarrow \pi^0 \pi^0$	> 3.4	90%
τ_{68}	$p p \rightarrow e^+ e^+$	> 5.8	90%
τ_{69}	$p p \rightarrow e^+ \mu^+$	> 3.6	90%
τ_{70}	$p p \rightarrow \mu^+ \mu^+$	> 1.7	90%
τ_{71}	$p n \rightarrow e^+ \bar{\nu}$	> 2.8	90%
τ_{72}	$p n \rightarrow \mu^+ \bar{\nu}$	> 1.6	90%
τ_{73}	$n n \rightarrow \nu_e \bar{\nu}_e$	> 0.000012	90%
τ_{74}	$n n \rightarrow \nu_\mu \bar{\nu}_\mu$	> 0.000006	90%
τ_{75}	$p p \rightarrow \text{neutrinos}$	> 0.00000055	90%

\bar{p} DECAY MODES

Mode	Partial mean life (years)	Confidence level
$\tau_{76} \bar{p} \rightarrow e^- \gamma$	$> 7 \times 10^5$	90%
$\tau_{77} \bar{p} \rightarrow \mu^- \gamma$	$> 5 \times 10^4$	90%
$\tau_{78} \bar{p} \rightarrow e^- \pi^0$	$> 4 \times 10^5$	90%
$\tau_{79} \bar{p} \rightarrow \mu^- \pi^0$	$> 5 \times 10^4$	90%
$\tau_{80} \bar{p} \rightarrow e^- \eta$	$> 2 \times 10^4$	90%
$\tau_{81} \bar{p} \rightarrow \mu^- \eta$	$> 8 \times 10^3$	90%
$\tau_{82} \bar{p} \rightarrow e^- K_S^0$	> 900	90%
$\tau_{83} \bar{p} \rightarrow \mu^- K_S^0$	$> 4 \times 10^3$	90%
$\tau_{84} \bar{p} \rightarrow e^- K_L^0$	$> 9 \times 10^3$	90%
$\tau_{85} \bar{p} \rightarrow \mu^- K_L^0$	$> 7 \times 10^3$	90%
$\tau_{86} \bar{p} \rightarrow e^- \gamma \gamma$	$> 2 \times 10^4$	90%
$\tau_{87} \bar{p} \rightarrow \mu^- \gamma \gamma$	$> 2 \times 10^4$	90%
$\tau_{88} \bar{p} \rightarrow e^- \rho$	—	
$\tau_{89} \bar{p} \rightarrow e^- \omega$	> 200	90%
$\tau_{90} \bar{p} \rightarrow e^- K^*(892)^0$	—	

p PARTIAL MEAN LIVES

The “partial mean life” limits tabulated here are the limits on τ/B_i , where τ is the total mean life for the proton and B_i is the branching fraction for the mode in question.

Decaying particle: p = proton, n = bound neutron. The same event may appear under more than one partial decay mode. Background estimates may be accurate to a factor of two.

— Antilepton + meson —

$\tau(N \rightarrow e^+ \pi)$

 τ_1

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
> 158	n	90	3	5	MCGREW	99 IMB3
> 1600	p	90	0	0.1	SHIOZAWA	98 SKAM

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 540	p	90	0	0.2	MCGREW	99 IMB3
> 70	p	90	0	0.5	BERGER	91 FREJ
> 70	n	90	0	≤ 0.1	BERGER	91 FREJ
> 550	p	90	0	0.7	²⁷ BECKER-SZ...	90 IMB3
> 260	p	90	0	<0.04	HIRATA	89C KAMI
> 130	n	90	0	<0.2	HIRATA	89C KAMI
> 310	p	90	0	0.6	SEIDEL	88 IMB
> 100	n	90	0	1.6	SEIDEL	88 IMB
> 1.3	n	90	0		BARTEL	87 SOUD
> 1.3	p	90	0		BARTEL	87 SOUD
> 250	p	90	0	0.3	HAINES	86 IMB
> 31	n	90	8	9	HAINES	86 IMB
> 64	p	90	0	<0.4	ARISAKA	85 KAMI
> 26	n	90	0	<0.7	ARISAKA	85 KAMI
> 82	p (free)	90	0	0.2	BLEWITT	85 IMB
> 250	p	90	0	0.2	BLEWITT	85 IMB
> 25	n	90	4	4	PARK	85 IMB
> 15	p, n	90	0		²⁸ BATTISTONI	84 NUSX
> 0.5	p	90	1	0.3	BARTEL	83 SOUD
> 0.5	n	90	1	0.3	²⁸ BARTEL	83 SOUD
> 5.8	p	90	2		²⁹ KRISHNA...	82 KOLR
> 5.8	n	90	2		²⁹ KRISHNA...	82 KOLR
> 0.1	n	90			³⁰ GURU	67 CNTR

²⁷ This BECKER-SZENDY 90 result includes data from SEIDEL 88.

²⁸ Limit based on zero events.

²⁹ We have calculated 90% CL limit from 1 confined event.

³⁰ We have converted half-life to 90% CL mean life.

$\tau(N \rightarrow \mu^+ \pi)$

 τ_2

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
> 473	p	90	0	0.6	MCGREW	99 IMB3
> 100	n	90	0	<0.2	HIRATA	89C KAMI

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 90	<i>n</i>	90	1	1.9	MCGREW	99	IMB3
> 81	<i>p</i>	90	0	0.2	BERGER	91	FREJ
> 35	<i>n</i>	90	1	1.0	BERGER	91	FREJ
>230	<i>p</i>	90	0	<0.07	HIRATA	89C	KAMI
>270	<i>p</i>	90	0	0.5	SEIDEL	88	IMB
> 63	<i>n</i>	90	0	0.5	SEIDEL	88	IMB
> 76	<i>p</i>	90	2	1	HAINES	86	IMB
> 23	<i>n</i>	90	8	7	HAINES	86	IMB
> 46	<i>p</i>	90	0	<0.7	ARISAKA	85	KAMI
> 20	<i>n</i>	90	0	<0.4	ARISAKA	85	KAMI
> 59	<i>p</i> (free)	90	0	0.2	BLEWITT	85	IMB
>100	<i>p</i>	90	1	0.4	BLEWITT	85	IMB
> 38	<i>n</i>	90	1	4	PARK	85	IMB
> 10	<i>p, n</i>	90	0		BATTISTONI	84	NUSX
> 1.3	<i>p, n</i>	90	0		ALEKSEEV	81	BAKS

$\tau(N \rightarrow \nu\pi)$

τ_3

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN	
>112	<i>n</i>	90	6	6.6	MCGREW	99	IMB3
> 25	<i>p</i>	90	32	32.8	HIRATA	89C	KAMI

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 16	<i>p</i>	90	6	6.7	WALL	00B	SOU2
> 39	<i>n</i>	90	4	3.8	WALL	00B	SOU2
> 10	<i>p</i>	90	15	20.3	MCGREW	99	IMB3
> 13	<i>n</i>	90	1	1.2	BERGER	89	FREJ
> 10	<i>p</i>	90	11	14	BERGER	89	FREJ
>100	<i>n</i>	90	1	3	HIRATA	89C	KAMI
> 6	<i>n</i>	90	73	60	HAINES	86	IMB
> 2	<i>p</i>	90	16	13	KAJITA	86	KAMI
> 40	<i>n</i>	90	0	1	KAJITA	86	KAMI
> 7	<i>n</i>	90	28	19	PARK	85	IMB
> 7	<i>n</i>	90	0		BATTISTONI	84	NUSX
> 2	<i>p</i>	90	≤ 3		BATTISTONI	84	NUSX
> 5.8	<i>p</i>	90	1		31 KRISHNA...	82	KOLR
> 0.3	<i>p</i>	90	2		32 CHERRY	81	HOME
> 0.1	<i>p</i>	90			33 GURU	67	CNTR

³¹We have calculated 90% CL limit from 1 confined event.

³²We have converted 2 possible events to 90% CL limit.

³³We have converted half-life to 90% CL mean life.

$\tau(p \rightarrow e^+ \eta)$

τ_4

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN	
>313	<i>p</i>	90	0	0.2	MCGREW	99	IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 81	<i>p</i>	90	1	1.7	WALL	00B	SOU2
> 44	<i>p</i>	90	0	0.1	BERGER	91	FREJ
>140	<i>p</i>	90	0	<0.04	HIRATA	89C	KAMI
>100	<i>p</i>	90	0	0.6	SEIDEL	88	IMB
>200	<i>p</i>	90	5	3.3	HAINES	86	IMB
> 64	<i>p</i>	90	0	<0.8	ARISAKA	85	KAMI
> 64	<i>p</i> (free)	90	5	6.5	BLEWITT	85	IMB
>200	<i>p</i>	90	5	4.7	BLEWITT	85	IMB
> 1.2	<i>p</i>	90	2		³⁴ CHERRY	81	HOME

³⁴ We have converted 2 possible events to 90% CL limit.

$\tau(p \rightarrow \mu^+ \eta)$

τ_5

<i>LIMIT</i> (10^{30} years)	<i>PARTICLE</i>	<i>CL%</i>	<i>EVTS</i>	<i>BKGD EST</i>	<i>DOCUMENT ID</i>	<i>TECN</i>
>126	<i>p</i>	90	3	2.8	MCGREW	99 IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 89	<i>p</i>	90	0	1.6	WALL	00B	SOU2
> 26	<i>p</i>	90	1	0.8	BERGER	91	FREJ
> 69	<i>p</i>	90	1	<0.08	HIRATA	89C	KAMI
> 1.3	<i>p</i>	90	0	0.7	PHILLIPS	89	HPW
> 34	<i>p</i>	90	1	1.5	SEIDEL	88	IMB
> 46	<i>p</i>	90	7	6	HAINES	86	IMB
> 26	<i>p</i>	90	1	<0.8	ARISAKA	85	KAMI
> 17	<i>p</i> (free)	90	6	6	BLEWITT	85	IMB
> 46	<i>p</i>	90	7	8	BLEWITT	85	IMB

$\tau(n \rightarrow \nu\eta)$

τ_6

<i>LIMIT</i> (10^{30} years)	<i>PARTICLE</i>	<i>CL%</i>	<i>EVTS</i>	<i>BKGD EST</i>	<i>DOCUMENT ID</i>	<i>TECN</i>
>158	<i>n</i>	90	0	1.2	MCGREW	99 IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 71	<i>n</i>	90	2	3.7	WALL	00B	SOU2
> 29	<i>n</i>	90	0	0.9	BERGER	89	FREJ
> 54	<i>n</i>	90	2	0.9	HIRATA	89C	KAMI
> 16	<i>n</i>	90	3	2.1	SEIDEL	88	IMB
> 25	<i>n</i>	90	7	6	HAINES	86	IMB
> 30	<i>n</i>	90	0	0.4	KAJITA	86	KAMI
> 18	<i>n</i>	90	4	3	PARK	85	IMB
> 0.6	<i>n</i>	90	2		³⁵ CHERRY	81	HOME

³⁵ We have converted 2 possible events to 90% CL limit.

$\tau(N \rightarrow e^+ \rho)$

τ_7

<i>LIMIT</i> (10^{30} years)	<i>PARTICLE</i>	<i>CL%</i>	<i>EVTS</i>	<i>BKGD EST</i>	<i>DOCUMENT ID</i>	<i>TECN</i>
>217	<i>n</i>	90	4	4.8	MCGREW	99 IMB3
> 75	<i>p</i>	90	2	2.7	HIRATA	89C KAMI

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 29	<i>p</i>	90	0	2.2	BERGER	91	FREJ
> 41	<i>n</i>	90	0	1.4	BERGER	91	FREJ
> 58	<i>n</i>	90	0	1.9	HIRATA	89C	KAMI
> 38	<i>n</i>	90	2	4.1	SEIDEL	88	IMB
> 1.2	<i>p</i>	90	0		BARTEL	87	SOU
> 1.5	<i>n</i>	90	0		BARTEL	87	SOU
> 17	<i>p</i>	90	7	7	HAINES	86	IMB
> 14	<i>n</i>	90	9	4	HAINES	86	IMB
> 12	<i>p</i>	90	0	<1.2	ARISAKA	85	KAMI
> 6	<i>n</i>	90	2	<1	ARISAKA	85	KAMI
> 6.7	<i>p</i> (free)	90	6	6	BLEWITT	85	IMB
> 17	<i>p</i>	90	7	7	BLEWITT	85	IMB
> 12	<i>n</i>	90	4	2	PARK	85	IMB
> 0.6	<i>n</i>	90	1	0.3	36 BARTEL	83	SOU
> 0.5	<i>p</i>	90	1	0.3	36 BARTEL	83	SOU
> 9.8	<i>p</i>	90	1		37 KRISHNA...	82	KOLR
> 0.8	<i>p</i>	90	2		38 CHERRY	81	HOME

36 Limit based on zero events.

37 We have calculated 90% CL limit from 0 confined events.

38 We have converted 2 possible events to 90% CL limit.

$\tau(N \rightarrow \mu^+ \rho)$

τ_8

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>228	<i>n</i>	90	3	9.5	MCGREW	99 IMB3
>110	<i>p</i>	90	0	1.7	HIRATA	89C KAMI

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 12	<i>p</i>	90	0	0.5	BERGER	91	FREJ
> 22	<i>n</i>	90	0	1.1	BERGER	91	FREJ
> 23	<i>n</i>	90	1	1.8	HIRATA	89C	KAMI
> 4.3	<i>p</i>	90	0	0.7	PHILLIPS	89	HPW
> 30	<i>p</i>	90	0	0.5	SEIDEL	88	IMB
> 11	<i>n</i>	90	1	1.1	SEIDEL	88	IMB
> 16	<i>p</i>	90	4	4.5	HAINES	86	IMB
> 7	<i>n</i>	90	6	5	HAINES	86	IMB
> 12	<i>p</i>	90	0	<0.7	ARISAKA	85	KAMI
> 5	<i>n</i>	90	1	<1.2	ARISAKA	85	KAMI
> 5.5	<i>p</i> (free)	90	4	5	BLEWITT	85	IMB
> 16	<i>p</i>	90	4	5	BLEWITT	85	IMB
> 9	<i>n</i>	90	1	2	PARK	85	IMB

$\tau(N \rightarrow \nu \rho)$

τ_9

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>162	<i>p</i>	90	18	21.7	MCGREW	99 IMB3
> 19	<i>n</i>	90	0	0.5	SEIDEL	88 IMB

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 9	<i>n</i>	90	4	2.4	BERGER	89	FREJ
> 24	<i>p</i>	90	0	0.9	BERGER	89	FREJ
> 27	<i>p</i>	90	5	1.5	HIRATA	89C	KAMI
> 13	<i>n</i>	90	4	3.6	HIRATA	89C	KAMI
> 13	<i>p</i>	90	1	1.1	SEIDEL	88	IMB
> 8	<i>p</i>	90	6	5	HAINES	86	IMB
> 2	<i>n</i>	90	15	10	HAINES	86	IMB
> 11	<i>p</i>	90	2	1	KAJITA	86	KAMI
> 4	<i>n</i>	90	2	2	KAJITA	86	KAMI
> 4.1	<i>p</i> (free)	90	6	7	BLEWITT	85	IMB
> 8.4	<i>p</i>	90	6	5	BLEWITT	85	IMB
> 2	<i>n</i>	90	7	3	PARK	85	IMB
> 0.9	<i>p</i>	90	2				
> 0.6	<i>n</i>	90	2				

³⁹We have converted 2 possible events to 90% CL limit.

$\tau(p \rightarrow e^+ \omega)$

τ_{10}

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>107	<i>p</i>	90	7	10.8	MCGREW	99 IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 17	<i>p</i>	90	0	1.1	BERGER	91	FREJ
> 45	<i>p</i>	90	2	1.45	HIRATA	89C	KAMI
> 26	<i>p</i>	90	1	1.0	SEIDEL	88	IMB
> 1.5	<i>p</i>	90	0		BARTEL	87	SOUD
> 37	<i>p</i>	90	6	5.3	HAINES	86	IMB
> 25	<i>p</i>	90	1	<1.4	ARISAKA	85	KAMI
> 12	<i>p</i> (free)	90	6	7.5	BLEWITT	85	IMB
> 37	<i>p</i>	90	6	5.7	BLEWITT	85	IMB
> 0.6	<i>p</i>	90	1	0.3	83 SOUD		
> 9.8	<i>p</i>	90	1		82 KOLR		
> 2.8	<i>p</i>	90	2				

⁴⁰Limit based on zero events.

⁴¹We have calculated 90% CL limit from 0 confined events.

⁴²We have converted 2 possible events to 90% CL limit.

$\tau(p \rightarrow \mu^+ \omega)$

τ_{11}

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>117	<i>p</i>	90	11	12.1	MCGREW	99 IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 11	<i>p</i>	90	0	1.0	BERGER	91	FREJ
> 57	<i>p</i>	90	2	1.9	HIRATA	89C	KAMI
> 4.4	<i>p</i>	90	0	0.7	PHILLIPS	89	HPW
> 10	<i>p</i>	90	2	1.3	SEIDEL	88	IMB
> 23	<i>p</i>	90	2	1	HAINES	86	IMB
> 6.5	<i>p</i> (free)	90	9	8.7	BLEWITT	85	IMB
> 23	<i>p</i>	90	8	7	BLEWITT	85	IMB

$\tau(n \rightarrow \nu\omega)$ τ_{12}

<i>LIMIT</i> (10^{-30} years)	<i>PARTICLE</i>	<i>CL%</i>	<i>EVTS</i>	<i>BKGD EST</i>	<i>DOCUMENT ID</i>	<i>TECN</i>
>108	<i>n</i>	90	12	22.5	MCGREW	99 IMB3
• • • We do not use the following data for averages, fits, limits, etc. • • •						
> 17	<i>n</i>	90	1	0.7	BERGER	89 FREJ
> 43	<i>n</i>	90	3	2.7	HIRATA	89C KAMI
> 6	<i>n</i>	90	2	1.3	SEIDEL	88 IMB
> 12	<i>n</i>	90	6	6	HAINES	86 IMB
> 18	<i>n</i>	90	2	2	KAJITA	86 KAMI
> 16	<i>n</i>	90	1	2	PARK	85 IMB
> 2.0	<i>n</i>	90	2		⁴³ CHERRY	81 HOME

43 We have converted 2 possible events to 90% CL limit.

 $\tau(N \rightarrow e^+ K)$ τ_{13}

<i>LIMIT</i> (10^{-30} years)	<i>PARTICLE</i>	<i>CL%</i>	<i>EVTS</i>	<i>BKGD EST</i>	<i>DOCUMENT ID</i>	<i>TECN</i>
> 17	<i>n</i>	90	35	29.4	MCGREW	99 IMB3
>150	<i>p</i>	90	0	<0.27	HIRATA	89C KAMI
• • • We do not use the following data for averages, fits, limits, etc. • • •						
> 85	<i>p</i>	90	3	4.9	WALL	00 SOU2
> 31	<i>p</i>	90	23	25.2	MCGREW	99 IMB3
> 60	<i>p</i>	90	0		BERGER	91 FREJ
> 70	<i>p</i>	90	0	1.8	SEIDEL	88 IMB
> 77	<i>p</i>	90	5	4.5	HAINES	86 IMB
> 38	<i>p</i>	90	0	<0.8	ARISAKA	85 KAMI
> 24	<i>p</i> (free)	90	7	8.5	BLEWITT	85 IMB
> 77	<i>p</i>	90	5	4	BLEWITT	85 IMB
> 1.3	<i>p</i>	90	0		ALEKSEEV	81 BAKS
> 1.3	<i>n</i>	90	0		ALEKSEEV	81 BAKS

 $\tau(p \rightarrow e^+ K_S^0)$ τ_{14}

<i>LIMIT</i> (10^{-30} years)	<i>PARTICLE</i>	<i>CL%</i>	<i>EVTS</i>	<i>BKGD EST</i>	<i>DOCUMENT ID</i>	<i>TECN</i>
>120	<i>p</i>	90	1	1.3	WALL	00 SOU2
• • • We do not use the following data for averages, fits, limits, etc. • • •						
> 76	<i>p</i>	90	0	0.5	BERGER	91 FREJ

 $\tau(p \rightarrow e^+ K_L^0)$ τ_{15}

<i>LIMIT</i> (10^{-30} years)	<i>PARTICLE</i>	<i>CL%</i>	<i>EVTS</i>	<i>BKGD EST</i>	<i>DOCUMENT ID</i>	<i>TECN</i>
>51	<i>p</i>	90	2	3.5	WALL	00 SOU2
• • • We do not use the following data for averages, fits, limits, etc. • • •						
>44	<i>p</i>	90	0	≤ 0.1	BERGER	91 FREJ

$\tau(N \rightarrow \mu^+ K)$ τ_{16}

<u>LIMIT</u> (10^{-30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>120	p	90	0	<1.2	WALL	00 SOU2
>120	p	90	4	7.2	MCGREW	99 IMB3
> 26	n	90	20	28.4	MCGREW	99 IMB3
>120	p	90	1	0.4	HIRATA	89C KAMI

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 54	p	90	0		BERGER	91 FREJ
> 3.0	p	90	0	0.7	PHILLIPS	89 HPW
> 19	p	90	3	2.5	SEIDEL	88 IMB
> 1.5	p	90	0		⁴⁴ BARTEL	87 SOUD
> 1.1	n	90	0		BARTEL	87 SOUD
> 40	p	90	7	6	HAINES	86 IMB
> 19	p	90	1	<1.1	ARISAKA	85 KAMI
> 6.7	p (free)	90	11	13	BLEWITT	85 IMB
> 40	p	90	7	8	BLEWITT	85 IMB
> 6	p	90	1		BATTISTONI	84 NUSX
> 0.6	p	90	0		⁴⁵ BARTEL	83 SOUD
> 0.4	n	90	0		⁴⁵ BARTEL	83 SOUD
> 5.8	p	90	2		⁴⁶ KRISHNA...	82 KOLR
> 2.0	p	90	0		CHERRY	81 HOME
> 0.2	n	90			⁴⁷ GURR	67 CNTR

⁴⁴ BARTEL 87 limit applies to $p \rightarrow \mu^+ K_S^0$.

⁴⁵ Limit based on zero events.

⁴⁶ We have calculated 90% CL limit from 1 confined event.

⁴⁷ We have converted half-life to 90% CL mean life.

 $\tau(p \rightarrow \mu^+ K_S^0)$ τ_{17}

<u>LIMIT</u> (10^{-30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>150	p	90	0	<0.8	WALL	00 SOU2

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 64	p	90	0	1.2	BERGER	91 FREJ
------	---	----	---	-----	--------	---------

 $\tau(p \rightarrow \mu^+ K_L^0)$ τ_{18}

<u>LIMIT</u> (10^{-30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>83	p	90	0	0.4	WALL	00 SOU2

• • • We do not use the following data for averages, fits, limits, etc. • • •

>44	p	90	0	≤ 0.1	BERGER	91 FREJ
-----	---	----	---	------------	--------	---------

 $\tau(N \rightarrow \nu K)$ τ_{19}

<u>LIMIT</u> (10^{-30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>670	p	90			HAYATO	99 SKAM
> 86	n	90	0	2.4	HIRATA	89C KAMI

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 26	<i>n</i>	90	16	9.1	WALL	00	SOU2
>151	<i>p</i>	90	15	21.4	MCGREW	99	IMB3
> 30	<i>n</i>	90	34	34.1	MCGREW	99	IMB3
> 43	<i>p</i>	90	1	1.54	48 ALLISON	98	SOU2
> 15	<i>n</i>	90	1	1.8	BERGER	89	FREJ
> 15	<i>p</i>	90	1	1.8	BERGER	89	FREJ
>100	<i>p</i>	90	9	7.3	HIRATA	89C	KAMI
> 0.28	<i>p</i>	90	0	0.7	PHILLIPS	89	HPW
> 0.3	<i>p</i>	90	0		BARTEL	87	SOU2
> 0.75	<i>n</i>	90	0		49 BARTEL	87	SOU2
> 10	<i>p</i>	90	6	5	HAINES	86	IMB
> 15	<i>n</i>	90	3	5	HAINES	86	IMB
> 28	<i>p</i>	90	3	3	KAJITA	86	KAMI
> 32	<i>n</i>	90	0	1.4	KAJITA	86	KAMI
> 1.8	<i>p</i> (free)	90	6	11	BLEWITT	85	IMB
> 9.6	<i>p</i>	90	6	5	BLEWITT	85	IMB
> 10	<i>n</i>	90	2	2	PARK	85	IMB
> 5	<i>n</i>	90	0		BATTISTONI	84	NUSX
> 2	<i>p</i>	90	0		BATTISTONI	84	NUSX
> 0.3	<i>n</i>	90	0		50 BARTEL	83	SOU2
> 0.1	<i>p</i>	90	0		50 BARTEL	83	SOU2
> 5.8	<i>p</i>	90	1		51 KRISHNA...	82	KOLR
> 0.3	<i>n</i>	90	2		52 CHERRY	81	HOME

⁴⁸ This ALLISON 98 limit is with no background subtraction; with subtraction the limit becomes $> 46 \times 10^{30}$ years.

⁴⁹ BARTEL 87 limit applies to $n \rightarrow \nu K_S^0$.

⁵⁰ Limit based on zero events.

⁵¹ We have calculated 90% CL limit from 1 confined event.

⁵² We have converted 2 possible events to 90% CL limit.

$\tau(n \rightarrow \nu K_S^0)$

τ_{20}

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN	
>51	<i>n</i>	90	16	9.1	WALL	00	SOU2

$\tau(p \rightarrow e^+ K^*(892)^0)$

τ_{21}

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN	
>84	<i>p</i>	90	38	52.0	MCGREW	99	IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

>10	<i>p</i>	90	0	0.8	BERGER	91	FREJ
>52	<i>p</i>	90	2	1.55	HIRATA	89C	KAMI
>10	<i>p</i>	90	1	<1	ARISAKA	85	KAMI

$\tau(N \rightarrow \nu K^*(892))$

τ_{22}

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN	
>51	<i>p</i>	90	7	9.1	MCGREW	99	IMB3
>78	<i>n</i>	90	40	50	MCGREW	99	IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

>22	<i>n</i>	90	0	2.1	BERGER	89	FREJ
>17	<i>p</i>	90	0	2.4	BERGER	89	FREJ
>20	<i>p</i>	90	5	2.1	HIRATA	89C	KAMI
>21	<i>n</i>	90	4	2.4	HIRATA	89C	KAMI
>10	<i>p</i>	90	7	6	HAINES	86	IMB
> 5	<i>n</i>	90	8	7	HAINES	86	IMB
> 8	<i>p</i>	90	3	2	KAJITA	86	KAMI
> 6	<i>n</i>	90	2	1.6	KAJITA	86	KAMI
> 5.8	<i>p</i> (free)	90	10	16	BLEWITT	85	IMB
> 9.6	<i>p</i>	90	7	6	BLEWITT	85	IMB
> 7	<i>n</i>	90	1	4	PARK	85	IMB
> 2.1	<i>p</i>	90	1		53 BATTISTONI	82	NUSX

⁵³We have converted 1 possible event to 90% CL limit.

— Antilepton + mesons —

$\tau(p \rightarrow e^+ \pi^+ \pi^-)$

τ_{23}

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>82	<i>p</i>	90	16	23.1	MCGREW	99

• • • We do not use the following data for averages, fits, limits, etc. • • •

>21	<i>p</i>	90	0	2.2	BERGER	91	FREJ
-----	----------	----	---	-----	--------	----	------

$\tau(p \rightarrow e^+ \pi^0 \pi^0)$

τ_{24}

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>147	<i>p</i>	90	2	0.8	MCGREW	99

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 38	<i>p</i>	90	1	0.5	BERGER	91	FREJ
------	----------	----	---	-----	--------	----	------

$\tau(n \rightarrow e^+ \pi^- \pi^0)$

τ_{25}

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>52	<i>n</i>	90	38	34.2	MCGREW	99

• • • We do not use the following data for averages, fits, limits, etc. • • •

>32	<i>n</i>	90	1	0.8	BERGER	91	FREJ
-----	----------	----	---	-----	--------	----	------

$\tau(p \rightarrow \mu^+ \pi^+ \pi^-)$

τ_{26}

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>133	<i>p</i>	90	25	38.0	MCGREW	99

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 17	<i>p</i>	90	1	2.6	BERGER	91	FREJ
> 3.3	<i>p</i>	90	0	0.7	PHILLIPS	89	HPW

$\tau(p \rightarrow \mu^+ \pi^0 \pi^0)$ τ_{27}

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>101	p	90	3	1.6	MCGREW	99 IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 33	p	90	1	0.9	BERGER	91 FREJ
------	---	----	---	-----	--------	---------

 $\tau(n \rightarrow \mu^+ \pi^- \pi^0)$ τ_{28}

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>74	n	90	17	20.8	MCGREW	99 IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

>33	n	90	0	1.1	BERGER	91 FREJ
-----	---	----	---	-----	--------	---------

 $\tau(n \rightarrow e^+ K^0 \pi^-)$ τ_{29}

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>18	n	90	1	0.2	BERGER	91 FREJ

 Lepton + meson

 $\tau(n \rightarrow e^- \pi^+)$ τ_{30}

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>65	n	90	0	1.6	SEIDEL	88 IMB

• • • We do not use the following data for averages, fits, limits, etc. • • •

>55	n	90	0	1.09	BERGER	91B FREJ
>16	n	90	9	7	HAINES	86 IMB
>25	n	90	2	4	PARK	85 IMB

 $\tau(n \rightarrow \mu^- \pi^+)$ τ_{31}

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>49	n	90	0	0.5	SEIDEL	88 IMB

• • • We do not use the following data for averages, fits, limits, etc. • • •

>33	n	90	0	1.40	BERGER	91B FREJ
> 2.7	n	90	0	0.7	PHILLIPS	89 HPW
>25	n	90	7	6	HAINES	86 IMB
>27	n	90	2	3	PARK	85 IMB

 $\tau(n \rightarrow e^- \rho^+)$ τ_{32}

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>62	n	90	2	4.1	SEIDEL	88 IMB

• • • We do not use the following data for averages, fits, limits, etc. • • •

>12	n	90	13	6	HAINES	86 IMB
>12	n	90	5	3	PARK	85 IMB

$\tau(n \rightarrow \mu^- \rho^+)$ τ_{33}

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>7	n	90	1	1.1	SEIDEL	88 IMB
• • • We do not use the following data for averages, fits, limits, etc. • • •						
>2.6	n	90	0	0.7	PHILLIPS	89 HPW
>9	n	90	7	5	HAINES	86 IMB
>9	n	90	2	2	PARK	85 IMB

 $\tau(n \rightarrow e^- K^+)$ τ_{34}

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>32	n	90	3	2.96	BERGER	91B FREJ
• • • We do not use the following data for averages, fits, limits, etc. • • •						
> 0.23	n	90	0	0.7	PHILLIPS	89 HPW

 $\tau(n \rightarrow \mu^- K^+)$ τ_{35}

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>57	n	90	0	2.18	BERGER	91B FREJ
• • • We do not use the following data for averages, fits, limits, etc. • • •						
> 4.7	n	90	0	0.7	PHILLIPS	89 HPW

Lepton + mesons $\tau(p \rightarrow e^- \pi^+ \pi^+)$ τ_{36}

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>30	p	90	1	2.50	BERGER	91B FREJ
• • • We do not use the following data for averages, fits, limits, etc. • • •						
> 2.0	p	90	0	0.7	PHILLIPS	89 HPW

 $\tau(n \rightarrow e^- \pi^+ \pi^0)$ τ_{37}

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>29	n	90	1	0.78	BERGER	91B FREJ

 $\tau(p \rightarrow \mu^- \pi^+ \pi^+)$ τ_{38}

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>17	p	90	1	1.72	BERGER	91B FREJ
• • • We do not use the following data for averages, fits, limits, etc. • • •						
> 7.8	p	90	0	0.7	PHILLIPS	89 HPW

 $\tau(n \rightarrow \mu^- \pi^+ \pi^0)$ τ_{39}

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>34	n	90	0	0.78	BERGER	91B FREJ

$\tau(p \rightarrow e^- \pi^+ K^+)$

T40

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>75	p	90	81	127.2	MCGREW	99 IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

>20 p 90 3 2.50

BERGER 91B FREJ

$\tau(p \rightarrow \mu^- \pi^+ K^+)$

T41

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>245	p	90	3	4.0	MCGREW	99 IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 5 p 90 2 0.78

BERGER 91B FREJ

— Antilepton + photon(s) —

$\tau(p \rightarrow e^+ \gamma)$

T42

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>670	p	90	0	0.1	MCGREW	99 IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

>133 p 90 0 0.3
 >460 p 90 0 0.6
 >360 p 90 0 0.3
 > 87 p (free) 90 0 0.2
 >360 p 90 0 0.2
 > 0.1 p 90

⁵⁴ GURR 67 CNTR

⁵⁴ We have converted half-life to 90% CL mean life.

$\tau(p \rightarrow \mu^+ \gamma)$

T43

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>478	p	90	0	0.1	MCGREW	99 IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

>155 p 90 0 0.1
 >380 p 90 0 0.5
 > 97 p 90 3 2
 > 61 p (free) 90 0 0.2
 >280 p 90 0 0.6
 > 0.3 p 90

⁵⁵ GURR 67 CNTR

⁵⁵ We have converted half-life to 90% CL mean life.

$\tau(n \rightarrow \nu \gamma)$

T44

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>28	n	90	163	144.7	MCGREW	99 IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

>24 n 90 10 6.86
 > 9 n 90 73 60
 >11 n 90 28 19

BERGER 91B FREJ
 HAINES 86 IMB
 PARK 85 IMB

$\tau(p \rightarrow e^+ \gamma\gamma)$ **T45**

<i>LIMIT</i> (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST
>100	p	90	1	0.8

DOCUMENT ID	TECN
BERGER	91 FREJ

 $\tau(n \rightarrow \nu\gamma\gamma)$ **T46**

<i>LIMIT</i> (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST
>219	n	90	5	7.5

DOCUMENT ID	TECN
MCGREW	99 IMB3

Three (or more) leptons $\tau(p \rightarrow e^+ e^+ e^-)$ **T47**

<i>LIMIT</i> (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST
>793	p	90	0	0.5

DOCUMENT ID	TECN
MCGREW	99 IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

>147	p	90	0	0.1	BERGER	91	FREJ
>510	p	90	0	0.3	HAINES	86	IMB
> 89	p (free)	90	0	0.5	BLEWITT	85	IMB
>510	p	90	0	0.7	BLEWITT	85	IMB

 $\tau(p \rightarrow e^+ \mu^+ \mu^-)$ **T48**

<i>LIMIT</i> (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST
>359	p	90	1	0.9

DOCUMENT ID	TECN
MCGREW	99 IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 81	p	90	0	0.16	BERGER	91	FREJ
> 5.0	p	90	0	0.7	PHILLIPS	89	HPW

 $\tau(p \rightarrow e^+ \nu\nu)$ **T49**

<i>LIMIT</i> (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST
>17	p	90	152	153.7

DOCUMENT ID	TECN
MCGREW	99 IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

>11	p	90	11	6.08	BERGER	91B	FREJ
-----	-----	----	----	------	--------	-----	------

 $\tau(n \rightarrow e^+ e^- \nu)$ **T50**

<i>LIMIT</i> (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST
>257	n	90	5	7.5

DOCUMENT ID	TECN
MCGREW	99 IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 74	n	90	0	< 0.1	BERGER	91B	FREJ
> 45	n	90	5	5	HAINES	86	IMB
> 26	n	90	4	3	PARK	85	IMB

 $\tau(n \rightarrow \mu^+ e^- \nu)$ **T51**

<i>LIMIT</i> (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST
>83	n	90	25	29.4

DOCUMENT ID	TECN
MCGREW	99 IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

>47	n	90	0	< 0.1	BERGER	91B	FREJ
-----	-----	----	---	-------	--------	-----	------

$\tau(n \rightarrow \mu^+ \mu^- \nu)$ **T52**

<i>LIMIT</i> (10^{30} years)	<i>PARTICLE</i>	<i>CL%</i>	<i>EVTS</i>	<i>BKGD EST</i>	<i>DOCUMENT ID</i>	<i>TECN</i>
>79	n	90	100	145	MCGREW	99 IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

>42	<i>n</i>	90	0	1.4	BERGER	91B FREJ
> 5.1	<i>n</i>	90	0	0.7	PHILLIPS	89 HPW
>16	<i>n</i>	90	14	7	HAINES	86 IMB
>19	<i>n</i>	90	4	7	PARK	85 IMB

 $\tau(p \rightarrow \mu^+ e^+ e^-)$ **T53**

<i>LIMIT</i> (10^{30} years)	<i>PARTICLE</i>	<i>CL%</i>	<i>EVTS</i>	<i>BKGD EST</i>	<i>DOCUMENT ID</i>	<i>TECN</i>
>529	p	90	0	1.0	MCGREW	99 IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 91	<i>p</i>	90	0	≤ 0.1	BERGER	91 FREJ
------	----------	----	---	------------	--------	---------

 $\tau(p \rightarrow \mu^+ \mu^+ \mu^-)$ **T54**

<i>LIMIT</i> (10^{30} years)	<i>PARTICLE</i>	<i>CL%</i>	<i>EVTS</i>	<i>BKGD EST</i>	<i>DOCUMENT ID</i>	<i>TECN</i>
>675	p	90	0	0.3	MCGREW	99 IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

>119	<i>p</i>	90	0	0.2	BERGER	91 FREJ
> 10.5	<i>p</i>	90	0	0.7	PHILLIPS	89 HPW
>190	<i>p</i>	90	1	0.1	HAINES	86 IMB
> 44	<i>p</i> (free)	90	1	0.7	BLEWITT	85 IMB
>190	<i>p</i>	90	1	0.9	BLEWITT	85 IMB
> 2.1	<i>p</i>	90	1		⁵⁶ BATTISTONI	82 NUSX

⁵⁶ We have converted 1 possible event to 90% CL limit.

 $\tau(p \rightarrow \mu^+ \nu \nu)$ **T55**

<i>LIMIT</i> (10^{30} years)	<i>PARTICLE</i>	<i>CL%</i>	<i>EVTS</i>	<i>BKGD EST</i>	<i>DOCUMENT ID</i>	<i>TECN</i>
>21	p	90	7	11.23	BERGER	91B FREJ

 $\tau(p \rightarrow e^- \mu^+ \mu^+)$ **T56**

<i>LIMIT</i> (10^{30} years)	<i>PARTICLE</i>	<i>CL%</i>	<i>EVTS</i>	<i>BKGD EST</i>	<i>DOCUMENT ID</i>	<i>TECN</i>
>6.0	p	90	0	0.7	PHILLIPS	89 HPW

 $\tau(n \rightarrow 3\nu)$ **T57**

<i>LIMIT</i> (10^{30} years)	<i>PARTICLE</i>	<i>CL%</i>	<i>EVTS</i>	<i>BKGD EST</i>	<i>DOCUMENT ID</i>	<i>TECN</i>
>0.00049	n	90	2	2	⁵⁷ SUZUKI	93B KAMI

• • • We do not use the following data for averages, fits, limits, etc. • • •

>0.0023	<i>n</i>	90			⁵⁸ GLICENSTEIN	97 KAMI
>0.00003	<i>n</i>	90	11	6.1	⁵⁹ BERGER	91B FREJ
>0.00012	<i>n</i>	90	7	11.2	⁵⁹ BERGER	91B FREJ
>0.0005	<i>n</i>	90	0		LEARNED	79 RVUE

$\Delta B = 2$ dinucleon modes $\tau(pp \rightarrow \pi^+ \pi^+)$ τ_{64}

<i>LIMIT</i> (10^{30} years)	<i>CL%</i>	<i>EVTS</i>	<i>BKGD EST</i>
>0.7	90	4	2.34

<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
BERGER	91B FREJ	τ per iron nucleus

 $\tau(pn \rightarrow \pi^+ \pi^0)$ τ_{65}

<i>LIMIT</i> (10^{30} years)	<i>CL%</i>	<i>EVTS</i>	<i>BKGD EST</i>
>2.0	90	0	0.31

<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
BERGER	91B FREJ	τ per iron nucleus

 $\tau(nn \rightarrow \pi^+ \pi^-)$ τ_{66}

<i>LIMIT</i> (10^{30} years)	<i>CL%</i>	<i>EVTS</i>	<i>BKGD EST</i>
>0.7	90	4	2.18

<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
BERGER	91B FREJ	τ per iron nucleus

 $\tau(nn \rightarrow \pi^0 \pi^0)$ τ_{67}

<i>LIMIT</i> (10^{30} years)	<i>CL%</i>	<i>EVTS</i>	<i>BKGD EST</i>
>3.4	90	0	0.78

<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
BERGER	91B FREJ	τ per iron nucleus

 $\tau(pp \rightarrow e^+ e^+)$ τ_{68}

<i>LIMIT</i> (10^{30} years)	<i>CL%</i>	<i>EVTS</i>	<i>BKGD EST</i>
>5.8	90	0	<0.1

<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
BERGER	91B FREJ	τ per iron nucleus

 $\tau(pp \rightarrow e^+ \mu^+)$ τ_{69}

<i>LIMIT</i> (10^{30} years)	<i>CL%</i>	<i>EVTS</i>	<i>BKGD EST</i>
>3.6	90	0	<0.1

<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
BERGER	91B FREJ	τ per iron nucleus

 $\tau(pp \rightarrow \mu^+ \mu^+)$ τ_{70}

<i>LIMIT</i> (10^{30} years)	<i>CL%</i>	<i>EVTS</i>	<i>BKGD EST</i>
>1.7	90	0	0.62

<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
BERGER	91B FREJ	τ per iron nucleus

 $\tau(pn \rightarrow e^+ \bar{\nu})$ τ_{71}

<i>LIMIT</i> (10^{30} years)	<i>CL%</i>	<i>EVTS</i>	<i>BKGD EST</i>
>2.8	90	5	9.67

<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
BERGER	91B FREJ	τ per iron nucleus

 $\tau(pn \rightarrow \mu^+ \bar{\nu})$ τ_{72}

<i>LIMIT</i> (10^{30} years)	<i>CL%</i>	<i>EVTS</i>	<i>BKGD EST</i>
>1.6	90	4	4.37

<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
BERGER	91B FREJ	τ per iron nucleus

$\tau(nn \rightarrow \nu_e \bar{\nu}_e)$ **T73**

<i>LIMIT</i> (10^{-30} years)	<i>CL%</i>	<i>EVTS</i>	<i>BKGD EST</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
>0.000012	90			64 BERNABEI	00B DAMA	
>0.000012	90	5	9.7	BERGER	91B FREJ	τ per iron nucleus

64 BERNABEI 00B looks for the decay of a $^{127}_{54}\text{Xe}$ nucleus following the disappearance of an nn pair in the otherwise-stable $^{129}_{54}\text{Xe}$ nucleus. The limit here applies as well to $nn \rightarrow \nu_\mu \bar{\nu}_\mu$, $nn \rightarrow \nu_\tau \bar{\nu}_\tau$, or any “disappearance” mode.

 $\tau(nn \rightarrow \nu_\mu \bar{\nu}_\mu)$ **T74**

<i>LIMIT</i> (10^{-30} years)	<i>CL%</i>	<i>EVTS</i>	<i>BKGD EST</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
>0.000006	90	4	4.4	BERGER	91B FREJ	τ per iron nucleus

 $\tau(pp \rightarrow \text{neutrinos})$ **T75**

<i>LIMIT</i> (10^{-30} years)	<i>CL%</i>	<i>EVTS</i>	<i>BKGD EST</i>	<i>DOCUMENT ID</i>	<i>TECN</i>
>0.00000055	90			65 BERNABEI	00B DAMA

65 BERNABEI 00B looks for the decay of a $^{127}_{52}\text{Te}$ nucleus following the disappearance of a pp pair in the otherwise-stable $^{129}_{54}\text{Xe}$ nucleus. Note that the decay doesn't conserve charge as well as baryon number.

 \bar{p} PARTIAL MEAN LIVES

The “partial mean life” limits tabulated here are the limits on $\bar{\tau}/B_i$, where $\bar{\tau}$ is the total mean life for the antiproton and B_i is the branching fraction for the mode in question.

 $\tau(\bar{p} \rightarrow e^- \gamma)$ **T76**

<i>VALUE</i> (years)	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
> 7×10^5	90	GEER	00	APEX 8.9 GeV/c \bar{p} beam
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>1848	95	GEER	94	CALO 8.9 GeV/c \bar{p} beam

 $\tau(\bar{p} \rightarrow \mu^- \gamma)$ **T77**

<i>VALUE</i> (years)	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
> 5×10^4	90	GEER	00	APEX 8.9 GeV/c \bar{p} beam
• • • We do not use the following data for averages, fits, limits, etc. • • •				
> 5.0×10^4	90	HU	98B APEX	8.9 GeV/c \bar{p} beam

 $\tau(\bar{p} \rightarrow e^- \pi^0)$ **T78**

<i>VALUE</i> (years)	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
> 4×10^5	90	GEER	00	APEX 8.9 GeV/c \bar{p} beam
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>554	95	GEER	94	CALO 8.9 GeV/c \bar{p} beam

$\tau(\bar{p} \rightarrow \mu^- \pi^0)$ **τ_{79}**

<u>VALUE</u> (years)	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$>5 \times 10^4$	90	GEER	00	APEX 8.9 GeV/c \bar{p} beam
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$>4.8 \times 10^4$	90	HU	98B	APEX 8.9 GeV/c \bar{p} beam

 $\tau(\bar{p} \rightarrow e^- \eta)$ **τ_{80}**

<u>VALUE</u> (years)	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$> 2 \times 10^4$	90	GEER	00	APEX 8.9 GeV/c \bar{p} beam
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
>171	95	GEER	94	CALO 8.9 GeV/c \bar{p} beam

 $\tau(\bar{p} \rightarrow \mu^- \eta)$ **τ_{81}**

<u>VALUE</u> (years)	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$>8 \times 10^3$	90	GEER	00	APEX 8.9 GeV/c \bar{p} beam
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$>7.9 \times 10^3$	90	HU	98B	APEX 8.9 GeV/c \bar{p} beam

 $\tau(\bar{p} \rightarrow e^- K_S^0)$ **τ_{82}**

<u>VALUE</u> (years)	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
>900	90	GEER	00	APEX 8.9 GeV/c \bar{p} beam
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
>29	95	GEER	94	CALO 8.9 GeV/c \bar{p} beam

 $\tau(\bar{p} \rightarrow \mu^- K_S^0)$ **τ_{83}**

<u>VALUE</u> (years)	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$>4 \times 10^3$	90	GEER	00	APEX 8.9 GeV/c \bar{p} beam
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$>4.3 \times 10^3$	90	HU	98B	APEX 8.9 GeV/c \bar{p} beam

 $\tau(\bar{p} \rightarrow e^- K_L^0)$ **τ_{84}**

<u>VALUE</u> (years)	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$>9 \times 10^3$	90	GEER	00	APEX 8.9 GeV/c \bar{p} beam
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
>9	95	GEER	94	CALO 8.9 GeV/c \bar{p} beam

 $\tau(\bar{p} \rightarrow \mu^- K_L^0)$ **τ_{85}**

<u>VALUE</u> (years)	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$>7 \times 10^3$	90	GEER	00	APEX 8.9 GeV/c \bar{p} beam
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$>6.5 \times 10^3$	90	HU	98B	APEX 8.9 GeV/c \bar{p} beam

 $\tau(\bar{p} \rightarrow e^- \gamma\gamma)$ **τ_{86}**

<u>VALUE</u> (years)	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$>2 \times 10^4$	90	GEER	00	APEX 8.9 GeV/c \bar{p} beam

$\tau(\bar{p} \rightarrow \mu^- \gamma\gamma)$ **T87**

<u>VALUE (years)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$>2 \times 10^4$	90	GEER	00	APEX 8.9 GeV/c \bar{p} beam
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$>2.3 \times 10^4$	90	HU	98B	APEX 8.9 GeV/c \bar{p} beam

 $\tau(\bar{p} \rightarrow e^- \rho)$ **T88**

<u>VALUE (years)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
>200	90	66 GEER	00	APEX 8.9 GeV/c \bar{p} beam
66 This GEER 00 measurement has been withdrawn; see GEER 00C.				

 $\tau(\bar{p} \rightarrow e^- \omega)$ **T89**

<u>VALUE (years)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
>200	90	GEER	00	APEX 8.9 GeV/c \bar{p} beam

 $\tau(\bar{p} \rightarrow e^- K^*(892)^0)$ **T90**

<u>VALUE (years)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$>1 \times 10^3$	90	67 GEER	00	APEX 8.9 GeV/c \bar{p} beam
67 This GEER 00 measurement has been withdrawn; see GEER 00C.				

p REFERENCES

BARANOV	01	PPN 32 376 Translated from FECAY	P.S. Baranov <i>et al.</i> 32 699.	
BLANPIED	01	PR C64 025203	G. Blanpied <i>et al.</i>	(BNL LEGS Collab.)
ESCHRICH	01	PL B522 233	I. Eschrich <i>et al.</i>	(FNAL SELEX Collab.)
HORI	01	PRL 87 093401	M. Hori <i>et al.</i>	(CERN ASACUSA Collab.)
OLMOSDEL...	01	EPJ A10 207	V. Olmos de Leon <i>et al.</i>	(MAMI TAPS Collab.)
TRETYAK	01	PL B505 59	V.I. Tretyak, Yu.G. Zdesenko	(KIEV)
BERNABEI	00B	PL B493 12	R. Bernabei <i>et al.</i>	(Gran Sasso DAMA Collab.)
GEER	00	PRL 84 590	S. Geer <i>et al.</i>	(FNAL APEX Collab.)
Also	00B	PR D62 052004	S. Geer <i>et al.</i>	(FNAL APEX Collab.)
Also	00C	PR 85 3546 (erratum)	S. Geer <i>et al.</i>	(FNAL APEX Collab.)
GEER	00C	PRL 85 3546 (erratum)	S. Geer <i>et al.</i>	(FNAL APEX Collab.)
GEER	00D	APJ 532 648	S.H. Geer, D.C. Kennedy	
MELNIKOV	00	PRL 84 1673	K. Melnikov <i>et al.</i>	(SLAC, KARL)
ROSENFELDR...	00	PL B479 381	R. Rosenfelder	
SENGUPTA	00	PL B484 275	S. Sengupta	
WALL	00	PR D61 072004	D. Wall <i>et al.</i>	(Soudan-2 Collab.)
WALL	00B	PR D62 092003	D. Wall <i>et al.</i>	(Soudan-2 Collab.)
GABRIELSE	99	PRL 82 3198	G. Gabrielse <i>et al.</i>	
HAYATO	99	PRL 83 1529	Y. Hayato <i>et al.</i>	(Super-Kamiokande Collab.)
MCGREW	99	PR D59 052004	C. McGrew <i>et al.</i>	(IMB-3 Collab.)
MOHR	99	JPCRD 28 1713	P.J. Mohr, B.N. Taylor	(NIST)
Also	00	RMP 72 351	P.J. Mohr, B.N. Taylor	(NIST)
TORII	99	PR A59 223	H.A. Torii <i>et al.</i>	(CERN-PS-205 Collab.)
ALLISON	98	PL B427 217	W.W.M. Allison <i>et al.</i>	(Soudan-2 Collab.)
HU	98B	PR D58 111101	M. Hu <i>et al.</i>	(FNAL APEX Collab.)
SHIOZAWA	98	PRL 81 3319	M. Shiozawa <i>et al.</i>	(Super-Kamiokande Collab.)
GLICENSTEIN	97	PL B411 326	J.F. Glicenstein	(SACL)
MERGELL	96	NP A596 367	P. Mergell <i>et al.</i>	(MANZ, BONN)
GABRIELSE	95	PRL 74 3544	G. Gabrielse <i>et al.</i>	(HARV, MANZ, SEOUL)
MACGIBBON	95	PR C52 2097	B.E. MacGibbon <i>et al.</i>	(ILL, SASK, INRM)
GEER	94	PRL 72 1596	S. Geer <i>et al.</i>	(FNAL, UCLA, PSU)
WONG	94	IJMP E3 821	C.W. Wong	(UCLA)
HALLIN	93	PR C48 1497	E.L. Hallin <i>et al.</i>	(SASK, BOST, ILL)
SUZUKI	93B	PL B311 357	Y. Suzuki <i>et al.</i>	(KAMIOKANDE Collab.)

HUGHES	92	PRL 69 578	R.J. Hughes, B.I. Deutch	(LANL, AARH)
ZIEGER	92	PL B278 34	A. Zieger <i>et al.</i>	(MPCM)
Also	92B	PL B281 417 (erratum)	A. Zieger <i>et al.</i>	(MPCM)
BERGER	91	ZPHY C50 385	C. Berger <i>et al.</i>	(FREJUS Collab.)
BERGER	91B	PL B269 227	C. Berger <i>et al.</i>	(FREJUS Collab.)
FEDERSPIEL	91	PRL 67 1511	F.J. Federspiel <i>et al.</i>	(ILL)
MCCORD	91	NIM B56/57 496	M. McCord <i>et al.</i>	
BECKER-SZ...	90	PR D42 2974	R.A. Becker-Szendy <i>et al.</i>	(IMB-3 Collab.)
ERICSON	90	EPL 11 295	T.E.O. Ericson, A. Richter	(CERN, DARM)
GABRIELSE	90	PRL 65 1317	G. Gabrielse <i>et al.</i>	(HARV, MANZ, WASH+)
BERGER	89	NP B313 509	C. Berger <i>et al.</i>	(FREJUS Collab.)
CHO	89	PRL 63 2559	D. Cho, K. Sangster, E.A. Hinds	(YALE)
HIRATA	89C	PL B220 308	K.S. Hirata <i>et al.</i>	(Kamiokande Collab.)
PHILLIPS	89	PL B224 348	T.J. Phillips <i>et al.</i>	(HPW Collab.)
KREISSL	88	ZPHY C37 557	A. Kreissl <i>et al.</i>	(CERN PS176 Collab.)
SEIDEL	88	PRL 61 2522	S. Seidel <i>et al.</i>	(IMB Collab.)
BARTELT	87	PR D36 1990	J.E. Bartelt <i>et al.</i>	(Soudan Collab.)
Also	89	PR D40 1701 erratum	J.E. Bartelt <i>et al.</i>	(Soudan Collab.)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
HAINES	86	PRL 57 1986	T.J. Haines <i>et al.</i>	(IMB Collab.)
KAJITA	86	JPSJ 55 711	T. Kajita <i>et al.</i>	(Kamiokande Collab.)
ARISAKA	85	JPSJ 54 3213	K. Arisaka <i>et al.</i>	(Kamiokande Collab.)
BLEWITT	85	PRL 55 2114	G.B. Blewitt <i>et al.</i>	(IMB Collab.)
DZUBA	85	PL 154B 93	V.A. Dzuba, V.V. Flambaum, P.G. Silvestrov	(NOVO)
PARK	85	PRL 54 22	H.S. Park <i>et al.</i>	(IMB Collab.)
BATTISTONI	84	PL 133B 454	G. Battistoni <i>et al.</i>	(NUSEX Collab.)
MARINELLI	84	PL 137B 439	M. Marinelli, G. Morpurgo	(GENO)
WILKENING	84	PR A29 425	D.A. Wilkening, N.F. Ramsey, D.J. Larson	(HARV+)
BARTELT	83	PRL 50 651	J.E. Bartelt <i>et al.</i>	(MINN, ANL)
BATTISTONI	82	PL 118B 461	G. Battistoni <i>et al.</i>	(NUSEX Collab.)
KRISHNA...	82	PL 115B 349	M.R. Krishnaswamy <i>et al.</i>	(TATA, OSKC+)
ALEKSEEV	81	JETPL 33 651	E.N. Alekseev <i>et al.</i>	(PNPI)
		Translated from ZETFP 33 664.		
CHERRY	81	PRL 47 1507	M.L. Cherry <i>et al.</i>	(PENN, BNL)
COWSIK	80	PR D22 2204	R. Cowsik, V.S. Narasimham	(TATA)
SIMON	80	NP A333 381	G.G. Simon <i>et al.</i>	
BELL	79	PL 86B 215	M. Bell <i>et al.</i>	(CERN)
GOLDEN	79	PRL 43 1196	R.L. Golden <i>et al.</i>	(NASA, PSLL)
LEARNED	79	PRL 43 907	J.G. Learned, F. Reines, A. Soni	(UCI)
BREGMAN	78	PL 78B 174	M. Bregman <i>et al.</i>	(CERN)
ROBERTS	78	PR D17 358	B.L. Roberts	(WILL, RHEL)
EVANS	77	Science 197 989	J.C. Evans Jr., R.I. Steinberg	(BNL, PENN)
HU	75	NP A254 403	E. Hu <i>et al.</i>	(COLU, YALE)
BORKOWSKI	74	NP A222 269	F. Borkowski <i>et al.</i>	
COHEN	73	JPCRD 2 664	E.R. Cohen, B.N. Taylor	(RISC, NBS)
DYLLA	73	PR A7 1224	H.F. Dylla, J.G. King	(MIT)
AKIMOV	72	JETP 35 651	Yu.K. Akimov <i>et al.</i>	(YERE)
		Translated from ZETF 62 1231.		
DIX	70	Thesis Case	F.W. Dix	(CASE)
HARRISON	69	PRL 22 1263	G.E. Harrison, P.G.H. Sandars, S.J. Wright	(OXF)
GURR	67	PR 158 1321	H.S. Gurr <i>et al.</i>	(CASE, WITW)
FREREJACQ...	66	PR 141 1308	D. Frerejacque <i>et al.</i>	
HAND	63	RMP 35 335	L.N. hand <i>et al.</i>	
FLEROV	58	DOKL 3 79	G.N. Flerov <i>et al.</i>	(ASCI)