

B^\pm/B^0 ADMIXTURE

B DECAY MODES

The branching fraction measurements are for an admixture of B mesons at the $\Upsilon(4S)$. The values quoted assume that $B(\Upsilon(4S) \rightarrow B\bar{B}) = 100\%$.

For inclusive branching fractions, e.g., $B \rightarrow D^\pm$ anything, the treatment of multiple D 's in the final state must be defined. One possibility would be to count the number of events with one-or-more D 's and divide by the total number of B 's. Another possibility would be to count the total number of D 's and divide by the total number of B 's, which is the definition of average multiplicity. The two definitions are identical when only one of the specified particles is allowed in the final state. Even though the "one-or-more" definition seems sensible, for practical reasons inclusive branching fractions are almost always measured using the multiplicity definition. For heavy final state particles, authors call their results inclusive branching fractions while for light particles some authors call their results multiplicities. In the B sections, we list all results as inclusive branching fractions, adopting a multiplicity definition. This means that inclusive branching fractions can exceed 100% and that inclusive partial widths can exceed total widths, just as inclusive cross sections can exceed total cross sections.

\bar{B} modes are charge conjugates of the modes below. Reactions indicate the weak decay vertex and do not include mixing.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Semileptonic and leptonic modes		
Γ_1 $B \rightarrow e^+ \nu_e$ anything	[a]	(10.2 ± 0.4) %
Γ_2 $B \rightarrow \bar{p} e^+ \nu_e$ anything	< 1.6	$\times 10^{-3}$ CL=90%
Γ_3 $B \rightarrow \mu^+ \nu_\mu$ anything	[a]	
Γ_4 $B \rightarrow \ell^+ \nu_\ell$ anything	[a,b]	(10.38 ± 0.32) %
Γ_5 $B \rightarrow D^- \ell^+ \nu_\ell$ anything	[b]	(2.7 ± 0.8) %
Γ_6 $B \rightarrow \bar{D}^0 \ell^+ \nu_\ell$ anything	[b]	(7.0 ± 1.4) %
Γ_7 $B \rightarrow D^{*-} \ell^+ \nu_\ell$ anything		
Γ_8 $B \rightarrow D^{*0} \ell^+ \nu_\ell$ anything		
Γ_9 $B \rightarrow \bar{D}^{**} \ell^+ \nu_\ell$	[b,c]	(2.7 ± 0.7) %
Γ_{10} $B \rightarrow \bar{D}_1(2420) \ell^+ \nu_\ell$ anything		(7.4 ± 1.6) $\times 10^{-3}$
Γ_{11} $B \rightarrow D \pi \ell^+ \nu_\ell$ anything + $D^* \pi \ell^+ \nu_\ell$ anything		(2.6 ± 0.5) % S=1.5
Γ_{12} $B \rightarrow D \pi \ell^+ \nu_\ell$ anything		(1.5 ± 0.6) %
Γ_{13} $B \rightarrow D^* \pi \ell^+ \nu_\ell$ anything		(1.9 ± 0.4) %
Γ_{14} $B \rightarrow \bar{D}_2^*(2460) \ell^+ \nu_\ell$ anything	< 6.5	$\times 10^{-3}$ CL=95%

Γ_{15}	$B \rightarrow D^{*-} \pi^+ \ell^+ \nu_\ell$ anything	(1.00 ± 0.34) %		
Γ_{16}	$B \rightarrow D_s^- \ell^+ \nu_\ell$ anything	[b] < 9 $\times 10^{-3}$	CL=90%	
Γ_{17}	$B \rightarrow D_s^- \ell^+ \nu_\ell K^+$ anything	[b] < 6 $\times 10^{-3}$	CL=90%	
Γ_{18}	$B \rightarrow D_s^- \ell^+ \nu_\ell K^0$ anything	[b] < 9 $\times 10^{-3}$	CL=90%	
Γ_{19}	$B \rightarrow \ell^+ \nu_\ell$ noncharmed	[b]		
Γ_{20}	$B \rightarrow K^+ \ell^+ \nu_\ell$ anything	[b] (6.0 ± 0.6) %		
Γ_{21}	$B \rightarrow K^- \ell^+ \nu_\ell$ anything	[b] (10 ± 4) $\times 10^{-3}$		
Γ_{22}	$B \rightarrow K^0/\overline{K}^0 \ell^+ \nu_\ell$ anything	[b] (4.4 ± 0.5) %		

D , D^* , or D_s modes

Γ_{23}	$B \rightarrow D^\pm$ anything	(23.9 ± 1.9) %		
Γ_{24}	$B \rightarrow D^0/\overline{D}^0$ anything	(63.9 ± 3.0) %	S=1.1	
Γ_{25}	$B \rightarrow D^*(2010)^\pm$ anything	(22.5 ± 1.5) %		
Γ_{26}	$B \rightarrow D^*(2007)^0$ anything	(26.0 ± 2.7) %		
Γ_{27}	$B \rightarrow D_s^\pm$ anything	[d] (10.5 ± 2.6) %		
Γ_{28}	$B \rightarrow D_s^{*\pm}$ anything	(7.9 ± 2.2) %		
Γ_{29}	$B \rightarrow D_s^{*\pm} \overline{D}^{(*)}$	(4.2 ± 1.2) %		
Γ_{30}	$B \rightarrow D^{(*)} \overline{D}^{(*)} K^0 + D^{(*)} \overline{D}^{(*)} K^\pm$	[d,e] (7.1 ± 2.7) %		
Γ_{31}	$b \rightarrow c \overline{c} s$	(22 ± 4) %		
Γ_{32}	$B \rightarrow D_s^{(*)} \overline{D}^{(*)}$	[d,e] (4.9 ± 1.2) %		
Γ_{33}	$B \rightarrow D^* D^*(2010)^\pm$	[d] < 5.9 $\times 10^{-3}$	CL=90%	
Γ_{34}	$B \rightarrow D D^*(2010)^\pm + D^* D^\pm$	[d] < 5.5 $\times 10^{-3}$	CL=90%	
Γ_{35}	$B \rightarrow D D^\pm$	[d] < 3.1 $\times 10^{-3}$	CL=90%	
Γ_{36}	$B \rightarrow D_s^{(*)\pm} \overline{D}^{(*)} X(n\pi^\pm)$	[d,e] ($9 \pm 5/4$) %		
Γ_{37}	$B \rightarrow D^*(2010) \gamma$	< 1.1 $\times 10^{-3}$	CL=90%	
Γ_{38}	$B \rightarrow D_s^+ \pi^-, D_s^{*+} \pi^-, D_s^+ \rho^-, [d] < 5 D_s^{*+} \rho^-, D_s^+ \pi^0, D_s^{*+} \pi^0,$ $D_s^+ \eta, D_s^{*+} \eta, D_s^+ \rho^0,$ $D_s^{*+} \rho^0, D_s^+ \omega, D_s^{*+} \omega$	$\times 10^{-4}$	CL=90%	
Γ_{39}	$B \rightarrow D_{s1}(2536)^+$ anything	< 9.5 $\times 10^{-3}$	CL=90%	

Charmonium modes

Γ_{40}	$B \rightarrow J/\psi(1S)$ anything	(1.15 ± 0.06) %		
Γ_{41}	$B \rightarrow J/\psi(1S)$ (direct) anything	(8.0 ± 0.8) $\times 10^{-3}$		
Γ_{42}	$B \rightarrow \psi(2S)$ anything	(3.5 ± 0.5) $\times 10^{-3}$		
Γ_{43}	$B \rightarrow \chi_{c1}(1P)$ anything	(3.6 ± 0.5) $\times 10^{-3}$		
Γ_{44}	$B \rightarrow \chi_{c1}(1P)$ (direct) anything	(3.3 ± 0.5) $\times 10^{-3}$		
Γ_{45}	$B \rightarrow \chi_{c2}(1P)$ anything	(7 ± 4) $\times 10^{-4}$		
Γ_{46}	$B \rightarrow \eta_c(1S)$ anything	< 9 $\times 10^{-3}$	CL=90%	

K or K^* modes

Γ_{47}	$B \rightarrow K^\pm$ anything	[d]	(78.9 \pm 2.5) %
Γ_{48}	$B \rightarrow K^+$ anything		(66 \pm 5) %
Γ_{49}	$B \rightarrow K^-$ anything		(13 \pm 4) %
Γ_{50}	$B \rightarrow K^0/\bar{K}^0$ anything	[d]	(64 \pm 4) %
Γ_{51}	$B \rightarrow K^*(892)^\pm$ anything		(18 \pm 6) %
Γ_{52}	$B \rightarrow K^*(892)^0/\bar{K}^*(892)^0$ anything	[d]	(14.6 \pm 2.6) %
Γ_{53}	$B \rightarrow K^*(892)\gamma$		(4.2 \pm 0.6) $\times 10^{-5}$
Γ_{54}	$B \rightarrow K_1(1400)\gamma$	<	1.27 $\times 10^{-4}$
Γ_{55}	$B \rightarrow K_2^*(1430)\gamma$		(1.7 \pm 0.6) $\times 10^{-5}$
Γ_{56}	$B \rightarrow K_2(1770)\gamma$	<	1.2 $\times 10^{-3}$
Γ_{57}	$B \rightarrow K_3^*(1780)\gamma$	<	3.0 $\times 10^{-3}$
Γ_{58}	$B \rightarrow K_4^*(2045)\gamma$	<	1.0 $\times 10^{-3}$
Γ_{59}	$B \rightarrow \eta'(958)K$		(8.3 \pm 1.1) $\times 10^{-5}$
Γ_{60}	$B \rightarrow \eta'(958)K^*(892)$	<	2.2 $\times 10^{-5}$
Γ_{61}	$B \rightarrow \eta K$	<	5.2 $\times 10^{-6}$
Γ_{62}	$B \rightarrow \eta K^*(892)$		(1.8 \pm 0.5) $\times 10^{-5}$
Γ_{63}	$B \rightarrow \bar{b} \rightarrow \bar{s}\gamma$		(3.3 \pm 0.4) $\times 10^{-4}$
Γ_{64}	$B \rightarrow \bar{b} \rightarrow \bar{s}$ gluon	<	6.8 %
Γ_{65}	$B \rightarrow \eta$ anything	<	4.4 $\times 10^{-4}$
Γ_{66}	$B \rightarrow \eta' \text{ anything}$		(6.2 \pm 2.1) $\times 10^{-4}$

Light unflavored meson modes

Γ_{67}	$B \rightarrow \rho\gamma$	<	1.4 $\times 10^{-5}$	CL=90%
Γ_{68}	$B \rightarrow \pi^\pm$ anything	[d,f]	(358 \pm 7) %	
Γ_{69}	$B \rightarrow \pi^0$ anything		(235 \pm 11) %	
Γ_{70}	$B \rightarrow \eta$ anything		(17.6 \pm 1.6) %	
Γ_{71}	$B \rightarrow \rho^0$ anything		(21 \pm 5) %	
Γ_{72}	$B \rightarrow \omega$ anything	<	81 %	CL=90%
Γ_{73}	$B \rightarrow \phi$ anything		(3.5 \pm 0.7) %	S=1.8
Γ_{74}	$B \rightarrow \phi K^*(892)$	<	2.2 $\times 10^{-5}$	CL=90%

Baryon modes

Γ_{75}	$B \rightarrow \Lambda_c^\pm$ anything		(6.4 \pm 1.1) %
Γ_{76}	$B \rightarrow \Lambda_c^+$ anything		
Γ_{77}	$B \rightarrow \bar{\Lambda}_c^-$ anything		
Γ_{78}	$B \rightarrow \bar{\Lambda}_c^- e^+$ anything	<	3.2 $\times 10^{-3}$
Γ_{79}	$B \rightarrow \bar{\Lambda}_c^- p$ anything		(3.6 \pm 0.7) %
Γ_{80}	$B \rightarrow \bar{\Lambda}_c^- p e^+ \nu_e$	<	1.5 $\times 10^{-3}$
Γ_{81}	$B \rightarrow \bar{\Sigma}_c^{--}$ anything		(4.2 \pm 2.4) $\times 10^{-3}$
Γ_{82}	$B \rightarrow \bar{\Sigma}_c^-$ anything	<	9.6 $\times 10^{-3}$
Γ_{83}	$B \rightarrow \bar{\Sigma}_c^0$ anything		(4.6 \pm 2.4) $\times 10^{-3}$

Γ_{84}	$B \rightarrow \overline{\Sigma}_c^0 N$ ($N = p$ or n)	$< 1.5 \times 10^{-3}$	CL=90%
Γ_{85}	$B \rightarrow \Xi_c^0$ anything $\times B(\Xi_c^0 \rightarrow \Xi^- \pi^+)$	$(1.4 \pm 0.5) \times 10^{-4}$	
Γ_{86}	$B \rightarrow \Xi_c^+ \text{anything}$ $\times B(\Xi_c^+ \rightarrow \Xi^- \pi^+ \pi^+)$	$(4.5 \pm 1.3) \times 10^{-4}$	
Γ_{87}	$B \rightarrow p/\bar{p}$ anything	[d] $(8.0 \pm 0.4) \%$	
Γ_{88}	$B \rightarrow p/\bar{p}$ (direct) anything	[d] $(5.5 \pm 0.5) \%$	
Γ_{89}	$B \rightarrow \Lambda/\bar{\Lambda}$ anything	[d] $(4.0 \pm 0.5) \%$	
Γ_{90}	$B \rightarrow \Lambda$ anything		
Γ_{91}	$B \rightarrow \bar{\Lambda}$ anything		
Γ_{92}	$B \rightarrow \Xi^-/\Xi^+$ anything	[d] $(2.7 \pm 0.6) \times 10^{-3}$	
Γ_{93}	$B \rightarrow \text{baryons}$ anything	$(6.8 \pm 0.6) \%$	
Γ_{94}	$B \rightarrow p\bar{p}$ anything	$(2.47 \pm 0.23) \%$	
Γ_{95}	$B \rightarrow \Lambda\bar{p}/\bar{\Lambda}p$ anything	[d] $(2.5 \pm 0.4) \%$	
Γ_{96}	$B \rightarrow \Lambda\bar{\Lambda}$ anything	$< 5 \times 10^{-3}$	CL=90%

**Lepton Family number (*LF*) violating modes or
 $\Delta B = 1$ weak neutral current (*B1*) modes**

Γ_{97}	$B \rightarrow e^+ e^- s$	<i>B1</i>	$< 5.7 \times 10^{-5}$	CL=90%
Γ_{98}	$B \rightarrow \mu^+ \mu^- s$	<i>B1</i>	$< 5.8 \times 10^{-5}$	CL=90%
Γ_{99}	$B \rightarrow K e^+ e^-$	<i>B1</i>	$< 1.3 \times 10^{-6}$	CL=90%
Γ_{100}	$B \rightarrow K^*(892) e^+ e^-$	<i>B1</i>	$< 5.6 \times 10^{-6}$	CL=90%
Γ_{101}	$B \rightarrow K \mu^+ \mu^-$	<i>B1</i>	$(9.9 \pm 4.2) \times 10^{-7}$	
Γ_{102}	$B \rightarrow K^*(892) \mu^+ \mu^-$	<i>B1</i>	$< 3.1 \times 10^{-6}$	CL=90%
Γ_{103}	$B \rightarrow K \ell^+ \ell^-$	<i>B1</i>	$(7.5 \pm 2.6) \times 10^{-7}$	
Γ_{104}	$B \rightarrow K^*(892) \ell^+ \ell^-$	<i>B1</i>	$< 3.3 \times 10^{-6}$	CL=90%
Γ_{105}	$B \rightarrow e^\pm \mu^\mp s$	<i>LF</i>	[d] $< 2.2 \times 10^{-5}$	CL=90%

[a] These values are model dependent. See ‘Note on Semileptonic Decays’ in the B^+ Particle Listings.

[b] An ℓ indicates an e or a μ mode, not a sum over these modes.

[c] D^{**} stands for the sum of the $D(1^1P_1)$, $D(1^3P_0)$, $D(1^3P_1)$, $D(1^3P_2)$, $D(2^1S_0)$, and $D(2^1S_1)$ resonances.

[d] The value is for the sum of the charge states or particle/antiparticle states indicated.

[e] $D^{(*)}\overline{D}^{(*)}$ stands for the sum of $D^*\overline{D}^*$, $D^*\overline{D}$, $D\overline{D}^*$, and $D\overline{D}$.

[f] Inclusive branching fractions have a multiplicity definition and can be greater than 100%.

B^\pm/B^0 ADMIXTURE BRANCHING RATIOS

$\Gamma(\ell^+ \nu_\ell \text{anything})/\Gamma_{\text{total}}$

Γ_4/Γ

These branching fraction values are model dependent. See the note on "Semileptonic Decays of B Mesons at the beginning of the B^+ Particle Listings.

VALUE	DOCUMENT ID	TECN	COMMENT
0.1038 ± 0.0032 OUR AVERAGE			Includes data from the 2 datablocks that follow this one.

$0.108 \pm 0.002 \pm 0.0056$ ¹HENDERSON 92 CLEO $e^+ e^- \rightarrow \Upsilon(4S)$
¹HENDERSON 92 measurement employs e and μ . The systematic error contains 0.004 in quadrature from model dependence. The authors average a variation of the Isgur, Scora, Grinstein, and Wise model with that of the Altarelli-Cabibbo-Corbò-Maiani-Martinelli model for semileptonic decays to correct the acceptance.

$\Gamma(e^+ \nu_e \text{anything})/\Gamma_{\text{total}}$

Γ_1/Γ

These branching fraction values are model dependent. See the note on "Semileptonic Decays of B Mesons at the beginning of the B^+ Particle Listings.

VALUE	DOCUMENT ID	TECN	COMMENT
The data in this block is included in the average printed for a previous datablock.			

0.102 ± 0.004 OUR AVERAGE

$0.1049 \pm 0.0017 \pm 0.0043$ ²BARISH 96B CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$
 $0.097 \pm 0.005 \pm 0.004$ ³ALBRECHT 93H ARG $e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •
 $0.100 \pm 0.004 \pm 0.003$ ⁴YANAGISAWA 91 CSB2 $e^+ e^- \rightarrow \Upsilon(4S)$
 $0.103 \pm 0.006 \pm 0.002$ ⁵ALBRECHT 90H ARG $e^+ e^- \rightarrow \Upsilon(4S)$
 $0.117 \pm 0.004 \pm 0.010$ ⁶WACHS 89 CBAL Direct e at $\Upsilon(4S)$
 $0.120 \pm 0.007 \pm 0.005$ CHEN 84 CLEO Direct e at $\Upsilon(4S)$
 $0.132 \pm 0.008 \pm 0.014$ ⁷KLOPFEN... 83B CUSB Direct e at $\Upsilon(4S)$

²BARISH 96B analysis performed using tagged semileptonic decays of the B . This technique is almost model independent for the lepton branching ratio.

³ALBRECHT 93H analysis performed using tagged semileptonic decays of the B . This technique is almost model independent for the lepton branching ratio.

⁴YANAGISAWA 91 also measures an average semileptonic branching ratio at the $\Upsilon(5S)$ of 9.6–10.5% depending on assumptions about the relative production of different B meson species.

⁵ALBRECHT 90H uses the model of ALTARELLI 82 to correct over all lepton momenta. 0.099 ± 0.006 is obtained using ISGUR 89B.

⁶Using data above $p(e) = 2.4$ GeV, WACHS 89 determine $\sigma(B \rightarrow e\nu \text{up})/\sigma(B \rightarrow e\nu \text{charm}) < 0.065$ at 90% CL.

⁷Ratio $\sigma(b \rightarrow e\nu \text{up})/\sigma(b \rightarrow e\nu \text{charm}) < 0.055$ at CL = 90%.

$\Gamma(\mu^+ \nu_\mu \text{anything})/\Gamma_{\text{total}}$

Γ_3/Γ

These branching fraction values are model dependent. See the note on "Semileptonic Decays of B Mesons at the beginning of the B^+ Particle Listings.

VALUE	DOCUMENT ID	TECN	COMMENT
The data in this block is included in the average printed for a previous datablock.			

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.100 \pm 0.006 \pm 0.002$ ⁸ALBRECHT 90H ARG $e^+ e^- \rightarrow \Upsilon(4S)$
 $0.108 \pm 0.006 \pm 0.01$ CHEN 84 CLEO Direct μ at $\Upsilon(4S)$
 $0.112 \pm 0.009 \pm 0.01$ LEVMAN 84 CUSB Direct μ at $\Upsilon(4S)$

⁸ ALBRECHT 90H uses the model of ALTARELLI 82 to correct over all lepton momenta.
 0.097 ± 0.006 is obtained using ISGUR 89B.

$\Gamma(\bar{p}e^+\nu_e\text{anything})/\Gamma_{\text{total}}$		Γ_2/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0016	90	ALBRECHT	90H ARG	$e^+e^- \rightarrow \gamma(4S)$

$\Gamma(D^-\ell^+\nu_\ell\text{anything})/\Gamma(\ell^+\nu_\ell\text{anything})$		Γ_5/Γ_4	
VALUE	DOCUMENT ID	TECN	COMMENT
$0.26 \pm 0.07 \pm 0.04$	⁹ FULTON	91 CLEO	$e^+e^- \rightarrow \gamma(4S)$

⁹ FULTON 91 uses $B(D^+ \rightarrow K^-\pi^+\pi^+) = (9.1 \pm 1.3 \pm 0.4)\%$ as measured by MARK III.

$\Gamma(\bar{D}^0\ell^+\nu_\ell\text{anything})/\Gamma(\ell^+\nu_\ell\text{anything})$		Γ_6/Γ_4	
VALUE	DOCUMENT ID	TECN	COMMENT
$0.67 \pm 0.09 \pm 0.10$	¹⁰ FULTON	91 CLEO	$e^+e^- \rightarrow \gamma(4S)$

¹⁰ FULTON 91 uses $B(D^0 \rightarrow K^-\pi^+) = (4.2 \pm 0.4 \pm 0.4)\%$ as measured by MARK III.

$\Gamma(D^{*-}\ell^+\nu_\ell\text{anything})/\Gamma_{\text{total}}$		Γ_7/Γ	
VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.6 \pm 0.3 \pm 0.1	¹¹ BARISH	95 CLE2	$e^+e^- \rightarrow \gamma(4S)$

¹¹ BARISH 95 use $B(D^0 \rightarrow K^-\pi^+) = (3.91 \pm 0.08 \pm 0.17)\%$ and $B(D^{*+} \rightarrow D^0\pi^+) = (68.1 \pm 1.0 \pm 1.3)\%$.

$\Gamma(D^{*0}\ell^+\nu_\ell\text{anything})/\Gamma_{\text{total}}$		Γ_8/Γ	
VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.6 \pm 0.6 \pm 0.1	¹² BARISH	95 CLE2	$e^+e^- \rightarrow \gamma(4S)$

¹² BARISH 95 use $B(D^0 \rightarrow K^-\pi^+) = (3.91 \pm 0.08 \pm 0.17)\%$, $B(D^{*+} \rightarrow D^0\pi^+) = (68.1 \pm 1.0 \pm 1.3)\%$, $B(D^{*0} \rightarrow D^0\pi^0) = (63.6 \pm 2.3 \pm 3.3)\%$.

$\Gamma(\bar{D}^{**}\ell^+\nu_\ell)/\Gamma_{\text{total}}$		Γ_9/Γ			
VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$0.027 \pm 0.005 \pm 0.005$	63	¹³ ALBRECHT	93 ARG	$e^+e^- \rightarrow \gamma(4S)$	

$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

<0.028	95	¹⁴ BARISH	95 CLE2	$e^+e^- \rightarrow \gamma(4S)$
--------	----	----------------------	---------	---------------------------------

¹³ ALBRECHT 93 assumes the GISW model to correct for unseen modes. Using the BHKT model, the result becomes $0.023 \pm 0.006 \pm 0.004$. Assumes $B(D^{*+} \rightarrow D^0\pi^+) = 68.1\%$, $B(D^0 \rightarrow K^-\pi^+) = 3.65\%$, $B(D^0 \rightarrow K^-\pi^+\pi^-\pi^+) = 7.5\%$. We have taken their average e and μ value.

¹⁴ BARISH 95 use $B(D^0 \rightarrow K^-\pi^+) = (3.91 \pm 0.08 \pm 0.17)\%$, assume all nonresonant channels are zero, and use GISW model for relative abundances of D^{**} states.

$\Gamma(\overline{D}_1(2420)\ell^+\nu_\ell\text{anything})/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0074±0.0016	15 BUSKULIC	97B ALEP	$e^+e^- \rightarrow Z$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
seen	16 BUSKULIC	95B ALEP	Repl. by BUSKULIC 97B
15 BUSKULIC 97B assumes $B(D_1(2420) \rightarrow D^*\pi) = 1$, $B(D_1(2420) \rightarrow D^*\pi^\pm) = 2/3$, and $B(b \rightarrow B) = 0.378 \pm 0.022$.			
16 BUSKULIC 95B reports $f_B \times B(B \rightarrow \overline{D}_1(2420)^0\ell^+\nu_\ell\text{anything}) \times B(\overline{D}_1(2420)^0 \rightarrow \overline{D}^*(2010)^-\pi^+) = (2.04 \pm 0.58 \pm 0.34)10^{-3}$, where f_B is the production fraction for a single B charge state.			

$[\Gamma(D\pi\ell^+\nu_\ell\text{anything}) + \Gamma(D^*\pi\ell^+\nu_\ell\text{anything})]/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.026 ±0.005 OUR AVERAGE			Error includes scale factor of 1.5.
0.0340±0.0052±0.0032	17 ABREU	00R DLPH	$e^+e^- \rightarrow Z$
0.0226±0.0029±0.0033	18 BUSKULIC	97B ALEP	$e^+e^- \rightarrow Z$
17 Assumes no contribution from B_s and b baryons. Further assumes contributions from single pion ($D\pi$ and $D^*\pi$) states only, allowing isospin conservation to relate the relative π^0 and π^+ rates.			
18 BUSKULIC 97B assumes $B(b \rightarrow B) = 0.378 \pm 0.022$ and uses isospin invariance by assuming that all observed $D^0\pi^+$, $D^{*0}\pi^+$, $D^+\pi^-$, and $D^{*+}\pi^-$ are from D^{**} states. A correction has been applied to account for the production of B_s^0 and Λ_b^0 .			

$\Gamma(D\pi\ell^+\nu_\ell\text{anything})/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0154±0.0061	ABREU	00R DLPH	$e^+e^- \rightarrow Z$

$\Gamma(D^*\pi\ell^+\nu_\ell\text{anything})/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0186±0.0038	ABREU	00R DLPH	$e^+e^- \rightarrow Z$

$\Gamma(\overline{D}_2^*(2460)\ell^+\nu_\ell\text{anything})/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0065	95	19 BUSKULIC	97B ALEP	$e^+e^- \rightarrow Z$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
not seen		20 BUSKULIC	95B ALEP	$e^+e^- \rightarrow Z$
19 A revised number based on BUSKULIC 97B which assumes $B(D_2^*(2460) \rightarrow D^*\pi^\pm) = 0.20$ and $B(b \rightarrow B) = 0.378 \pm 0.022$.				
20 BUSKULIC 95B reports $f_B \times B(B \rightarrow \overline{D}_2^*(2460)^0\ell^+\nu_\ell\text{anything}) \times B(\overline{D}_2^*(2460)^0 \rightarrow \overline{D}^*(2010)^-\pi^+) \leq 0.81 \times 10^{-3}$ at CL=95%, where f_B is the production fraction for a single B charge state.				

$\Gamma(D^{*-}\pi^+\ell^+\nu_\ell\text{anything})/\Gamma_{\text{total}}$ Γ_{15}/Γ

Includes resonant and nonresonant contributions.

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
10.0±2.7±2.1	21 BUSKULIC	95B ALEP	$e^+e^- \rightarrow Z$
21 BUSKULIC 95B reports $f_B \times B(B \rightarrow \overline{D}^*(2010)^-\pi^+\ell^+\nu_\ell\text{anything}) = (3.7 \pm 1.0 \pm 0.7)10^{-3}$. Above value assumes $f_B = 0.37 \pm 0.03$.			

$\Gamma(D_s^- \ell^+ \nu_\ell \text{anything})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{16}/Γ
<0.009	90	22 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \gamma(4S)$	

22 ALBRECHT 93E reports < 0.012 for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$.

$\Gamma(D_s^- \ell^+ \nu_\ell K^+ \text{anything})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{17}/Γ
<0.006	90	23 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \gamma(4S)$	

23 ALBRECHT 93E reports < 0.008 for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$.

$\Gamma(D_s^- \ell^+ \nu_\ell K^0 \text{anything})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{18}/Γ
<0.009	90	24 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \gamma(4S)$	

24 ALBRECHT 93E reports < 0.012 for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$.

$\Gamma(\ell^+ \nu_\ell \text{noncharmed})/\Gamma(\ell^+ \nu_\ell \text{anything})$

Γ_{19}/Γ_4

ℓ denotes e or μ , not the sum. These experiments measure this ratio in very limited momentum intervals.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
			25 ALBRECHT	94C ARG	$e^+ e^- \rightarrow \gamma(4S)$
107			26 BARTEL	93B CLE2	$e^+ e^- \rightarrow \gamma(4S)$
77			27 ALBRECHT	91C ARG	$e^+ e^- \rightarrow \gamma(4S)$
76			28 FULTON	90 CLEO	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

	41	29 ALBRECHT	90 ARG	$e^+ e^- \rightarrow \gamma(4S)$
<0.04	90	30 BEHRENDS	87 CLEO	$e^+ e^- \rightarrow \gamma(4S)$
<0.04	90	CHEN	84 CLEO	Direct e at $\gamma(4S)$
<0.055	90	KLOPFEN...	83B CUSB	Direct e at $\gamma(4S)$

25 ALBRECHT 94C find $\Gamma(b \rightarrow c)/\Gamma(b \rightarrow \text{all}) = 0.99 \pm 0.02 \pm 0.04$.

26 BARTEL 93B (CLEO II) measures an excess of $107 \pm 15 \pm 11$ leptons in the lepton momentum interval $2.3\text{--}2.6 \text{ GeV}/c$ which is attributed to $b \rightarrow u\ell\nu_\ell$. This corresponds to a model-dependent partial branching ratio ΔB_{ub} between $(1.15 \pm 0.16 \pm 0.15) \times 10^{-4}$, as evaluated using the KS model (KOERNER 88), and $(1.54 \pm 0.22 \pm 0.20) \times 10^{-4}$ using the ACCMM model (ARTUSO 93). The corresponding values of $|V_{ub}|/|V_{cb}|$ are 0.056 ± 0.006 and 0.076 ± 0.008 , respectively.

27 ALBRECHT 91C result supersedes ALBRECHT 90. Two events are fully reconstructed providing evidence for the $b \rightarrow u$ transition. Using the model of ALTARELLI 82, they obtain $|V_{ub}/V_{cb}| = 0.11 \pm 0.012$ from 77 leptons in the $2.3\text{--}2.6 \text{ GeV}$ momentum range.

28 FULTON 90 observe 76 ± 20 excess e and μ (lepton) events in the momentum interval $p = 2.4\text{--}2.6 \text{ GeV}$ signaling the presence of the $b \rightarrow u$ transition. The average branching ratio, $(1.8 \pm 0.4 \pm 0.3) \times 10^{-4}$, corresponds to a model-dependent measurement of approximately $|V_{ub}/V_{cb}| = 0.1$ using $B(b \rightarrow c\ell\nu) = 10.2 \pm 0.2 \pm 0.7\%$.

29 ALBRECHT 90 observes 41 ± 10 excess e and μ (lepton) events in the momentum interval $p = 2.3\text{--}2.6 \text{ GeV}$ signaling the presence of the $b \rightarrow u$ transition. The events correspond to a model-dependent measurement of $|V_{ub}/V_{cb}| = 0.10 \pm 0.01$.

30 The quoted possible limits range from 0.018 to 0.04 for the ratio, depending on which model or momentum range is chosen. We select the most conservative limit they have calculated. This corresponds to a limit on $|V_{ub}|/|V_{cb}| < 0.20$. While the endpoint technique employed is more robust than their previous results in CHEN 84, these results do not provide a numerical improvement in the limit.

$\Gamma(K^+\ell^+\nu_\ell \text{anything})/\Gamma(\ell^+\nu_\ell \text{anything})$

ℓ denotes e or μ , not the sum.

Γ_{20}/Γ_4

VALUE	DOCUMENT ID	TECN	COMMENT
0.58 ±0.05 OUR AVERAGE			
0.594±0.021±0.056	ALBRECHT	94C ARG	$e^+e^- \rightarrow \gamma(4S)$
0.54 ±0.07 ±0.06	31 ALAM	87B CLEO	$e^+e^- \rightarrow \gamma(4S)$

31 ALAM 87B measurement relies on lepton-kaon correlations.

$\Gamma(K^-\ell^+\nu_\ell \text{anything})/\Gamma(\ell^+\nu_\ell \text{anything})$

ℓ denotes e or μ , not the sum.

Γ_{21}/Γ_4

VALUE	DOCUMENT ID	TECN	COMMENT
0.092±0.035 OUR AVERAGE			
0.086±0.011±0.044	ALBRECHT	94C ARG	$e^+e^- \rightarrow \gamma(4S)$
0.10 ±0.05 ±0.02	32 ALAM	87B CLEO	$e^+e^- \rightarrow \gamma(4S)$

32 ALAM 87B measurement relies on lepton-kaon correlations.

$\Gamma(K^0/\bar{K}^0 \ell^+\nu_\ell \text{anything})/\Gamma(\ell^+\nu_\ell \text{anything})$

Γ_{22}/Γ_4

ℓ denotes e or μ , not the sum. Sum over K^0 and \bar{K}^0 states.

VALUE	DOCUMENT ID	TECN	COMMENT
0.42 ±0.05 OUR AVERAGE			
0.452±0.038±0.056	33 ALBRECHT	94C ARG	$e^+e^- \rightarrow \gamma(4S)$
0.39 ±0.06 ±0.04	34 ALAM	87B CLEO	$e^+e^- \rightarrow \gamma(4S)$

33 ALBRECHT 94C assume a K^0/\bar{K}^0 multiplicity twice that of K_S^0 .

34 ALAM 87B measurement relies on lepton-kaon correlations.

$\langle n_c \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
1.10±0.05			
35 GIBBONS	97B CLE2	$e^+e^- \rightarrow \gamma(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •			

35 GIBBONS 97B from charm counting using $B(D_s^+ \rightarrow \phi\pi) = 0.036 \pm 0.009$ and $B(\Lambda_c^+ \rightarrow pK^-\pi^+) = 0.044 \pm 0.006$.

36 From the difference between K^- and K^+ widths. ALAM 87B measurement relies on lepton-kaon correlations. It does not consider the possibility of $B\bar{B}$ mixing. We have thus removed it from the average.

$\Gamma(D^\pm \text{anything})/\Gamma_{\text{total}}$

Γ_{23}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.239±0.019 OUR AVERAGE				
0.238±0.013±0.015	37 GIBBONS	97B CLE2	$e^+e^- \rightarrow \gamma(4S)$	
0.25 ±0.04 ±0.02	38 BORTOLETTO92	CLEO	$e^+e^- \rightarrow \gamma(4S)$	
0.23 ±0.05 ±0.01	39 ALBRECHT	91H ARG	$e^+e^- \rightarrow \gamma(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.21 ±0.05 ±0.01	20k	40 BORTOLETTO87	CLEO	Sup. by BORTOLETTO 92

- ³⁷ GIBBONS 97B reports $[B(B \rightarrow D^\pm \text{anything}) \times B(D^+ \rightarrow K^- \pi^+ \pi^+)] = 0.0216 \pm 0.0008 \pm 0.00082$. We divide by our best value $B(D^+ \rightarrow K^- \pi^+ \pi^+) = (9.1 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ³⁸ BORTOLETTO 92 reports $[B(B \rightarrow D^\pm \text{anything}) \times B(D^+ \rightarrow K^- \pi^+ \pi^+)] = 0.0226 \pm 0.0030 \pm 0.0018$. We divide by our best value $B(D^+ \rightarrow K^- \pi^+ \pi^+) = (9.1 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ³⁹ ALBRECHT 91H reports $[B(B \rightarrow D^\pm \text{anything}) \times B(D^+ \rightarrow K^- \pi^+ \pi^+)] = 0.0209 \pm 0.0027 \pm 0.0040$. We divide by our best value $B(D^+ \rightarrow K^- \pi^+ \pi^+) = (9.1 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ⁴⁰ BORTOLETTO 87 reports $[B(B \rightarrow D^\pm \text{anything}) \times B(D^+ \rightarrow K^- \pi^+ \pi^+)] = 0.019 \pm 0.004 \pm 0.002$. We divide by our best value $B(D^+ \rightarrow K^- \pi^+ \pi^+) = (9.1 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D^0/\bar{D}^0 \text{anything})/\Gamma_{\text{total}}$

Γ_{24}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.639 ± 0.030 OUR AVERAGE				Error includes scale factor of 1.1.
0.660 ± 0.025	$+0.016$	41 GIBBONS	97B CLE2	$e^+ e^- \rightarrow \gamma(4S)$
-0.015				
0.61 ± 0.05	± 0.01	42 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
0.51 ± 0.08	± 0.01	43 ALBRECHT	91H ARG	$e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
0.55 ± 0.07	± 0.01	21k	44 BORTOLETTO87	CLEO $e^+ e^- \rightarrow \gamma(4S)$
0.63 ± 0.19	$+0.02$		45 GREEN	83 CLEO Repl. by BORTOLETTO 87
-0.01				
41 GIBBONS 97B reports $[B(B \rightarrow D^0/\bar{D}^0 \text{anything}) \times B(D^0 \rightarrow K^- \pi^+)] = 0.0251 \pm 0.0006 \pm 0.00075$. We divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.80 \pm 0.09) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
42 BORTOLETTO 92 reports $[B(B \rightarrow D^0/\bar{D}^0 \text{anything}) \times B(D^0 \rightarrow K^- \pi^+)] = 0.0233 \pm 0.0012 \pm 0.0014$. We divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.80 \pm 0.09) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
43 ALBRECHT 91H reports $[B(B \rightarrow D^0/\bar{D}^0 \text{anything}) \times B(D^0 \rightarrow K^- \pi^+)] = 0.0194 \pm 0.0015 \pm 0.0025$. We divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.80 \pm 0.09) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
44 BORTOLETTO 87 reports $[B(B \rightarrow D^0/\bar{D}^0 \text{anything}) \times B(D^0 \rightarrow K^- \pi^+)] = 0.0210 \pm 0.0015 \pm 0.0021$. We divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.80 \pm 0.09) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
45 GREEN 83 reports $[B(B \rightarrow D^0/\bar{D}^0 \text{anything}) \times B(D^0 \rightarrow K^- \pi^+)] = 0.024 \pm 0.006 \pm 0.004$. We divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.80 \pm 0.09) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(D^*(2010)^{\pm} \text{anything})/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.225±0.015 OUR AVERAGE				
0.247±0.019±0.01		46 GIBBONS	97B CLE2	$e^+ e^- \rightarrow \gamma(4S)$
0.205±0.019±0.007		47 ALBRECHT	96D ARG	$e^+ e^- \rightarrow \gamma(4S)$
0.230±0.028±0.009		48 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.283±0.053±0.002		49 ALBRECHT	91H ARG	Sup. by ALBRECHT 96D
0.22 ± 0.04 ± 0.07 - 0.04	5200	50 BORTOLETTO87	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
0.27 ± 0.06 ± 0.08 - 0.06	510	51 CSORNA	85 CLEO	Repl. by BORTOLETTO 87

46 GIBBONS 97B reports $B(B \rightarrow D^*(2010)^+ \text{anything}) = 0.239 \pm 0.015 \pm 0.014 \pm 0.009$ using CLEO measured D and D^* branching fractions. We rescale to our PDG 96 values of D and D^* branching ratios. Our first error is their experiment's error and our second error is the systematic error from using our best value.

47 ALBRECHT 96D reports $B(B \rightarrow D^*(2010)^+ \text{anything}) = 0.196 \pm 0.019$ using CLEO measured $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.681 \pm 0.01 \pm 0.013$, $B(D^0 \rightarrow K^- \pi^+) = 0.0401 \pm 0.0014$, $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-) = 0.081 \pm 0.005$. We rescale to our PDG 96 values of D and D^* branching ratios. Our first error is their experiment's error and our second error is the systematic error from using our best value.

48 BORTOLETTO 92 reports $B(B \rightarrow D^*(2010)^+ \text{anything}) = 0.25 \pm 0.03 \pm 0.04$ using MARK II $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.57 \pm 0.06$ and $B(D^0 \rightarrow K^- \pi^+) = 0.042 \pm 0.008$. We rescale to our PDG 96 values of D and D^* branching ratios. Our first error is their experiment's error and our second error is the systematic error from using our best value.

49 ALBRECHT 91H reports $0.348 \pm 0.060 \pm 0.035$ for $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.55 \pm 0.04$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Uses the PDG 90 $B(D^0 \rightarrow K^- \pi^+) = 0.0371 \pm 0.0025$.

50 BORTOLETTO 87 uses old MARK III (BALTRUSAITIS 86E) branching ratios $B(D^0 \rightarrow K^- \pi^+) = 0.056 \pm 0.004 \pm 0.003$ and also assumes $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.60^{+0.08}_{-0.15}$. The product branching ratio for $B(B \rightarrow D^*(2010)^+)$ $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ is $0.13 \pm 0.02 \pm 0.012$. Superseded by BORTOLETTO 92.

51 $V-A$ momentum spectrum used to extrapolate below $p = 1$ GeV. We correct the value assuming $B(D^0 \rightarrow K^- \pi^+) = 0.042 \pm 0.006$ and $B(D^{*+} \rightarrow D^0 \pi^+) = 0.6^{+0.08}_{-0.15}$. The product branching fraction is $B(B \rightarrow D^{*+} X) \cdot B(D^{*+} \rightarrow \pi^+ D^0) \cdot B(D^0 \rightarrow K^- \pi^+) = (68 \pm 15 \pm 9) \times 10^{-4}$.

 $\Gamma(D^*(2007)^0 \text{anything})/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.260±0.023±0.015	52 GIBBONS	97B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

52 GIBBONS 97B reports $B(B \rightarrow D^*(2007)^0 \text{anything}) = 0.247 \pm 0.012 \pm 0.018 \pm 0.018$ using CLEO measured D and D^* branching fractions. We rescale to our PDG 96 values of D and D^* branching ratios. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_s^\pm \text{anything})/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{27}/Γ
0.105±0.026 OUR AVERAGE					
0.109±0.006	+0.026 -0.027	53 AUBERT	02G BABR	$e^+ e^- \rightarrow \gamma(4S)$	
0.117±0.009	+0.028 -0.029	54 GIBAUT	96 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
0.081±0.014	+0.019 -0.020	55 ALBRECHT	92G ARG	$e^+ e^- \rightarrow \gamma(4S)$	
0.085±0.013	+0.020 -0.021	257 56 BORTOLETTO90	CLEO	$e^+ e^- \rightarrow \gamma(4S)$	
0.105±0.028	+0.025 -0.026	57 HAAS	86 CLEO	$e^+ e^- \rightarrow \gamma(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.116±0.030±0.028		58 ALBRECHT	87H ARG	$e^+ e^- \rightarrow \gamma(4S)$	
53 AUBERT 02G reports $[B(B \rightarrow D_s^\pm \text{anything}) \times B(D_s^\pm \rightarrow \phi\pi^+)] = 0.00393 \pm 0.00007 \pm 0.00021$. We divide by our best value $B(D_s^\pm \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
54 GIBAUT 96 reports $0.1211 \pm 0.0039 \pm 0.0088$ for $B(D_s^\pm \rightarrow \phi\pi^+) = 0.035$. We rescale to our best value $B(D_s^\pm \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
55 ALBRECHT 92G reports $[B(B \rightarrow D_s^\pm \text{anything}) \times B(D_s^\pm \rightarrow \phi\pi^+)] = 0.00292 \pm 0.00039 \pm 0.00031$. We divide by our best value $B(D_s^\pm \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
56 BORTOLETTO 90 reports $[B(B \rightarrow D_s^\pm \text{anything}) \times B(D_s^\pm \rightarrow \phi\pi^+)] = 0.00306 \pm 0.00047$. We divide by our best value $B(D_s^\pm \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
57 HAAS 86 reports $[B(B \rightarrow D_s^\pm \text{anything}) \times B(D_s^\pm \rightarrow \phi\pi^+)] = 0.0038 \pm 0.0010$. We divide by our best value $B(D_s^\pm \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. $64 \pm 22\%$ decays are 2-body.					
58 ALBRECHT 87H reports $[B(B \rightarrow D_s^\pm \text{anything}) \times B(D_s^\pm \rightarrow \phi\pi^+)] = 0.0042 \pm 0.0009 \pm 0.0006$. We divide by our best value $B(D_s^\pm \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. $46 \pm 16\%$ of $B \rightarrow D_s X$ decays are 2-body. Superseded by ALBRECHT 92G.					

 $\Gamma(D_s^{*\pm} \text{anything})/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{28}/Γ
0.079±0.011±0.019	59 AUBERT	02G BABR	$e^+ e^- \rightarrow \gamma(4S)$	
59 AUBERT 02G reports $[B(B \rightarrow D_s^{*\pm} \text{anything}) \times B(D_s^{*\pm} \rightarrow \phi\pi^+)] = 0.00284 \pm 0.00029 \pm 0.00025$. We divide by our best value $B(D_s^{*\pm} \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(D_s^{*\pm} \bar{D}^{(*)})/\Gamma(D_s^{*\pm} \text{anything})$
 Sum over modes
 Γ_{29}/Γ_{28}

VALUE	DOCUMENT ID	TECN	COMMENT
0.533±0.037±0.037	AUBERT	02G BABR	$e^+ e^- \rightarrow \gamma(4S)$

 $[\Gamma(D^{(*)} \bar{D}^{(*)} K^0) + \Gamma(D^{(*)} \bar{D}^{(*)} K^\pm)]/\Gamma_{\text{total}}$
 Γ_{30}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.071^{+0.025}_{-0.015}^{+0.010}_{-0.009}	60 BARATE	98Q ALEP	$e^+ e^- \rightarrow Z$

60 The systematic error includes the uncertainties due to the charm branching ratios.

 $\Gamma(c\bar{c}s)/\Gamma_{\text{total}}$
 Γ_{31}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.219±0.037	61 COAN	98 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

61 COAN 98 uses D - ℓ correlation.

 $\Gamma(D_s^{(*)} \bar{D}^{(*)})/\Gamma(D_s^\pm \text{anything})$
 Sum over modes.
 Γ_{32}/Γ_{27}

VALUE	DOCUMENT ID	TECN	COMMENT
0.469±0.017 OUR AVERAGE			

$0.464\pm 0.013\pm 0.015$

DOCUMENT ID	TECN	COMMENT
AUBERT	02G BABR	$e^+ e^- \rightarrow \gamma(4S)$

$0.56\begin{array}{l} +0.21 \\ -0.15 \end{array}\begin{array}{l} +0.09 \\ -0.08 \end{array}$

DOCUMENT ID	TECN	COMMENT
62 BARATE	98Q ALEP	$e^+ e^- \rightarrow Z$

$0.457\pm 0.019\pm 0.037$

DOCUMENT ID	TECN	COMMENT
GIBAUT	96 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

$0.58\pm 0.07\pm 0.09$

DOCUMENT ID	TECN	COMMENT
ALBRECHT	92G ARG	$e^+ e^- \rightarrow \gamma(4S)$

0.56 ± 0.10

DOCUMENT ID	TECN	COMMENT
BORTOLETTO90	CLEO	$e^+ e^- \rightarrow \gamma(4S)$

62 BARATE 98Q measures $B(B \rightarrow D_s^{(*)} \bar{D}^{(*)}) = 0.056^{+0.021}_{-0.015}{}^{+0.009}_{-0.008}{}^{+0.019}_{-0.011}$, where the third error results from the uncertainty on the different D branching ratios and is dominated by the uncertainty on $B(D_s^{+} \rightarrow \phi\pi^{+})$. We divide $B(B \rightarrow D_s^{(*)} \bar{D}^{(*)})$ by our best value of $B(B \rightarrow D_s \text{anything}) = 0.1 \pm 0.025$.

 $\Gamma(D^* D^*(2010)^\pm)/\Gamma_{\text{total}}$
 Γ_{33}/Γ

VALUE	CL%
$<5.9 \times 10^{-3}$	90

DOCUMENT ID	TECN	COMMENT
BARATE	98Q ALEP	$e^+ e^- \rightarrow Z$

 $[\Gamma(DD^*(2010)^\pm) + \Gamma(D^* D^\pm)]/\Gamma_{\text{total}}$
 Γ_{34}/Γ

VALUE	CL%
$<5.5 \times 10^{-3}$	90

DOCUMENT ID	TECN	COMMENT
BARATE	98Q ALEP	$e^+ e^- \rightarrow Z$

 $\Gamma(DD^\pm)/\Gamma_{\text{total}}$
 Γ_{35}/Γ

VALUE	CL%
$<3.1 \times 10^{-3}$	90

DOCUMENT ID	TECN	COMMENT
BARATE	98Q ALEP	$e^+ e^- \rightarrow Z$

 $\Gamma(D_s^{(*)\pm} \bar{D}^{(*)} X(n\pi^\pm))/\Gamma_{\text{total}}$
 Γ_{36}/Γ

VALUE	CL%
$0.094\begin{array}{l} +0.040 \\ -0.031 \end{array}\begin{array}{l} +0.034 \\ -0.024 \end{array}$	63

DOCUMENT ID	TECN	COMMENT
BARATE	98Q ALEP	$e^+ e^- \rightarrow Z$

63 The systematic error includes the uncertainties due to the charm branching ratios.

$\Gamma(D^*(2010)\gamma)/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.1 \times 10^{-3}$	90	64 LESIAK	92 CBAL	$e^+ e^- \rightarrow \gamma(4S)$

⁶⁴ LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s -quark hadronization.

$\Gamma(D_s^+\pi^-, D_s^{*+}\pi^-, D_s^+\rho^-, D_s^{*+}\rho^-, D_s^+\pi^0, D_s^{*+}\pi^0, D_s^+\eta, D_s^{*+}\eta, D_s^+\rho^0, D_s^{*+}\rho^0, D_s^+\omega, D_s^{*+}\omega)/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0005	90	65 ALEXANDER	93B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

⁶⁵ ALEXANDER 93B reports $< 4.8 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$. This branching ratio limit provides a model-dependent upper limit $|V_{ub}|/|V_{cb}| < 0.16$ at CL=90%.

$\Gamma(D_{s1}(2536)^+ \text{anything})/\Gamma_{\text{total}}$ Γ_{39}/Γ

$D_{s1}(2536)^+$ is the narrow P -wave D_s^+ meson with $J^P = 1^+$.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0095	90	66 BISHAI	98 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

⁶⁶ Assuming factorization, the decay constant $f_{D_{s1}^+}$ is at least a factor of 2.5 times smaller than $f_{D_s^+}$.

$\Gamma(J/\psi(1S)\text{anything})/\Gamma_{\text{total}}$ Γ_{40}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.15 ± 0.06 OUR AVERAGE				
1.13 $\pm 0.06 \pm 0.02$	1489	67 BAlest	95B CLE2	$e^+ e^- \rightarrow \gamma(4S)$
1.30 $\pm 0.45 \pm 0.02$	27	68 MASCHMANN	90 CBAL	$e^+ e^- \rightarrow \gamma(4S)$
1.24 $\pm 0.27 \pm 0.02$	120	69 ALBRECHT	87D ARG	$e^+ e^- \rightarrow \gamma(4S)$
1.37 $\pm 0.25 \pm 0.02$	52	70 ALAM	86 CLEO	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.4 $\begin{array}{l} +0.6 \\ -0.5 \end{array}$	7	71 ALBRECHT	85H ARG	$e^+ e^- \rightarrow \gamma(4S)$
1.1 $\pm 0.21 \pm 0.23$	46	72 HAAS	85 CLEO	Repl. by ALAM 86

⁶⁷ BAlest 95B reports $1.12 \pm 0.04 \pm 0.06$ for $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0599 \pm 0.0025$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.93 \pm 0.10) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.. They measure $J/\psi(1S) \rightarrow e^+ e^-$ and $\mu^+ \mu^-$ and use PDG 1994 values for the branching fractions. The rescaling is the same for either mode so we use $e^+ e^-$.

⁶⁸ MASCHMANN 90 reports $1.12 \pm 0.33 \pm 0.25$ for $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.93 \pm 0.10) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁶⁹ ALBRECHT 87D reports $1.07 \pm 0.16 \pm 0.22$ for $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.93 \pm 0.10) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using

- our best value. ALBRECHT 87D find the branching ratio for J/ψ not from $\psi(2S)$ to be 0.0081 ± 0.0023 .
- 70 ALAM 86 reports $1.09 \pm 0.16 \pm 0.21$ for $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.074 \pm 0.012$. We rescale to our best value $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = (5.88 \pm 0.10) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- 71 Statistical and systematic errors were added in quadrature. ALBRECHT 85H also report a CL = 90% limit of 0.007 for $B \rightarrow J/\psi(1S) + X$ where $m_X < 1$ GeV.
- 72 Dimuon and dielectron events used.

$\Gamma(J/\psi(1S)(\text{direct}) \text{ anything})/\Gamma_{\text{total}}$ Γ_{41}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0080±0.0008	73 BAEST	95B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

73 BAEST 95B assume PDG 1994 values for sub mode branching ratios. $J/\psi(1S)$ mesons are reconstructed in $J/\psi(1S) \rightarrow e^+ e^-$ and $J/\psi(1S) \rightarrow \mu^+ \mu^-$. The $B \rightarrow J/\psi(1S)X$ branching ratio contains $J/\psi(1S)$ mesons directly from B decays and also from feeddown through $\psi(2S) \rightarrow J/\psi(1S)$, $\chi_{c1}(1P) \rightarrow J/\psi(1S)$, or $\chi_{c2}(1P) \rightarrow J/\psi(1S)$. Using the measured inclusive rates, BAEST 95B corrects for the feeddown and finds the $B \rightarrow J/\psi(1S)(\text{direct}) X$ branching ratio.

$\Gamma(\psi(2S) \text{ anything})/\Gamma_{\text{total}}$ Γ_{42}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0035±0.0005 OUR AVERAGE				
0.0034±0.0004±0.0003	240	74 BAEST	95B CLE2	$e^+ e^- \rightarrow \gamma(4S)$
0.0046±0.0017±0.0011	8	ALBRECHT	87D ARG	$e^+ e^- \rightarrow \gamma(4S)$

74 BAEST 95B assume PDG 1994 values for sub mode branching ratios. They find $B(B \rightarrow \psi(2S)X, \psi(2S) \rightarrow \ell^+ \ell^-) = 0.30 \pm 0.05 \pm 0.04$ and $B(B \rightarrow \psi(2S)X, \psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = 0.37 \pm 0.05 \pm 0.05$. Weighted average is quoted for $B(B \rightarrow \psi(2S)X)$.

$\Gamma(\chi_{c1}(1P) \text{ anything})/\Gamma_{\text{total}}$ Γ_{43}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0036±0.0004±0.0004		75 CHEN	01 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0040±0.0006±0.0004	112	76 BAEST	95B CLE2	Repl. by CHEN 01
0.0105±0.0035±0.0025		77 ALBRECHT	92E ARG	$e^+ e^- \rightarrow \gamma(4S)$

75 CHEN 01 reports $0.00414 \pm 0.00031 \pm 0.00040$ for $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$. We rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (31.6 \pm 3.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

76 BAEST 95B assume $B(\chi_{c1}(1P) \rightarrow J/\psi(1S)\gamma) = (27.3 \pm 1.6) \times 10^{-2}$, the PDG 1994 value. Fit to ψ -photon invariant mass distribution allows for a $\chi_{c1}(1P)$ and a $\chi_{c2}(1P)$ component.

77 ALBRECHT 92E assumes no $\chi_{c2}(1P)$ production.

$\Gamma(\chi_{c1}(1P)(\text{direct}) \text{ anything})/\Gamma_{\text{total}}$ Γ_{44}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0033±0.0004±0.0003	78 CHEN	01 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.0037±0.0007	79 BAEST	95B CLE2	Repl. by CHEN 01

⁷⁸ CHEN 01 reports $0.00383 \pm 0.00031 \pm 0.00040$ for $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$. We rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (31.6 \pm 3.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

⁷⁹ BALEST 95B assume PDG 1994 values. $J/\psi(1S)$ mesons are reconstructed in the $e^+ e^-$ and $\mu^+ \mu^-$ modes. The $B \rightarrow \chi_{c1}(1P)X$ branching ratio contains $\chi_{c1}(1P)$ mesons directly from B decays and also from feeddown through $\psi(2S) \rightarrow \chi_{c1}(1P)\gamma$. Using the measured inclusive rates, BALEST 95B corrects for the feeddown and finds the $B \rightarrow \chi_{c1}(1P)$ (direct) X branching ratio.

$\Gamma(\chi_{c2}(1P)\text{anything})/\Gamma_{\text{total}}$

Γ_{45}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$7.1 \pm 3.6 \pm 0.8$		80	CHEN	01 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<38 90 35 81 BALEST 95B CLE2 Repl. by CHEN 01

⁸⁰ CHEN 01 reports $9.8 \pm 4.8 \pm 1.5$ for $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$. We rescale to our best value $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (18.7 \pm 2.0) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

⁸¹ BALEST 95B assume $B(\chi_{c2}(1P) \rightarrow J/\psi(1S)\gamma) = (13.5 \pm 1.1) \times 10^{-2}$, the PDG 1994 value. $J/\psi(1S)$ mesons are reconstructed in the $e^+ e^-$ and $\mu^+ \mu^-$ modes, and PDG 1994 branching fractions are used. If interpreted as signal, the 35 ± 13 events correspond to $B(B \rightarrow \chi_{c2}(1P)X) = (0.25 \pm 0.10 \pm 0.03) \times 10^{-2}$.

$\Gamma(\eta_c(1S)\text{anything})/\Gamma_{\text{total}}$

Γ_{46}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.009	90	82 BALEST	95B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

⁸² BALEST 95B assume PDG 1994 values for sub mode branching ratios. $J/\psi(1S)$ mesons are reconstructed in $J/\psi(1S) \rightarrow e^+ e^-$ and $J/\psi(1S) \rightarrow \mu^+ \mu^-$. Search region $2960 < m_{\eta_c(1S)} < 3010$ MeV/ c^2 .

$\Gamma(K^\pm\text{anything})/\Gamma_{\text{total}}$

Γ_{47}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.789 ± 0.025 OUR AVERAGE			
0.82 $\pm 0.01 \pm 0.05$	ALBRECHT 94C ARG	$e^+ e^- \rightarrow \Upsilon(4S)$	
$0.775 \pm 0.015 \pm 0.025$	83 ALBRECHT 93I ARG	$e^+ e^- \rightarrow \Upsilon(4S)$	
0.85 $\pm 0.07 \pm 0.09$	ALAM 87B CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen 84 BRODY 82 CLEO $e^+ e^- \rightarrow \Upsilon(4S)$

seen 85 GIANNINI 82 CUSB $e^+ e^- \rightarrow \Upsilon(4S)$

⁸³ ALBRECHT 93I value is not independent of the sum of $B \rightarrow K^+$ anything and $B \rightarrow K^-$ anything ALBRECHT 94C values.

⁸⁴ Assuming $\Upsilon(4S) \rightarrow B\bar{B}$, a total of $3.38 \pm 0.34 \pm 0.68$ kaons per $\Upsilon(4S)$ decay is found (the second error is systematic). In the context of the standard B -decay model, this leads to a value for $(b\text{-quark} \rightarrow c\text{-quark})/(b\text{-quark} \rightarrow \text{all})$ of $1.09 \pm 0.33 \pm 0.13$.

⁸⁵ GIANNINI 82 at CESR-CUSB observed 1.58 ± 0.35 K^0 per hadronic event much higher than 0.82 ± 0.10 below threshold. Consistent with predominant $b \rightarrow cX$ decay.

$\Gamma(K^+ \text{anything})/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.66 ± 0.05	86 ALBRECHT	94C ARG	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.620 ± 0.013 ± 0.038	87 ALBRECHT	94C ARG	$e^+ e^- \rightarrow \gamma(4S)$
0.66 ± 0.05 ± 0.07	87 ALAM	87B CLEO	$e^+ e^- \rightarrow \gamma(4S)$
86 Measurement relies on lepton-kaon correlations. It is for the weak decay vertex and does not include mixing of the neutral B meson. Mixing effects were corrected for by assuming a mixing parameter r of $(18.1 \pm 4.3)\%$.			
87 Measurement relies on lepton-kaon correlations. It includes production through mixing of the neutral B meson.			

 $\Gamma(K^- \text{anything})/\Gamma_{\text{total}}$ Γ_{49}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.13 ± 0.04	88 ALBRECHT	94C ARG	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.165 ± 0.011 ± 0.036	89 ALBRECHT	94C ARG	$e^+ e^- \rightarrow \gamma(4S)$
0.19 ± 0.05 ± 0.02	89 ALAM	87B CLEO	$e^+ e^- \rightarrow \gamma(4S)$
88 Measurement relies on lepton-kaon correlations. It is for the weak decay vertex and does not include mixing of the neutral B meson. Mixing effects were corrected for by assuming a mixing parameter r of $(18.1 \pm 4.3)\%$.			
89 Measurement relies on lepton-kaon correlations. It includes production through mixing of the neutral B meson.			

 $\Gamma(K^0/\bar{K}^0 \text{anything})/\Gamma_{\text{total}}$ Γ_{50}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.64 ± 0.04 OUR AVERAGE	90 ALBRECHT	94C ARG	$e^+ e^- \rightarrow \gamma(4S)$
0.642 ± 0.010 ± 0.042	ALAM	87B CLEO	$e^+ e^- \rightarrow \gamma(4S)$
0.63 ± 0.06 ± 0.06	90 ALBRECHT 94C assume a K^0/\bar{K}^0 multiplicity twice that of K_S^0 .		

 $\Gamma(K^*(892)^{\pm} \text{anything})/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.182 ± 0.054 ± 0.024	ALBRECHT	94J ARG	$e^+ e^- \rightarrow \gamma(4S)$

 $\Gamma(K^*(892)^0/\bar{K}^*(892)^0 \text{anything})/\Gamma_{\text{total}}$ Γ_{52}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.146 ± 0.016 ± 0.020	ALBRECHT	94J ARG	$e^+ e^- \rightarrow \gamma(4S)$

 $\Gamma(K^*(892)\gamma)/\Gamma_{\text{total}}$ Γ_{53}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
4.24 ± 0.54 ± 0.32		91 COAN	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<150 90 92 LESIAK 92 CBAL $e^+ e^- \rightarrow \gamma(4S)$ < 24 90 ALBRECHT 88H ARG $e^+ e^- \rightarrow \gamma(4S)$ 91 An average of $B(B^+ \rightarrow K^*(892)^+\gamma)$ and $B(B^0 \rightarrow K^*(892)^0\gamma)$ measurements reported in COAN 00 by assuming full correlated systematic errors.92 LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s -quark hadronization.

$\Gamma(K_1(1400)\gamma)/\Gamma_{\text{total}}$ Γ_{54}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<12.7 \times 10^{-5}$	90	93 COAN	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$< 1.6 \times 10^{-3}$	90	94 LESIAK	92 CBAL	$e^+ e^- \rightarrow \Upsilon(4S)$
$< 4.1 \times 10^{-4}$	90	ALBRECHT	88H ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

93 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.94 LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s -quark hadronization. $\Gamma(K_2^*(1430)\gamma)/\Gamma_{\text{total}}$ Γ_{55}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.66^{+0.59}_{-0.53} \pm 0.13$	95 COAN	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$	

 $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ <83 90 ALBRECHT 88H ARG $e^+ e^- \rightarrow \Upsilon(4S)$ 95 COAN 00 obtains a fitted signal yield of $15.9^{+5.7}_{-5.2}$ events. A search for contamination by $K^*(1410)$ yielded a rate consistent with 0; the central value assumes no contamination. $\Gamma(K_2(1770)\gamma)/\Gamma_{\text{total}}$ Γ_{56}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.2 \times 10^{-3}$	90	96 LESIAK	92 CBAL	$e^+ e^- \rightarrow \Upsilon(4S)$

96 LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s -quark hadronization. $\Gamma(K_3^*(1780)\gamma)/\Gamma_{\text{total}}$ Γ_{57}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.0 \times 10^{-3}$	90	ALBRECHT	88H ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

 $\Gamma(K_4^*(2045)\gamma)/\Gamma_{\text{total}}$ Γ_{58}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.0 \times 10^{-3}$	90	97 LESIAK	92 CBAL	$e^+ e^- \rightarrow \Upsilon(4S)$

97 LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s -quark hadronization. $\Gamma(\eta'(958)K)/\Gamma_{\text{total}}$ Γ_{59}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$(8.3^{+0.9}_{-0.8} \pm 0.7) \times 10^{-5}$	98 RICHICHI	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$	

98 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. $\Gamma(\eta'(958)K^*(892))/\Gamma_{\text{total}}$ Γ_{60}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.2 \times 10^{-5}$	90	99 RICHICHI	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

99 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta K)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{61}/Γ
$<5.2 \times 10^{-6}$	90	100	RICHICHI	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
100 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.					

 $\Gamma(\eta K^*(892))/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{62}/Γ
$(1.80^{+0.49}_{-0.43} \pm 0.18) \times 10^{-5}$	101	RICHICHI	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
101 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.				

 $\Gamma(\bar{b} \rightarrow \bar{s}\gamma)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{63}/Γ
$(3.3 \pm 0.4) \times 10^{-4}$ OUR AVERAGE				
$(3.36 \pm 0.53^{+0.65}_{-0.68}) \times 10^{-4}$	102	ABE	01F BELL	$e^+ e^- \rightarrow \gamma(4S)$
$(3.21 \pm 0.43^{+0.32}_{-0.29}) \times 10^{-4}$	103	CHEN	01C CLE2	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$(2.32 \pm 0.57 \pm 0.35) \times 10^{-4}$	ALAM	95	CLE2	Repl. by CHEN 01C
102 ABE 01F reports their systematic errors $\pm 0.42^{+0.50}_{-0.54}$, where the second error is due to the theoretical uncertainty. We combine them in quadrature.				
103 We have combined the experimental systematic theoretical uncertainties in quadrature. Also determined the first and second moments of the photon energy spectrum above 2.0 GeV: $\langle E_\gamma \rangle = 2.346 \pm 0.032 \pm 0.011$ GeV and $\langle E_\gamma^2 \rangle - \langle E_\gamma \rangle^2 = 0.0226 \pm 0.0066 \pm 0.0020$ GeV ² .				

 $\Gamma(\bar{b} \rightarrow \bar{s}\text{gluon})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{64}/Γ
<0.068	90	104	COAN	98 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.08	2	105	ALBRECHT	95D ARG	$e^+ e^- \rightarrow \gamma(4S)$	
104 COAN 98 uses D - ℓ correlation.						
105 ALBRECHT 95D use full reconstruction of one B decay as tag. Two candidate events for charmless B decay can be interpreted as either $b \rightarrow s\text{gluon}$ or $b \rightarrow u$ transition. If interpreted as $b \rightarrow s\text{gluon}$ they find a branching ratio of ~ 0.026 or the upper limit quoted above. Result is highly model dependent.						

 $\Gamma(\eta \text{anything})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{65}/Γ
$<4.4 \times 10^{-4}$	90	106	BROWDER	98 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
106 BROWDER 98 search for high momentum $B \rightarrow \eta X_s$ between 2.1 and 2.7 GeV/c.					

 $\Gamma(\eta' \text{anything})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{66}/Γ
$(6.2 \pm 1.6^{+1.3}_{-2.0}) \times 10^{-4}$	107	BROWDER	98 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
107 BROWDER 98 observed a signal of 39.0 ± 11.6 events in high momentum $B \rightarrow \eta' X_s$ production between 2.0 and 2.7 GeV/c. The branching fraction is based on the interpretation of $b \rightarrow sg$, where the last error includes additional uncertainties due to the color-suppressed $b \rightarrow$ backgrounds.				

$\Gamma(\rho\gamma)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{67}/Γ
$<1.4 \times 10^{-5}$	90	108 COAN	00	CLE2 $e^+ e^- \rightarrow \gamma(4S)$	

108 COAN 00 reports $B(B \rightarrow \rho\gamma)/B(B \rightarrow K^*(892)\gamma) < 0.32$ at 90%CL and scaled by the central value of $B(B \rightarrow K^*(892)\gamma) = (4.24 \pm 0.54 \pm 0.32) \times 10^{-5}$.

 $\Gamma(\pi^\pm \text{ anything})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{68}/Γ
$3.585 \pm 0.025 \pm 0.070$	109 ALBRECHT	93I ARG	$e^+ e^- \rightarrow \gamma(4S)$	

109 ALBRECHT 93 excludes π^\pm from K_S^0 and Λ decays. If included, they find $4.105 \pm 0.025 \pm 0.080$.

 $\Gamma(\pi^0 \text{ anything})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{69}/Γ
$2.35 \pm 0.02 \pm 0.11$	110 ABE	01J BELL	$e^+ e^- \rightarrow \gamma(4S)$	

110 From fully inclusive π^0 yield with no corrections from decays of K_S^0 or other particles.

 $\Gamma(\eta \text{ anything})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{70}/Γ
$0.176 \pm 0.011 \pm 0.012$	KUBOTA	96	CLE2 $e^+ e^- \rightarrow \gamma(4S)$	

 $\Gamma(\rho^0 \text{ anything})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{71}/Γ
$0.208 \pm 0.042 \pm 0.032$	ALBRECHT	94J ARG	$e^+ e^- \rightarrow \gamma(4S)$	

 $\Gamma(\omega \text{ anything})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{72}/Γ
<0.81	90	ALBRECHT	94J ARG	$e^+ e^- \rightarrow \gamma(4S)$	

 $\Gamma(\phi \text{ anything})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{73}/Γ
0.035 ± 0.007 OUR AVERAGE			Error includes scale factor of 1.8.	
0.0390 $\pm 0.0030 \pm 0.0035$	ALBRECHT	94J ARG	$e^+ e^- \rightarrow \gamma(4S)$	
0.023 $\pm 0.006 \pm 0.005$	BORTOLETTO86	CLEO	$e^+ e^- \rightarrow \gamma(4S)$	

 $\Gamma(\phi K^*(892))/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
$<2.2 \times 10^{-5}$	90	111 BERGFELD	98 CLE2

111 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\Lambda_c^\pm \text{ anything})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{75}/Γ
0.064 $\pm 0.008 \pm 0.008$		112 CRAWFORD	92 CLEO	$e^+ e^- \rightarrow \gamma(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.14 ± 0.09		113 ALBRECHT	88E ARG	$e^+ e^- \rightarrow \gamma(4S)$	
<0.112	90	114 ALAM	87 CLEO	$e^+ e^- \rightarrow \gamma(4S)$	

112 CRAWFORD 92 result derived from lepton baryon correlations. Assumes all charmed baryons in B^0 and B^\pm decay are Λ_c .

113 ALBRECHT 88E measured $B(B \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (0.30 \pm 0.12 \pm 0.06)\%$ and used $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (2.2 \pm 1.0)\%$ from ABRAMS 80 to obtain above number.

114 Assuming all baryons result from charmed baryons, ALAM 86 conclude the branching fraction is $7.4 \pm 2.9\%$. The limit given above is model independent.

$\Gamma(\Lambda_c^+ \text{ anything})/\Gamma(\bar{\Lambda}_c^- \text{ anything})$ Γ_{76}/Γ_{77}

VALUE	DOCUMENT ID	TECN	COMMENT
0.19±0.13±0.04	115 AMMAR	97 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

115 AMMAR 97 uses a high-momentum lepton tag ($P_\ell > 1.4 \text{ GeV}/c^2$).

$\Gamma(\bar{\Lambda}_c^- e^+ \text{ anything})/\Gamma(\Lambda_c^\pm \text{ anything})$ Γ_{78}/Γ_{75}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.05	90	116 BONVICINI	98 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

116 BONVICINI 98 uses the electron with momentum above $0.6 \text{ GeV}/c$.

$\Gamma(\bar{\Lambda}_c^- p \text{ anything})/\Gamma(\Lambda_c^\pm \text{ anything})$ Γ_{79}/Γ_{75}

VALUE	DOCUMENT ID	TECN	COMMENT
0.57±0.05±0.05	BONVICINI	98 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

$\Gamma(\bar{\Lambda}_c^- p e^+ \nu_e)/\Gamma(\bar{\Lambda}_c^- p \text{ anything})$ Γ_{80}/Γ_{79}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.04	90	117 BONVICINI	98 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

117 BONVICINI 98 uses the electron with momentum above $0.6 \text{ GeV}/c$.

$\Gamma(\bar{\Sigma}_c^{--} \text{ anything})/\Gamma_{\text{total}}$ Γ_{81}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0042±0.0021±0.0011	77	118 PROCARIO	94 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

118 PROCARIO 94 reports $[B(B \rightarrow \bar{\Sigma}_c^{--} \text{ anything}) \times B(\Lambda_c^+ \rightarrow p K^- \pi^+)] = 0.00021 \pm 0.00008 \pm 0.00007$. We divide by our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\bar{\Sigma}_c^- \text{ anything})/\Gamma_{\text{total}}$ Γ_{82}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.010	90	119 PROCARIO	94 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

119 PROCARIO 94 reports $[B(B \rightarrow \bar{\Sigma}_c^- \text{ anything}) \times B(\Lambda_c^+ \rightarrow p K^- \pi^+)] = < 0.00048$. We divide by our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.050$.

$\Gamma(\bar{\Sigma}_c^0 \text{ anything})/\Gamma_{\text{total}}$ Γ_{83}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0046±0.0021±0.0012	76	120 PROCARIO	94 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

120 PROCARIO 94 reports $[B(B \rightarrow \bar{\Sigma}_c^0 \text{ anything}) \times B(\Lambda_c^+ \rightarrow p K^- \pi^+)] = 0.00023 \pm 0.00008 \pm 0.00007$. We divide by our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Sigma_c^0 N(N=p \text{ or } n))/\Gamma_{\text{total}}$ Γ_{84}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0015	90	121 PROCARIO	94 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

121 PROCARIO 94 reports < 0.0017 for $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.043$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.050$.

$\Gamma(\Xi_c^0 \text{anything} \times B(\Xi_c^0 \rightarrow \Xi^- \pi^+))/\Gamma_{\text{total}}$ Γ_{85}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.144 ± 0.048 ± 0.021	122 BARISH	97 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

122 BARISH 97 find $79 \pm 27 \Xi_c^0$ events.

$\Gamma(\Xi_c^+ \text{anything} \times B(\Xi_c^+ \rightarrow \Xi^- \pi^+ \pi^+))/\Gamma_{\text{total}}$ Γ_{86}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.453 ± 0.096 ± 0.085	123 BARISH	97 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

123 BARISH 97 find $125 \pm 28 \Xi_c^+$ events.

$\Gamma(p/\bar{p}\text{anything})/\Gamma_{\text{total}}$ Γ_{87}/Γ

Includes p and \bar{p} from Λ and $\bar{\Lambda}$ decay.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.080 ± 0.004 OUR AVERAGE				

0.080 ± 0.005 ± 0.005		ALBRECHT	93I ARG	$e^+ e^- \rightarrow \gamma(4S)$
0.080 ± 0.005 ± 0.003		CRAWFORD	92 CLEO	$e^+ e^- \rightarrow \gamma(4S)$
0.082 ± 0.005 ± 0.013	2163	124 ALBRECHT	89K ARG	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

>0.021		125 ALAM	83B CLEO	$e^+ e^- \rightarrow \gamma(4S)$
--------	--	----------	----------	----------------------------------

124 ALBRECHT 89K include direct and nondirect protons.

125 ALAM 83B reported their result as $> 0.036 \pm 0.006 \pm 0.009$. Data are consistent with equal yields of p and \bar{p} . Using assumed yields below cut, $B(B \rightarrow p + X) = 0.03$ not including protons from Λ decays.

$\Gamma(p/\bar{p}(\text{direct})\text{anything})/\Gamma_{\text{total}}$ Γ_{88}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.055 ± 0.005 OUR AVERAGE				

0.055 ± 0.005 ± 0.0035		ALBRECHT	93I ARG	$e^+ e^- \rightarrow \gamma(4S)$
0.056 ± 0.006 ± 0.005		CRAWFORD	92 CLEO	$e^+ e^- \rightarrow \gamma(4S)$
0.055 ± 0.016	1220	126 ALBRECHT	89K ARG	$e^+ e^- \rightarrow \gamma(4S)$

126 ALBRECHT 89K subtract contribution of Λ decay from the inclusive proton yield.

$\Gamma(\Lambda/\bar{\Lambda}\text{anything})/\Gamma_{\text{total}}$ Γ_{89}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.040 ± 0.005 OUR AVERAGE				

0.038 ± 0.004 ± 0.006	2998	CRAWFORD	92 CLEO	$e^+ e^- \rightarrow \gamma(4S)$
0.042 ± 0.005 ± 0.006	943	ALBRECHT	89K ARG	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.022 ± 0.003 ± 0.0022		127 ACKERSTAFF	97N OPAL	$e^+ e^- \rightarrow Z$
------------------------	--	----------------	----------	-------------------------

>0.011		128 ALAM	83B CLEO	$e^+ e^- \rightarrow \gamma(4S)$
--------	--	----------	----------	----------------------------------

127 ACKERSTAFF 97N assures $B(b \rightarrow B) = 0.868 \pm 0.041$, i.e., an admixture of B^0 , B^\pm , and B_s .

128 ALAM 83B reported their result as $> 0.022 \pm 0.007 \pm 0.004$. Values are for $(B(\Lambda X) + B(\bar{\Lambda} X))/2$. Data are consistent with equal yields of p and \bar{p} . Using assumed yields below cut, $B(B \rightarrow \Lambda X) = 0.03$.

$\Gamma(\Lambda \text{anything})/\Gamma(\bar{\Lambda} \text{anything})$

Γ_{90}/Γ_{91}

VALUE	DOCUMENT ID	TECN	COMMENT
0.43±0.09±0.07	129 AMMAR 97	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

129 AMMAR 97 uses a high-momentum lepton tag ($P_\ell > 1.4 \text{ GeV}/c^2$).

$\Gamma(\Xi^-/\Xi^+ \text{anything})/\Gamma_{\text{total}}$

Γ_{92}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0027±0.0006 OUR AVERAGE				
0.0027±0.0005±0.0004	147	CRAWFORD 92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$

0.0028±0.0014 54 ALBRECHT 89K ARG $e^+ e^- \rightarrow \gamma(4S)$

$\Gamma(\text{baryons anything})/\Gamma_{\text{total}}$

Γ_{93}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.068±0.005±0.003	130 ALBRECHT 920 ARG	$e^+ e^- \rightarrow \gamma(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.076±0.014 131 ALBRECHT 89K ARG $e^+ e^- \rightarrow \gamma(4S)$

130 ALBRECHT 920 result is from simultaneous analysis of p and Λ yields, $p\bar{p}$ and $\Lambda\bar{p}$ correlations, and various lepton-baryon and lepton-baryon-antibaryon correlations. Supersedes ALBRECHT 89K.

131 ALBRECHT 89K obtain this result by adding their measurements ($5.5 \pm 1.6\%$) for direct protons and ($4.2 \pm 0.5 \pm 0.6\%$) for inclusive Λ production. They then assume ($5.5 \pm 1.6\%$) for neutron production and add it in also. Since each B decay has two baryons, they divide by 2 to obtain ($7.6 \pm 1.4\%$).

$\Gamma(p\bar{p}\text{anything})/\Gamma_{\text{total}}$

Γ_{94}/Γ

Includes p and \bar{p} from Λ and $\bar{\Lambda}$ decay.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0247±0.0023 OUR AVERAGE				

0.024 ± 0.001 ± 0.004 CRAWFORD 92 CLEO $e^+ e^- \rightarrow \gamma(4S)$

0.025 ± 0.002 ± 0.002 918 ALBRECHT 89K ARG $e^+ e^- \rightarrow \gamma(4S)$

$\Gamma(p\bar{p}\text{anything})/\Gamma(p/\bar{p}\text{anything})$

Γ_{94}/Γ_{87}

Includes p and \bar{p} from Λ and $\bar{\Lambda}$ decay.

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			

0.30±0.02±0.05 132 CRAWFORD 92 CLEO $e^+ e^- \rightarrow \gamma(4S)$

132 CRAWFORD 92 value is not independent of their $\Gamma(p\bar{p}\text{anything})/\Gamma_{\text{total}}$ value.

$\Gamma(\Lambda\bar{p}/\bar{\Lambda}p\text{anything})/\Gamma_{\text{total}}$

Γ_{95}/Γ

Includes p and \bar{p} from Λ and $\bar{\Lambda}$ decay.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.025±0.004 OUR AVERAGE				

0.029±0.005±0.005 CRAWFORD 92 CLEO $e^+ e^- \rightarrow \gamma(4S)$

0.023±0.004±0.003 165 ALBRECHT 89K ARG $e^+ e^- \rightarrow \gamma(4S)$

$\Gamma(\Lambda\bar{p}/\bar{\Lambda}p\text{anything})/\Gamma(\Lambda/\bar{\Lambda}\text{anything})$ Γ_{95}/Γ_{89} Includes p and \bar{p} from Λ and $\bar{\Lambda}$ decay.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$0.76 \pm 0.11 \pm 0.08$	133 CRAWFORD	92 CLEO	$e^+ e^- \rightarrow \gamma(4S)$
133 CRAWFORD	92	value	is not independent of their $[\Gamma(\Lambda\bar{p}\text{anything}) + \Gamma(\bar{\Lambda}p\text{anything})]/\Gamma_{\text{total}}$ value.

 $\Gamma(\Lambda\bar{\Lambda}\text{anything})/\Gamma_{\text{total}}$ Γ_{96}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.005	90		CRAWFORD	92 CLEO	$e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<0.0088	90	12	ALBRECHT	89K ARG	$e^+ e^- \rightarrow \gamma(4S)$

 $\Gamma(\Lambda\bar{\Lambda}\text{anything})/\Gamma(\Lambda/\bar{\Lambda}\text{anything})$ Γ_{96}/Γ_{89}

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<0.13	90	134 CRAWFORD	92 CLEO	$e^+ e^- \rightarrow \gamma(4S)$
134 CRAWFORD	92	value	is not independent of their $\Gamma(\Lambda\bar{\Lambda}\text{anything})/\Gamma_{\text{total}}$ value.	

 $\Gamma(e^+ e^- s)/\Gamma_{\text{total}}$ Γ_{97}/Γ Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5.7 $\times 10^{-5}$	90	GLENN	98 CLEO	$e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<0.05	90	BEBEK	81 CLEO	$e^+ e^- \rightarrow \gamma(4S)$

 $\Gamma(\mu^+ \mu^- s)/\Gamma_{\text{total}}$ Γ_{98}/Γ Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5.8 $\times 10^{-5}$	90	GLENN	98 CLEO	$e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<0.017	90	CHADWICK	81 CLEO	$e^+ e^- \rightarrow \gamma(4S)$

 $\Gamma(K e^+ e^-)/\Gamma_{\text{total}}$ Γ_{99}/Γ Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.3 $\times 10^{-6}$	90	ABE	02 BELL	$e^+ e^- \rightarrow \gamma(4S)$

 $\Gamma(K^*(892)e^+ e^-)/\Gamma_{\text{total}}$ Γ_{100}/Γ Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5.6 $\times 10^{-6}$	90	ABE	02 BELL	$e^+ e^- \rightarrow \gamma(4S)$

$\Gamma(K\mu^+\mu^-)/\Gamma_{\text{total}}$

Γ_{101}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	DOCUMENT ID	TECN	COMMENT
$(0.99 \pm 0.40 \pm 0.13) \times 10^{-6}$	ABE	02	BELL $e^+ e^- \rightarrow \gamma(4S)$

$\Gamma(K^*(892)\mu^+\mu^-)/\Gamma_{\text{total}}$

Γ_{102}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.1 \times 10^{-6}$	90	ABE	02	BELL $e^+ e^- \rightarrow \gamma(4S)$

$\Gamma(K\ell^+\ell^-)/\Gamma_{\text{total}}$

Γ_{103}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$0.75 \pm 0.25 \pm 0.06$	135	ABE	02	BELL $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.7	90	136 ANDERSON	01B CLE2	$e^+ e^- \rightarrow \gamma(4S)$
--------	----	--------------	----------	----------------------------------

135 Assumes lepton universality.

136 The result is for di-lepton masses above 0.5 GeV.

$\Gamma(K^*(892)\ell^+\ell^-)/\Gamma_{\text{total}}$

Γ_{104}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.3 \times 10^{-6}$	90	137 ANDERSON	01B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

137 The result is for di-lepton masses above 0.5 GeV.

$[\Gamma(e^+e^-s) + \Gamma(\mu^+\mu^-s)]/\Gamma_{\text{total}}$

$(\Gamma_{97} + \Gamma_{98})/\Gamma$

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.2 \times 10^{-5}$	90	GLENN	98	CLEO $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0024	90	138 BEAN	87	CLEO Repl. by GLENN 98
-----------	----	----------	----	------------------------

<0.0062	90	139 AVERY	84	CLEO Repl. by BEAN 87
-----------	----	-----------	----	-----------------------

138 BEAN 87 reports $[(\mu^+\mu^-) + (e^+e^-)]/2$ and we converted it.

139 Determine ratio of B^+ to B^0 semileptonic decays to be in the range 0.25–2.9.

$\Gamma(e^\pm\mu^\mp s)/\Gamma_{\text{total}}$

Γ_{105}/Γ

Test for lepton family number conservation. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.2 \times 10^{-5}$	90	GLENN	98	CLEO $e^+ e^- \rightarrow \gamma(4S)$

CP VIOLATION

A_{CP} is defined as

$$\frac{B(B \rightarrow \bar{f}) - B(\bar{B} \rightarrow f)}{B(B \rightarrow \bar{f}) + B(\bar{B} \rightarrow f)},$$

the CP-violation charge asymmetry of inclusive B^\pm and B^0 decay.

$A_{CP}(B \rightarrow K^*(892)\gamma)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.01 ± 0.07 OUR AVERAGE			
-0.044 ± 0.076 ± 0.012	140 AUBERT	02C BABR	$e^+ e^- \rightarrow \gamma(4S)$
+0.08 ± 0.13 ± 0.03	141 COAN	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
140 A 90% CL range is $-0.170 < A_{CP} < 0.082$.			
141 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.			

$A_{CP}(B \rightarrow s\gamma)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.079 ± 0.108 ± 0.022			
142 COAN	01 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
142 Corresponds to $-0.27 < A_{CP} < 0.10$ at 90% CL.			

$B \rightarrow X_c \ell \nu$ HADRONIC MASS MOMENTS

$\langle M_X^2 - \bar{M}_D^2 \rangle$ (First Moments)

VALUE (GeV ²)	DOCUMENT ID	TECN	COMMENT
0.251 ± 0.023 ± 0.062	143 CRONIN-HEN..01B	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

143 The leptons are required to have $P_1 > 1.5$ GeV/c.

$\langle M_X^2 - \bar{M}_X^2 \rangle$ (Second Moments)

VALUE (GeV ⁴)	DOCUMENT ID	TECN	COMMENT
0.576 ± 0.048 ± 0.168	144 CRONIN-HEN..01B	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

144 The leptons are required to have $P_1 > 1.5$ GeV/c.

$\langle (M_X^2 - \bar{M}_D^2)^2 \rangle$ (Second Moments)

VALUE (GeV ⁴)	DOCUMENT ID	TECN	COMMENT
0.639 ± 0.056 ± 0.178	145 CRONIN-HEN..01B	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

145 The leptons are required to have $P_1 > 1.5$ GeV/c.

B^\pm/B^0 ADMIXTURE REFERENCES

ABE	02	PRL 88 021801	K. Abe <i>et al.</i>	(Belle Collab.)
AUBERT	02C	PRL 88 101805	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	02G	PR D (to be publ.)	B. Aubert <i>et al.</i>	(Babar Collab.)
	hep-ex/0201041			
ABE	01F	PL B511 151	K. Abe <i>et al.</i>	(Belle Collab.)
ABE	01J	PR D64 072001	K. Abe <i>et al.</i>	(Belle Collab.)
ANDERSON	01B	PRL 87 181803	S. Anderson <i>et al.</i>	(CLEO Collab.)
CHEN	01	PR D63 031102	S. Chen <i>et al.</i>	(CLEO Collab.)
CHEN	01C	PRL 87 251807	S. Chen <i>et al.</i>	(CLEO Collab.)
COAN	01	PRL 86 5661	T.E. Coan <i>et al.</i>	(CLEO Collab.)
CRONIN-HEN...	01B	PRL 87 251808	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
ABREU	00R	PL B475 407	P. Abreu <i>et al.</i>	(DELPHI Collab.)
COAN	00	PR 84 5283	T.E. Coan <i>et al.</i>	(CLEO Collab.)
RICHICHI	00	PRL 85 520	S.J. Richichi <i>et al.</i>	(CLEO Collab.)
BARATE	98Q	EPJ C4 387	R. Barate <i>et al.</i>	(ALEPH Collab.)
BERGFELD	98	PRL 81 272	T. Bergfeld <i>et al.</i>	(CLEO Collab.)
BISHAI	98	PR D57 3847	M. Bishai <i>et al.</i>	(CLEO Collab.)
BONVICINI	98	PR D57 6604	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
BROWDER	98	PRL 81 1786	T.E. Browder <i>et al.</i>	(CLEO Collab.)
COAN	98	PRL 80 1150	T.E. Coan <i>et al.</i>	(CLEO Collab.)
GLENN	98	PRL 80 2289	S. Glenn <i>et al.</i>	(CLEO Collab.)
ACKERSTAFF	97N	ZPHY C74 423	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
AMMAR	97	PR D55 13	R. Ammar <i>et al.</i>	(CLEO Collab.)
BARISH	97	PRL 79 3599	B. Barish <i>et al.</i>	(CLEO Collab.)
BUSKULIC	97B	ZPHY C73 601	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
GIBBONS	97B	PR D56 3783	L. Gibbons <i>et al.</i>	(CLEO Collab.)
ALBRECHT	96D	PL B374 256	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BARISH	96B	PRL 76 1570	B.C. Barish <i>et al.</i>	(CLEO Collab.)
GIBAUT	96	PR D53 4734	D. Gibaut <i>et al.</i>	(CLEO Collab.)
KUBOTA	96	PR D53 6033	Y. Kubota <i>et al.</i>	(CLEO Collab.)
PDG	96	PR D54 1	R. M. Barnett <i>et al.</i>	
ALAM	95	PRL 74 2885	M.S. Alam <i>et al.</i>	(CLEO Collab.)
ALBRECHT	95D	PL B353 554	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BALEST	95B	PR D52 2661	R. Balest <i>et al.</i>	(CLEO Collab.)
BARISH	95	PR D51 1014	B.C. Barish <i>et al.</i>	(CLEO Collab.)
BUSKULIC	95B	PL B345 103	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ALBRECHT	94C	ZPHY C62 371	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	94J	ZPHY C61 1	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
PROCARIO	94	PRL 73 1472	M. Procario <i>et al.</i>	(CLEO Collab.)
ALBRECHT	93	ZPHY C57 533	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	93E	ZPHY C60 11	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	93H	PL B318 397	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	93I	ZPHY C58 191	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	93B	PL B319 365	J. Alexander <i>et al.</i>	(CLEO Collab.)
ARTUSO	93	PL B311 307	M. Artuso	(SYRA)
BARTEL	93B	PRL 71 4111	J.E. Bartelt <i>et al.</i>	(CLEO Collab.)
ALBRECHT	92E	PL B277 209	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	92G	ZPHY C54 1	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	92O	ZPHY C56 1	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BORTOLETTO	92	PR D45 21	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
CRAWFORD	92	PR D45 752	G. Crawford <i>et al.</i>	(CLEO Collab.)
HENDERSON	92	PR D45 2212	S. Henderson <i>et al.</i>	(CLEO Collab.)
LESIAK	92	ZPHY C55 33	T. Lesiak <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	91C	PL B255 297	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	91H	ZPHY C52 353	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
FULTON	91	PR D43 651	R. Fulton <i>et al.</i>	(CLEO Collab.)
YANAGISAWA	91	PRL 66 2436	C. Yanagisawa <i>et al.</i>	(CUSB II Collab.)
ALBRECHT	90	PL B234 409	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	90H	PL B249 359	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BORTOLETTO	90	PRL 64 2117	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
Also	92	PR D45 21	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
FULTON	90	PRL 64 16	R. Fulton <i>et al.</i>	(CLEO Collab.)
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)
PDG	90	PL B239	J.J. Hernandez <i>et al.</i>	(IFIC, BOST, CIT+)
ALBRECHT	89K	ZPHY C42 519	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ISGUR	89B	PR D39 799	N. Isgur <i>et al.</i>	(TNTO, CIT)
WACHS	89	ZPHY C42 33	K. Wachs <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	88E	PL B210 263	H. Albrecht <i>et al.</i>	(ARGUS Collab.)

ALBRECHT	88H	PL B210 258	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
KOERNER	88	ZPHY C38 511	J.G. Korner, G.A. Schuler	(MANZ, DESY)
ALAM	87	PRL 59 22	M.S. Alam <i>et al.</i>	(CLEO Collab.)
ALAM	87B	PRL 58 1814	M.S. Alam <i>et al.</i>	(CLEO Collab.)
ALBRECHT	87D	PL B199 451	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	87H	PL B187 425	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BEAN	87	PR D35 3533	A. Bean <i>et al.</i>	(CLEO Collab.)
BEHRENDS	87	PRL 59 407	S. Behrends <i>et al.</i>	(CLEO Collab.)
BORTOLETTO	87	PR D35 19	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
ALAM	86	PR D34 3279	M.S. Alam <i>et al.</i>	(CLEO Collab.)
BALTRUSAIT...	86E	PRL 56 2140	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BORTOLETTO	86	PRL 56 800	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
HAAS	86	PRL 56 2781	J. Haas <i>et al.</i>	(CLEO Collab.)
ALBRECHT	85H	PL 162B 395	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
CSORNA	85	PRL 54 1894	S.E. Csorna <i>et al.</i>	(CLEO Collab.)
HAAS	85	PRL 55 1248	J. Haas <i>et al.</i>	(CLEO Collab.)
AVERY	84	PRL 53 1309	P. Avery <i>et al.</i>	(CLEO Collab.)
CHEN	84	PRL 52 1084	A. Chen <i>et al.</i>	(CLEO Collab.)
LEVMAN	84	PL 141B 271	G.M. Levman <i>et al.</i>	(CUSB Collab.)
ALAM	83B	PRL 51 1143	M.S. Alam <i>et al.</i>	(CLEO Collab.)
GREEN	83	PRL 51 347	J. Green <i>et al.</i>	(CLEO Collab.)
KLOPFEN...	83B	PL 130B 444	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)
ALTARELLI	82	NP B208 365	G. Altarelli <i>et al.</i>	(ROMA, INFN, FRAS)
BRODY	82	PRL 48 1070	A.D. Brody <i>et al.</i>	(CLEO Collab.)
GIANNINI	82	NP B206 1	G. Giannini <i>et al.</i>	(CUSB Collab.)
BEBEK	81	PRL 46 84	C. Bebek <i>et al.</i>	(CLEO Collab.)
CHADWICK	81	PRL 46 88	K. Chadwick <i>et al.</i>	(CLEO Collab.)
ABRAMS	80	PRL 44 10	G.S. Abrams <i>et al.</i>	(SLAC, LBL)