

# $V_{cb}$ and $V_{ub}$ CKM Matrix Elements

OMITTED FROM SUMMARY TABLE

---

## DETERMINATION OF $|V_{cb}|$

Written April 2002 by M. Artuso (Syracuse University)  
and E. Barberio (CERN).

### *I. Introduction*

In the framework of the Standard Model, the quark sector is characterized by a rich pattern of flavor-changing transitions, described by the Cabibbo-Kobayashi-Maskawa (CKM) matrix (See CKM review [1]). This report focuses on the quark mixing parameter  $|V_{cb}|$ .

Two different methods have been used to extract this parameter from data: the **exclusive** measurement, where  $|V_{cb}|$  is extracted by studying exclusive  $B \rightarrow D^* \ell \nu$  and  $B \rightarrow D \ell \nu$  decay processes; and the **inclusive** measurement, which uses the semileptonic width of  $b$ -hadron decays. Theoretical estimates play a crucial role in extracting  $|V_{cb}|$ , and an understanding of their uncertainties is very important.

### *II. Exclusive $|V_{cb}|$ determination*

The exclusive  $|V_{cb}|$  determination is obtained studying the  $B \rightarrow D^* \ell \nu$  and  $B \rightarrow D \ell \nu$  decays, using Heavy Quark Effective Theory (HQET), an exact theory in the limit of infinite quark masses. Presently the  $B \rightarrow D \ell \nu$  transition provides a less precise value, and is used as a check.

***The decay  $B \rightarrow D^* \ell \nu$  in HQET:*** HQET predicts that the differential partial decay width for this process,  $d\Gamma/dw$ , is related to  $|V_{cb}|$  through:

$$\frac{d\Gamma}{dw}(B \rightarrow D^* \ell \nu) = \frac{G_F^2 |V_{cb}|^2}{48\pi^3} \mathcal{K}(w) \mathcal{F}(w)^2, \quad (1)$$

where  $w$  is the inner product of the  $B$  and  $D^*$  meson 4-velocities,  $\mathcal{K}(w)$  is a known phase-space factor, and the form factor  $\mathcal{F}(w)$  is generally expressed as the product of a normalization constant,  $\mathcal{F}(1)$ , and a function,  $g(w)$ , constrained by dispersion relations [2].

There are several different corrections to the infinite mass value  $\mathcal{F}(1) = 1$  [3]:

$$\mathcal{F}(1) = \eta_{\text{QED}}\eta_A \left[ 1 + \delta_{1/m_Q^2} + \dots \right] \quad (2)$$

where  $Q = c$  or  $b$ . By virtue of Luke's theorem [4], the first term in the non-perturbative expansion in powers of  $1/m_Q$  vanishes. QED corrections up to leading-logarithmic order give  $\eta_{\text{QED}} \approx 1.007$  [3] and QCD radiative corrections to two loops give  $\eta_A = 0.960 \pm 0.007$  [5]. Different estimates of the  $1/m_Q^2$  corrections, involving terms proportional to  $1/m_b^2$ ,  $1/m_c^2$ , and  $1/(m_b m_c)$ , have been performed in a quark model [6,7], with OPE sum rules [8], and, more recently, with an HQET based lattice gauge calculation [9]. The value from this quenched lattice HQET calculation is  $\mathcal{F}(1) = 0.913_{-0.017}^{+0.024} \pm 0.016_{-0.014}^{+0.003} \begin{smallmatrix} +0.000 & +0.006 \\ -0.016 & -0.014 \end{smallmatrix}$ . The errors quoted reflect the statistical accuracy, the matching error, the lattice finite size, the uncertainty in the quark masses, and an estimate of the error induced by the quenched approximation, respectively. The central value obtained with OPE sum rules is similar, with an error of  $\pm 0.04$  [10]. Consequently,  $\mathcal{F}(1) = 0.91 \pm 0.04$  [10] will be used here.

The analytical expression of  $\mathcal{F}(w)$  is not known a-priori, and this introduces an additional uncertainty in the determination of  $\mathcal{F}(1)|V_{cb}|$ . First measurements of  $|V_{cb}|$  were performed assuming a linear approximation for  $\mathcal{F}(w)$ . It has been shown [11]

that this assumption is not justified, and that linear fits systematically underestimate the extrapolation at zero recoil ( $w = 1$ ) by about 3%. Most of this effect is related to the curvature of the form factor, and does not depend strongly upon the details of the non-linear shape chosen [11]. All recent published results use a non-linear shape for  $\mathcal{F}(w)$ , approximated with an expansion near  $w = 1$  [12].  $\mathcal{F}(w)$  is parameterized in terms of the variable  $\rho^2$ , which is the slope of the form factor at zero recoil given in Ref. 12.

***Experimental techniques to study the decay  $B \rightarrow D^*\ell\nu$ :***

The decay  $B \rightarrow D^*\ell\nu$  has been studied in experiments performed at center-of-mass energies equal to the  $\Upsilon(4S)$  mass and the  $Z^0$  mass. At the  $\Upsilon(4S)$ , experiments have the advantage that the  $w$  resolution is quite good. However, they have more limited statistics near  $w = 1$  in the decay  $\overline{B}^0 \rightarrow D^{*+}\ell\nu$ , because of the lower reconstruction efficiency of the slow pion, from the  $D^{*+} \rightarrow \pi^+D^0$  decay. The decay  $B^- \rightarrow D^{*0}\ell\overline{\nu}$  is not affected by this problem, and CLEO [13] studies both channels. In addition, kinematic constraints enable  $\Upsilon(4S)$  experiments to identify the final state, including the  $D^*$ , without a large contamination from the poorly known semileptonic decays including a hadronic system heavier than  $D^*$ , commonly identified as ‘ $D^{**}$ .’ At LEP,  $B$ ’s are produced with a large momentum (about 30 GeV on average). This makes the determination of  $w$  dependent upon the neutrino four-momentum reconstruction, thus giving a relatively poor resolution and limited physics background rejection capabilities. By contrast, LEP experiments benefit from an efficiency that is only mildly dependent upon  $w$ .

Experiments determine the product  $(\mathcal{F}(1) \cdot |V_{cb}|)^2$  by fitting the measured  $d\Gamma/dw$  distribution. Measurements have been

performed by CLEO [13], Belle [14], DELPHI [15], ALEPH [16], and OPAL [17]. At LEP, the dominant source of systematic error is the uncertainty on the contribution to  $d\Gamma/dw$  from semileptonic  $B$  decays with final states including a hadron system heavier than the  $D^*$ . This component includes both narrow orbitally excited charmed mesons and non-resonant or broad species. The existence of narrow resonant states is well established [1], and a signal of a broad resonance has been seen by CLEO [18], but the decay characteristics of these states in  $b$ -hadron semileptonic decays have large uncertainties. The average of ALEPH [19], CLEO [20], and DELPHI [21] narrow state branching fractions show that the ratio  $R_{\star\star} = \frac{\text{B}(\overline{B} \rightarrow D_2^* \ell \overline{\nu})}{\text{B}(\overline{B} \rightarrow D_1 \ell \overline{\nu})}$  is smaller than one ( $< 0.6$  at 95% C.L. [22]), in disagreement with HQET models where an infinite quark mass is assumed [23], but in agreement with models which take into account finite quark mass corrections [24]. Hence, LEP experiments use the treatment of narrow  $D^{\star\star}$  proposed in Ref. 24, which accounts for  $\mathcal{O}(1/m_c)$  corrections. Ref. 24 provides several possible approximations of the form factors that depend on five different expansion schemes, and on three input parameters. To calculate the systematic errors, each proposed scheme is tested, with the relevant input parameters varied over a range consistent with the experimental limit on  $R_{\star\star}$ . The quoted systematic error is the maximal difference from the central value obtained with this method. Broad resonances or other non-resonant terms may not be modelled correctly with this approach.

To combine the published data, the central values and the errors of  $\mathcal{F}(1)|V_{cb}|$  and  $\rho^2$  are re-scaled to the same set of input parameters and their quoted uncertainties. The  $\mathcal{F}(1)|V_{cb}|$

values used for this average are extracted using the parametrization in Ref. 13, based on the experimental determinations of the vector and axial form factor ratios  $R_1$  and  $R_2$  [26]. The LEP data, which originally used theoretical values for these ratios, are re-scaled accordingly [25]. Table 1 summarizes the corrected data. The averaging procedure [25] takes into account statistical and systematic correlations between  $\mathcal{F}(1)|V_{cb}|$  and  $\rho^2$ . Averaging the measurements in Table 1, we get:

$$\mathcal{F}(1)|V_{cb}| = (38.3 \pm 1.0) \times 10^{-3}$$


and

$$\rho^2 = 1.5 \pm 0.13, \quad (3)$$

with a confidence level \* The  $\chi^2$  per degree of freedom is less than 2, and we do not scale the error. of 5.1%. The error ellipses for the corrected measurements and for the world average are shown in Figure 1.

The main contributions to the  $\mathcal{F}(1)|V_{cb}|$  systematic error are from the uncertainty on the  $B \rightarrow D^{*\ell}\nu$  shape and  $B(b \rightarrow B_d)$ , ( $0.57 \times 10^{-3}$ ), fully correlated among the LEP experiments, the branching fraction of  $D$  and  $D^*$  decays, ( $0.4 \times 10^{-3}$ ), fully correlated among all the experiments, and the slow pion reconstruction from Belle and CLEO which are uncorrelated, ( $0.28 \times 10^{-3}$ ). The main contribution to the  $\rho^2$  systematic error is from the uncertainties in the measured values of  $R_1$  and  $R_2$  (0.13), fully correlated among experiments. Because of the large contribution of this uncertainty to the non-diagonal terms of the covariance matrix, the averaged  $\rho^2$  is higher than one would naively expect.

Using  $\mathcal{F}(1) = 0.91 \pm 0.04$  [10], we get  $|V_{cb}| = (42.1 \pm 1.1_{\text{exp}} \pm 1.9_{\text{theo}}) \times 10^{-3}$ . The dominant error is theoretical, but there



**Figure 1:** The error ellipses for the corrected measurements and world average for  $\mathcal{F}(1)|V_{cb}|$  vs  $\rho^2$ . The ellipses are the product between the  $1\sigma$  error of  $\mathcal{F}(1)|V_{cb}|$ ,  $\rho^2$ , and the correlation between the two. Consequently the ellipses correspond to about 37% CL.

are good prospects that lattice gauge calculations will improve significantly the accuracy of their estimate.

**The decay  $B \rightarrow D\ell\nu$ :** The study of the decay  $B \rightarrow D\ell\nu$  poses new challenges both from the theoretical and experimental point of view.

**Table 1:** Experimental results (from  $B \rightarrow D^* \ell \nu$  analyses) after the correction to common inputs and world average. The LEP numbers are corrected to use  $R_1$  and  $R_2$  from CLEO data.  $\rho^2$  is the slope of the form factor at zero recoil as defined in Ref. 12.

Exp.	$\mathcal{F}(1) V_{cb} (\times 10^3)$	$\rho^2$	Corr <sub>stat</sub>
ALEPH	$33.8 \pm 2.1 \pm 1.6$	$0.74 \pm 0.25 \pm 0.41$	94%
DELPHI	$36.1 \pm 1.4 \pm 2.5$	$1.42 \pm 0.14 \pm 0.37$	94%
OPAL	$38.5 \pm 0.9 \pm 1.8$	$1.35 \pm 0.12 \pm 0.31$	89%
Belle	$36.0 \pm 1.9 \pm 1.8$	$1.45 \pm 0.16 \pm 0.20$	90%
CLEO	$43.3 \pm 1.3 \pm 1.8$	$1.61 \pm 0.09 \pm 0.21$	86%
World average	$38.3 \pm 0.5 \pm 0.9$	$1.51 \pm 0.05 \pm 0.12$	86%

The differential decay rate for  $B \rightarrow D \ell \nu$  can be expressed as:

$$\frac{d\Gamma_D}{dw}(B \rightarrow D \ell \nu) = \frac{G_F^2 |V_{cb}|^2}{48\pi^3} \mathcal{K}_D(w) \mathcal{G}(w)^2, \quad (4)$$

where  $w$  is the inner product of the  $B$  and  $D$  meson 4-velocities,  $\mathcal{K}_D(w)$  is the phase space, and the form factor  $\mathcal{G}(w)$  is generally expressed as the product of a normalization factor,  $\mathcal{G}(1)$ , and a function,  $g_D(w)$ , constrained by dispersion relations [2].

The strategy to extract  $\mathcal{G}(1)|V_{cb}|$  is identical to that used for the  $B \rightarrow D^* \ell \nu$  decay. However, in this case there is no suppression of  $1/m_Q$  (*i.e.*, no Luke theorem) and corrections and QCD effects on  $\mathcal{G}(1)$  are calculated with less accuracy than  $\mathcal{F}(1)$  [27,28]. Moreover,  $d\Gamma_D/dw$  is more heavily suppressed near  $w = 1$  than  $d\Gamma_{D^*}/dw$ , due to the helicity mismatch between initial and final states. This channel is also much

more challenging from the experimental point of view as it is hard to isolate from the dominant background  $B \rightarrow D^* \ell \nu$ , as well as from fake  $D\text{-}\ell$  combinations. Thus, the extraction of  $|V_{cb}|$  from this channel is less precise than the one from the  $B \rightarrow D^* \ell \nu$  decay. Nevertheless, the  $B \rightarrow D \ell \nu$  channel provides a consistency check, and allows a test of heavy-quark symmetry [28] through the measurement of the form factor  $\mathcal{G}(w)$ , as HQET predicts the ratio  $\mathcal{G}(w)/\mathcal{F}(w)$  to be very close to one.

Belle [29] and ALEPH [16] studied the  $\overline{B}^0 \rightarrow D^+ \ell^- \overline{\nu}$  channel, while CLEO [30] studied both  $B^+ \rightarrow D^0 \ell^+ \overline{\nu}$  and  $\overline{B}^0 \rightarrow D^+ \ell^- \overline{\nu}$  decays. Averaging the data in Table 2 [25], we get  $\mathcal{G}(1)|V_{cb}| = (41.3 \pm 4.0) \times 10^{-3}$  and  $\rho_D^2 = 1.19 \pm 0.19$ , where  $\rho_D^2$  is the slope of the form factor at zero recoil given in Ref. 12.

**Table 2:** Experimental results after the correction to common inputs and world average.  $\rho_D^2$  is the slope of the form factor at zero recoil given in Ref. 12.

Exp.	$\mathcal{G}(1) V_{cb} (\times 10^3)$	$\rho_D^2$
ALEPH	$37.7 \pm 9.9 \pm 6.5$	$0.90 \pm 0.98 \pm 0.38$
Belle	$41.2 \pm 4.4 \pm 5.1$	$1.12 \pm 0.22 \pm 0.14$
CLEO	$44.6 \pm 5.8 \pm 3.5$	$1.27 \pm 0.25 \pm 0.14$
World average	$41.3 \pm 2.9 \pm 2.7$	$1.19 \pm 0.15 \pm 0.12$



The theoretical predictions for  $\mathcal{G}(1)$  are consistent:  $1.03 \pm 0.07$  [31], and  $0.98 \pm 0.07$  [28]. A quenched lattice calculation gives  $\mathcal{G}(1) = 1.058_{-0.017}^{+0.020}$  [32], where the errors do not include the uncertainties induced by the quenching approximation and lattice spacing. Using  $\mathcal{G}(1) = 1.0 \pm 0.07$ , we get  $|V_{cb}| = (41.3 \pm 4.0_{\text{exp}} \pm 2.9_{\text{theo}}) \times 10^{-3}$ , consistent with the value extracted from  $B \rightarrow D^* \ell \nu$  decay, but with a larger uncertainty.

The experiments have also measured the differential decay rate distribution to extract the ratio  $\mathcal{G}(w)/\mathcal{F}(w)$ . The data are compatible with a universal form factor as predicted by HQET. From the measured values of  $\mathcal{G}(1)|V_{cb}|$  and  $\mathcal{F}(1)|V_{cb}|$ , we get  $\mathcal{G}(1)/\mathcal{F}(1) = 1.08 \pm 0.09$ , consistent with the form-factor values we used.

### ***III. $|V_{cb}|$ determination from inclusive $B$ semileptonic decays***

Alternatively,  $|V_{cb}|$  can be extracted from the inclusive branching fraction for semileptonic  $b$  hadron decays  $B(B \rightarrow X_c \ell \nu)$  [33,34]. Several studies have shown that the spectator model decay rate is the leading term in a well-defined expansion controlled by the parameter  $\Lambda_{\text{QCD}}/m_b$ . Non-perturbative corrections to this leading approximation arise only to order  $1/m_b^2$ . The key issue in this approach is the ability to separate non-perturbative corrections, that can be expressed as a series in powers of  $1/m_b$ , and perturbative corrections, expressed in powers of  $\alpha_s$ . Quark-hadron duality is an important *ab initio* assumption in these calculations. While several authors [35] argue that this ansatz does not introduce appreciable errors, as they expect that duality violations affect the semileptonic width only in high powers in the non-perturbative expansion, other authors recognize that a presently unknown

correction may be associated with this assumption [36]. Arguments supporting a possible sizeable source of errors related to the assumption of quark-hadron duality have been proposed [37]. This issue needs to be resolved with further experimental studies. At present, no explicit additional error has been added to account for possible quark-hadron duality violation.

The coefficients of the  $1/m_b$  power terms are expectations values of operators that include non-perturbative physics. Relationships that are valid up to  $1/m_b^2$  include four such parameters: the expectation value of the kinetic operator, corresponding to the average of the square of the heavy-quark momentum inside the hadron, the expectation value of the chromomagnetic operator, and the heavy-quark masses ( $m_b$  and  $m_c$ ). The expectation value of the kinetic operator is introduced in the literature as  $-\lambda_1$  [38,39] or  $\mu_\pi^2$  [33,34], whereas the expectation value of the chromomagnetic operator is defined as  $\lambda_2$  [38,39] or  $\mu_G^2$  [33,34]. The two notations reflect a difference in the approach used to handle the energy scale  $\mu$  used to separate long-distance from short-distance physics. HQET is most commonly renormalized in a mass-independent scheme, thus making the quark masses the pole masses of the underlying theory (QCD). The second group of authors prefer the definition of the non-perturbative operators using a mass scale  $\mu \approx 1$  GeV.

The corresponding equations for the semileptonic width can be found in Refs. 33,40. Ref. 40 has been used to extract  $|V_{cb}|$  from the semileptonic branching fraction measured by CLEO, and to measure the heavy-quark expansion (HQE) parameters  $\overline{\Lambda}$  and  $\lambda_1$ , as discussed below.

The quark masses are related to the corresponding meson masses through [6]:

$$m_b = \overline{M}_B - \overline{\Lambda} + \frac{\lambda_1}{2\overline{M}_B}, \quad (5)$$

where  $\overline{M}_B$  is the spin averaged  $B-B^*$  mass ( $\overline{M}_B = 5.3134$  GeV/ $c^2$ ). A similar equation relates  $m_c$  and  $\overline{M}_D$ . The parameter  $\overline{\Lambda}$  represents the energy of the light quark and gluons.

**HQE and moments in semileptonic decays:** Experimental determinations of the HQE parameters are important in several respects. In particular, redundant determinations of these parameters may uncover inconsistencies, or point to violation of some important assumptions inherent in these calculations. The parameter  $\lambda_2$  can be extracted from the  $B^*-B$  mass splitting, whereas the other parameters need more elaborate measurements.

The first stage of this experimental program has been completed recently. The CLEO collaboration has measured the shape of the photon spectrum in  $b \rightarrow s\gamma$  inclusive decays. Its first moment, giving the average energy of the  $\gamma$  emitted in this transition, is related to the  $b$  quark mass. In the formalism of Ref. 40, this corresponds to the measurement of the parameter  $\overline{\Lambda} = 0.35 \pm 0.07 \pm 0.10$  GeV [41].

The parameter  $\lambda_1$  is determined experimentally through a measurement of the first moment of the mass  $M_X$  of the hadronic system recoiling against the  $\ell-\bar{\nu}$  pair. The relationship between the first moment of  $M_1 = \langle M_X^2 - M_D^2 \rangle / M_B^2$  and the parameters  $\overline{\Lambda}$  and  $\lambda_1$  is given in Ref. 42.

The measured value for  $\langle M_X^2 - M_D^2 \rangle$  [42] is  $0.251 \pm 0.066$  GeV<sup>2</sup>. This constraint, combined with the measurement of the mean photon energy in  $b \rightarrow s\gamma$ , implies a value of

$\lambda_1 = -0.24 \pm 0.11 \text{ GeV}^2$ , to order  $1/M_B^3$  and  $\beta_0\alpha_s^2$  in  $(\overline{\text{MS}})$ . The quoted theoretical uncertainty of 2% accounts for the  $1/M_B^3$  and  $\alpha_s$  uncertainties, but not for possible violations of quark-hadron duality.

***Experimental determination of the semileptonic branching fraction:*** The value of  $B(B \rightarrow X_c\ell\nu)$  has been measured both at the  $\Upsilon(4S)$  and LEP.

The most recent CLEO data, published in 1996 and based on a subset of the data sample accumulated now, obtains this branching fraction using a lepton tagged sample [43]. In this approach, a di-lepton sample is studied, and the charge correlation between the two leptons is used to disentangle leptons coming from the direct decay  $B \rightarrow X_c\ell\nu$  and the dominant background at low lepton momenta, the cascade decay  $B \rightarrow X_c \rightarrow X_s\ell\nu$ . This method was pioneered by the ARGUS collaboration [44] to measure the electron spectrum from  $B \rightarrow X_c\ell\nu$  down to 0.6 GeV/ $c$ . Thus, it reduces the model dependence of the extracted semileptonic branching fraction very substantially. They obtain  $B(B \rightarrow X_c e \nu) = (10.49 \pm 0.17 \pm 0.43)\%$ . The systematic error (4%) is dominated by experimental uncertainties. Lepton identification efficiency, fake rate determination, and tracking efficiencies contribute to 3% of this overall error. The remaining error is a sum of several small corrections associated with the uncertainty in the mixing parameter, and additional background estimates [43].

Combining  $\Upsilon(4S)$  results [1], we obtain:  $B(b \rightarrow X\ell\nu) = (10.38 \pm 0.32)\%$ . Using  $\tau_{B^+}, \tau_{B^0}$  [1],  $f_{+-}/f_{00} = 1.04 \pm 0.08$  [45], and subtracting  $B(b \rightarrow u\ell\nu) = (0.17 \pm 0.05)\%$ , we get:  $B(b \rightarrow X_c\ell\nu) = (10.21 \pm 0.32)\%$  and  $\Gamma(b \rightarrow X_c\ell\nu) = (0.419 \pm 0.013 \pm 0.003) \times 10^{-10} \text{ MeV}$ , where  $0.003 \times 10^{-10} \text{ MeV}$  includes the

uncertainties from  $B(b \rightarrow u\ell\nu)$ , and the model dependence correlated with LEP.

At LEP,  $B^0$ ,  $B^-$ ,  $B_s$ , and  $b$  baryon are produced, so the measured inclusive semileptonic branching ratio is an average over the different hadron species. Assuming that the semileptonic widths of all  $b$  hadrons are equal, the following relation holds:

$$\begin{aligned}
 B(b \rightarrow X_c\ell\nu)_{\text{LEP}} &= \\
 & f_{B^0} \frac{\Gamma(B^0 \rightarrow X_c\ell\nu)}{\Gamma(B^0)} + f_{B^-} \frac{\Gamma(B^- \rightarrow X_c\ell\nu)}{\Gamma(B^-)} \\
 & + f_{B_s} \frac{\Gamma(B_s \rightarrow X_c\ell\nu)}{\Gamma(B_s)} + f_{\Lambda_b} \frac{\Gamma(\Lambda_b \rightarrow X_c\ell\nu)}{\Gamma(\Lambda_b)} \\
 & = \Gamma(B \rightarrow X_c\ell\nu)\tau_b, \tag{6}
 \end{aligned}$$

where  $\tau_b$  is the average  $b$ -hadron lifetime. Taking into account the present precision of LEP measurements of  $b$ -baryon semileptonic branching ratios and lifetimes, the estimate uncertainty for a possible difference for the width of  $b$  baryons is 0.13%.

At LEP,  $B(b \rightarrow X\ell\nu)$  is measured with dedicated analyses [47–50](Table 3). The average LEP value for  $B(b \rightarrow X\ell\nu) = (10.59 \pm 0.09 \pm 0.30)\%$  is taken from a fit [46], which combines the semileptonic branching ratios, the  $B^0 - \bar{B}^0$  mixing parameter  $\bar{\chi}_b$ , and  $R_b = \Gamma(Z \rightarrow b\bar{b})/\Gamma(Z \rightarrow \text{had})$ . Ref. 47 shows that the main contribution to the modelling error is the uncertainty in the composition of the semileptonic width, including the narrow, wide and non-resonant  $D^{*\star}$  states.  $B_s$  and  $b$  baryons are about 20% of the total signal, and their contribution to the uncertainty of the spectrum is small. In this average, we use the modelling error quoted by Ref. 47, rather than the error from the combined fit, as the ALEPH procedure is based on

more recent information. The dominant errors in the combined branching fraction are the modelling of semileptonic decays (2.6%) and the detector related items (1.3%).

**Table 3:**  $B(b \rightarrow \ell \nu)$  measurement from LEP and their average. The errors quoted reflect statistical, systematic, and modelling uncertainties respectively.

Experiment	$B(b \rightarrow \ell \nu)\%$
ALEPH	$10.70 \pm 0.10 \pm 0.23 \pm 0.26$
DELPHI	$10.70 \pm 0.08 \pm 0.21 \pm_{-0.30}^{+0.44}$
L3	$10.85 \pm 0.12 \pm 0.38 \pm 0.26$
L3 (double-tag)	$10.16 \pm 0.13 \pm 0.20 \pm 0.22$
OPAL	$10.83 \pm 0.10 \pm 0.20 \pm_{-0.13}^{+0.20}$
LEP Average	$10.59 \pm 0.09 \pm 0.15 \pm 0.26$

Subtracting  $B(b \rightarrow u \ell \nu)$  from the LEP semileptonic branching fraction, we get:  $B(b \rightarrow X_c \ell \nu) = (10.42 \pm 0.34)\%$ , and using  $\tau_b$  [1]:  $\Gamma(b \rightarrow X_c \ell \nu) = (0.439 \pm 0.010 \pm 0.011) \times 10^{-10}$  MeV, where the systematic error  $0.011 \times 10^{-10}$  MeV reflects the  $B(b \rightarrow u \ell \nu)$  uncertainty and the model dependence, correlated with the  $\Upsilon(4S)$  result.

Combining the LEP and the  $\Upsilon(4S)$  semileptonic widths, we get:  $\Gamma(b \rightarrow X_c \ell \nu) = (0.43 \pm 0.01) \times 10^{-10}$  MeV, which is used in the formula of Ref. 42 to get:

$$|V_{cb}|_{\text{incl}} = (40.4 \pm 0.5_{\text{exp}} \pm 0.5_{\lambda_1, \bar{\Lambda}} \pm 0.8_{\text{theo}}) \times 10^{-3}, \quad (7)$$

where the first error is experimental, and the second is from the measured value of  $\lambda_1$  and  $\bar{\Lambda}$ , assumed to be universal up

to higher orders. The third error is from  $1/m_b^3$  corrections and from the ambiguity in the  $\alpha_s$  scale definition. The error on the average  $b$ -hadron lifetime is assumed to be uncorrelated with the error on the semileptonic branching ratio.

#### IV. Conclusions

The values of  $|V_{cb}|$  obtained both from the inclusive and exclusive method agree within errors. The value of  $|V_{cb}|$  obtained from the analysis of the  $B \rightarrow D^* \ell \nu$  decay is:

$$|V_{cb}|_{\text{exclusive}} = (42.1 \pm 1.1_{\text{exp}} \pm 1.9_{\text{theo}}) \times 10^{-3}, \quad (8)$$

where the first error is experimental and the second error is from the  $1/m_Q^2$  corrections to  $\mathcal{F}(1)$ . The value of  $|V_{cb}|$ , obtained from inclusive semileptonic branching fractions is:

$$|V_{cb}|_{\text{incl}} = (40.4 \pm 0.5_{\text{exp}} \pm 0.5_{\lambda_1, \bar{\Lambda}} \pm 0.8_{\text{theo}}) \times 10^{-3}, \quad (9)$$

where the first error is experimental, the second error is from the measured values of  $\lambda_1$  and  $\bar{\Lambda}$ , assumed to be universal up to higher orders, and the last is from  $1/m_b^3$  corrections and  $\alpha_s$ . Non-quantified uncertainties are associated with a possible quark-hadron duality violation. For this reason, we chose not to average the two numbers.

While experimental errors have reached 2.7% and 1.2% levels respectively, the dominant uncertainties remain of theoretical origin. The theoretical errors are difficult to assign and may not correspond to a Gaussian probability distribution function. High precision tests of HQET, checks on possible violations of quark-hadron duality in semileptonic decays, and experimental determination of  $m_b$ ,  $m_b - m_c$ , and  $\mu_\pi^2$  are needed to complete this challenging experimental program.

## References

1. See the review on “CKM quark-mixing matrix” by F.J. Gilman, K. Kleinknecht, and B. Renk in this *Review*.
2. C. Glenn Boyd, B. Grinstein, and R.F. Lebed, Phys. Lett. **B353**, 306 (1995).
3. P.F. Harrison and H.R. Quinn, editors *The BaBar Physics Book* (1998) and references therein.
4. M. Luke, Phys. Lett. **B252** (1990) 447.
5. A. Czarnecki, Phys. Rev. Lett. **76**, 4124 (1993).
6. A.F. Falk and M. Neubert, Phys. Rev. **D47**, 2965 (1993) and Phys. Rev. **D47**, 2982 (1993).
7. T. Mannel, Phys. Rev. **D50**, 428 (1994).
8. M. Shifman, N.G. Uraltsev, and A. Vainshtein, Phys. Rev. **D51**, 2217 (1995).
9. J.N. Simone *et al.*, Nucl. Phys. Proc. Suppl. **83**, 334 (2000); [hep-ph/0110253](http://hep-ph/0110253).
10. Working Group 1 Summary, CKM Workshop, CERN, CH (2002); <http://ckm-workshop.web.cern.ch/ckm-workshop/>.
11. S. Stone, in *B Decays, 2nd Edition*, S. Stone editor (1994), 283.
12. I. Caprini, L. Lellouch, and M. Neubert, Nucl. Phys. **B530**, 153 (1998);  
C.G. Boyd, B. Grinstein, R.F. Lebed, Phys. Rev. **56**, 6895 (1997).
13. The CLEO Collaboration, LNS Preprint CLNS 01-1776 (2002); [hep-ex/0203032](http://hep-ex/0203032).
14. K. Abe *et al.* (Belle), Phys. Lett. **B526**, 247 (2002).
15. P. Abreu *et al.* (DELPHI), Phys. Lett. **B510**, 55 (2001).
16. D. Buskulic *et al.* (ALEPH), Phys. Lett. **B395**, 373 (1997).
17. G. Abbiendi *et al.* (OPAL), Phys. Lett. **B482**, 15 (2000).
18. S. Anderson *et al.* (CLEO), Nucl. Phys. **A663**, 647 (2000).
19. D. Buskulic *et al.* (ALEPH), Z. Phys. **C73**, 601 (1997).



20. A. Anastassov *et al.* (CLEO), Phys. Rev. Lett. **80**, 4127 (1998).
21. D. Block *et al.* (DELPHI), Contributed Paper to *ICHEP 2000*, DELPHI 2000-106 Conf. 45 (2000).
22. ALEPH, CDF, DELPHI, L3, OPAL, SLD, CERN-EP/2001-050.
23. V. Morenas *et al.*, Phys. Rev. **D56**, 5668 (1997);  
M.Q. Huang and Y.B. Dai, Phys. Rev. **D59**, 34018 (1999);  
M.Oda *et al.*, [hep-ph/0005102](http://hep-ph/0005102) (2000).
24. A.K. Leibovich *et al.*, Phys. Rev. **D57**, 308 (1998);  
Phys. Rev. Lett. **78**, 3995 (1997).
25. LEP  $V_{cb}$  Working Group, Internal Note,  
<http://lepvcb.web.cern.ch/LEPVCB/>.
26. J.E. Duboscq *et al.* (CLEO), Phys. Rev. Lett. **76**, 3898 (1996).
27. M. Neubert, Phys. Lett. **B264**, 455 (1991).
28. Z. Ligeti, Y. Nir, and M. Neubert, Phys. Rev. **D49**, 1302 (1994).
29. K. Abe *et al.* (Belle), Phys. Lett. **B526**, 258 (2002).
30. J. Bartelt *et al.* (CLEO), Phys. Rev. Lett. **82**, 3746 (1999).
31. D. Scora and N. Isgur, Phys. Rev. **D52**, 2783 (1995).
32. S. Hashimoto *et al.*, Phys. Rev. **D61**, 014502 (1999).
33. I. Bigi, M. Shifman, and N. Uraltsev, Ann. Rev. Nuc. Part. Sci. **47**, 591 (1997).
34. A.V. Manohar and M.B. Wise, Phys. Rev. **D49**, 110 (1994);  
A.F. Falk, Lectures presented at TASI-2000,  
[hep-ph/0007339](http://hep-ph/0007339).
35. I.I. Bigi and N. Uraltsev, Int. J. Mod. Phys. **A16**, 5201(2001);  
A. Le Yaouanc *et al.*, Phys. Lett. **B488**, 153 (2000);  
A. Le Yaouanc *et al.*, Phys. Rev. **D62**, 74007 (2000);  
A. Le Yaouanc *et al.*, Phys. Lett. **B517**, 135 (2001).
36. G. Buchalla, [hep-ph/0202092](http://hep-ph/0202092) (2002)  
and references therein.

37. N. Isgur, Phys. Lett. **B448**, 111 (1999).
38. M. Gremm and N. Kapustin, Phys. Rev. **D55**, 6924 (1997).
39. A. Falk, M. Luke, and M.J. Savage, Phys. Rev. **D33**, 2491 (1996).
40. A. Falk and M. Luke, Phys. Rev. **D57**, 424 (1998).
41. S. Chen *et al.*, Phys. Rev. Lett. **87**, 251807 (2001).
42. D. Cronin-Hennessy *et al.* (CLEO), Phys. Rev. Lett. **87**, 251808 (2001).
43. B.C. Barish *et al.* (CLEO), Phys. Rev. Lett. **76**, 1570 (1996);  
S. Henderson *et al.* (CLEO), Phys. Rev. **D45**, 2212 (1992).
44. H. Albrecht *et al.* (ARGUS), Phys. Lett. **B318**, 397 (1993).
45. J.P. Alexander *et al.* (CLEO), Phys. Rev. Lett. **86**, 2737 (2001).
46. LEP/SLD Electroweak Heavy Flavor Results Winter 2001 Conferences;  
<http://lepewwg.web.cern.ch/LEPEWWG/heavy/>.
47. A. Heister *et al.* (ALEPH), Eur. Phys. J. **C22**, 613 (2002).
48. P. Abreu *et al.* (DELPHI), Eur. Phys. J. **C20**, 455 (2001).
49. M. Acciari *et al.* (L3), Z. Phys. **C71**, 379 (1996);  
M. Acciari *et al.* (L3), Eur. Phys. J. **C13**, 47 (2000).
50. G. Abbiendi *et al.* (OPAL), Eur. Phys. J. **C13**, 225 (2000).

---

### **$V_{cb}$ MEASUREMENTS**

For the discussion of  $V_{cb}$  measurements, which is not repeated here, see the review on "Determination of  $|V_{cb}|$ ."

The CKM matrix element  $|V_{cb}|$  can be determined by studying the rate of the semileptonic decay  $B \rightarrow D^{(*)} \ell \nu$  as a function of the recoil kinematics of  $D^{(*)}$  mesons. Taking advantage of theoretical constraints on the normalization and a linear  $\omega$  dependence of the form factors provided by Heavy Quark Effective Theory (HQET), the  $|V_{cb}| \times F(\omega)$  and  $\rho^2$  ( $a^2$ ) can be simultaneously extracted from data, where  $\omega$  is the scalar product of the two-meson four velocities,  $F(1)$  is the form factor at zero recoil ( $\omega=1$ ) and  $\rho^2$  is the slope, sometimes denoted as  $a^2$ . Using the theoretical input of  $F(1)$ , a value of  $|V_{cb}|$  can be obtained.

"OUR EVALUATION" is an average of the data listed below performed by the LEP Heavy Flavor Working Group. The average procedure takes into account correlations between the measurements.

**$|V_{cb}| \times F(1)$  (from  $B^0 \rightarrow D^{*-} \ell^+ \nu$ )**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

**0.0382 ± 0.0012 OUR EVALUATION** with  $\rho^2=1.81 \pm 0.16$  and a correlation 0.86. The fitted  $\chi^2$  is 13 for 10 degrees of freedom.

**0.0351 ± 0.0011 OUR AVERAGE**

0.0354 ± 0.0019 ± 0.0018	<sup>1</sup> ABE	02F BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
0.0355 ± 0.0014 <sup>+0.0023</sup> <sub>-0.0024</sub>	<sup>2</sup> ABREU	01H DLPH	$e^+ e^- \rightarrow Z$
0.0371 ± 0.0010 ± 0.0020	<sup>3</sup> ABBIENDI	00Q OPAL	$e^+ e^- \rightarrow Z$
0.0319 ± 0.0018 ± 0.0019	<sup>4</sup> BUSKULIC	97 ALEP	$e^+ e^- \rightarrow Z$
0.0351 ± 0.0019 ± 0.0020	<sup>5</sup> BARISH	95 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0328 ± 0.0019 ± 0.0022	ACKERSTAFF	97G OPAL	Repl. by ABBIENDI 00Q
0.0350 ± 0.0019 ± 0.0023	<sup>6</sup> ABREU	96P DLPH	Repl. by ABREU 01H
0.0314 ± 0.0023 ± 0.0025	BUSKULIC	95N ALEP	Repl. by BUSKULIC 97

<sup>1</sup> Measured using exclusive  $B^0 \rightarrow D^{*}(892)^- e^+ \nu$  decays with  $\rho^2 = 1.35 \pm 0.17 \pm 0.19$  and a correlation of 0.91.

<sup>2</sup> ABREU 01H measured using about 5000 partial reconstructed  $D^*$  sample with a  $\rho^2 = 1.34 \pm 0.14$ <sup>+0.24</sup><sub>-0.22</sub>.

<sup>3</sup> ABBIENDI 00Q: measured using both inclusively and exclusively reconstructed  $D^{*\pm}$  samples with a  $\rho^2 = 1.21 \pm 0.12 \pm 0.20$ . The statistical and systematic correlations between  $|V_{cb}| \times F(1)$  and  $\rho^2$  are 0.90 and 0.54 respectively.

<sup>4</sup> BUSKULIC 97: measured using exclusively reconstructed  $D^{*\pm}$  with a  $a^2 = 0.31 \pm 0.17 \pm 0.08$ . The statistical correlation is 0.92.

<sup>5</sup> BARISH 95: measured using both exclusive reconstructed  $B^0 \rightarrow D^{*-} \ell^+ \nu$  and  $B^+ \rightarrow D^{*0} \ell^+ \nu$  samples. They report their experiment's uncertainties  $\pm 0.0019 \pm 0.0018 \pm 0.0008$ , where the first error is statistical, the second is systematic, and the third is the uncertainty in the lifetimes. We combine the last two in quadrature.

<sup>6</sup> ABREU 96P: measured using both inclusively and exclusively reconstructed  $D^{*\pm}$  samples.

**$|V_{cb}| \times F(1)$  (from  $B \rightarrow D^- \ell^+ \nu$ )**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

**0.0413 ± 0.0040 OUR EVALUATION** with  $\rho^2=1.19 \pm 0.19$  and a correlation of 0.94. The fitted  $\chi^2$  is 0.4 for 4 degrees of freedom.

**0.039 ± 0.004 OUR AVERAGE**

0.0411 ± 0.0044 ± 0.0052	<sup>7</sup> ABE	02E BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
0.0416 ± 0.0047 ± 0.0037	<sup>8</sup> BARTELT	99 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
0.0278 ± 0.0068 ± 0.0065	<sup>9</sup> BUSKULIC	97 ALEP	$e^+ e^- \rightarrow Z$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0337 ± 0.0044 <sup>+0.0072</sup> <sub>-0.0049</sub>	<sup>10</sup> ATHANAS	97 CLE2	Repl. by BARTELT 99
---	-----------------------	---------	---------------------

<sup>7</sup> Using the missing energy and momentum to extract kinematic information about the undetected neutrino in the  $B^0 \rightarrow D^- \ell^+ \nu$  decay.

<sup>8</sup> BARTELT 99: measured using both exclusive reconstructed  $B^0 \rightarrow D^- \ell^+ \nu$  and  $B^+ \rightarrow D^0 \ell^+ \nu$  samples.

<sup>9</sup> BUSKULIC 97: measured using exclusively reconstructed  $D^\pm$  with a  $a^2 = -0.05 \pm 0.53 \pm 0.38$ . The statistical correlation is 0.99.

<sup>10</sup> ATHANAS 97: measured using both exclusive reconstructed  $B^0 \rightarrow D^- \ell^+ \nu$  and  $B^+ \rightarrow D^0 \ell^+ \nu$  samples with a  $\rho^2 = 0.59 \pm 0.22 \pm 0.12_{-0}^{+0.59}$ . They report their experiment's uncertainties  $\pm 0.0044 \pm 0.0048_{-0.0012}^{+0.0053}$ , where the first error is statistical, the second is systematic, and the third is the uncertainty due to the form factor model variations. We combine the last two in quadrature.

## DETERMINATION OF $|V_{ub}|$

Written April 2002 by M. Battaglia (CERN) and L. Gibbons (Cornell University).

The precise determination of the magnitude of  $|V_{ub}|$  with a robust, well-understood uncertainty remains one of the key goals of the heavy flavor physics programs, both experimentally and theoretically. Because  $|V_{ub}|$ , the smallest element in the CKM mixing matrix, provides a bound on the upper vertex of one of the triangles representing the unitarity property of the CKM matrix, it plays a crucial role in the examination of the unitarity constraints and the fundamental questions on which the constraints can bear [1–2]. Investigation of these issues requires measurements that are precise and that have well-understood uncertainties.

Since the initial observation of the  $b \rightarrow u$  transition by CLEO [3] and ARGUS [4] over a decade ago, we have made great strides both in defining and performing new measurements, and in evaluating the related uncertainties in the extraction of  $|V_{ub}|$ .

The charmless semi-leptonic (S.L.) decay channel  $b \rightarrow u \ell \bar{\nu}$  provides the cleanest path for the determination of  $|V_{ub}|$ . However, the theory for the heavy-to-light  $b \rightarrow u$  transition cannot be as well-constrained as that for the heavy-to-heavy  $b \rightarrow c$  transition used in the determination of  $|V_{cb}|$  [5]. The extraction

of  $|V_{ub}|$  and the interplay between experimental measurements and their theoretical interpretation are further complicated by the large background from  $b \rightarrow c\ell\bar{\nu}$  decay, which has a rate about 60 times higher than that for charmless S.L. decay. Measurements based both on exclusive decay channels and on inclusive techniques have been, and are being, pursued.

We will review the current determinations of  $|V_{ub}|$  by CLEO and the LEP experiments within this overall context, and outline the potential for precise determinations of  $|V_{ub}|$  at the  $B$  factories.

***Exclusive determinations:*** Reconstruction of exclusive  $b \rightarrow u\ell\bar{\nu}$  channels provides powerful kinematic constraints for suppression of the  $b \rightarrow c\ell\bar{\nu}$  background. For this suppression to be effective, an estimate of the four momenta of the undetected neutrino must be provided. The measurements to date have made use of detector hermeticity and the well-determined beam parameters to define a missing momentum that is defined as the neutrino momentum. Signal-to-background ratios (S/B) of order one have been obtained in these channels.

To extract  $|V_{ub}|$  from an exclusive channel, the form factors for that channel must be known. The form factor normalization dominates the uncertainty on  $|V_{ub}|$ . The  $q^2$  dependence of the form factors, which is needed to determine the experimental efficiency, also contributes to the uncertainty, but at a much reduced level. For example, the requirement of a stiff lepton for background reduction in these analyses introduces a  $q^2$  dependence to the efficiency. In the limit of a massless charged lepton (a reasonable limit for the electron and muon decay channels), the  $B \rightarrow \pi\ell\nu$  decay depends on one form factor  $f_1(q^2)$ :

$$\frac{d\Gamma(B^0 \rightarrow \pi^- \ell^+ \nu)}{dy d \cos \theta_\ell} = |V_{ub}|^2 \frac{G_F^2 p_\pi^3 M_B^2}{32\pi^3} \sin^2 \theta_\ell |f_1(q^2)|^2, \quad (1)$$

where  $y = q^2/M_B^2$ , and  $\theta_\ell$  is the angle between the charged lepton direction in the virtual  $W$  ( $\ell + \nu$ ) rest frame and the direction of the virtual  $W$ . For the vector meson final states  $\rho$  and  $\omega$ , three form factors,  $A_1$ ,  $A_2$  and  $V$ , are necessary (see *e.g.*, Ref. 6).

Calculation of these form factors constitutes a considerable theoretical industry, with a variety of techniques now being employed. Form factors based on lattice calculations [11–23], and on light cone sum rules [24–32], currently have uncertainties in the 15% to 20% range. A variety of quark model calculations exist [33–47]. Finally, a number of other approaches [48–53], such as dispersive bounds and experimentally-constrained models based on Heavy Quark Symmetry, seek to improve the  $q^2$  range over which the form factors can be estimated without introduction of a significant model dependence. Unfortunately, all these calculations currently have contributions to the uncertainty that remain uncontrolled. The light cone sum rules calculations assume quark-hadron duality, offering a “canonical” contribution to the uncertainty of 10%, but with no known means of rigorously limiting that uncertainty. The lattice calculations to date remain in the “quenched” approximation (no light quark loops in the propagators), which limits the ultimate precision to the 15% to 20% range. For the quark model calculations, there exists no means for systematic evaluation of the uncertainties.

There have been two exclusive  $|V_{ub}|$  analyses by the CLEO Collaboration: a simultaneous measurement of the  $B \rightarrow \pi \ell \bar{\nu}$  and the  $B \rightarrow \rho \ell \bar{\nu}$  transitions [9], and a second measurement of the  $B \rightarrow \rho \ell \bar{\nu}$  rate [10]. Both measurements employ the missing energy and momentum to estimate the neutrino momentum. With that technique, the major background results from  $b \rightarrow$

$c\bar{\ell}\bar{\nu}$  decays in events that cannot be properly reconstructed (for example, because of additional neutrinos in the event). Both measurements also employ the isospin relations

$$\Gamma(B^0 \rightarrow \pi^- \ell^+ \nu) = 2\Gamma(B^+ \rightarrow \pi^0 \ell^+ \nu)$$

and

$$\Gamma(B^0 \rightarrow \rho^- \ell^+ \nu) = 2\Gamma(B^+ \rightarrow \rho^0 \ell^+ \nu) \quad (2)$$

to combine the charged and neutral decays. In the original method, strict event quality requirements were made that resulted in a low efficiency, but a relatively low background to signal ratio over a fairly broad lepton momentum range. The branching fractions obtained were

$$B(B^0 \rightarrow \pi^- \ell^+ \nu) = (1.8 \pm 0.4 \pm 0.3 \pm 0.2) \times 10^{-4}$$

and

$$B(B^0 \rightarrow \rho^- \ell^+ \nu) = (2.5 \pm 0.4_{-0.7}^{+0.5} \pm 0.5) \times 10^{-4} . \quad (3)$$

The second analysis loosened the event cleanliness requirements, resulting in a much higher efficiency. The efficiency gain comes at the price of an increased background, and the analysis was primarily sensitive to signal with lepton momenta above 2.3 GeV/ $c$ , which is near (and beyond) the kinematic endpoint for  $b \rightarrow c\bar{\ell}\bar{\nu}$  decays, which are therefore highly suppressed. This analysis obtained

$$B(B^0 \rightarrow \rho^- \ell^+ \nu) = (2.69 \pm 0.41_{-0.40}^{+0.35} \pm 0.50) \times 10^{-4} . \quad (4)$$

The results of the two analyses are largely statistically independent, and they have been combined, accounting for correlated uncertainties, to obtain:

$$|V_{ub}| = (3.25 \pm 0.14_{-0.29}^{+0.21} \pm 0.55) \times 10^{-3} , \quad (5)$$

where the errors arise from statistical, experimental systematic, and form factor uncertainties, respectively. The last term has been estimated by comparing a large number of available models and, for this average, the earlier analysis was updated to the set of models used in the later analysis. A potential non-resonant  $\pi\pi\ell\nu$  contribution (assumed to be zero in the analyses) results in the asymmetric systematic uncertainty. The model dependence uncertainty is dominated by the overall normalization. Evaluation of the systematic considered both the spread among individual models and calculations, as well as the uncertainties claimed for the calculations. The central value spread and the estimated uncertainties were of the same order (roughly 15%).

The  $B$  factories have recently released very preliminary results of their first analyses of these exclusive modes. Belle has produced a  $B \rightarrow \pi\ell\nu$  analysis [54] that is very similar to the original CLEO exclusive analysis [9]. They find

$$B(B^0 \rightarrow \pi^- \ell^+ \nu) = (1.28 \pm 0.20 \pm 0.26 \pm \sigma_{\text{model}}) \times 10^{-4} . \quad (6)$$

BABAR has recently presented preliminary results for a  $B \rightarrow \rho\ell\nu$  analysis [55] that is quite similar to the second CLEO analysis [10], for which they obtained

$$B(B^0 \rightarrow \rho^- \ell^+ \nu) = (2.97 \pm 0.56_{-0.56}^{+0.48} \pm \sigma_{\text{model}}) \times 10^{-4} . \quad (7)$$

Both experiments use neutrino reconstruction, but have not yet advanced the detailed event cleanup (see Ref. 56) to the level of CLEO. The uncertainties, which are comparable to the original CLEO errors despite the much larger integrated luminosity, reflect this preliminary situation. This situation will certainly improve as the experiments mature.



The future for exclusive determinations of  $|V_{ub}|$  appears promising. Unquenched lattice calculations begin to be feasible, and this will eliminate the primary source of uncontrolled uncertainty in these calculations. Simultaneously, the  $B$  factories are performing very well, and very large samples of events in which one  $B$  meson has been fully reconstructed will be available. This will allow a more robust determination of the neutrino momentum, and should allow a significant reduction of backgrounds and experimental systematic uncertainties. The high statistics should also allow detailed measurements of  $d\Gamma/dq^2$ , which will provide a sorely-needed litmus test for the form factor calculations, and will make the form factor shape contribution to the uncertainty on  $|V_{ub}|$  negligible. Should theory allow use of the full range of  $q^2$  in the extraction of  $|V_{ub}|$ , the  $B$  factories have already logged data sufficient for a 5% statistical determination of  $|V_{ub}|$ . If the data must be restricted to low hadronic recoil momentum (large  $q^2$ ), an order of magnitude more data would be necessary.

For both lattice and the  $B$  factories,  $\pi\ell\nu$  appears to be a golden mode for future precise determination of  $|V_{ub}|$ . The one caveat is management of contributions from the  $B^*$  pole, but recent work [21] suggests that this problem can be successfully overcome.  $B \rightarrow \eta\ell\nu$  will provide a valuable cross-check. The  $\rho\ell\nu$  mode will be more problematic for high precision: the broad width introduces both experimental and theoretical difficulties. Experiments must, for example, assess potential nonresonant  $\pi\pi$  contributions, but only crude arguments based on isospin and quark-popping have been brought to bear to date. Theoretically, no calculation, including lattice, has dealt with the width of the  $\rho$ . Even worse, when the lattice calculations become unquenched, the  $\rho$  will become unstable, and the  $\pi\pi$  final

state must be faced by the calculations. The methodology for accommodation of two particle final states on the lattice remains quite primitive, very costly, and works only for low-energy states, so it may be unsuitable for  $\rho$  decay. One could put the  $\rho$  “in a box” to prevent its decay, but this introduces uncertainties of order width/mass [57]. Fortunately, the  $\omega\ell\nu$  mode provides an excellent alternative to the  $\rho$  mode, though it has remained elusive to date. Agreement between accurate  $|V_{ub}|$  determinations from  $\pi\ell\nu$  and from  $\omega\ell\nu$  will provide added confidence in both.

***Inclusive determinations:*** In principle, the fully inclusive rate  $\text{BR}(b \rightarrow X_u\ell\bar{\nu})$  can be calculated quite reliably within the OPE framework [58–61], with a  $\simeq 6\%$  theoretical uncertainty in  $|V_{ub}|$  attainable. The calculations find

$$|V_{ub}| = 0.00445 \times \left( \frac{\text{B}(b \rightarrow u\ell\bar{\nu})}{0.002} \frac{1.55\text{ps}}{\tau_b} \right)^{1/2} \times (1 \pm 0.020 \pm 0.052), \quad (8)$$

where the first error arises from uncertainties in the OPE expansion, and the second from uncertainty in the  $b$  quark mass, for which  $m_b^{\text{kin}}(1 \text{ GeV}) = (4.58 \pm 0.09) \text{ GeV}$  has been assumed. With the large number of final states available over a broad mass range, deviations from global quark-hadron duality are expected to be small for the total charmless S.L. rate.

Unfortunately, realizing this accuracy is extremely difficult in practice. The background from  $b \rightarrow c\ell\bar{\nu}$  decays forces experiments to limit their sensitivity within some restricted region of the total phase space. These regions are the lepton energy endpoint  $E_\ell > (M_B^2 - M_D^2)/(2M_B)$ , the low hadronic mass region  $M_X < M_D$ , and the large dilepton mass  $q^2 > (M_B - M_D)^2$ . They select  $\simeq 15\%$ ,  $70\%$ , and  $20\%$  of the charmless S.L. rate, respectively. The typical S/B ratios achieved within these regions

are a factor of 5–10 smaller than those for the exclusive analyses. These restrictions introduce additional uncertainties in the calculation of the total charmless branching fraction that are difficult to quantify. In addition, they may end up moving the quark-duality assumption away from the well-grounded global assumption towards a local assumption. While the limitations of the quark-hadron duality assumption are expected to be quite small for fully integrated rates, they may become more pronounced in partially integrated rates. The applicability of general results for the inclusive OPE approach, and the control of the theoretical uncertainties for measurements restricted within these limited regions, remain the subject of ongoing debate within the community.

The original observations of the  $b \rightarrow u$  transition at the  $\Upsilon(4S)$  [3,4] were inclusive analyses that focused on leptons in the endpoint region of the single lepton spectrum, beyond the kinematical limit for compatibility with  $b \rightarrow c\ell\bar{\nu}$  transitions. The ACCMM [7] and ISGW [35] models were used to estimate the rate into the endpoint, from which  $|V_{ub}/V_{cb}| = (0.08 \pm 0.02)$  was obtained, where the 25% error is dominated by the theoretical uncertainty. That the error “guesstimated” with those models happened to give an error that was reasonably appropriate was a historical accident. Had the ISGW II model [36] been available at that time (and used), the model dependence would have been significantly underestimated [8]. The theoretical uncertainty for an endpoint analysis has been very difficult to quantify. Because the endpoint region extends beyond the partonic endpoint, and the size of the endpoint is of order  $\Lambda_{\text{QCD}}$ , an infinite series of terms in the OPE rate calculation become equally important. The leading twist singularities can be resummed into a structure function [75–78]. This shape

function encapsulates the “Fermi motion” of the  $b$  quark within the hadron, and must be evaluated when experimental determinations of  $|V_{ub}|$  are forced near the boundary of a kinematic distribution [73].

A number of authors observed that selection of S.L. decays  $b \rightarrow X\ell\bar{\nu}$ , with hadronic mass  $M_X$  below that of the  $D$  meson, provides a separation of the charmless  $X_u\ell\bar{\nu}$  signal from the  $X_c\ell\bar{\nu}$  background, with an efficiency that can be reliably estimated [62,68,69,73]. These observations motivated an intense effort into such inclusive analyses at LEP. The significant  $B$ -hadron boost, and the containment of its decay products into a narrow jet in  $Z^0 \rightarrow b\bar{b}$  events, make these measurements at the  $Z^0$  pole interestingly complementary to those performed at the  $\Upsilon(4S)$  peak. Over the past several years, the ALEPH [63], DELPHI [64], L3 [65], and, most recently, OPAL [66] collaborations, have published inclusive measurements of the  $b \rightarrow u\ell\bar{\nu}$  rate. Three separate methods have been employed. The ALEPH and OPAL analyses use neural networks that take as input a large number (twenty in the case of ALEPH, and seven in the OPAL analysis) of kinematic variables which provide separation between  $b \rightarrow c\ell\nu$  and  $b \rightarrow u\ell\nu$  decays. The signal is extracted in both cases from a fit to the network output discriminant, restricted to a region enriched in signal decays. L3 applies a sequential-cut analysis, using the kinematics of the lepton and of the leading hadron in the same jet for discrimination of the signal events. The DELPHI analysis performs an inclusive reconstruction of the hadronic mass of the system emitted together with the lepton in the  $b$ -hadron decay. The S.L.  $B$  sample is split into  $b \rightarrow u$ -enriched and -depleted samples, based on the separation between tertiary and secondary vertices (making use of the

finite charm lifetime), and on the presence of tagged kaons in the final state. The mass of the hadronic system  $M_X$  is used to further subdivide the sample into a  $b \rightarrow X_u \ell \nu$ -favored region ( $M_X < 1.6$  GeV) and a  $b \rightarrow X_c \ell \nu$ -dominated region. The signal is extracted from a simultaneous fit to the number of decays classified according to the four different categories and the distributions of the lepton energy in the reconstructed  $B$ -rest frame. While the approaches of the various LEP analyses differ, they tend to be sensitive to  $b \rightarrow u \ell \bar{\nu}$ , primarily when the mass of the hadronic system is in the region  $M_X \lesssim M_D$ : DELPHI explicitly so, ALEPH and OPAL implicitly, in that after their neural net requirement, the efficiency falls noticeably with increasing hadronic mass (they gain some additional sensitivity at higher masses when stiff leptons or hadrons are present in the event). These analyses are sensitive to a significantly larger portion of the phase space than the endpoint analyses, but at the cost of larger backgrounds from  $b \rightarrow c \ell \bar{\nu}$  decays (see Table 1).

The uncertainty in the determination of the fraction of the total charmless rate selected by a given cut in  $M_X$  has been studied by several authors. A major source of uncertainty is represented by the value of the  $b$ -quark mass  $m_b$ . A relative error of  $\pm 15$ – $30\%$  on the inclusive charmless branching fraction, obtained with  $M_X$  cut values from  $M_D$  down to 1.5 GeV, has been estimated from the uncertainty on the mass (assuming  $\pm 150$  MeV) and the kinetic energy of the  $b$  quark [73]. Other estimates are compatible with this range [68,69]. For restrictions in the range  $M_X^2 < m_b \Lambda_{\text{QCD}} \approx M_D^2$ , the calculation of the inclusive charmless branching fraction is also sensitive to the shape function uncertainties that affect the endpoint region, particularly as the cut is lowered much below

**Table 1:** Summary of  $|V_{ub}|$  determinations by experiment. The method, the S/B ratio of the analyses, the result with the statistical+experimental, the  $b \rightarrow c$  and the published  $b \rightarrow u$  uncertainties, the fractional systematic uncertainty for the non  $b \rightarrow u$  contributions, and our evaluation for the range of the estimated theoretical uncertainty (including uncertainty in the shape functions from  $b \rightarrow s\gamma$ ) are given.

Exp.	Method	S/B	$ V_{ub} $ [ $10^{-3}$ ]	$\sigma_{b \rightarrow c}$ ( $ V_{ub} $ )	$\sigma_{th}$ ( $ V_{ub} $ )
ALEPH	Neural Net	0.07	$4.12 \pm .67 \pm .62 \pm 0.35$	15%	9%
OPAL	Neural Net	0.05	$4.00 \pm .71 \pm .59 \pm 0.40$	15%	10%
DELPHI	$M_X$	0.10	$4.07 \pm .65 \pm .47 \pm 0.39$	12%	10%
L3	$\pi - \ell$ Cut	0.22	$5.7 \pm 1.0 \pm 1.3 \pm 0.5$	22%	10%
<b>LEP</b>	<b>Average</b>		<b><math>4.09 \pm 0.37 \pm 0.44 \pm 0.34</math></b>		<b>9–15%</b>
<b>CLEO</b>	<b><math>E_\ell</math> endpoint</b>	<b>0.39</b>	<b><math>4.12 \pm 0.34 \pm 0.44 \pm 0.33</math></b>	<b>7%</b>	<b>10–15%</b>
CLEO	$\pi(\rho)\ell\bar{\nu}+$ strict $\nu$ -rec	2.1 (0.6)	$3.30 \pm 0.4 \pm 0.7$	8%	
CLEO	$\rho\ell\bar{\nu}+$ loose $\nu$ -rec	0.7–1.5	$3.23 \pm 0.35 \pm 0.58$	5%	
<b>CLEO</b>	<b><math>\pi + \rho\ell\bar{\nu}</math></b> <b>Average</b>		<b><math>3.25^{+0.25}_{-0.32} \pm 0.55</math></b>		<b>15–20%</b>

$M_D$  [68,69,73](though model-dependent studies suggest that the importance of the shape function may be reduced in the  $M_X$  case). Higher-twist contributions and unknown power corrections of order  $\Lambda_{\text{QCD}}/M_B \approx 10\%$  (for example, corrections to the method of convolution of the parton-level spectra with a shape function [70,71]) also contribute to the uncertainty on  $|V_{ub}|$ . This leads to an estimate of the overall  $b \rightarrow u$  systematics on  $|V_{ub}|$  extracted with these methods of order 10–15%, which still allows for a largely model-independent determination.

The DELPHI analysis follows this framework with the theoretical uncertainties evaluated within the framework outlined, for example, in Ref. 69. At that time, however, no detailed information regarding the shape function existed, and the experiment relied on models. The neural net analyses are somewhat more difficult to interpret directly, and the experiments rely more heavily on model estimates to gauge the uncertainty. Given that those measurements tend to have sensitivity primarily in the regions affected by the shape functions, a theoretical uncertainty within the 10%–15% window seems likely. An average by the LEP Heavy Flavour Group [72] results in

$$V_{ub} = (4.09_{-0.39-0.47}^{+0.36+0.42} \pm 0.25 \pm 0.23) \times 10^{-3} . \quad (9)$$

The first error includes the statistical and detector-level systematic uncertainties, and the second the systematics from the  $b \rightarrow c$  background. The third error reflects the uncertainty in the extrapolation of the yields measured in the restricted kinematic region to the total charmless S.L. branching fraction determination. The above was obtained based on model studies for each analysis, accounting for the contributions that could be quantified. It profits from the partially uncorrelated sources of systematics that result from the different techniques adopted by the four experiments. The discussion above suggests the more conservative range of  $\pm 10$ – $15\%$ . The final error reflects the  $\pm 6\%$  uncertainty for extraction of  $|V_{ub}|$  from the total charmless S.L. branching fraction.

Observation of semileptonic  $b \rightarrow u$  decays at LEP has been an experimental tour de force. The successful realization of those analyses is due to some advantages offered by the kinematics at the  $Z^0$  pole, and to the performance of the detectors. While these studies have pioneered a path towards

new approaches for extracting  $|V_{ub}|$ , they have exposed the drawbacks of analyses with S/B ratios that require control of the background level to better than 5% of itself. Some areas of concern here, discussed within the community, include the modeling uncertainties of the non- $D$  and  $D^*$  components of the background from  $B$  decay, the modeling of the  $B_s$  and  $b$ -baryon S.L. decays, and the estimate of the  $b \rightarrow u$  modeling uncertainties due to the uneven sampling of the decay phase space.

CLEO has recently submitted for publication [74] a new measurement based on the lepton endpoint fraction that makes a significant step away from reliance on models for the theoretical uncertainty. It has been known for some time [77,76] that, at leading twist, the same shape function corrects the parton level  $b \rightarrow s\gamma$  photon spectrum and the  $b \rightarrow u\ell\nu$  lepton spectrum. While measurements of the non-perturbative  $1/m_b^2$  parameters have been made for the heavy-to-heavy  $b \rightarrow c\ell\bar{\nu}$  transition [79], differences in the higher order corrections in the OPE spoil their application to heavy-to-light transitions. Since both  $b \rightarrow s\gamma$  and  $b \rightarrow u\ell\bar{\nu}$  are heavy-to-light transitions, however, one can relate the parameters determined in one system to the other, up to power corrections of order  $\Lambda_{\text{QCD}}/M_B$  arising from nonlocal operators [71]. It has been suggested that the stability of the extracted  $|V_{ub}|$  under variation of the lepton momentum endpoint region can limit the uncertainty due to these corrections [71].

Because the CLEO experiment must account for the distortion of the endpoint spectrum due to the motion of the  $B$  mesons, initial-state radiation, and experimental resolution, it fits the observed data using a theoretical momentum spectrum



to which the distortions are applied. Several ansatz [82,83] for the form of the shape function were employed. CLEO finds

$$|V_{ub}| = (4.12 \pm 0.34 \pm 0.44 \pm 0.23 \pm 0.24) \times 10^{-3}, \quad (10)$$

based on the lepton momentum range 2.2–2.6 GeV/ $c$ . The first error is the combined statistical and experimental uncertainty on the rate into the accepted momentum range. The second error is the uncertainty on the fraction of leptons expected to lie within this range based on the uncertainty, statistical and systematic, in the parameters derived from the  $b \rightarrow s\gamma$  photon spectrum. The third uncertainty is the same HQE uncertainty on the extraction of  $|V_{ub}|$  from the total rate as above. The final error is an estimate of the scale of the uncertainty that results from the unknown power corrections in applying the  $b \rightarrow s\gamma$  shape function to  $b \rightarrow ul\nu$ . In the limit of integration over the full lepton spectrum, this uncertainty would vanish: in fact, as one moves away from the phase space boundary, the importance of the shape function diminishes. To evaluate this uncertainty, the parameters of the shape function were varied by the expected order of the corrections:  $\Lambda_{\text{QCD}}/M_B \approx 10\%$ . This sets the *scale* of the uncertainty, but is not a precise evaluation of the uncertainty—we do not know if the true uncertainty is a factor of two larger or smaller, for example. Variation of the size of the endpoint region results in consistent determinations of  $|V_{ub}|$ , but the experimental uncertainties limit our ability to make more precise statements regarding the power corrections at this time.

Finally, a method of extracting  $|V_{ub}|$  inclusively in a restricted region of  $q^2$ , the mass of the leptonic system, has been proposed [84]. This region is free from uncertainty due to the shape functions, but does receive order  $1/m_c^3$  corrections [85].

Given the orthogonal uncertainties, this method will provide a valuable crosscheck to other inclusive determinations of  $|V_{ub}|$ . While this method has been found to be unsuitable for experiments at higher energy, application at the  $\Upsilon(4S)$  facilities should be feasible, where resolutions on  $q^2$  of approximately 1  $\text{GeV}^2$  can already be obtained, though with large tails from mis-reconstructed events.

As the  $B$  factories bring their full data samples to bear on inclusive measurements of  $|V_{ub}|$ , there is a potential for considerable progress in  $|V_{ub}|$ . More precise evaluation of the  $b \rightarrow s\gamma$  photon spectrum will lead to more precisely determined effective shape functions. With the potential for large samples of events with one fully reconstructed  $B$ , reconstruction of the hadronic recoil mass with much reduced contamination from  $b \rightarrow c\ell\bar{\nu}$  decays and of  $q^2$  should be possible. Hence, statistical and systematic experimental uncertainties should be reduced as well. As long as the various determinations remain in agreement, while their precision improves, we can enhance our confidence that the uncontrolled theoretical uncertainties are not biasing  $|V_{ub}|$  beyond whatever level of precision has been reached in the individual measurements.

**Summary and outlook:** There is considerable debate (even between the authors) regarding the soundest use of the various measurements. While our knowledge regarding  $|V_{ub}|$  is far more robust than it was ten years ago, the uncertainty on  $|V_{ub}|$  from each method still receives contributions from some uncontrolled sources. To validate a given level of precision in this situation, measurements based on complementary techniques that agree within that precision are needed. The present results from inclusive and exclusive determinations display a promising agreement (see Table 1). However, there is a fairly widespread

consensus that the inclusive and exclusive measurements cannot be reliably combined until we can quantify all of their yet-uncontrolled uncertainties.

Restricting to the inclusive determinations, several results have been already obtained, all within the same HQE framework, and with comparable estimated accuracies. Their averaging would improve the overall  $|V_{ub}|$  accuracy, since the uncorrelated uncertainties are sizeable. The LEP Heavy Flavour Group has already performed the exercise of such averaging for the four LEP measurements. The result has been obtained in the framework of OPE, and no additional error has been added to account for additional corrections to the  $1/m_b$  expansion, and of the quark-hadron duality assumption [72]. These issues need to be addressed and tested by further experimental studies.

As discussed above, there remain uncertainties that cannot be precisely quantified in the different analyses. Furthermore, potential violation of local quark-hadron duality might affect each kinematic region differently, though future study of the end-point spectrum for  $B_u$  and  $B_d$  separately may help in constraining these effects. Hence, there exist sources of uncertainty that we cannot yet quantify that may be as large as the current statistical uncertainties, though, in the end, they could also be found to be small. The agreement among the current results limits the uncontrolled uncertainties to the order of 20%, the current precision of the comparison. Without some independent means of controlling such uncertainties, an average of the different methods may underestimate the uncertainty with which we have truly determined  $|V_{ub}|$ .

Currently, we have a variety of measurements that individually approach a 15% uncertainty, and that all agree within that uncertainty. Any of the individual measurements can, therefore, be used as representative of  $|V_{ub}|$ .

The prospects for improved precision on  $|V_{ub}|$  are excellent. With the large data samples now becoming available at the  $\Upsilon(4S)$  facilities, the experimental uncertainties will continue to shrink. With these improved uncertainties will come more stringent comparisons of techniques to “stress test” the theory, and either continued confidence in the increased precision on  $|V_{ub}|$ , or an indication of where the shortcomings lie. With the continued advancement of lattice QCD, exclusive determinations of  $|V_{ub}|$  from  $B \rightarrow \pi \ell \nu$  and  $B \rightarrow \omega \ell \nu$  well below 10%, appear feasible within the decade. We now have event samples that allow model independent extractions of  $|V_{ub}|$  from a variety of inclusive techniques. As the event samples from the  $B$  factories increase, the precision of these techniques will continue to improve, and agreement among them can limit the uncontrolled uncertainties. Barring disagreements as the measurements improve, sub 10% precisions also appear feasible from the inclusive techniques.

## References

1. See the review on “CKM Quark Mixing Matrix” by F.J. Gilman, K. Kleinknecht, and B. Renk in this *Review*.
2. See the review on “ $CP$  Violation in  $B$  decays” by H. Quinn and A. Sanda in this *Review*.
3. R. Fulton *et al.* (CLEO Collab.), Phys. Rev. Lett. **64**, 16, (1990);  
J. Bartelt *et al.*, Phys. Rev. Lett. **71**, 4111 (1993).
4. H. Albrecht *et al.* (ARGUS Collab.), Phys. Lett. **B234**, 409 (1990); and Phys. Lett. **B255**, 297 (1991).
5. See the review on “Determination of  $|V_{cb}|$ ” by M. Artuso and E. Barberio in this *Review*.
6. F.J. Gilman and R.L. Singleton, Phys. Rev. **D41**, 142 (1990).
7. G. Altarelli *et al.*, Nucl. Phys. **B208**, 365 (1982).

8. R.A. Poling, private communication.
9. J.P. Alexander *et al.* (CLEO Collab.), Phys. Rev. Lett. **77**, 5000 (1996).
10. B.H. Behrens *et al.* (CLEO Collab.), Phys. Rev. **D61**, 052001 (2000).
11. A. Abada *et al.*, Nucl. Phys. **B416**, 675 (1994).
12. C.R. Allton *et al.* (APE Collab.), Phys. Lett. **B345**, 513 (1995).
13. L. Del Debbio *et al.* (UKQCD Collab.), Phys. Lett. **B416**, 392 (1998).
14. S. Hashimoto *et al.*, Phys. Rev. **D58**, 014502 (1998).
15. S. Ryan *et al.*, Nucl. Phys. Proc. Suppl. **73**, 390 (1999).
16. S.M. Ryan *et al.*, Nucl. Phys. Proc. Suppl. **83**, 328 (2000).
17. L. Lellouch, [[arXiv:hep-ph/9912353](https://arxiv.org/abs/hep-ph/9912353)].
18. K.C. Bowler *et al.* (UKQCD Collab.), Phys. Lett. **B486**, 111 (2000).
19. D. Becirevic and A.B. Kaidalov, Phys. Lett. **B478**, 417 (2000).
20. S. Aoki *et al.* (JLQCD Collab.), Nucl. Phys. Proc. Suppl. **94**, 329 (2001).
21. A.X. El-Khadra *et al.*, Phys. Rev. **D64**, 014502 (2001).
22. S. Aoki *et al.* (JLQCD Collab.), Phys. Rev. **D64**, 114505 (2001).
23. A. Abada *et al.*, Nucl. Phys. **B619**, 565 (2001).
24. P. Ball and V.M. Braun, Phys. Rev. **D55**, 5561 (1997).
25. P. Ball and V.M. Braun, Phys. Rev. **D58**, 094016 (1998).
26. A. Khodjamirian *et al.*, Phys. Lett. **B410**, 275 (1997).
27. A. Khodjamirian *et al.*, Phys. Rev. **D62**, 114002 (2000).
28. A.P. Bakulev, S.V. Mikhailov, and R. Ruskov, [[arXiv:hep-ph/0006216](https://arxiv.org/abs/hep-ph/0006216)].
29. T. Huang, Z. Li, and X. Wu, [[arXiv:hep-ph/0011161](https://arxiv.org/abs/hep-ph/0011161)].
30. W.Y. Wang and Y.L. Wu, Phys. Lett. **B515**, 57 (2001).
31. W.Y. Wang and Y.L. Wu, Phys. Lett. **B519**, 219 (2001).
32. P. Ball and R. Zwicky, JHEP **0110**, 019 (2001).

33. M. Wirbel, B. Stech, and M. Bauer, Z. Phys. **C29**, 637 (1985).
34. J.G. Korner and G.A. Schuler, Z. Phys. **C38**, 511 (1988) [Erratum: *ibid.* **C41**, 690 (1988)].
35. N. Isgur *et al.*, Phys. Rev. **D39**, 799 (1989).
36. D. Scora and N. Isgur, Phys. Rev. **D52**, 2783 (1995).
37. D. Melikhov, Phys. Rev. **D53**, 2460 (1996).
38. M. Beyer and D. Melikhov, Phys. Lett. **B436**, 344 (1998).
39. R.N. Faustov, V.O. Galkin, and A.Y. Mishurov, Phys. Rev. **D53**, 6302 (1996).
40. N.B. Demchuk *et al.*, Phys. Atom. Nucl. **60**, 1292 (1997) [Yad. Fiz. **60N8**, 1429 (1997)].
41. I.L. Grach, I.M. Narodetsky, and S. Simula, Phys. Lett. **B385**, 317 (1996).
42. Riazuddin, T.A. Al-Aithan, and A.H. Gilani, [arXiv:hep-ph/0007164].
43. D. Melikhov and B. Stech, Phys. Rev. **D62**, 014006 (2000).
44. T. Feldmann and P. Kroll, Eur. Phys. J. C **12**, 99 (2000).
45. J.M. Flynn and J. Nieves, Phys. Lett. **B505**, 82 (2001).
46. M. Beneke and T. Feldmann, Nucl. Phys. **B592**, 3 (2001).
47. H.M. Choi and C.R. Ji, Phys. Lett. **B460**, 461 (1999).
48. T. Kurimoto, H. Li, and A.I. Sanda, Phys. Rev. **D65**, 014007 (2002).
49. Z. Ligeti and M.B. Wise, Phys. Rev. **D53**, 4937 (1996).
50. E.M. Aitala *et al.* (E791 Collab.), Phys. Rev. Lett. **80**, 1393 (1998).
51. G. Burdman and J. Kambor, Phys. Rev. **D55**, 2817 (1997).
52. L. Lellouch, Nucl. Phys. **B479**, 353 (1996).
53. T. Mannel and B. Postler, Nucl. Phys. **B535**, 372 (1998).
54. K. Abe *et al.* (BELLE Collab.), BELLE-CONF-0124, 2001.
55. B. Serfass, talk presented at "Workshop on the CKM Unitarity Triangle," CERN, Geneva, Feb., 2002.
56. L. Gibbons, Ann. Rev. Nucl. Sci. **48**, 121 (1998).
57. G.P. Lepage, private communication.

58. I.I. Bigi, M.A. Shifman, and N. Uraltsev, Ann. Rev. Nucl. Part. Sci. **47**, 591 (1997).
59. A.H. Hoang, Z. Ligeti, and A.V. Manohar, Phys. Rev. Lett. **82**, 277 (1999).
60. I.I. Bigi, UND-HEP-BIG-99-05, [hep-ph/9907270](http://arxiv.org/abs/hep-ph/9907270).
61. Z. Ligeti, FERMILAB-Conf-99/213-T, [hep-ph/9908432](http://arxiv.org/abs/hep-ph/9908432).
62. V. Barger, C.S. Kim, and R.J.N. Phillips, Phys. Lett. **B251**, 629 (1990).
63. R. Barate *et al.* (ALEPH Collab.), Eur. Phys. J. **C6**, 555 (1999).
64. P. Abreu *et al.* (DELPHI Collab.), Phys. Lett. **B478**, 14 (2000).
65. M. Acciarri *et al.* (L3 Collab.), Phys. Lett. **B436**, 174 (1998).
66. G. Abbiendi *et al.* (OPAL Collab.), Eur. Phys. J. **C21** 399 (2001).
67. R.D. Dikeman, M. Shifman, and N.G. Uraltsev, Int. J. Mod. Phys. **A11**, 571 (1996).
68. A.F. Falk, Z. Ligeti, and M.B. Wise, Phys. Lett. **B406**, 225 (1997).
69. I.I. Bigi, R.D. Dikeman, and N. Uraltsev, Eur. Phys. J. **C4**, 453 (1998).
70. A.K. Leibovich, in *Proc. of the 5th International Symposium on Radiative Corrections (RADCOR 2000)* ed. Howard E. Haber, [[arXiv:hep-ph/0011181](http://arxiv.org/abs/hep-ph/0011181)].
71. M. Neubert, Phys. Lett. **B513**, 88 (2001).
72. The LEP VUB Working Group, Note LEPVUB-01/01.
73. F. De Fazio and M. Neubert, JHEP **9906**, 017 (1999).
74. A. Bornheim *et al.* (CLEO Collab.), CLNS 01/1767, [[arXiv:hep-ex/0202019](http://arxiv.org/abs/hep-ex/0202019)].
75. M. Neubert, Phys. Rev. **D49**, 3392 (1994).
76. I. Bigi *et al.*, Int. J. Mod. Phys. **A9**, 2467 (1994).
77. M. Neubert, Phys. Rev. **D49**, 4623 (1994).
78. U. Aglietti and G. Ricciardi, Nucl. Phys. **B587**, 363 (2000).

79. D. Cronin-Hennessy *et al.* (CLEO Collab.), Phys. Rev. Lett. **87**, 251808 (2001).
80. A.K. Leibovich, I. Low, I.Z. Rothstein, Phys. Rev. **D61**, 053006 (2000).
81. A.K. Leibovich, I. Low, and I.Z. Rothstein, Phys. Lett. **B513**, 83 (2001).
82. I.I. Bigi *et al.*, Phys. Lett. **B328**, 431 (1994).
83. A.L. Kagan and M. Neubert, Eur. Phys. J. **C7**, 5 (1999).
84. C.W. Bauer, Z. Ligeti, and M. Luke, Phys. Lett. **B479**, 395 (2000).
85. M. Neubert, JHEP **0007**, 022 (2000). M. Neubert and T. Becher, CLNS-01-1737, [hep-ph/0105217](http://hep-ph/0105217).
86. A.K. Leibovich, I. Low, and I.Z. Rothstein, Phys. Lett. **B486**, 86 (2000).

### $V_{ub}$ MEASUREMENTS

For the discussion of  $V_{ub}$  measurements, which is not repeated here, see the review on "Determination of  $|V_{ub}|$ ."

The CKM matrix element  $|V_{ub}|$  can be determined by studying the rate of the charmless semileptonic decay  $b \rightarrow u \ell \nu$ . Measurements based on on exclusive decay channels and on inclusive techniques can be found in the previous  $B$  Listings, which will not repeat here.

### $V_{cb}$ and $V_{ub}$ CKM Matrix Elements REFERENCES

ABE	02E	PL B526 258	K. Abe <i>et al.</i>	(Belle Collab.)
ABE	02F	PL B526 247	K. Abe <i>et al.</i>	(Belle Collab.)
ABREU	01H	PL B510 55	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABBIENDI	00Q	PL B482 15	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
BARTELT	99	PRL 82 3746	J. Bartelt <i>et al.</i>	(CLEO Collab.)
ACKERSTAFF	97G	PL B395 128	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ATHANAS	97	PRL 79 2208	M. Athanas <i>et al.</i>	(CLEO Collab.)
BUSKULIC	97	PL B395 373	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ABREU	96P	ZPHY C71 539	P. Abreu <i>et al.</i>	(DELPHI Collab.)
BARISH	95	PR D51 1014	B.C. Barish <i>et al.</i>	(CLEO Collab.)
BUSKULIC	95N	PL B359 236	D. Buskulic <i>et al.</i>	(ALEPH Collab.)