

# Quark and Lepton Compositeness, Searches for

## SEARCHES FOR QUARK AND LEPTON COMPOSITENESS

Revised 2001 by K. Hagiwara (KEK), and K. Hikasa and M. Tanabashi (Tohoku University).

If quarks and leptons are made of constituents, then at the scale of constituent binding energies, there should appear new interactions among quarks and leptons. At energies much below the compositeness scale ( $\Lambda$ ), these interactions are suppressed by inverse powers of  $\Lambda$ . The dominant effect should come from the lowest dimensional interactions with four fermions (contact terms), whose most general chirally invariant form reads [1]

$$L = \frac{g^2}{2\Lambda^2} \left[ \eta_{LL} \bar{\psi}_L \gamma_\mu \psi_L \bar{\psi}_L \gamma^\mu \psi_L + \eta_{RR} \bar{\psi}_R \gamma_\mu \psi_R \bar{\psi}_R \gamma^\mu \psi_R + 2\eta_{LR} \bar{\psi}_L \gamma_\mu \psi_L \bar{\psi}_R \gamma^\mu \psi_R \right]. \quad (1)$$

Chiral invariance provides a natural explanation why quark and lepton masses are much smaller than their inverse size  $\Lambda$ . We may determine the scale  $\Lambda$  unambiguously by using the above form of the effective interactions; the conventional method [1] is to fix its scale by setting  $g^2/4\pi = g^2(\Lambda)/4\pi = 1$  for the new strong interaction coupling and by setting the largest magnitude of the coefficients  $\eta_{\alpha\beta}$  to be unity. In the following, we denote

$$\begin{aligned} \Lambda &= \Lambda_{LL}^\pm \quad \text{for } (\eta_{LL}, \eta_{RR}, \eta_{LR}) = (\pm 1, 0, 0), \\ \Lambda &= \Lambda_{RR}^\pm \quad \text{for } (\eta_{LL}, \eta_{RR}, \eta_{LR}) = (0, \pm 1, 0), \\ \Lambda &= \Lambda_{VV}^\pm \quad \text{for } (\eta_{LL}, \eta_{RR}, \eta_{LR}) = (\pm 1, \pm 1, \pm 1), \\ \Lambda &= \Lambda_{AA}^\pm \quad \text{for } (\eta_{LL}, \eta_{RR}, \eta_{LR}) = (\pm 1, \pm 1, \mp 1), \end{aligned} \quad (2)$$

as typical examples. Such interactions can arise by constituent interchange (when the fermions have common constituents, *e.g.*, for  $ee \rightarrow ee$ ) and/or by exchange of the binding quanta (whenever binding quanta couple to constituents of both particles).

Another typical consequence of compositeness is the appearance of excited leptons and quarks ( $\ell^*$  and  $q^*$ ). Phenomenologically, an excited lepton is defined to be a heavy lepton which shares leptonic quantum number with one of the existing leptons (an excited quark is defined similarly). For example, an excited electron  $e^*$  is characterized by a nonzero transition-magnetic coupling with electrons. Smallness of the lepton mass and the success of QED prediction for  $g-2$  suggest chirality conservation, *i.e.*, an excited lepton should not couple to both left- and right-handed components of the corresponding lepton.

Excited leptons may be classified by  $SU(2) \times U(1)$  quantum numbers. Typical examples are:

1. Sequential type

$$\begin{pmatrix} \nu^* \\ \ell^* \end{pmatrix}_L, \quad [\nu_R^*], \quad \ell_R^*.$$

$\nu_R^*$  is necessary unless  $\nu^*$  has a Majorana mass.

2. Mirror type

$$[\nu_L^*], \quad \ell_L^*, \quad \begin{pmatrix} \nu^* \\ \ell^* \end{pmatrix}_R.$$

3. Homodoublet type

$$\begin{pmatrix} \nu^* \\ \ell^* \end{pmatrix}_L, \quad \begin{pmatrix} \nu^* \\ \ell^* \end{pmatrix}_R.$$

Similar classification can be made for excited quarks.

Excited fermions can be pair produced via their gauge couplings. The couplings of excited leptons with  $Z$  are listed

	Sequential type	Mirror type	Homodoublet type
$V^{\ell^*}$	$-\frac{1}{2} + 2 \sin^2 \theta_W$	$-\frac{1}{2} + 2 \sin^2 \theta_W$	$-1 + 2 \sin^2 \theta_W$
$A^{\ell^*}$	$-\frac{1}{2}$	$+\frac{1}{2}$	0
$V^{\nu_D^*}$	$+\frac{1}{2}$	$+\frac{1}{2}$	+1
$A^{\nu_D^*}$	$+\frac{1}{2}$	$-\frac{1}{2}$	0
$V^{\nu_M^*}$	0	0	—
$A^{\nu_M^*}$	+1	-1	—

in the following table (for notation see Eq. (1) in “Standard Model of Electroweak Interactions”):

Here  $\nu_D^*$  ( $\nu_M^*$ ) stands for Dirac (Majorana) excited neutrino. The corresponding couplings of excited quarks can be easily obtained. Although form factor effects can be present for the gauge couplings at  $q^2 \neq 0$ , they are usually neglected.

In addition, transition magnetic type couplings with a gauge boson are expected. These couplings can be generally parameterized as follows:

$$\begin{aligned}
 \mathcal{L} = & \frac{\lambda_\gamma^{(f^*)}}{2m_{f^*}} e \bar{f}^* \sigma^{\mu\nu} (\eta_L \frac{1-\gamma_5}{2} + \eta_R \frac{1+\gamma_5}{2}) f F_{\mu\nu} \\
 & + \frac{\lambda_Z^{(f^*)}}{2m_{f^*}} e \bar{f}^* \sigma^{\mu\nu} (\eta_L \frac{1-\gamma_5}{2} + \eta_R \frac{1+\gamma_5}{2}) f Z_{\mu\nu} \\
 & + \frac{\lambda_W^{(\ell^*)}}{2m_{\ell^*}} g \bar{\ell}^* \sigma^{\mu\nu} \frac{1-\gamma_5}{2} \nu W_{\mu\nu} \\
 & + \frac{\lambda_W^{(\nu^*)}}{2m_{\nu^*}} g \bar{\nu}^* \sigma^{\mu\nu} (\eta_L \frac{1-\gamma_5}{2} + \eta_R \frac{1+\gamma_5}{2}) \ell W_{\mu\nu}^\dagger \\
 & + \text{h.c.} , \tag{3}
 \end{aligned}$$

where  $g = e/\sin\theta_W$ ,  $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$  is the photon field strength,  $Z_{\mu\nu} = \partial_\mu Z_\nu - \partial_\nu Z_\mu$ , *etc.* The normalization of the coupling is chosen such that

$$\max(|\eta_L|, |\eta_R|) = 1 .$$

Chirality conservation requires

$$\eta_L \eta_R = 0 . \tag{4}$$

Some experimental analyses assume the relation  $\eta_L = \eta_R = 1$ , which violates chiral symmetry. We encode the results of such analyses if the crucial part of the cross section is proportional to the factor  $\eta_L^2 + \eta_R^2$  and the limits can be reinterpreted as those for chirality conserving cases  $(\eta_L, \eta_R) = (1, 0)$  or  $(0, 1)$  after rescaling  $\lambda$ .

These couplings in Eq. (3) can arise from  $SU(2) \times U(1)$ -invariant higher-dimensional interactions. A well-studied model is the interaction of homodoublet type  $\ell^*$  with the Lagrangian [2,3]

$$\mathcal{L} = \frac{1}{2\Lambda} \bar{L}^* \sigma^{\mu\nu} (gf \frac{\tau^a}{2} W_{\mu\nu}^a + g' f' Y B_{\mu\nu}) \frac{1-\gamma_5}{2} L + \text{h.c.} , \tag{5}$$

where  $L$  denotes the lepton doublet  $(\nu, \ell)$ ,  $\Lambda$  is the compositeness scale,  $g, g'$  are  $SU(2)$  and  $U(1)_Y$  gauge couplings, and  $W_{\mu\nu}^a$  and  $B_{\mu\nu}$  are the field strengths for  $SU(2)$  and  $U(1)_Y$  gauge fields. The same interaction occurs for mirror-type excited leptons. For sequential-type excited leptons, the  $\ell^*$  and  $\nu^*$  couplings become unrelated, and the couplings receive the extra suppression of  $(250 \text{ GeV})/\Lambda$  or  $m_{L^*}/\Lambda$ . In any case, these couplings satisfy the relation

$$\lambda_W = -\sqrt{2} \sin^2 \theta_W (\lambda_Z \cot \theta_W + \lambda_\gamma) . \tag{6}$$

Additional coupling with gluons is possible for excited quarks:

$$\mathcal{L} = \frac{1}{2\Lambda} \bar{Q}^* \sigma^{\mu\nu} \left( g_s f_s \frac{\lambda^a}{2} G_{\mu\nu}^a + g f \frac{\tau^a}{2} W_{\mu\nu}^a + g' f' Y B_{\mu\nu} \right) \times \frac{1-\gamma_5}{2} Q + \text{h.c.}, \quad (7)$$

where  $Q$  denotes a quark doublet,  $g_s$  is the QCD gauge coupling, and  $G_{\mu\nu}^a$  the gluon field strength.

It should be noted that the electromagnetic radiative decay of  $\ell^*(\nu^*)$  is forbidden if  $f = -f'$  ( $f = f'$ ). These two possibilities ( $f = f'$  and  $f = -f'$ ) are investigated in many analyses of the LEP experiments above the Z pole.

Several different conventions are used by LEP experiments on Z pole to express the transition magnetic couplings. To facilitate comparison, we re-express these in terms of  $\lambda_Z$  and  $\lambda_\gamma$  using the following relations and taking  $\sin^2\theta_W = 0.23$ . We assume chiral couplings, *i.e.*,  $|c| = |d|$  in the notation of Ref. 2.

1. ALEPH (charged lepton and neutrino)

$$\lambda_Z^{\text{ALEPH}} = \frac{1}{2} \lambda_Z \quad (1990 \text{ papers}) \quad (8a)$$

$$\frac{2c}{\Lambda} = \frac{\lambda_Z}{m_{\ell^*} [\text{or } m_{\nu^*}]} \quad (\text{for } |c| = |d|) \quad (8b)$$

2. ALEPH (quark)

$$\lambda_u^{\text{ALEPH}} = \frac{\sin\theta_W \cos\theta_W}{\sqrt{\frac{1}{4} - \frac{2}{3}\sin^2\theta_W + \frac{8}{9}\sin^4\theta_W}} \lambda_Z = 1.11\lambda_Z \quad (9)$$

3. L3 and DELPHI (charged lepton)

$$\lambda^{\text{L3}} = \lambda_Z^{\text{DELPHI}} = -\frac{\sqrt{2}}{\cot\theta_W - \tan\theta_W} \lambda_Z = -1.10\lambda_Z \quad (10)$$

4. L3 (neutrino)

$$f_Z^{\text{L3}} = \sqrt{2}\lambda_Z \quad (11)$$

5. OPAL (charged lepton)

$$\frac{f^{\text{OPAL}}}{\Lambda} = -\frac{2}{\cot\theta_W - \tan\theta_W} \frac{\lambda_Z}{m_{\ell^*}} = -1.56 \frac{\lambda_Z}{m_{\ell^*}} \quad (12)$$

6. OPAL (quark)

$$\frac{f^{\text{OPAL}}_c}{\Lambda} = \frac{\lambda_Z}{2m_{q^*}} \quad (\text{for } |c| = |d|) \quad (13)$$

7. DELPHI (charged lepton)

$$\lambda_\gamma^{\text{DELPHI}} = -\frac{1}{\sqrt{2}} \lambda_\gamma \quad (14)$$

If leptons are made of color triplet and antitriplet constituents, we may expect their color-octet partners. Transitions between the octet leptons ( $\ell_8$ ) and the ordinary lepton ( $\ell$ ) may take place via the dimension-five interactions

$$\mathcal{L} = \frac{1}{2\Lambda} \sum_{\ell} \left\{ \bar{\ell}_8^\alpha g_S F_{\mu\nu}^\alpha \sigma^{\mu\nu} (\eta_L \ell_L + \eta_R \ell_R) + h.c. \right\} \quad (15)$$

where the summation is over charged leptons and neutrinos. The leptonic chiral invariance implies  $\eta_L \eta_R = 0$  as before.

## References

1. E.J. Eichten, K.D. Lane, and M.E. Peskin, Phys. Rev. Lett. **50**, 811 (1983).
2. K. Hagiwara, S. Komamiya, and D. Zeppenfeld, Z. Phys. **C29**, 115 (1985).
3. N. Cabibbo, L. Maiani, and Y. Srivastava, Phys. Lett. **139B**, 459 (1984).

## SCALE LIMITS for Contact Interactions: $\Lambda(eeee)$

Limits are for  $\Lambda_{LL}^{\pm}$  only. For other cases, see each reference.

$\Lambda_{LL}^+$ (TeV)	$\Lambda_{LL}^-$ (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;8.3</b>	<b>&gt;10.3</b>	95	<sup>1</sup> BOURILKOV 01	RVUE	$E_{cm} = 192\text{--}208$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
>3.8	>5.6	95	ABBIENDI 00R	OPAL	$E_{cm} = 189$ GeV
>4.4	>5.4	95	ABREU 00S	DLPH	$E_{cm} = 183\text{--}189$ GeV
>4.3	>4.9	95	ACCIARRI 00P	L3	$E_{cm} = 130\text{--}189$ GeV
>3.5	>3.2	95	BARATE 00I	ALEP	$E_{cm} = 130\text{--}183$ GeV
>6.0	>7.7	95	<sup>2</sup> BOURILKOV 00	RVUE	$E_{cm} = 183\text{--}189$ GeV
>3.1	>3.8	95	ABBIENDI 99	OPAL	$E_{cm} = 130\text{--}136, 161\text{--}172, 183$ GeV
>2.2	>2.8	95	ABREU 99A	DLPH	$E_{cm} = 130\text{--}172$ GeV
>2.7	>2.4	95	ACCIARRI 98J	L3	$E_{cm} = 130\text{--}172$ GeV
>3.0	>2.5	95	ACKERSTAFF 98V	OPAL	$E_{cm} = 130\text{--}172$ GeV
>2.4	>2.2	95	ACKERSTAFF 97C	OPAL	$E_{cm} = 130\text{--}136, 161$ GeV
>1.7	>2.3	95	ARIMA 97	VNS	$E_{cm} = 57.77$ GeV
>1.6	>2.0	95	<sup>3</sup> BUSKULIC 93Q	ALEP	$E_{cm} = 88.25\text{--}94.25$ GeV
>1.6		95	<sup>3,4</sup> BUSKULIC 93Q	RVUE	
	>2.2	95	BUSKULIC 93Q	RVUE	
	>3.6	95	<sup>5</sup> KROHA 92	RVUE	
>1.3		95	<sup>5</sup> KROHA 92	RVUE	
>0.7	>2.8	95	BEHREND 91C	CELL	$E_{cm} = 35$ GeV
>1.3	>1.3	95	KIM 89	AMY	$E_{cm} = 50\text{--}57$ GeV
>1.4	>3.3	95	<sup>6</sup> BRAUNSCH... 88	TASS	$E_{cm} = 12\text{--}46.8$ GeV
>1.0	>0.7	95	<sup>7</sup> FERNANDEZ 87B	MAC	$E_{cm} = 29$ GeV
>1.1	>1.4	95	<sup>8</sup> BARTEL 86C	JADE	$E_{cm} = 12\text{--}46.8$ GeV
>1.17	>0.87	95	<sup>9</sup> DERRICK 86	HRS	$E_{cm} = 29$ GeV
>1.1	>0.76	95	<sup>10</sup> BERGER 85B	PLUT	$E_{cm} = 34.7$ GeV

<sup>1</sup> A combined analysis of the data from ALEPH, DELPHI, L3, and OPAL.

<sup>2</sup> A combined analysis of the data from ALEPH, L3, and OPAL.

<sup>3</sup> BUSKULIC 93Q uses the following prescription to obtain the limit: when the naive 95%CL limit is better than the statistically expected sensitivity for the limit, the latter is adopted for the limit.

<sup>4</sup> This BUSKULIC 93Q value is from ALEPH data plus PEP/PETRA/TRISTAN data re-analyzed by KROHA 92.

<sup>5</sup> KROHA 92 limit is from fit to BERGER 85B, BARTEL 86C, DERRICK 86B, FERNANDEZ 87B, BRAUNSCHWEIG 88, BEHREND 91B, and BEHREND 91C. The fit gives  $\eta/\Lambda_{LL}^2 = +0.230 \pm 0.206$  TeV<sup>-2</sup>.

<sup>6</sup> BRAUNSCHWEIG 88 assumed  $m_Z = 92$  GeV and  $\sin^2\theta_W = 0.23$ .

<sup>7</sup> FERNANDEZ 87B assumed  $\sin^2\theta_W = 0.22$ .

<sup>8</sup> BARTEL 86C assumed  $m_Z = 93$  GeV and  $\sin^2\theta_W = 0.217$ .

<sup>9</sup> DERRICK 86 assumed  $m_Z = 93$  GeV and  $g_V^2 = (-1/2 + 2\sin^2\theta_W)^2 = 0.004$ .

<sup>10</sup> BERGER 85B assumed  $m_Z = 93$  GeV and  $\sin^2\theta_W = 0.217$ .

### SCALE LIMITS for Contact Interactions: $\Lambda(ee\mu\mu)$

Limits are for  $\Lambda_{LL}^{\pm}$  only. For other cases, see each reference.

$\Lambda_{LL}^+$ (TeV)	$\Lambda_{LL}^-$ (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
>6.6	> <b>6.3</b>	95	ABREU	00S DLPH	$E_{cm} = 183\text{--}189$ GeV
> <b>8.5</b>	>3.8	95	ACCIARRI	00P L3	$E_{cm} = 130\text{--}189$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
>7.3	>4.6	95	ABBIENDI	00R OPAL	$E_{cm} = 189$ GeV
>4.0	>4.7	95	BARATE	00I ALEP	$E_{cm} = 130\text{--}183$ GeV
>4.5	>4.3	95	ABBIENDI	99 OPAL	$E_{cm} = 130\text{--}136, 161\text{--}172, 183$ GeV
>3.4	>2.7	95	ABREU	99A DLPH	$E_{cm} = 130\text{--}172$ GeV
>3.6	>2.4	95	ACCIARRI	98J L3	$E_{cm} = 130\text{--}172$ GeV
>2.9	>3.4	95	ACKERSTAFF	98V OPAL	$E_{cm} = 130\text{--}172$ GeV
>3.1	>2.0	95	MIURA	98 VNS	$E_{cm} = 57.77$ GeV
>2.4	>2.9	95	ACKERSTAFF	97C OPAL	$E_{cm} = 130\text{--}136, 161$ GeV
>1.7	>2.2	95	<sup>11</sup> VELISSARIS	94 AMY	$E_{cm} = 57.8$ GeV
>1.3	>1.5	95	<sup>11</sup> BUSKULIC	93Q ALEP	$E_{cm} = 88.25\text{--}94.25$ GeV
>2.6	>1.9	95	<sup>11,12</sup> BUSKULIC	93Q RVUE	
>2.3	>2.0	95	HOWELL	92 TOPZ	$E_{cm} = 52\text{--}61.4$ GeV
	>1.7	95	<sup>13</sup> KROHA	92 RVUE	
>2.5	>1.5	95	BEHREND	91C CELL	$E_{cm} = 35\text{--}43$ GeV
>1.6	>2.0	95	<sup>14</sup> ABE	90I VNS	$E_{cm} = 50\text{--}60.8$ GeV
>1.9	>1.0	95	KIM	89 AMY	$E_{cm} = 50\text{--}57$ GeV
>2.3	>1.3	95	BRAUNSCH...	88D TASS	$E_{cm} = 30\text{--}46.8$ GeV
>4.4	>2.1	95	<sup>15</sup> BARTEL	86C JADE	$E_{cm} = 12\text{--}46.8$ GeV
>2.9	>0.86	95	<sup>16</sup> BERGER	85 PLUT	$E_{cm} = 34.7$ GeV

<sup>11</sup>BUSKULIC 93Q and VELISSARIS 94 use the following prescription to obtain the limit: when the naive 95%CL limit is better than the statistically expected sensitivity for the limit, the latter is adopted for the limit.

<sup>12</sup>This BUSKULIC 93Q value is from ALEPH data plus PEP/PETRA/TRISTAN data re-analyzed by KROHA 92.

<sup>13</sup>KROHA 92 limit is from fit to BARTEL 86C, BEHREND 87C, BRAUNSCHWEIG 88D, BRAUNSCHWEIG 89C, ABE 90I, and BEHREND 91C. The fit gives  $\eta/\Lambda_{LL}^2 = -0.155 \pm 0.095 \text{ TeV}^{-2}$ .

<sup>14</sup>ABE 90I assumed  $m_Z = 91.163$  GeV and  $\sin^2\theta_W = 0.231$ .

<sup>15</sup>BARTEL 86C assumed  $m_Z = 93$  GeV and  $\sin^2\theta_W = 0.217$ .

<sup>16</sup>BERGER 85 assumed  $m_Z = 93$  GeV and  $\sin^2\theta_W = 0.217$ .

### SCALE LIMITS for Contact Interactions: $\Lambda(ee\tau\tau)$

Limits are for  $\Lambda_{LL}^{\pm}$  only. For other cases, see each reference.

$\Lambda_{LL}^+$ (TeV)	$\Lambda_{LL}^-$ (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
>3.9	> <b>6.5</b>	95	ABBIENDI	00R OPAL	$E_{cm} = 189$ GeV
> <b>5.4</b>	>4.7	95	ACCIARRI	00P L3	$E_{cm} = 130\text{--}189$ GeV



• • • We do not use the following data for averages, fits, limits, etc. • • •

>5.2	>5.4	95	ABREU	00S	DLPH	$E_{\text{cm}} = 183\text{--}189$ GeV
>3.9	>3.7	95	BARATE	00I	ALEP	$E_{\text{cm}} = 130\text{--}183$ GeV
>3.8	>4.0	95	ABBIENDI	99	OPAL	$E_{\text{cm}} = 130\text{--}136, 161\text{--}172, 183$ GeV
>2.8	>2.6	95	ABREU	99A	DLPH	$E_{\text{cm}} = 130\text{--}172$ GeV
>2.4	>2.8	95	ACCIARRI	98J	L3	$E_{\text{cm}} = 130\text{--}172$ GeV
>2.3	>3.7	95	ACKERSTAFF	98V	OPAL	$E_{\text{cm}} = 130\text{--}172$ GeV
>1.9	>3.0	95	ACKERSTAFF	97C	OPAL	$E_{\text{cm}} = 130\text{--}136, 161$ GeV
>1.4	>2.0	95	<sup>17</sup> VELISSARIS	94	AMY	$E_{\text{cm}} = 57.8$ GeV
>1.0	>1.5	95	<sup>17</sup> BUSKULIC	93Q	ALEP	$E_{\text{cm}} = 88.25\text{--}94.25$ GeV
>1.8	>2.3	95	<sup>17,18</sup> BUSKULIC	93Q	RVUE	
>1.9	>1.7	95	HOWELL	92	TOPZ	$E_{\text{cm}} = 52\text{--}61.4$ GeV
>1.9	>2.9	95	<sup>19</sup> KROHA	92	RVUE	
>1.6	>2.3	95	BEHREND	91C	CELL	$E_{\text{cm}} = 35\text{--}43$ GeV
>1.8	>1.3	95	<sup>20</sup> ABE	90I	VNS	$E_{\text{cm}} = 50\text{--}60.8$ GeV
>2.2	>3.2	95	<sup>21</sup> BARTEL	86	JADE	$E_{\text{cm}} = 12\text{--}46.8$ GeV

<sup>17</sup> BUSKULIC 93Q and VELISSARIS 94 use the following prescription to obtain the limit: when the naive 95%CL limit is better than the statistically expected sensitivity for the limit, the latter is adopted for the limit.

<sup>18</sup> This BUSKULIC 93Q value is from ALEPH data plus PEP/PETRA/TRISTAN data re-analyzed by KROHA 92.

<sup>19</sup> KROHA 92 limit is from fit to BARTEL 86C BEHREND 89B, BRAUNSCHWEIG 89C, ABE 90I, and BEHREND 91C. The fit gives  $\eta/\Lambda_{LL}^2 = +0.095 \pm 0.120 \text{ TeV}^{-2}$ .

<sup>20</sup> ABE 90I assumed  $m_Z = 91.163$  GeV and  $\sin^2\theta_W = 0.231$ .

<sup>21</sup> BARTEL 86 assumed  $m_Z = 93$  GeV and  $\sin^2\theta_W = 0.217$ .

## SCALE LIMITS for Contact Interactions: $\Lambda(\ell\ell\ell\ell)$

Lepton universality assumed. Limits are for  $\Lambda_{LL}^{\pm}$  only. For other cases, see each reference.

$\Lambda_{LL}^+(\text{TeV})$	$\Lambda_{LL}^-(\text{TeV})$	CL%	DOCUMENT ID	TECN	COMMENT
>7.3	<b>&gt; 7.8</b>	95	ABREU	00S DLPH	$E_{\text{cm}} = 183\text{--}189$ GeV
<b>&gt; 9.0</b>	>5.2	95	ACCIARRI	00P L3	$E_{\text{cm}} = 130\text{--}189$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

>6.4	>7.2	95	ABBIENDI	00R	OPAL	$E_{\text{cm}} = 189$ GeV
>5.3	>5.5	95	BARATE	00I	ALEP	$E_{\text{cm}} = 130\text{--}183$ GeV
>5.2	>5.3	95	ABBIENDI	99	OPAL	$E_{\text{cm}} = 130\text{--}136, 161\text{--}172, 183$ GeV
>4.4	>4.2	95	ABREU	99A	DLPH	$E_{\text{cm}} = 130\text{--}172$ GeV
>4.0	>3.1	95	<sup>22</sup> ACCIARRI	98J	L3	$E_{\text{cm}} = 130\text{--}172$ GeV
>3.4	>4.4	95	ACKERSTAFF	98V	OPAL	$E_{\text{cm}} = 130\text{--}172$ GeV
>2.7	>3.8	95	ACKERSTAFF	97C	OPAL	$E_{\text{cm}} = 130\text{--}136, 161$ GeV
>3.0	>2.3	95	<sup>22,23</sup> BUSKULIC	93Q	ALEP	$E_{\text{cm}} = 88.25\text{--}94.25$ GeV
>3.5	>2.8	95	<sup>23,24</sup> BUSKULIC	93Q	RVUE	
>2.5	>2.2	95	<sup>25</sup> HOWELL	92	TOPZ	$E_{\text{cm}} = 52\text{--}61.4$ GeV
>3.4	>2.7	95	<sup>26</sup> KROHA	92	RVUE	

<sup>22</sup> From  $e^+e^- \rightarrow e^+e^-, \mu^+\mu^-,$  and  $\tau^+\tau^-$ .

<sup>23</sup> BUSKULIC 93Q uses the following prescription to obtain the limit: when the naive 95%CL limit is better than the statistically expected sensitivity for the limit, the latter is adopted for the limit.

<sup>24</sup> This BUSKULIC 93Q value is from ALEPH data plus PEP/PETRA/TRISTAN data re-analyzed by KROHA 92.

<sup>25</sup> HOWELL 92 limit is from  $e^+e^- \rightarrow \mu^+\mu^-$  and  $\tau^+\tau^-$ .

<sup>26</sup> KROHA 92 limit is from fit to most PEP/PETRA/TRISTAN data. The fit gives  $\eta/\Lambda_{LL}^2 = -0.0200 \pm 0.0666 \text{ TeV}^{-2}$ .

## SCALE LIMITS for Contact Interactions: $\Lambda(eeqq)$

Limits are for  $\Lambda_{LL}^\pm$  only. For other cases, see each reference.

$\Lambda_{LL}^+$ (TeV)	$\Lambda_{LL}^-$ (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;23.3</b>	<b>&gt;12.5</b>	95	27 CHEUNG	01B RVUE	( <i>eeuu</i> )
<b>&gt;11.1</b>	<b>&gt;26.4</b>	95	27 CHEUNG	01B RVUE	( <i>eedd</i> )
<b>&gt; 5.6</b>	<b>&gt;4.9</b>	95	28 BARATE	00I ALEP	( <i>eebb</i> )
<b>&gt; 1.0</b>	<b>&gt;2.1</b>	95	29 ABREU	99A DLPH	( <i>eecc</i> )
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
> 5.5	>3.1	95	30 ABBIENDI	00R OPAL	( <i>eeqq</i> )
> 4.9	>6.1	95	30 ABBIENDI	00R OPAL	( <i>eeuu</i> )
> 5.7	>4.5	95	30 ABBIENDI	00R OPAL	( <i>eedd</i> )
> 4.2	>2.8	95	31 ACCIARRI	00P L3	( <i>eeqq</i> )
> 2.4	>1.3	95	32 ADLOFF	00 H1	( <i>eeqq</i> )
> 5.4	>6.2	95	33 BARATE	00I ALEP	( <i>eeqq</i> )
			34 BREITWEG	00B ZEUS	
> 4.4	>2.8	95	35 ABBIENDI	99 OPAL	( <i>eeqq</i> )
> 4.0	>4.8	95	36 ABBIENDI	99 OPAL	( <i>eebb</i> )
> 3.3	>4.2	95	37 ABBOTT	99D D0	( <i>eeqq</i> )
> 2.4	>2.8	95	29 ABREU	99A DLPH	( <i>eeqq</i> ) ( <i>d</i> or <i>s</i> quark)
> 4.4	>3.9	95	29 ABREU	99A DLPH	( <i>eebb</i> )
> 1.0	>2.4	95	29 ABREU	99A DLPH	( <i>eeuu</i> )
> 4.0	>3.4	95	38 ZARNECKI	99 RVUE	( <i>eedd</i> )
> 4.3	>5.6	95	38 ZARNECKI	99 RVUE	( <i>eeuu</i> )
> 3.0	>2.1	95	39 ACCIARRI	98J L3	( <i>eeqq</i> )
> 3.4	>2.2	95	40 ACKERSTAFF	98V OPAL	( <i>eeqq</i> )
> 4.0	>2.8	95	41 ACKERSTAFF	98V OPAL	( <i>eebb</i> )
> 9.3	>12.0	95	42 BARGER	98E RVUE	( <i>eeuu</i> )
> 8.8	>11.9	95	42 BARGER	98E RVUE	( <i>eedd</i> )
> 2.5	>3.7	95	43 ABE	97T CDF	( <i>eeqq</i> ) (isosinglet)
> 2.5	>2.1	95	44 ACKERSTAFF	97C OPAL	( <i>eeqq</i> )
> 3.1	>2.9	95	45 ACKERSTAFF	97C OPAL	( <i>eebb</i> )
> 7.4	>11.7	95	46 DEANDREA	97 RVUE	<i>eeuu</i> , atomic parity violation
> 2.3	>1.0	95	47 AID	95 H1	( <i>eeqq</i> ) ( <i>u, d</i> quarks)
1.7	>2.2	95	48 ABE	91D CDF	( <i>eeqq</i> ) ( <i>u, d</i> quarks)
> 1.2		95	49 ADACHI	91 TOPZ	( <i>eeqq</i> ) (flavor-universal)
	>1.6	95	49 ADACHI	91 TOPZ	( <i>eeqq</i> ) (flavor-universal)

> 0.6	>1.7	95	50	BEHREND	91C CELL	( <i>eecc</i> )
> 1.1	>1.0	95	50	BEHREND	91C CELL	( <i>eebb</i> )
> 0.9		95	51	ABE	89L VNS	( <i>eeqq</i> ) (flavor-universal)
	>1.7	95	51	ABE	89L VNS	( <i>eeqq</i> ) (flavor-universal)
> 1.05	>1.61	95	52	HAGIWARA	89 RVUE	( <i>eecc</i> )
> 1.21	>0.53	95	53	HAGIWARA	89 RVUE	( <i>eebb</i> )

27 CHEUNG 01B is an update of BARGER 98E.

28 BARATE 00I limits are from  $R_b$  and jet-charge asymmetry at 130–183 GeV.

29 ABREU 99A limits are from flavor-tagged  $e^+e^- \rightarrow q\bar{q}$  cross section at 130–172 GeV.

30 ABBIENDI 00R limits are from  $e^+e^- \rightarrow q\bar{q}$  cross section at  $\sqrt{s}=130$ –189 GeV.

31 ACCIARRI 00P limit is from  $e^+e^- \rightarrow qq$  cross section at  $\sqrt{s}=130$ –189 GeV.

32 ADLOFF 00 limits are from the  $Q^2$  spectrum measurement of  $e^+p \rightarrow e^+X$ .

33 BARATE 00I limits are from  $e^+e^- \rightarrow q\bar{q}$  cross section and jet-charge asymmetry at 130–183 GeV.

34 BREITWEG 00B limits are from  $Q^2$  spectrum measurement of  $e^+p$  collisions. See their Table 3 for the limits of various models.

35 ABBIENDI 99 limits are from  $e^+e^- \rightarrow q\bar{q}$  cross section at 130–136, 161–172, 183 GeV.

36 ABBIENDI 99 limits are from  $R_b$  at 130–136, 161–172, 183 GeV.

37 ABBOTT 99D limits are from  $e^+e^-$  mass distribution in  $p\bar{p} \rightarrow e^+e^-X$  at  $E_{cm}=1.8$  TeV.

38 ZARNECKI 99 use data from HERA, LEP, Tevatron, and various low-energy experiments.

39 ACCIARRI 98J limits are from  $e^+e^- \rightarrow q\bar{q}$  cross section at  $E_{cm}=130$ –172 GeV.

40 ACKERSTAFF 98V limits are from  $e^+e^- \rightarrow q\bar{q}$  at  $E_{cm}=130$ –172 GeV.

41 ACKERSTAFF 98V limits are from  $R_b$  measurements at  $E_{cm}=130$ –172 GeV.

42 BARGER 98E use data from HERA, LEP, Tevatron, and various low-energy experiments.

43 ABE 97T limits are from  $e^+e^-$  mass distribution in  $p\bar{p} \rightarrow e^+e^-X$  at  $E_{cm}=1.8$  TeV.

44 ACKERSTAFF 97C limits are from  $e^+e^- \rightarrow q\bar{q}$  cross section at  $E_{cm}=130$ –136 GeV and 161 GeV.

45 ACKERSTAFF 97C limits are  $R_b$  measurements at  $E_{cm}=133$  GeV and 161 GeV.

46 DEANDREA 97 limit is from atomic parity violation of cesium. The limit is eluded if the contact interactions are parity conserving.

47 AID 95 limits are from the  $Q^2$  spectrum measurement of  $ep \rightarrow eX$ .

48 ABE 91D limits are from  $e^+e^-$  mass distribution in  $p\bar{p} \rightarrow e^+e^-X$  at  $E_{cm}=1.8$  TeV.

49 ADACHI 91 limits are from differential jet cross section. Universality of  $\Lambda(eeqq)$  for five flavors is assumed.

50 BEHREND 91C is from data at  $E_{cm}=35$ –43 GeV.

51 ABE 89L limits are from jet charge asymmetry. Universality of  $\Lambda(eeqq)$  for five flavors is assumed.

52 The HAGIWARA 89 limit is derived from forward-backward asymmetry measurements of  $D/D^*$  mesons by ALTHOFF 83C, BARTEL 84E, and BARINGER 88.

53 The HAGIWARA 89 limit is derived from forward-backward asymmetry measurement of  $b$  hadrons by BARTEL 84D.

### SCALE LIMITS for Contact Interactions: $\Lambda(\mu\mu qq)$

$\Lambda_{LL}^+$ (TeV)	$\Lambda_{LL}^-$ (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
> <b>2.9</b>	> <b>4.2</b>	95	54 ABE	97T CDF	( $\mu\mu qq$ ) (isosinglet)
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
>1.4	>1.6	95	ABE	92B CDF	( $\mu\mu qq$ ) (isosinglet)

<sup>54</sup> ABE 97T limits are from  $\mu^+\mu^-$  mass distribution in  $\bar{p}p \rightarrow \mu^+\mu^-X$  at  $E_{\text{cm}}=1.8$  TeV.

### SCALE LIMITS for Contact Interactions: $\Lambda(\ell\nu\ell\nu)$

VALUE (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;3.10</b>	90	55 JODIDIO	86 SPEC	$\Lambda_{LR}^{\pm}(\nu_{\mu}\nu_e\mu e)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

>3.8		56 DIAZCRUZ	94 RVUE	$\Lambda_{LL}^+(\tau\nu_{\tau}e\nu_e)$
>8.1		56 DIAZCRUZ	94 RVUE	$\Lambda_{LL}^-(\tau\nu_{\tau}e\nu_e)$
>4.1		57 DIAZCRUZ	94 RVUE	$\Lambda_{LL}^+(\tau\nu_{\tau}\mu\nu_{\mu})$
>6.5		57 DIAZCRUZ	94 RVUE	$\Lambda_{LL}^-(\tau\nu_{\tau}\mu\nu_{\mu})$

<sup>55</sup> JODIDIO 86 limit is from  $\mu^+ \rightarrow \bar{\nu}_{\mu} e^+ \nu_e$ . Chirality invariant interactions  $L = (g^2/\Lambda^2) [\eta_{LL} (\bar{\nu}_{\mu} L \gamma^{\alpha} \mu_L) (\bar{e} L \gamma^{\alpha} \nu_e) + \eta_{LR} (\bar{\nu}_{\mu} L \gamma^{\alpha} \nu_e) (\bar{e} R \gamma^{\alpha} \mu_R)]$  with  $g^2/4\pi = 1$  and  $(\eta_{LL}, \eta_{LR}) = (0, \pm 1)$  are taken. No limits are given for  $\Lambda_{LL}^{\pm}$  with  $(\eta_{LL}, \eta_{LR}) = (\pm 1, 0)$ . For more general constraints with right-handed neutrinos and chirality nonconserving contact interactions, see their text.

<sup>56</sup> DIAZCRUZ 94 limits are from  $\Gamma(\tau \rightarrow e\nu\nu)$  and assume flavor-dependent contact interactions with  $\Lambda(\tau\nu_{\tau}e\nu_e) \ll \Lambda(\mu\nu_{\mu}e\nu_e)$ .

<sup>57</sup> DIAZCRUZ 94 limits are from  $\Gamma(\tau \rightarrow \mu\nu\nu)$  and assume flavor-dependent contact interactions with  $\Lambda(\tau\nu_{\tau}\mu\nu_{\mu}) \ll \Lambda(\mu\nu_{\mu}e\nu_e)$ .

### SCALE LIMITS for Contact Interactions: $\Lambda(e\nu qq)$

VALUE (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;2.81</b>	95	58 AFFOLDER	01i CDF	

<sup>58</sup> AFFOLDER 00i bound is for a scalar interaction  $\bar{q}_R q_L \bar{\nu} e_L$ .

### SCALE LIMITS for Contact Interactions: $\Lambda(qqqq)$

Limits are for  $\Lambda_{LL}^{\pm}$  with color-singlet isoscalar exchanges among  $u_L$ 's and  $d_L$ 's only, unless otherwise noted. See EICHTEN 84 for details.

VALUE (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;2.7</b>	95	59 ABBOTT	99C D0	$p\bar{p} \rightarrow$ dijet mass. $\Lambda_{LL}^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

>2.0	95	60 ABBOTT	00E D0	$H_T$ distribution; $\Lambda_{LL}^+$
>2.1	95	61 ABBOTT	98G D0	$p\bar{p} \rightarrow$ dijet angl. $\Lambda_{LL}^+$
		62 BERTRAM	98 RVUE	$p\bar{p} \rightarrow$ dijet mass
		63 ABE	96 CDF	$p\bar{p} \rightarrow$ jets inclusive
>1.6	95	64 ABE	96S CDF	$p\bar{p} \rightarrow$ dijet angl.; $\Lambda_{LL}^+$
>1.3	95	65 ABE	93G CDF	$p\bar{p} \rightarrow$ dijet mass
>1.4	95	66 ABE	92D CDF	$p\bar{p} \rightarrow$ jets inclusive
>1.0	99	67 ABE	92M CDF	$p\bar{p} \rightarrow$ dijet angl.

>0.825	95	68 ALITTI	91B UA2	$p\bar{p} \rightarrow$ jets inclusive
>0.700	95	66 ABE	89 CDF	$p\bar{p} \rightarrow$ jets inclusive
>0.330	95	69 ABE	89H CDF	$p\bar{p} \rightarrow$ dijet angl.
>0.400	95	70 ARNISON	86C UA1	$p\bar{p} \rightarrow$ jets inclusive
>0.415	95	71 ARNISON	86D UA1	$p\bar{p} \rightarrow$ dijet angl.
>0.370	95	72 APPEL	85 UA2	$p\bar{p} \rightarrow$ jets inclusive
>0.275	95	73 BAGNAIA	84C UA2	Repl. by APPEL 85

<sup>59</sup> The quoted limit is from inclusive dijet mass spectrum in  $p\bar{p}$  collisions at  $E_{\text{cm}}=1.8$  TeV.

ABBOTT 99C also obtain  $\Lambda_{LL}^- > 2.4$  TeV. All quarks are assumed composite.

<sup>60</sup> The quoted limit for ABBOTT 00E is from  $H_T$  distribution in  $p\bar{p}$  collisions at  $E_{\text{cm}}=1.8$  TeV. CTEQ4M PDF and  $\mu=E_T^{\text{max}}$  are assumed. For limits with different assumptions, see their Tables 2 and 3. All quarks are assumed composite.

<sup>61</sup> ABBOTT 98G limit is from dijet angular distribution in  $p\bar{p}$  collisions at  $E_{\text{cm}}=1.8$  TeV. All quarks are assumed composite.

<sup>62</sup> BERTRAM 98 obtain limit on the scale of color-octet axial-vector flavor-universal contact interactions:  $\Lambda_{A8} > 2.1$  TeV. They also obtain a limit  $\Lambda_{V8} > 2.4$  TeV on a color-octet flavor-universal vectorial contact interaction.

<sup>63</sup> ABE 96 finds that the inclusive jet cross section for  $E_T > 200$  GeV is significantly higher than the  $\mathcal{O}(\alpha_s^3)$  perturbative QCD prediction. This could be interpreted as the effect of a contact interaction with  $\Lambda_{LL} \sim 1.6$  TeV. However, ABE 96 state that uncertainty in the parton distribution functions, higher-order QCD corrections, and the detector calibration may possibly account for the effect.

<sup>64</sup> ABE 96S limit is from dijet angular distribution in  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.8$  TeV. The limit for  $\Lambda_{LL}^-$  is  $> 1.4$  TeV. ABE 96S also obtain limits for flavor symmetric contact interactions among all quark flavors:  $\Lambda_{LL}^+ > 1.8$  TeV and  $\Lambda_{LL}^- > 1.6$  TeV.

<sup>65</sup> ABE 93G limit is from dijet mass distribution in  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.8$  TeV. The limit is the weakest from several choices of structure functions and renormalization scale.

<sup>66</sup> Limit is from inclusive jet cross-section data in  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.8$  TeV. The limit takes into account uncertainties in choice of structure functions and in choice of process scale.

<sup>67</sup> ABE 92M limit is from dijet angular distribution for  $m_{\text{dijet}} > 550$  GeV in  $p\bar{p}$  collisions at  $E_{\text{cm}}=1.8$  TeV.

<sup>68</sup> ALITTI 91B limit is from inclusive jet cross section in  $p\bar{p}$  collisions at  $E_{\text{cm}} = 630$  GeV. The limit takes into account uncertainties in choice of structure functions and in choice of process scale.

<sup>69</sup> ABE 89H limit is from dijet angular distribution for  $m_{\text{dijet}} > 200$  GeV at the Fermilab Tevatron Collider with  $E_{\text{cm}} = 1.8$  TeV. The QCD prediction is quite insensitive to choice of structure functions and choice of process scale.

<sup>70</sup> ARNISON 86C limit is from the study of inclusive high- $p_T$  jet distributions at the CERN  $\bar{p}p$  collider ( $E_{\text{cm}} = 546$  and 630 GeV). The QCD prediction renormalized to the low- $p_T$  region gives a good fit to the data.

<sup>71</sup> ARNISON 86D limit is from the study of dijet angular distribution in the range  $240 < m(\text{dijet}) < 300$  GeV at the CERN  $\bar{p}p$  collider ( $E_{\text{cm}} = 630$  GeV). QCD prediction using EHLQ structure function (EICHTEN 84) with  $\Lambda_{\text{QCD}} = 0.2$  GeV for the choice of  $Q^2 = p_T^2$  gives the best fit to the data.

<sup>72</sup> APPEL 85 limit is from the study of inclusive high- $p_T$  jet distributions at the CERN  $\bar{p}p$  collider ( $E_{\text{cm}} = 630$  GeV). The QCD prediction renormalized to the low- $p_T$  region gives a good description of the data.

<sup>73</sup> BAGNAIA 84C limit is from the study of jet  $p_T$  and dijet mass distributions at the CERN  $\bar{p}p$  collider ( $E_{\text{cm}} = 540$  GeV). The limit suffers from the uncertainties in comparing the data with the QCD prediction.

## SCALE LIMITS for Contact Interactions: $\Lambda(\nu\nu qq)$

Limits are for  $\Lambda_{LL}^{\pm}$  only. For other cases, see each reference.

$\Lambda_{LL}^+$ (TeV)	$\Lambda_{LL}^-$ (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
>5.0	>5.4	95	<sup>74</sup> MCFARLAND 98	CCFR	$\nu N$ scattering

<sup>74</sup> MCFARLAND 98 assumed a flavor universal interaction. Neutrinos were mostly of muon type.

## MASS LIMITS for Excited $e$ ( $e^*$ )

Most  $e^+e^-$  experiments assume one-photon or  $Z$  exchange. The limits from some  $e^+e^-$  experiments which depend on  $\lambda$  have assumed transition couplings which are chirality violating ( $\eta_L = \eta_R$ ). However they can be interpreted as limits for chirality-conserving interactions after multiplying the coupling value  $\lambda$  by  $\sqrt{2}$ ; see Note.

Excited leptons have the same quantum numbers as other ortholeptons. See also the searches for ortholeptons in the "Searches for Heavy Leptons" section.

## Limits for Excited $e$ ( $e^*$ ) from Pair Production

These limits are obtained from  $e^+e^- \rightarrow e^{*+}e^{*-}$  and thus rely only on the (electroweak) charge of  $e^*$ . Form factor effects are ignored unless noted. For the case of limits from  $Z$  decay, the  $e^*$  coupling is assumed to be of sequential type. Possible  $t$  channel contribution from transition magnetic coupling is neglected. All limits assume a dominant  $e^* \rightarrow e\gamma$  decay except the limits from  $\Gamma(Z)$ .

For limits prior to 1987, see our 1992 edition (Physical Review **D45**, 1 June, Part II (1992)).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>100.0	95	<sup>75</sup> ACCIARRI	01D L3	$e^+e^- \rightarrow e^*e^*$ Homodoublet type
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
> 91.3	95	<sup>76</sup> ABBIENDI	00I OPAL	$e^+e^- \rightarrow e^*e^*$ Homodoublet type
> 94.2	95	<sup>77</sup> ACCIARRI	00E L3	$e^+e^- \rightarrow e^*e^*$ Homodoublet type
> 90.7	95	<sup>78</sup> ABREU	99O DLPH	Homodoublet type
> 85.0	95	<sup>79</sup> ACKERSTAFF	98C OPAL	$e^+e^- \rightarrow e^*e^*$ Homodoublet type
		<sup>80</sup> BARATE	98U ALEP	$Z \rightarrow e^*e^*$
> 79.6	95	<sup>81,82</sup> ABREU	97B DLPH	$e^+e^- \rightarrow e^*e^*$ Homodoublet type
> 77.9	95	<sup>81,83</sup> ABREU	97B DLPH	$e^+e^- \rightarrow e^*e^*$ Sequential type
> 79.7	95	<sup>81</sup> ACCIARRI	97G L3	$e^+e^- \rightarrow e^*e^*$ Sequential type
> 79.9	95	<sup>81,84</sup> ACKERSTAFF	97 OPAL	$e^+e^- \rightarrow e^*e^*$ Homodoublet type
> 62.5	95	<sup>85</sup> ABREU	96K DLPH	$e^+e^- \rightarrow e^*e^*$ Homodoublet type
> 64.7	95	<sup>86</sup> ACCIARRI	96D L3	$e^+e^- \rightarrow e^*e^*$ Sequential type
> 66.5	95	<sup>86</sup> ALEXANDER	96Q OPAL	$e^+e^- \rightarrow e^*e^*$ Homodoublet type
> 65.2	95	<sup>86</sup> BUSKULIC	96W ALEP	$e^+e^- \rightarrow e^*e^*$ Sequential type
> 45.6	95	ADRIANI	93M L3	$Z \rightarrow e^*e^*$
> 45.6	95	ABREU	92C DLPH	$Z \rightarrow e^*e^*$

> 29.8	95	87	BARDADIN-...	92	RVUE	$\Gamma(Z)$
> 26.1	95	88	DECAMP	92	ALEP	$Z \rightarrow e^* e^*; \Gamma(Z)$
> 46.1	95		DECAMP	92	ALEP	$Z \rightarrow e^* e^*$
> 33	95	88	ABREU	91F	DLPH	$Z \rightarrow e^* e^*; \Gamma(Z)$
> 45.0	95	89	ADEVA	90F	L3	$Z \rightarrow e^* e^*$
> 44.9	95		AKRAWY	90i	OPAL	$Z \rightarrow e^* e^*$
> 44.6	95	90	DECAMP	90G	ALEP	$e^+ e^- \rightarrow e^* e^*$
> 30.2	95		ADACHI	89B	TOPZ	$e^+ e^- \rightarrow e^* e^*$
> 28.3	95		KIM	89	AMY	$e^+ e^- \rightarrow e^* e^*$
> 27.9	95	91	ABE	88B	VNS	$e^+ e^- \rightarrow e^* e^*$

75 From  $e^+ e^-$  collisions at  $\sqrt{s} = 192\text{--}202$  GeV.  $f=f'$  is assumed. ACCIARRI 01D also obtain limit for  $f=-f'$ :  $m_{e^*} > 93.4$  GeV.

76 From  $e^+ e^-$  collisions at  $\sqrt{s}=161\text{--}183$  GeV.  $f=f'$  is assumed. ABBIENDI 00i also obtain limit for  $f=-f'$  ( $e^* \rightarrow \nu W$ ):  $m_{e^*} > 86.0$  GeV.

77 From  $e^+ e^-$  collisions at  $\sqrt{s}=189$  GeV.  $f=f'$  is assumed. ACCIARRI 00E also obtain limit for  $f=-f'$  ( $e^* \rightarrow \nu W$ ):  $m_{e^*} > 92.6$  GeV.

78 From  $e^+ e^-$  collisions at  $\sqrt{s}= 183$  GeV.  $f=f'$  is assumed. ABREU 990 also obtain limit for  $f=-f'$  ( $e^* \rightarrow \nu W$ ):  $m_{e^*} > 81.3$  GeV.

79 From  $e^+ e^-$  collisions at  $\sqrt{s}=170\text{--}172$  GeV. ACKERSTAFF 98C also obtain limit from  $e^* \rightarrow \nu W$  decay mode:  $m_{e^*} > 81.3$  GeV.

80 BARATE 98U obtain limits on the form factor. See their Fig. 14 for limits in mass-form factor plane.

81 From  $e^+ e^-$  collisions at  $\sqrt{s}= 161$  GeV.

82 ABREU 97B also obtain limit from charged current decay mode  $e^* \rightarrow \nu W$ ,  $m_{e^*} > 70.9$  GeV.

83 ABREU 97B also obtain limit from charged current decay mode  $e^* \rightarrow \nu W$ ,  $m_{e^*} > 44.6$  GeV.

84 ACKERSTAFF 97 also obtain limit from charged current decay mode  $e^* \rightarrow \nu W$ ,  $m_{\nu_e^*} > 77.1$  GeV.

85 From  $e^+ e^-$  collisions at  $\sqrt{s}= 130\text{--}136$  GeV.

86 From  $e^+ e^-$  collisions at  $\sqrt{s}= 130\text{--}140$  GeV.

87 BARDADIN-OTWINOWSKA 92 limit is independent of decay modes. Based on  $\Delta\Gamma(Z)<36$  MeV.

88 Limit is independent of  $e^*$  decay mode.

89 ADEVA 90F is superseded by ADRIANI 93M.

90 Superseded by DECAMP 92.

91 ABE 88B limits assume  $e^+ e^- \rightarrow e^{*+} e^{*-}$  with one photon exchange only and  $e^* \rightarrow e\gamma$  giving  $e e \gamma \gamma$ .

### Limits for Excited $e$ ( $e^*$ ) from Single Production

These limits are from  $e^+e^- \rightarrow e^*e$ ,  $W \rightarrow e^*\nu$ , or  $ep \rightarrow e^*X$  and depend on transition magnetic coupling between  $e$  and  $e^*$ . All limits assume  $e^* \rightarrow e\gamma$  decay except as noted. Limits from LEP, UA2, and H1 are for chiral coupling, whereas all other limits are for nonchiral coupling,  $\eta_L = \eta_R = 1$ . In most papers, the limit is expressed in the form of an excluded region in the  $\lambda - m_{e^*}$  plane. See the original papers.

For limits prior to 1987, see our 1992 edition (Physical Review **D45**, 1 June, Part II (1992)).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>223	95	92 ADLOFF	00E H1	$ep \rightarrow e^*X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
>202		93 ACCIARRI	01D L3	$e^+e^- \rightarrow ee^*$
		94 ABBIENDI	00I OPAL	$e^+e^- \rightarrow ee^*$
		95 ACCIARRI	00E L3	$e^+e^- \rightarrow ee^*$
		96 ABREU	99O DLPH	$e^+e^- \rightarrow ee^*$
none 20–170	95	97 ACCIARRI	98T L3	$e\gamma \rightarrow e^* \rightarrow e\gamma$
		98 ACKERSTAFF	98C OPAL	$e^+e^- \rightarrow ee^*$
		99 BARATE	98U ALEP	$e^+e^- \rightarrow ee^*$
		100,101 ABREU	97B DLPH	$e^+e^- \rightarrow ee^*$
		100,102 ACCIARRI	97G L3	$e^+e^- \rightarrow ee^*$
		103 ACKERSTAFF	97 OPAL	$e^+e^- \rightarrow ee^*$
		104 ADLOFF	97 H1	Lepton-flavor violation
none 30–200	95	105 BREITWEG	97C ZEUS	$ep \rightarrow e^*X$
		106 ABREU	96K DLPH	$e^+e^- \rightarrow ee^*$
		107 ACCIARRI	96D L3	$e^+e^- \rightarrow ee^*$
		108 ALEXANDER	96Q OPAL	$e^+e^- \rightarrow ee^*$
		109 BUSKULIC	96W ALEP	$e^+e^- \rightarrow ee^*$
		110 DERRICK	95B ZEUS	$ep \rightarrow e^*X$
		111 ABT	93 H1	$ep \rightarrow e^*X$
> 86	95	ADRIANI	93M L3	$\lambda_\gamma > 0.04$
> 89	95	ADRIANI	93M L3	$Z \rightarrow ee^*$ , $\lambda_Z > 0.5$
		112 DERRICK	93B ZEUS	Superseded by DERRICK 95B
> 88	95	ABREU	92C DLPH	$Z \rightarrow ee^*$ , $\lambda_Z > 0.5$
> 86	95	ABREU	92C DLPH	$e^+e^- \rightarrow ee^*$ , $\lambda_\gamma > 0.1$
> 91	95	DECAMP	92 ALEP	$Z \rightarrow ee^*$ , $\lambda_Z > 1$
> 88	95	113 ADEVA	90F L3	$Z \rightarrow ee^*$ , $\lambda_Z > 0.5$
> 86	95	113 ADEVA	90F L3	$Z \rightarrow ee^*$ , $\lambda_Z > 0.04$
> 87	95	AKRAWY	90I OPAL	$Z \rightarrow ee^*$ , $\lambda_Z > 0.5$
> 81	95	114 DECAMP	90G ALEP	$Z \rightarrow ee^*$ , $\lambda_Z > 1$
> 50	95	ADACHI	89B TOPZ	$e^+e^- \rightarrow ee^*$ , $\lambda_\gamma > 0.04$
> 56	95	KIM	89 AMY	$e^+e^- \rightarrow ee^*$ , $\lambda_\gamma > 0.03$
none 23–54	95	115 ABE	88B VNS	$e^+e^- \rightarrow ee^*$ , $\lambda_\gamma > 0.04$
> 75	95	116 ANSARI	87D UA2	$W \rightarrow e^*\nu$ ; $\lambda_W > 0.7$
> 63	95	116 ANSARI	87D UA2	$W \rightarrow e^*\nu$ ; $\lambda_W > 0.2$
> 40	95	116 ANSARI	87D UA2	$W \rightarrow e^*\nu$ ; $\lambda_W > 0.09$



- 92 ADLOFF 00E search for single  $e^*$  production in  $ep$  collisions with the decays  $e^* \rightarrow e\gamma, eZ, \nu W$ .  $f=f'=\Lambda/m_{e^*}$  is assumed for the  $e^*$  coupling. See their Fig. 9 for the exclusion plot in the mass-coupling plane.
- 93 ACCIARRI 01D result is from  $e^+e^-$  collisions at  $\sqrt{s} = 192\text{--}202$  GeV.  $f=f'=\Lambda/m_{e^*}$  is assumed for the  $e^*$  coupling. See their Fig. 4 for limits in the mass-coupling plane.
- 94 ABBIENDI 00I result is from  $e^+e^-$  collisions at  $\sqrt{s}=161\text{--}183$  GeV. See their Fig. 7 for limits in mass-coupling plane.
- 95 ACCIARRI 00E result is from  $e^+e^-$  collisions at  $\sqrt{s}=189$  GeV. See their Fig. 3 for limits in mass-coupling plane.
- 96 ABREU 99O result is from  $e^+e^-$  collisions at  $\sqrt{s}=183$  GeV. See their Figs. 4 and 5 for the exclusion limit in the mass-coupling plane.
- 97 ACCIARRI 98T search for single  $e^*$  production in quasi-real Compton scattering. The limit is for  $|\lambda| > 1.0 \times 10^{-1}$  and non-chiral coupling of  $e^*$ . See their Fig. 7 for the exclusion plot in the mass-coupling plane.
- 98 ACKERSTAFF 98C from  $e^+e^-$  collisions at  $\sqrt{s}=170\text{--}172$  GeV. See their Fig. 11 for the exclusion limit in the mass-coupling plane.
- 99 BARATE 98U is from  $e^+e^-$  collision at  $\sqrt{s}=M_Z$ . See their Fig. 12 for limits in mass-coupling plane
- 100 From  $e^+e^-$  collisions at  $\sqrt{s}=161$  GeV.
- 101 See Fig. 4a and Fig. 5a of ABREU 97B for the exclusion limit in the mass-coupling plane.
- 102 See Fig. 2 and Fig. 3 of ACCIARRI 97G for the exclusion limit in the mass-coupling plane.
- 103 ACKERSTAFF 97 result is from  $e^+e^-$  collisions at  $\sqrt{s}=161$  GeV. See their Fig. 3 for the exclusion limit in the mass-coupling plane.
- 104 ADLOFF 97 search for single  $e^*$  production in  $ep$  collisions with the decays  $e^* \rightarrow e\gamma, eZ, \nu W$ . See their Fig. 4 for the rejection limits on the product of the production cross section and the branching ratio into a specific decay channel.
- 105 BREITWEG 97C search for single  $e^*$  production in  $ep$  collisions with the decays  $e^* \rightarrow e\gamma, eZ, \nu W$ .  $f=f'=2\Lambda/m_{e^*}$  is assumed for the  $e^*$  coupling. See their Fig. 9 for the exclusion plot in the mass-coupling plane.
- 106 ABREU 96K result is from  $e^+e^-$  collisions at  $\sqrt{s}=130\text{--}136$  GeV. See their Fig. 4 for the exclusion limit in the mass-coupling plane.
- 107 ACCIARRI 96D result is from  $e^+e^-$  collisions at  $\sqrt{s}=130\text{--}140$  GeV. See their Fig. 2 for the exclusion limit in the mass-coupling plane.
- 108 ALEXANDER 96Q result is from  $e^+e^-$  collisions at  $\sqrt{s}=130\text{--}140$  GeV. See their Fig. 3a for the exclusion limit in the mass-coupling plane.
- 109 BUSKULIC 96W result is from  $e^+e^-$  collisions at  $\sqrt{s}=130\text{--}140$  GeV. See their Fig. 3 for the exclusion limit in the mass-coupling plane.
- 110 DERRICK 95B search for single  $e^*$  production via  $e^*e\gamma$  coupling in  $ep$  collisions with the decays  $e^* \rightarrow e\gamma, eZ, \nu W$ . See their Fig. 13 for the exclusion plot in the  $m_{e^*}-\lambda\gamma$  plane.
- 111 ABT 93 search for single  $e^*$  production via  $e^*e\gamma$  coupling in  $ep$  collisions with the decays  $e^* \rightarrow e\gamma, eZ, \nu W$ . See their Fig. 4 for exclusion plot in the  $m_{e^*}-\lambda\gamma$  plane.
- 112 DERRICK 93B search for single  $e^*$  production via  $e^*e\gamma$  coupling in  $ep$  collisions with the decays  $e^* \rightarrow e\gamma, eZ, \nu W$ . See their Fig. 3 for exclusion plot in the  $m_{e^*}-\lambda\gamma$  plane.
- 113 Superseded by ADRIANI 93M.
- 114 Superseded by DECAMP 92.
- 115 ABE 88B limits use  $e^+e^- \rightarrow ee^*$  where t-channel photon exchange dominates giving  $e\gamma(e)$  (quasi-real compton scattering).
- 116 ANSARI 87D is at  $E_{\text{cm}} = 546\text{--}630$  GeV.

## Limits for Excited $e$ ( $e^*$ ) from $e^+ e^- \rightarrow \gamma\gamma$

These limits are derived from indirect effects due to  $e^*$  exchange in the  $t$  channel and depend on transition magnetic coupling between  $e$  and  $e^*$ . All limits are for  $\lambda_\gamma = 1$ . All limits except ABE 89J and ACHARD 02D are for nonchiral coupling with  $\eta_L = \eta_R = 1$ . We choose the chiral coupling limit as the best limit and list it in the Summary Table.

For limits prior to 1987, see our 1992 edition (Physical Review **D45**, 1 June, Part II (1992)).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>310	95	ACHARD	02D L3	$\sqrt{s}=192\text{--}209$ GeV
>311	95	ABREU	00A DLPH	$\sqrt{s}=189\text{--}202$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
>283	95	<sup>117</sup> ACCIARRI	00G L3	$\sqrt{s}=183\text{--}189$ GeV
>306	95	ABBIENDI	99P OPAL	$\sqrt{s}=189$ GeV
>231	95	ABREU	98J DLPH	$\sqrt{s}=130\text{--}183$ GeV
>194	95	ACKERSTAFF	98 OPAL	$\sqrt{s}=130\text{--}172$ GeV
>227	95	ACKER...,K...	98B OPAL	$\sqrt{s}=183$ GeV
>250	95	BARATE	98J ALEP	$\sqrt{s}=183$ GeV
>160	95	BARATE	98U ALEP	
>210	95	<sup>119</sup> ACCIARRI	97W L3	$\sqrt{s}=161, 172$ GeV
>129	95	ACCIARRI	96L L3	$\sqrt{s}=133$ GeV
>147	95	ALEXANDER	96K OPAL	
>136	95	BUSKULIC	96Z ALEP	$\sqrt{s}=130, 136$ GeV
>146	95	ACCIARRI	95G L3	
		<sup>120</sup> BUSKULIC	93Q ALEP	
>127	95	<sup>121</sup> ADRIANI	92B L3	
>114	95	<sup>122</sup> BARDADIN-...	92 RVUE	
> 99	95	DECAMP	92 ALEP	
		<sup>123</sup> SHIMOZAWA	92 TOPZ	
>100	95	ABREU	91E DLPH	
>116	95	AKRAWY	91F OPAL	
> 83	95	ADEVA	90K L3	
> 82	95	AKRAWY	90F OPAL	
> 68	95	<sup>124</sup> ABE	89J VNS	$\eta_L=1, \eta_R=0$
> 90.2	95	ADACHI	89B TOPZ	
> 65	95	KIM	89 AMY	

<sup>117</sup> ACCIARRI 00G also obtain a limit on  $e^*$  with chiral coupling,  $m_{e^*} > 213$  GeV.

<sup>118</sup> BARATE 98U is from  $e^+ e^-$  collision at  $\sqrt{s}=M_Z$ . See their Fig. 5 for limits in mass-coupling plane

<sup>119</sup> ACCIARRI 97W also obtain a limit on  $e^*$  with chiral coupling,  $m_{e^*} > 157$  GeV (95%CL).

<sup>120</sup> BUSKULIC 93Q obtain  $\Lambda^+ > 121$  GeV (95%CL) from ALEPH experiment and  $\Lambda^+ > 135$  GeV from combined TRISTAN and ALEPH data. These limits roughly correspond to limits on  $m_{e^*}$ .

<sup>121</sup> ADRIANI 92B superseded by ACCIARRI 95G.

<sup>122</sup> BARDADIN-OTWINOWSKA 92 limit from fit to the combined data of DECAMP 92, ABREU 91E, ADEVA 90K, AKRAWY 91F.

<sup>123</sup> SHIMOZAWA 92 fit the data to the limiting form of the cross section with  $m_{e^*} \gg E_{cm}$  and obtain  $m_{e^*} > 168$  GeV at 95%CL. Use of the full form would reduce this limit by a few GeV. The statistically unexpected large value is due to fluctuation in the data.

<sup>124</sup> The ABE 89J limit assumes chiral coupling. This corresponds to  $\lambda_\gamma = 0.7$  for nonchiral coupling.

## Indirect Limits for Excited $e$ ( $e^*$ )

These limits make use of loop effects involving  $e^*$  and are therefore subject to theoretical uncertainty.

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
125	DORENBOS...	89 CHRM	$\bar{\nu}_\mu e \rightarrow \bar{\nu}_\mu e$ and $\nu_\mu e \rightarrow \nu_\mu e$
126	GRIFOLS	86 THEO	$\nu_\mu e \rightarrow \nu_\mu e$
127	RENARD	82 THEO	$g-2$ of electron
125	DORENBOSCH 89 obtain the limit $\lambda_\gamma^2 \Lambda_{\text{cut}}^2 / m_{e^*}^2 < 2.6$ (95% CL), where $\Lambda_{\text{cut}}$ is the cutoff scale, based on the one-loop calculation by GRIFOLS 86. If one assumes that $\Lambda_{\text{cut}} = 1$ TeV and $\lambda_\gamma = 1$ , one obtains $m_{e^*} > 620$ GeV. However, one generally expects $\lambda_\gamma \approx m_{e^*} / \Lambda_{\text{cut}}$ in composite models.		
126	GRIFOLS 86 uses $\nu_\mu e \rightarrow \nu_\mu e$ and $\bar{\nu}_\mu e \rightarrow \bar{\nu}_\mu e$ data from CHARM Collaboration to derive mass limits which depend on the scale of compositeness.		
127	RENARD 82 derived from $g-2$ data limits on mass and couplings of $e^*$ and $\mu^*$ . See figures 2 and 3 of the paper.		

## MASS LIMITS for Excited $\mu$ ( $\mu^*$ )

### Limits for Excited $\mu$ ( $\mu^*$ ) from Pair Production

These limits are obtained from  $e^+ e^- \rightarrow \mu^{*+} \mu^{*-}$  and thus rely only on the (electroweak) charge of  $\mu^*$ . Form factor effects are ignored unless noted. For the case of limits from  $Z$  decay, the  $\mu^*$  coupling is assumed to be of sequential type. All limits assume a dominant  $\mu^* \rightarrow \mu\gamma$  decay except the limits from  $\Gamma(Z)$ .

For limits prior to 1987, see our 1992 edition (Physical Review **D45**, 1 June, Part II (1992)).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;100.2</b>	95	128 ACCIARRI	01D L3	$e^+ e^- \rightarrow \mu^* \mu^*$ Homodoublet type
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
> 91.3	95	129 ABBIENDI	00I OPAL	$e^+ e^- \rightarrow \mu^* \mu^*$ Homodoublet type
> 94.2	95	130 ACCIARRI	00E L3	$e^+ e^- \rightarrow \mu^* \mu^*$ Homodoublet type
> 90.7	95	131 ABREU	99O DLPH	Homodoublet type
> 85.3	95	132 ACKERSTAFF	98C OPAL	$e^+ e^- \rightarrow \mu^* \mu^*$ Homodoublet type
		133 BARATE	98U ALEP	$Z \rightarrow \mu^* \mu^*$
> 79.6	95	134,135 ABREU	97B DLPH	$e^+ e^- \rightarrow \mu^* \mu^*$ Homodoublet type
> 78.4	95	134,136 ABREU	97B DLPH	$e^+ e^- \rightarrow \mu^* \mu^*$ Sequential type
> 79.9	95	134 ACCIARRI	97G L3	$e^+ e^- \rightarrow \mu^* \mu^*$ Sequential type
> 80.0	95	134,137 ACKERSTAFF	97 OPAL	$e^+ e^- \rightarrow \mu^* \mu^*$ Homodoublet type
> 62.6	95	138 ABREU	96K DLPH	$e^+ e^- \rightarrow \mu^* \mu^*$ Homodoublet type
> 64.9	95	139 ACCIARRI	96D L3	$e^+ e^- \rightarrow \mu^* \mu^*$ Sequential type
> 66.8	95	139 ALEXANDER	96Q OPAL	$e^+ e^- \rightarrow \mu^* \mu^*$ Homodoublet type
> 65.4	95	139 BUSKULIC	96W ALEP	$e^+ e^- \rightarrow \mu^* \mu^*$ Sequential type
> 45.6	95	ADRIANI	93M L3	$Z \rightarrow \mu^* \mu^*$

> 45.6	95	ABREU	92C DLPH	$Z \rightarrow \mu^* \mu^*$
> 29.8	95	140 BARDADIN-...	92 RVUE	$\Gamma(Z)$
> 26.1	95	141 DECAMP	92 ALEP	$Z \rightarrow \mu^* \mu^*; \Gamma(Z)$
> 46.1	95	DECAMP	92 ALEP	$Z \rightarrow \mu^* \mu^*$
> 33	95	141 ABREU	91F DLPH	$Z \rightarrow \mu^* \mu^*; \Gamma(Z)$
> 45.3	95	142 ADEVA	90F L3	$Z \rightarrow \mu^* \mu^*$
> 44.9	95	AKRAWY	90i OPAL	$Z \rightarrow \mu^* \mu^*$
> 44.6	95	143 DECAMP	90G ALEP	$e^+ e^- \rightarrow \mu^* \mu^*$
> 29.9	95	ADACHI	89B TOPZ	$e^+ e^- \rightarrow \mu^* \mu^*$
> 28.3	95	KIM	89 AMY	$e^+ e^- \rightarrow \mu^* \mu^*$

128 From  $e^+ e^-$  collisions at  $\sqrt{s} = 192\text{--}202$  GeV.  $f=f'$  is assumed. ACCIARRI 01D also obtain limit for  $f=-f'$ :  $m_{\mu^*} > 93.4$  GeV.

129 From  $e^+ e^-$  collisions at  $\sqrt{s}=161\text{--}183$  GeV.  $f=f'$  is assumed. ABBIENDI 00i also obtain limit for  $f=-f'$  ( $\mu^* \rightarrow \nu W$ ):  $m_{\mu^*} > 86.0$  GeV.

130 From  $e^+ e^-$  collisions at  $\sqrt{s}=189$  GeV.  $f=f'$  is assumed. ACCIARRI 00E also obtain limit for  $f=-f'$  ( $\mu^* \rightarrow \nu W$ ):  $m_{\mu^*} > 92.6$  GeV.

131 From  $e^+ e^-$  collisions at  $\sqrt{s}= 183$  GeV.  $f=f'$  is assumed. ABREU 990 also obtain limit for  $f=-f'$  ( $\mu^* \rightarrow \nu W$ ):  $m_{\mu^*} > 81.3$  GeV.

132 From  $e^+ e^-$  collisions at  $\sqrt{s}=170\text{--}172$  GeV. ACKERSTAFF 98C also obtain limit from  $\mu^* \rightarrow \nu W$  decay mode:  $m_{\mu^*} > 81.3$  GeV.

133 BARATE 98U obtain limits on the form factor. See their Fig. 14 for limits in mass-form factor plane.

134 From  $e^+ e^-$  collisions at  $\sqrt{s}= 161$  GeV.

135 ABREU 97B also obtain limit from charged current decay mode  $\mu^* \rightarrow \nu W$ ,  $m_{\mu^*} > 70.9$  GeV.

136 ABREU 97B also obtain limit from charged current decay mode  $\mu^* \rightarrow \nu W$ ,  $m_{\mu^*} > 44.6$  GeV.

137 ACKERSTAFF 97 also obtain limit from charged current decay mode  $\mu^* \rightarrow \nu W$ ,  $m_{\nu^*} > 77.1$  GeV.

138 From  $e^+ e^-$  collisions at  $\sqrt{s}= 130\text{--}136$  GeV.

139 From  $e^+ e^-$  collisions at  $\sqrt{s}= 130\text{--}140$  GeV.

140 BARDADIN-OTWINOWSKA 92 limit is independent of decay modes. Based on  $\Delta\Gamma(Z)<36$  MeV.

141 Limit is independent of  $\mu^*$  decay mode.

142 Superseded by ADRIANI 93M.

143 Superseded by DECAMP 92.

## Limits for Excited $\mu$ ( $\mu^*$ ) from Single Production

These limits are from  $e^+e^- \rightarrow \mu^*\mu$  and depend on transition magnetic coupling between  $\mu$  and  $\mu^*$ . All limits assume  $\mu^* \rightarrow \mu\gamma$  decay. Limits from LEP are for chiral coupling, whereas all other limits are for nonchiral coupling,  $\eta_L = \eta_R = 1$ . In most papers, the limit is expressed in the form of an excluded region in the  $\lambda-m_{\mu^*}$  plane. See the original papers.

For limits prior to 1987, see our 1992 edition (Physical Review **D45**, 1 June, Part II (1992)).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>178	95	144 ACCIARRI	01D L3	$e^+e^- \rightarrow \mu\mu^*$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
		145 ABBIENDI	00I OPAL	$e^+e^- \rightarrow \mu\mu^*$
		146 ACCIARRI	00E L3	$e^+e^- \rightarrow \mu\mu^*$
		147 ABREU	99O DLPH	$e^+e^- \rightarrow \mu\mu^*$
		148 ACKERSTAFF	98C OPAL	$e^+e^- \rightarrow \mu\mu^*$
		149 BARATE	98U ALEP	$Z \rightarrow \mu\mu^*$
		150,151 ABREU	97B DLPH	$e^+e^- \rightarrow \mu\mu^*$
		150,152 ACCIARRI	97G L3	$e^+e^- \rightarrow \mu\mu^*$
		153 ACKERSTAFF	97 OPAL	$e^+e^- \rightarrow \mu\mu^*$
		154 ABREU	96K DLPH	$e^+e^- \rightarrow \mu\mu^*$
		155 ACCIARRI	96D L3	$e^+e^- \rightarrow \mu\mu^*$
		156 ALEXANDER	96Q OPAL	$e^+e^- \rightarrow \mu\mu^*$
		157 BUSKULIC	96W ALEP	$e^+e^- \rightarrow \mu\mu^*$
> 89	95	ADRIANI	93M L3	$Z \rightarrow \mu\mu^*$ , $\lambda_Z > 0.5$
> 88	95	ABREU	92C DLPH	$Z \rightarrow \mu\mu^*$ , $\lambda_Z > 0.5$
> 91	95	DECAMP	92 ALEP	$Z \rightarrow \mu\mu^*$ , $\lambda_Z > 1$
> 85	95	158 ADEVA	90F L3	$Z \rightarrow \mu\mu^*$ , $\lambda_Z > 1$
> 75	95	158 ADEVA	90F L3	$Z \rightarrow \mu\mu^*$ , $\lambda_Z > 0.1$
> 87	95	AKRAWY	90I OPAL	$Z \rightarrow \mu\mu^*$ , $\lambda_Z > 1$
> 80	95	159 DECAMP	90G ALEP	$e^+e^- \rightarrow \mu\mu^*$ , $\lambda_Z=1$
> 50	95	ADACHI	89B TOPZ	$e^+e^- \rightarrow \mu\mu^*$ , $\lambda_\gamma=0.7$
> 46	95	KIM	89 AMY	$e^+e^- \rightarrow \mu\mu^*$ , $\lambda_\gamma=0.2$

144 ACCIARRI 01D result is from  $e^+e^-$  collisions at  $\sqrt{s} = 192\text{--}202$  GeV.  $f=f'=1/m_{\mu^*}$  is assumed for the  $\mu^*$  coupling. See their Fig. 4 for limits in the mass-coupling plane.

145 ABBIENDI 00I result is from  $e^+e^-$  collisions at  $\sqrt{s}=161\text{--}183$  GeV. See their Fig. 7 for limits in mass-coupling plane.

146 ACCIARRI 00E result is from  $e^+e^-$  collisions at  $\sqrt{s}=189$  GeV. See their Fig. 3 for limits in mass-coupling plane.

147 ABREU 99O result is from  $e^+e^-$  collisions at  $\sqrt{s}= 183$  GeV. See their Figs. 4 and 5 for the exclusion limit in the mass-coupling plane.

148 ACKERSTAFF 98C from  $e^+e^-$  collisions at  $\sqrt{s}=170\text{--}172$  GeV. See their Fig. 11 for the exclusion limit in the mass-coupling plane.

149 BARATE 98U obtain limits on the  $Z\mu\mu^*$  coupling. See their Fig. 12 for limits in mass-coupling plane

150 From  $e^+e^-$  collisions at  $\sqrt{s}= 161$  GeV.

151 See Fig. 4a and Fig. 5a of ABREU 97B for the exclusion limit in the mass-coupling plane.

152 See Fig. 2 and Fig. 3 of ACCIARRI 97G for the exclusion limit in the mass-coupling plane.

153 ACKERSTAFF 97 result is from  $e^+e^-$  collisions at  $\sqrt{s}= 161$  GeV. See their Fig. 3 for the exclusion limit in the mass-coupling plane.

- 154 ABREU 96K result is from  $e^+e^-$  collisions at  $\sqrt{s}=130\text{--}136$  GeV. See their Fig. 4 for the exclusion limit in the mass-coupling plane.
- 155 ACCIARRI 96D result is from  $e^+e^-$  collisions at  $\sqrt{s}=130\text{--}140$  GeV. See their Fig. 2 for the exclusion limit in the mass-coupling plane.
- 156 ALEXANDER 96Q result is from  $e^+e^-$  collisions at  $\sqrt{s}=130\text{--}140$  GeV. See their Fig. 3a for the exclusion limit in the mass-coupling plane.
- 157 BUSKULIC 96W result is from  $e^+e^-$  collisions at  $\sqrt{s}=130\text{--}140$  GeV. See their Fig. 3 for the exclusion limit in the mass-coupling plane.
- 158 Superseded by ADRIANI 93M.
- 159 Superseded by DECAMP 92.

### Indirect Limits for Excited $\mu$ ( $\mu^*$ )

These limits make use of loop effects involving  $\mu^*$  and are therefore subject to theoretical uncertainty.

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
• • •	We do not use the following data for averages, fits, limits, etc. • • •		
	160 RENARD	82 THEO	$g-2$ of muon
160	RENARD 82 derived from $g-2$ data limits on mass and couplings of $e^*$ and $\mu^*$ . See figures 2 and 3 of the paper.		

## MASS LIMITS for Excited $\tau$ ( $\tau^*$ )

### Limits for Excited $\tau$ ( $\tau^*$ ) from Pair Production

These limits are obtained from  $e^+e^- \rightarrow \tau^{*+}\tau^{*-}$  and thus rely only on the (electroweak) charge of  $\tau^*$ . Form factor effects are ignored unless noted. For the case of limits from  $Z$  decay, the  $\tau^*$  coupling is assumed to be of sequential type. All limits assume a dominant  $\tau^* \rightarrow \tau\gamma$  decay except the limits from  $\Gamma(Z)$ .

For limits prior to 1987, see our 1992 edition (Physical Review **D45**, 1 June, Part II (1992)).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>99.8	95	161 ACCIARRI	01D L3	$e^+e^- \rightarrow \tau^*\tau^*$ Homodoublet type
• • •	We do not use the following data for averages, fits, limits, etc. • • •			
>91.2	95	162 ABBIENDI	00I OPAL	$e^+e^- \rightarrow \tau^*\tau^*$ Homodoublet type
>94.2	95	163 ACCIARRI	00E L3	$e^+e^- \rightarrow \tau^*\tau^*$ Homodoublet type
>89.7	95	164 ABREU	99O DLPH	Homodoublet type
>84.6	95	165 ACKERSTAFF	98C OPAL	$e^+e^- \rightarrow \tau^*\tau^*$ Homodoublet type
		166 BARATE	98U ALEP	$Z \rightarrow \tau^*\tau^*$
>79.4	95	167,168 ABREU	97B DLPH	$e^+e^- \rightarrow \tau^*\tau^*$ Homodoublet type
>77.4	95	167,169 ABREU	97B DLPH	$e^+e^- \rightarrow \tau^*\tau^*$ Sequential type
>79.3	95	167 ACCIARRI	97G L3	$e^+e^- \rightarrow \tau^*\tau^*$ Sequential type
>79.1	95	167,170 ACKERSTAFF	97 OPAL	$e^+e^- \rightarrow \tau^*\tau^*$ Homodoublet type
>62.2	95	171 ABREU	96K DLPH	$e^+e^- \rightarrow \tau^*\tau^*$ Homodoublet type
>64.2	95	172 ACCIARRI	96D L3	$e^+e^- \rightarrow \tau^*\tau^*$ Sequential type
>65.3	95	172 ALEXANDER	96Q OPAL	$e^+e^- \rightarrow \tau^*\tau^*$ Homodoublet type
>64.8	95	172 BUSKULIC	96W ALEP	$e^+e^- \rightarrow \tau^*\tau^*$ Sequential type

>45.6	95	ADRIANI	93M L3	$Z \rightarrow \tau^* \tau^*$
>45.3	95	ABREU	92C DLPH	$Z \rightarrow \tau^* \tau^*$
>29.8	95	173 BARDADIN-...	92 RVUE	$\Gamma(Z)$
>26.1	95	174 DECAMP	92 ALEP	$Z \rightarrow \tau^* \tau^*; \Gamma(Z)$
>46.0	95	DECAMP	92 ALEP	$Z \rightarrow \tau^* \tau^*$
>33	95	174 ABREU	91F DLPH	$Z \rightarrow \tau^* \tau^*; \Gamma(Z)$
>45.5	95	175 ADEVA	90L L3	$Z \rightarrow \tau^* \tau^*$
>44.9	95	AKRAWY	90I OPAL	$Z \rightarrow \tau^* \tau^*$
>41.2	95	176 DECAMP	90G ALEP	$e^+ e^- \rightarrow \tau^* \tau^*$
>29.0	95	ADACHI	89B TOPZ	$e^+ e^- \rightarrow \tau^* \tau^*$

161 From  $e^+ e^-$  collisions at  $\sqrt{s} = 192\text{--}202$  GeV.  $f=f'$  is assumed. ACCIARRI 01D also obtain limit for  $f=-f'$ :  $m_{\tau^*} > 93.4$  GeV.

162 From  $e^+ e^-$  collisions at  $\sqrt{s}=161\text{--}183$  GeV.  $f=f'$  is assumed. ABBIENDI 00I also obtain limit for  $f=-f'$  ( $\tau^* \rightarrow \nu W$ ):  $m_{\tau^*} > 86.0$  GeV.

163 From  $e^+ e^-$  collisions at  $\sqrt{s}=189$  GeV.  $f=f'$  is assumed. ACCIARRI 00E also obtain limit for  $f=-f'$  ( $\tau^* \rightarrow \nu W$ ):  $m_{\tau^*} > 92.6$  GeV.

164 From  $e^+ e^-$  collisions at  $\sqrt{s}= 183$  GeV.  $f=f'$  is assumed. ABREU 990 also obtain limit for  $f=-f'$  ( $\tau^* \rightarrow \nu W$ ):  $m_{\tau^*} > 81.3$  GeV.

165 From  $e^+ e^-$  collisions at  $\sqrt{s}=170\text{--}172$  GeV. ACKERSTAFF 98C also obtain limit from  $\tau^* \rightarrow \nu W$  decay mode:  $m_{\tau^*} > 81.3$  GeV.

166 BARATE 98U obtain limits on the form factor. See their Fig. 14 for limits in mass-form factor plane.

167 From  $e^+ e^-$  collisions at  $\sqrt{s}= 161$  GeV.

168 ABREU 97B also obtain limit from charged current decay mode  $\tau^* \rightarrow \nu W$ ,  $m_{\tau^*} > 70.9$  GeV.

169 ABREU 97B also obtain limit from charged current decay mode  $\tau^* \rightarrow \nu W$ ,  $m_{\tau^*} > 44.6$  GeV.

170 ACKERSTAFF 97 also obtain limit from charged current decay mode  $\tau^* \rightarrow \nu W$ ,  $m_{\nu_{\tau^*}} > 77.1$  GeV.

171 From  $e^+ e^-$  collisions at  $\sqrt{s}= 130\text{--}136$  GeV.

172 From  $e^+ e^-$  collisions at  $\sqrt{s}= 130\text{--}140$  GeV.

173 BARDADIN-OTWINOWSKA 92 limit is independent of decay modes. Based on  $\Delta\Gamma(Z)<36$  MeV.

174 Limit is independent of  $\tau^*$  decay mode.

175 Superseded by ADRIANI 93M.

176 Superseded by DECAMP 92.

## Limits for Excited $\tau$ ( $\tau^*$ ) from Single Production

These limits are from  $e^+e^- \rightarrow \tau^*\tau$  and depend on transition magnetic coupling between  $\tau$  and  $\tau^*$ . All limits assume  $\tau^* \rightarrow \tau\gamma$  decay. Limits from LEP are for chiral coupling, whereas all other limits are for nonchiral coupling,  $\eta_L = \eta_R = 1$ . In most papers, the limit is expressed in the form of an excluded region in the  $\lambda-m_{\tau^*}$  plane. See the original papers.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;173</b>	95	177 ACCIARRI	01D L3	$e^+e^- \rightarrow \tau\tau^*$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
		178 ABBIENDI	00I OPAL	$e^+e^- \rightarrow \tau\tau^*$
		179 ACCIARRI	00E L3	$e^+e^- \rightarrow \tau\tau^*$
		180 ABREU	99O DLPH	$e^+e^- \rightarrow \tau\tau^*$
		181 ACKERSTAFF	98C OPAL	$e^+e^- \rightarrow \tau\tau^*$
		182 BARATE	98U ALEP	$Z \rightarrow \tau\tau^*$
		183,184 ABREU	97B DLPH	$e^+e^- \rightarrow \tau\tau^*$
		183,185 ACCIARRI	97G L3	$e^+e^- \rightarrow \tau\tau^*$
		186 ACKERSTAFF	97 OPAL	$e^+e^- \rightarrow \tau\tau^*$
		187 ABREU	96K DLPH	$e^+e^- \rightarrow \tau\tau^*$
		188 ACCIARRI	96D L3	$e^+e^- \rightarrow \tau\tau^*$
		189 ALEXANDER	96Q OPAL	$e^+e^- \rightarrow \tau\tau^*$
		190 BUSKULIC	96W ALEP	$e^+e^- \rightarrow \tau\tau^*$
> 88	95	ADRIANI	93M L3	$Z \rightarrow \tau\tau^*$ , $\lambda_Z > 0.5$
> 87	95	ABREU	92C DLPH	$Z \rightarrow \tau\tau^*$ , $\lambda_Z > 0.5$
> 90	95	DECAMP	92 ALEP	$Z \rightarrow \tau\tau^*$ , $\lambda_Z > 0.18$
> 88	95	191 ADEVA	90L L3	$Z \rightarrow \tau\tau^*$ , $\lambda_Z > 1$
> 86.5	95	AKRAWY	90I OPAL	$Z \rightarrow \tau\tau^*$ , $\lambda_Z > 1$
> 59	95	192 DECAMP	90G ALEP	$Z \rightarrow \tau\tau^*$ , $\lambda_Z = 1$
> 40	95	193 BARTEL	86 JADE	$e^+e^- \rightarrow \tau\tau^*$ , $\lambda_\gamma = 1$
> 41.4	95	194 BEHREND	86 CELL	$e^+e^- \rightarrow \tau\tau^*$ , $\lambda_\gamma = 1$
> 40.8	95	194 BEHREND	86 CELL	$e^+e^- \rightarrow \tau\tau^*$ , $\lambda_\gamma = 0.7$

177 ACCIARRI 01D result is from  $e^+e^-$  collisions at  $\sqrt{s} = 192-202$  GeV.  $f=f'=\Lambda/m_{\tau^*}$  is assumed for the  $\tau^*$  coupling. See their Fig. 4 for limits in the mass-coupling plane.

178 ABBIENDI 00I result is from  $e^+e^-$  collisions at  $\sqrt{s}=161-183$  GeV. See their Fig. 7 for limits in mass-coupling plane.

179 ACCIARRI 00E result is from  $e^+e^-$  collisions at  $\sqrt{s}=189$  GeV. See their Fig. 3 for limits in mass-coupling plane.

180 ABREU 99O result is from  $e^+e^-$  collisions at  $\sqrt{s}= 183$  GeV. See their Figs. 4 and 5 for the exclusion limit in the mass-coupling plane.

181 ACKERSTAFF 98C from  $e^+e^-$  collisions at  $\sqrt{s}=170-172$  GeV. See their Fig. 11 for the exclusion limit in the mass-coupling plane.

182 BARATE 98U obtain limits on the  $Z\tau\tau^*$  coupling. See their Fig. 12 for limits in mass-coupling plane

183 From  $e^+e^-$  collisions at  $\sqrt{s}= 161$  GeV.

184 See Fig. 4a and Fig. 5a of ABREU 97B for the exclusion limit in the mass-coupling plane.

185 See Fig. 2 and Fig. 3 of ACCIARRI 97G for the exclusion limit in the mass-coupling plane.

186 ACKERSTAFF 97 result is from  $e^+e^-$  collisions at  $\sqrt{s}= 161$  GeV. See their Fig. 3 for the exclusion limit in the mass-coupling plane.

187 ABREU 96K result is from  $e^+e^-$  collisions at  $\sqrt{s}= 130-136$  GeV. See their Fig. 4 for the exclusion limit in the mass-coupling plane.



- 188 ACCIARRI 96D result is from  $e^+e^-$  collisions at  $\sqrt{s}=130\text{--}140$  GeV. See their Fig. 2 for the exclusion limit in the mass-coupling plane.
- 189 ALEXANDER 96Q result is from  $e^+e^-$  collisions at  $\sqrt{s}=130\text{--}140$  GeV. See their Fig. 3a for the exclusion limit in the mass-coupling plane.
- 190 BUSKULIC 96W result is from  $e^+e^-$  collisions at  $\sqrt{s}=130\text{--}140$  GeV. See their Fig. 3 for the exclusion limit in the mass-coupling plane.
- 191 Superseded by ADRIANI 93M.
- 192 Superseded by DECAMP 92.
- 193 BARTEL 86 is at  $E_{\text{cm}} = 30\text{--}46.78$  GeV.
- 194 BEHREND 86 limit is at  $E_{\text{cm}} = 33\text{--}46.8$  GeV.

## MASS LIMITS for Excited Neutrino ( $\nu^*$ )

### Limits for Excited $\nu$ ( $\nu^*$ ) from Pair Production

These limits are obtained from  $e^+e^- \rightarrow \nu^*\nu^*$  and thus rely only on the (electroweak) charge of  $\nu^*$ . Form factor effects are ignored unless noted. The  $\nu^*$  coupling is assumed to be of sequential type unless otherwise noted. All limits assume a dominant  $\nu^* \rightarrow \nu\gamma$  decay except the limits from  $\Gamma(Z)$ .

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;99.4</b>	95	195 ACCIARRI	01D L3	$e^+e^- \rightarrow \nu^*\nu^*$ Homodoublet type
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
>91.2	95	196 ABBIENDI	00I OPAL	$e^+e^- \rightarrow \nu^*\nu^*$ Homodoublet type
		197 ABBIENDI,G	00D OPAL	
>94.1	95	198 ACCIARRI	00E L3	$e^+e^- \rightarrow \nu^*\nu^*$ Homodoublet type
		199 ABBIENDI	99F OPAL	
>90.0	95	200 ABREU	99O DLPH	Homodoublet type
>84.9	95	201 ACKERSTAFF	98C OPAL	$e^+e^- \rightarrow \nu^*\nu^*$ Homodoublet type
		202 BARATE	98U ALEP	$Z \rightarrow \nu^*\nu^*$
>77.6	95	203,204 ABREU	97B DLPH	$e^+e^- \rightarrow \nu^*\nu^*$ Homodoublet type
>64.4	95	203,205 ABREU	97B DLPH	$e^+e^- \rightarrow \nu^*\nu^*$ Sequential type
>71.2	95	203,206 ACCIARRI	97G L3	$e^+e^- \rightarrow \nu^*\nu^*$ Sequential type
>77.8	95	203,207 ACKERSTAFF	97 OPAL	$e^+e^- \rightarrow \nu^*\nu^*$ Homodoublet type
>61.4	95	208,209 ACCIARRI	96D L3	$e^+e^- \rightarrow \nu^*\nu^*$ Sequential type
>65.0	95	210,211 ALEXANDER	96Q OPAL	$e^+e^- \rightarrow \nu^*\nu^*$ Homodoublet type
>63.6	95	208 BUSKULIC	96W ALEP	$e^+e^- \rightarrow \nu^*\nu^*$ Sequential type
>43.7	95	212 BARDADIN-...	92 RVUE	$\Gamma(Z)$
>47	95	213 DECAMP	92 ALEP	
>42.6	95	214 DECAMP	92 ALEP	$\Gamma(Z)$
>35.4	95	215,216 DECAMP	90O ALEP	$\Gamma(Z)$
>46	95	216,217 DECAMP	90O ALEP	

195 From  $e^+e^-$  collisions at  $\sqrt{s} = 192\text{--}202$  GeV.  $f=f'$  is assumed. ACCIARRI 01D also obtain limit for  $f=-f'$ :  $m_{\nu_e^*} > 99.1$  GeV,  $m_{\nu_\mu^*} > 99.3$  GeV,  $m_{\nu_\tau^*} > 90.5$  GeV.

196 From  $e^+e^-$  collisions at  $\sqrt{s}=161\text{--}183$  GeV.  $f=-f'$  (photonic decay) is assumed. ABBIENDI 00I also obtain limit for  $f=f'$  ( $\nu^* \rightarrow \ell W$ ):  $m_{\nu_e^*} > 91.1$  GeV,  $m_{\nu_\mu^*} > 91.1$  GeV,  $m_{\nu_\tau^*} > 83.1$  GeV.

- 197 From  $e^+e^-$  collisions at  $\sqrt{s}=189$  GeV. ABBIENDI, G 00D obtain limit on  $\sigma(e^+e^- \rightarrow \nu^*\nu^*)B(\nu^* \rightarrow \nu\gamma)^2$ . See their Fig. 14. The limit ranges from 50 to 80 fb for  $\sqrt{s}/2 > m_{\nu^*} > 45$  GeV.
- 198 From  $e^+e^-$  collisions at  $\sqrt{s}=189$  GeV.  $f=-f'$  (photonic decay) is assumed. ACCIARRI 00E also obtain limit for  $f=f'$  ( $\nu^* \rightarrow \ell W$ ):  $m_{\nu_e^*} > 93.9$  GeV,  $m_{\nu_\mu^*} > 94.0$  GeV,  $m_{\nu_\tau^*} > 91.5$  GeV.
- 199 From  $e^+e^-$  collisions at  $\sqrt{s}=130-183$  GeV, ABBIENDI 99F obtain limit on  $\sigma(e^+e^- \rightarrow \nu^*\nu^*)B(\nu^* \rightarrow \nu\gamma)^2$ . See their Fig. 13. The limit ranges from 0.094 to 0.14 pb for  $\sqrt{s}/2 > m_{\nu^*} > 45$  GeV.
- 200 From  $e^+e^-$  collisions at  $\sqrt{s}=183$  GeV.  $f=-f'$  is assumed. ABREU 99O also obtain limit for  $f=f'$ :  $m_{\nu_{e^*}} > 87.3$  GeV,  $m_{\nu_{\mu^*}} > 88.0$  GeV,  $m_{\nu_{\tau^*}} > 81.0$  GeV.
- 201 From  $e^+e^-$  collisions at  $\sqrt{s}=170-172$  GeV. ACKERSTAFF 98C also obtain limit from charged decay modes:  $m_{\nu_e^*} > 84.1$  GeV,  $m_{\nu_\mu^*} > 83.9$  GeV, and  $m_{\nu_\tau^*} > 79.4$  GeV.
- 202 BARATE 98U obtain limits on the form factor. See their Fig. 14 for limits in mass-form factor plane.
- 203 From  $e^+e^-$  collisions at  $\sqrt{s}=161$  GeV.
- 204 ABREU 97B also obtain limits from charged current decay modes,  $m_{\nu^*} > 56.4$  GeV.
- 205 ABREU 97B also obtain limits from charged current decay modes,  $m_{\nu^*} > 44.9$  GeV.
- 206 ACCIARRI 97G also obtain limits from charged current decay mode  $\nu_e^* \rightarrow eW$ ,  $m_{\nu^*} > 64.5$  GeV.
- 207 ACKERSTAFF 97 also obtain limits from charged current decay modes  $m_{\nu_e^*} > 78.3$  GeV,  $m_{\nu_\mu^*} > 78.9$  GeV,  $m_{\nu_\tau^*} > 76.2$  GeV.
- 208 From  $e^+e^-$  collisions at  $\sqrt{s}=130-140$  GeV.
- 209 ACCIARRI 96D also obtain limit from  $\nu^* \rightarrow eW$  decay mode:  $m_{\nu^*} > 57.3$  GeV.
- 210 From  $e^+e^-$  collisions at  $\sqrt{s}=130-136$  GeV.
- 211 ALEXANDER 96Q also obtain limits from charged current decay modes:  $m_{\nu_e^*} > 66.2$  GeV,  $m_{\nu_\mu^*} > 66.5$  GeV,  $m_{\nu_\tau^*} > 64.7$  GeV.
- 212 BARDADIN-OTWINOWSKA 92 limit is for Dirac  $\nu^*$ . Based on  $\Delta\Gamma(Z) < 36$  MeV. The limit is 36.4 GeV for Majorana  $\nu^*$ , 45.4 GeV for homodoublet  $\nu^*$ .
- 213 Limit is based on  $B(Z \rightarrow \nu^*\bar{\nu}^*) \times B(\nu^* \rightarrow \nu\gamma)^2 < 5 \times 10^{-5}$  (95%CL) assuming Dirac  $\nu^*$ ,  $B(\nu^* \rightarrow \nu\gamma) = 1$ .
- 214 Limit is for Dirac  $\nu^*$ . The limit is 34.6 GeV for Majorana  $\nu^*$ , 45.4 GeV for homodoublet  $\nu^*$ .
- 215 DECAMP 90O limit is from excess  $\Delta\Gamma(Z) < 89$  MeV. The above value is for Dirac  $\nu^*$ ; 26.6 GeV for Majorana  $\nu^*$ ; 44.8 GeV for homodoublet  $\nu^*$ .
- 216 Superseded by DECAMP 92.
- 217 DECAMP 90O limit based on  $B(Z \rightarrow \nu^*\nu^*) \cdot B(\nu^* \rightarrow \nu\gamma)^2 < 7 \times 10^{-5}$  (95%CL), assuming Dirac  $\nu^*$ ,  $B(\nu^* \rightarrow \nu\gamma) = 1$ .

### Limits for Excited $\nu$ ( $\nu^*$ ) from Single Production

These limits are from  $e^+e^- \rightarrow \nu\nu^*$ ,  $Z \rightarrow \nu\nu^*$ , or  $ep \rightarrow \nu^*X$  and depend on transition magnetic coupling between  $\nu/e$  and  $\nu^*$ . Assumptions about  $\nu^*$  decay mode are given in footnotes.

<u>VALUE (GeV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
>171	95	218 ACCIARRI	01D L3	$e^+e^- \rightarrow \nu\nu^*$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
none 50–150	95	219 ADLOFF	02 H1	$ep \rightarrow \nu^*X$
		220 ABBIENDI	00I OPAL	$e^+e^- \rightarrow \nu\nu^*$
		221 ABBIENDI,G	00D OPAL	
		222 ACCIARRI	00E L3	$e^+e^- \rightarrow \nu\nu^*$
>114	95	223 ADLOFF	00E H1	$ep \rightarrow \nu^*X$
		224 ABBIENDI	99F OPAL	
		225 ABREU	99O DLPH	$e^+e^- \rightarrow \nu\nu^*$
		226 ACKERSTAFF	98C OPAL	$e^+e^- \rightarrow \nu^*\nu^*$ Homodoublet type
		227 BARATE	98U ALEP	$Z \rightarrow \nu\nu^*$
	228,229	ABREU	97B DLPH	$e^+e^- \rightarrow \nu\nu^*$
		230 ABREU	97I DLPH	$\nu^* \rightarrow \ell W, \nu Z$
		231 ABREU	97J DLPH	$\nu^* \rightarrow \nu\gamma$
	228,232	ACCIARRI	97G L3	$e^+e^- \rightarrow \nu\nu^*$
		233 ACKERSTAFF	97 OPAL	$e^+e^- \rightarrow \nu\nu^*$
		234 ADLOFF	97 H1	Lepton-flavor violation
none 40–96	95	235 BREITWEG	97C ZEUS	$ep \rightarrow \nu^*X$
		236 ACCIARRI	96D L3	$e^+e^- \rightarrow \nu\nu^*$
		237 ALEXANDER	96Q OPAL	$e^+e^- \rightarrow \nu\nu^*$
		238 BUSKULIC	96W ALEP	$e^+e^- \rightarrow \nu\nu^*$
		239 DERRICK	95B ZEUS	$ep \rightarrow \nu^*X$
		240 ABT	93 H1	$ep \rightarrow \nu^*X$
> 91	95	ADRIANI	93M L3	$\lambda_Z > 1, \nu^* \rightarrow \nu\gamma$
> 89	95	ADRIANI	93M L3	$\lambda_Z > 1, \nu_e^* \rightarrow eW$
> 87	95	ADRIANI	93M L3	$\lambda_Z > 0.1, \nu^* \rightarrow \nu\gamma$
> 74	95	ADRIANI	93M L3	$\lambda_Z > 0.1, \nu_e^* \rightarrow eW$
		241 BARDADIN-...	92 RVUE	
> 91	95	242 DECAMP	92 ALEP	$\lambda_Z > 1$
> 74	95	242 DECAMP	92 ALEP	$\lambda_Z > 0.034$
> 91	95	243,244 ADEVA	90O L3	$\lambda_Z > 1$
> 83	95	244 ADEVA	90O L3	$\lambda_Z > 0.1, \nu^* \rightarrow \nu\gamma$
> 74	95	244 ADEVA	90O L3	$\lambda_Z > 0.1, \nu_e^* \rightarrow eW$
> 90	95	245,246 DECAMP	90O ALEP	$\lambda_Z > 1$
> 74.7	95	245,246 DECAMP	90O ALEP	$\lambda_Z > 0.06$

218 ACCIARRI 01D search for  $\nu\nu^*$  production in  $e^+e^-$  collisions at  $\sqrt{s} = 192\text{--}202$  GeV with decays  $\nu^* \rightarrow \nu\gamma, \nu^* \rightarrow eW$ .  $f = -f' = \Lambda/m_{\nu^*}$  is assumed for the  $\nu^*$  coupling. See their Fig. 4 for limits in the mass-coupling plane.

219 ADLOFF 02 search for single  $\nu^*$  production in  $ep$  collisions with the decays  $\nu^* \rightarrow \nu\gamma, \nu Z, eW$ . The quoted limit assumes  $f = -f' = \Lambda/m_{\nu^*}$ . See their Fig. 1 for the exclusion plots in the mass-coupling plane.

220 ABBIENDI 00I result is from  $e^+e^-$  collisions at  $\sqrt{s}=161\text{--}183$  GeV. See their Fig. 7 for limits in mass-coupling plane.

- 221 From  $e^+e^-$  collisions at  $\sqrt{s}=189$  GeV. ABBIENDI, G 00D obtain limit on  $\sigma(e^+e^- \rightarrow \nu^*\nu^*)B(\nu^* \rightarrow \nu\gamma)^2$ . See their Fig. 11.
- 222 ACCIARRI 00E result is from  $e^+e^-$  collisions at  $\sqrt{s}=189$  GeV. See their Fig. 3 for limits in mass-coupling plane.
- 223 ADLOFF 00E search for single  $\nu^*$  production in  $ep$  collisions with the decays  $\nu^* \rightarrow \nu\gamma, \nu Z, eW$ . The quoted limit assumes  $f=-f'=\Lambda/m_{\nu^*}$ . See their Fig. 10 for the exclusion plot in the mass-coupling plane.
- 224 From  $e^+e^-$  collisions at  $\sqrt{s}=130-183$  GeV, ABBIENDI 99F obtain limit on  $\sigma(e^+e^- \rightarrow \nu\nu^*)B(\nu^* \rightarrow \nu\gamma)$ . See their Fig. 8.
- 225 ABREU 99O result is from  $e^+e^-$  collisions at  $\sqrt{s}=183$  GeV. See their Figs. 4 and 5 for the exclusion limit in the mass-coupling plane.
- 226 ACKERSTAFF 98C from  $e^+e^-$  collisions at  $\sqrt{s}=170-172$  GeV. See their Fig. 11 for the exclusion limit in the mass-coupling plane.
- 227 BARATE 98U obtain limits on the  $Z\nu\nu^*$  coupling. See their Fig. 13 for limits in mass-coupling plane
- 228 From  $e^+e^-$  collisions at  $\sqrt{s}=161$  GeV.
- 229 See Fig. 4b and Fig. 5b of ABREU 97B for the exclusion limit in the mass-coupling plane.
- 230 ABREU 97I limit is from  $Z \rightarrow \nu\nu^*$ . See their Fig. 12 for the exclusion limit in the mass-coupling plane.
- 231 ABREU 97J limit is from  $Z \rightarrow \nu\nu^*$ . See their Fig. 5 for the exclusion limit in the mass-coupling plane.
- 232 See Fig. 2 and Fig. 3 of ACCIARRI 97G for the exclusion limit in the mass-coupling plane.
- 233 ACKERSTAFF 97 result is from  $e^+e^-$  collisions at  $\sqrt{s}=161$  GeV, for homodoublet  $\nu^*$ . See their Fig. 3 for the exclusion limit in the mass-coupling plane.
- 234 ADLOFF 97 search for single  $e^*$  production in  $ep$  collisions with the decays  $e^* \rightarrow e\gamma, eZ, \nu W$ . See their Fig. 4 for the rejection limits on the product of the production cross section and the branching ratio.
- 235 BREITWEG 97C search for single  $\nu^*$  production in  $ep$  collisions with the decay  $\nu^* \rightarrow \nu\gamma$ .  $f=-f'=2\Lambda/m_{\nu^*}$  is assumed for the  $\nu^*$  coupling. See their Fig. 10 for the exclusion plot in the mass-coupling plane.
- 236 ACCIARRI 96D result is from  $e^+e^-$  collisions at  $\sqrt{s}=130-140$  GeV. See their Fig. 2 for the exclusion limit in the mass-coupling plane.
- 237 ALEXANDER 96Q result is from  $e^+e^-$  collisions at  $\sqrt{s}=130-140$  GeV for homedoublet  $\nu^*$ . See their Fig. 3b and Fig. 3c for the exclusion limit in the mass-coupling plane.
- 238 BUSKULIC 96W result is from  $e^+e^-$  collisions at  $\sqrt{s}=130-140$  GeV. See their Fig. 4 for the exclusion limit in the mass-coupling plane.
- 239 DERRICK 95B search for single  $\nu^*$  production via  $\nu^*eW$  coupling in  $ep$  collisions with the decays  $\nu^* \rightarrow \nu\gamma, \nu Z, eW$ . See their Fig. 14 for the exclusion plot in the  $m_{\nu^*}-\lambda\gamma$  plane.
- 240 ABT 93 search for single  $\nu^*$  production via  $\nu^*eW$  coupling in  $ep$  collisions with the decays  $\nu^* \rightarrow \nu\gamma, \nu Z, eW$ . See their Fig. 4 for exclusion plot in the  $m_{\nu^*}-\lambda_W$  plane.
- 241 See Fig. 5 of BARDADIN-OTWINOWSKA 92 for combined limit of ADEVA 900, DECAMP 900, and DECAMP 92.
- 242 DECAMP 92 limit is based on  $B(Z \rightarrow \nu^*\bar{\nu}) \times B(\nu^* \rightarrow \nu\gamma) < 2.7 \times 10^{-5}$  (95%CL) assuming Dirac  $\nu^*$ ,  $B(\nu^* \rightarrow \nu\gamma) = 1$ .
- 243 Limit is either for  $\nu^* \rightarrow \nu\gamma$  or  $\nu^* \rightarrow eW$ .
- 244 Superseded by ADRIANI 93M.
- 245 DECAMP 900 limit based on  $B(Z \rightarrow \nu\nu^*) \cdot B(\nu^* \rightarrow \nu\gamma) < 6 \times 10^{-5}$  (95%CL), assuming  $B(\nu^* \rightarrow \nu\gamma) = 1$ .
- 246 Superseded by DECAMP 92.

## MASS LIMITS for Excited $q$ ( $q^*$ )

### Limits for Excited $q$ ( $q^*$ ) from Pair Production

These limits are obtained from  $e^+ e^- \rightarrow q^* \bar{q}^*$  and thus rely only on the (electroweak) charge of the  $q^*$ . Form factor effects are ignored unless noted. Assumptions about the  $q^*$  decay are given in the comments and footnotes.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;45.6</b>	95	<sup>247</sup> ADRIANI	93M L3	$u$ or $d$ type, $Z \rightarrow q^* q^*$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
		<sup>248</sup> BARATE	98U ALEP	$Z \rightarrow q^* q^*$
		<sup>249</sup> ADRIANI	92F L3	$Z \rightarrow q^* q^*$
>41.7	95	<sup>250</sup> BARDADIN-...	92 RVUE	$u$ -type, $\Gamma(Z)$
>44.7	95	<sup>250</sup> BARDADIN-...	92 RVUE	$d$ -type, $\Gamma(Z)$
>40.6	95	<sup>251</sup> DECAMP	92 ALEP	$u$ -type, $\Gamma(Z)$
>44.2	95	<sup>251</sup> DECAMP	92 ALEP	$d$ -type, $\Gamma(Z)$
>45	95	<sup>252</sup> DECAMP	92 ALEP	$u$ or $d$ type, $Z \rightarrow q^* q^*$
>45	95	<sup>251</sup> ABREU	91F DLPH	$u$ -type, $\Gamma(Z)$
>45	95	<sup>251</sup> ABREU	91F DLPH	$d$ -type, $\Gamma(Z)$
>21.1	95	<sup>253</sup> BEHREND	86C CELL	$e(q^*) = -1/3$ , $q^* \rightarrow$ $qg$
>22.3	95	<sup>253</sup> BEHREND	86C CELL	$e(q^*) = 2/3$ , $q^* \rightarrow qg$
>22.5	95	<sup>253</sup> BEHREND	86C CELL	$e(q^*) = -1/3$ , $q^* \rightarrow$ $q\gamma$
>23.2	95	<sup>253</sup> BEHREND	86C CELL	$e(q^*) = 2/3$ , $q^* \rightarrow q\gamma$

<sup>247</sup> ADRIANI 93M limit is valid for  $B(q^* \rightarrow qg) > 0.25$  (0.17) for up (down) type.

<sup>248</sup> BARATE 98U obtain limits on the form factor. See their Fig. 16 for limits in mass-form factor plane.

<sup>249</sup> ADRIANI 92F search for  $Z \rightarrow q^* \bar{q}^*$  followed with  $q^* \rightarrow q\gamma$  decays and give the limit  $\sigma_Z \cdot B(Z \rightarrow q^* \bar{q}^*) \cdot B^2(q^* \rightarrow q\gamma) < 2$  pb at 95%CL. Assuming five flavors of degenerate  $q^*$  of homodoublet type,  $B(q^* \rightarrow q\gamma) < 4\%$  is obtained for  $m_{q^*} < 45$  GeV.

<sup>250</sup> BARDADIN-OTWINOWSKA 92 limit based on  $\Delta\Gamma(Z) < 36$  MeV.

<sup>251</sup> These limits are independent of decay modes.

<sup>252</sup> Limit is for  $B(q^* \rightarrow qg) + B(q^* \rightarrow q\gamma) = 1$ .

<sup>253</sup> BEHREND 86C search for  $e^+ e^- \rightarrow q^* \bar{q}^*$  for  $m_{q^*} > 5$  GeV. But  $m < 5$  GeV excluded by total hadronic cross section. The limits are for point-like photon couplings of excited quarks.

### Limits for Excited $q$ ( $q^*$ ) from Single Production

These limits are from  $e^+ e^- \rightarrow q^* \bar{q}$  or  $p\bar{p} \rightarrow q^* X$  and depend on transition magnetic couplings between  $q$  and  $q^*$ . Assumptions about  $q^*$  decay mode are given in the footnotes and comments.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt; 570, none 580–760 (CL = 95%) OUR EVALUATION</b>				
none 200–520 and <b>580–760</b>	95	<sup>254</sup> ABE	97G CDF	$p\bar{p} \rightarrow q^* X$ , $q^* \rightarrow 2$ jets
<b>none 80–570</b>	95	<sup>255</sup> ABE	95N CDF	$p\bar{p} \rightarrow q^* X$ , $q^* \rightarrow qg$ $q\gamma$ , $qW$
<b>&gt;288</b>	90	<sup>256</sup> ALITTI	93 UA2	$p\bar{p} \rightarrow q^* X$ , $q^* \rightarrow qg$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

>188	95	257 ADLOFF	00E H1	$e p \rightarrow q^* X$
		258 ABREU	990 DLPH	$e^+ e^- \rightarrow q q^*$
		259 BARATE	98U ALEP	$Z \rightarrow q q^*$
		260 ADLOFF	97 H1	Lepton-flavor violation
none 40–169	95	261 BREITWEG	97C ZEUS	$e p \rightarrow q^* X$
		262 DERRICK	95B ZEUS	$e p \rightarrow q^* X$
none 80–540	95	263 ABE	94 CDF	$p\bar{p} \rightarrow q^* X, q^* \rightarrow q\gamma, qW$
> 79	95	264 ADRIANI	93M L3	$\lambda_Z(L3) > 0.06$
		265 ABREU	92D DLPH	$Z \rightarrow q q^*$
		266 ADRIANI	92F L3	$Z \rightarrow q q^*$
> 75	95	264 DECAMP	92 ALEP	$Z \rightarrow q q^*, \lambda_Z > 1$
> 88	95	267 DECAMP	92 ALEP	$Z \rightarrow q q^*, \lambda_Z > 1$
> 86	95	267 AKRAWY	90J OPAL	$Z \rightarrow q q^*, \lambda_Z > 1.2$
		268 ALBAJAR	89 UA1	$p\bar{p} \rightarrow q^* X, q^* \rightarrow qW$
> 39	95	269 BEHREND	86C CELL	$e^+ e^- \rightarrow q^* \bar{q} (q^* \rightarrow qg, q\gamma), \lambda_\gamma=1$

254 ABE 97G search for new particle decaying to dijets.

255 ABE 95N assume a degenerate  $u^*$  and  $d^*$  with  $f_s=f=f'= \Lambda/m_{q^*}$ . See their Fig. 4 for the excluded region in  $m_{q^*} - f$  plane.

256 ALITTI 93 search for resonances in the two-jet invariant mass. The limit is for  $f_s = f = f' = \Lambda/m_{q^*}$ .  $u^*$  and  $d^*$  are assumed to be degenerate. If not, the limit for  $u^*$  ( $d^*$ ) is 277 (247) GeV if  $m_{d^*} \gg m_{u^*}$  ( $m_{u^*} \gg m_{d^*}$ ).

257 ADLOFF 00E search for single  $q^*$  production in  $e p$  collisions with the decays  $q^* \rightarrow q\gamma, qZ, qW$ .  $f_s=0$  and  $f=f'= \Lambda/m_{q^*}$  is assumed for the  $q^*$  coupling. See their Fig. 11 for the exclusion plot in the mass-coupling plane.

258 ABREU 990 result is from  $e^+ e^-$  collisions at  $\sqrt{s}= 183$  GeV. See their Fig. 6 for the exclusion limit in the mass-coupling plane.

259 BARATE 98U obtain limits on the  $Z q q^*$  coupling. See their Fig. 16 for limits in mass-coupling plane

260 ADLOFF 97 search for single  $q^*$  production in  $e p$  collisions with the decay  $q^* \rightarrow q\gamma$ . See their Fig. 6 for the rejection limits on the product of the production cross section and the branching ratio.

261 BREITWEG 97C search for single  $q^*$  production in  $e p$  collisions with the decays  $q^* \rightarrow q\gamma, qW$ .  $f_s=0$ , and  $f=-f'=2\Lambda/m_{q^*}$  is assumed for the  $q^*$  coupling. See their Fig. 11 for the exclusion plot in the mass-coupling plane.

262 DERRICK 95B search for single  $q^*$  production via  $q^* q\gamma$  coupling in  $e p$  collisions with the decays  $q^* \rightarrow qW, qZ, qg, q\gamma$ . See their Fig. 15 for the exclusion plot in the  $m_{q^*} - \lambda_\gamma$  plane.

263 ABE 94 search for resonances in jet- $\gamma$  and jet- $W$  invariant mass in  $p\bar{p}$  collisions at  $E_{CM} = 1.8$  TeV. The limit is for  $f_s = f = f' = \Lambda/m_{q^*}$  and  $u^*$  and  $d^*$  are assumed to be degenerate. See their Fig. 4 for the excluded region in  $m_{q^*} - f$  plane.

264 Assumes  $B(q^* \rightarrow qg) = 1$ .

265 ABREU 92D give  $\sigma(e^+ e^- \rightarrow Z \rightarrow q^* \bar{q} \text{ or } q \bar{q}^*) \times B(q^* \rightarrow q\gamma) < 15$  pb (95% CL) for  $m_{q^*} < 80$  GeV.

- 266 ADRIANI 92F search for  $Z \rightarrow qq^*$  with  $q^* \rightarrow q\gamma$  and give the limit  $\sigma_Z \cdot B(Z \rightarrow qq^*) \cdot B(q^* \rightarrow q\gamma) < (2-10) \text{ pb}$  (95%CL) for  $m_{q^*} = (46-82) \text{ GeV}$ .
- 267 Assumes  $B(q^* \rightarrow q\gamma) = 0.1$ .
- 268 ALBAJAR 89 give  $\sigma(q^* \rightarrow W + \text{jet})/\sigma(W) < 0.019$  (90% CL) for  $m_{q^*} > 220 \text{ GeV}$ .
- 269 BEHREND 86C has  $E_{\text{cm}} = 42.5-46.8 \text{ GeV}$ . See their Fig. 3 for excluded region in the  $m_{q^*} - (\lambda_\gamma/m_{q^*})^2$  plane. The limit is for  $\lambda_\gamma = 1$  with  $\eta_L = \eta_R = 1$ .

### MASS LIMITS for Color Sextet Quarks ( $q_6$ )

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;84</b>	95	270 ABE	89D CDF	$p\bar{p} \rightarrow q_6\bar{q}_6$

- 270 ABE 89D look for pair production of unit-charged particles which leave the detector before decaying. In the above limit the color sextet quark is assumed to fragment into a unit-charged or neutral hadron with equal probability and to have long enough lifetime not to decay within the detector. A limit of 121 GeV is obtained for a color decuplet.

### MASS LIMITS for Color Octet Charged Leptons ( $l_8$ )

$$\lambda \equiv m_{l_8}/\Lambda$$

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;86</b>	95	271 ABE	89D CDF	Stable $l_8: p\bar{p} \rightarrow l_8\bar{l}_8$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
		272 ABT	93 H1	$e_8: ep \rightarrow e_8 X$
none 3.0-30.3	95	273 KIM	90 AMY	$e_8: e^+e^- \rightarrow ee + \text{jets}$
none 3.5-30.3	95	273 KIM	90 AMY	$\mu_8: e^+e^- \rightarrow \mu\mu + \text{jets}$
		274 KIM	90 AMY	$e_8: e^+e^- \rightarrow gg; R$
>19.8	95	275 BARTEL	87B JADE	$e_8, \mu_8, \tau_8: e^+e^-; R$
none 5-23.2	95	275 BARTEL	87B JADE	$\mu_8: e^+e^- \rightarrow \mu\mu + \text{jets}$
		276 BARTEL	85K JADE	$e_8: e^+e^- \rightarrow gg; R$

- 271 ABE 89D look for pair production of unit-charged particles which leave the detector before decaying. In the above limit the color octet lepton is assumed to fragment into a unit-charged or neutral hadron with equal probability and to have long enough lifetime not to decay within the detector. The limit improves to 99 GeV if it always fragments into a unit-charged hadron.
- 272 ABT 93 search for  $e_8$  production via  $e$ -gluon fusion in  $ep$  collisions with  $e_8 \rightarrow eg$ . See their Fig. 3 for exclusion plot in the  $m_{e_8} - \Lambda$  plane for  $m_{e_8} = 35-220 \text{ GeV}$ .
- 273 KIM 90 is at  $E_{\text{cm}} = 50-60.8 \text{ GeV}$ . The same assumptions as in BARTEL 87B are used.
- 274 KIM 90 result  $(m_{e_8} \Lambda_M)^{1/2} > 178.4 \text{ GeV}$  (95%CL,  $\alpha_s = 0.16$  used) is subject to the same restriction as for BARTEL 85K.
- 275 BARTEL 87B is at  $E_{\text{cm}} = 46.3-46.78 \text{ GeV}$ . The limits assume  $l_8$  pair production cross sections to be eight times larger than those of the corresponding heavy lepton pair production.
- 276 In BARTEL 85K,  $R$  can be affected by  $e^+e^- \rightarrow gg$  via  $e_q$  exchange. Their limit  $m_{e_8} > 173 \text{ GeV}$  (CL=95%) at  $\lambda = m_{e_8}/\Lambda_M = 1$  ( $\eta_L = \eta_R = 1$ ) is not listed above because the cross section is sensitive to the product  $\eta_L\eta_R$ , which should be absent in ordinary theory with electronic chiral invariance.

## MASS LIMITS for Color Octet Neutrinos ( $\nu_8$ )

$$\lambda \equiv m_{\ell_8}/\Lambda$$

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;110</b>	90	277 BARGER	89 RVUE	$\nu_8: p\bar{p} \rightarrow \nu_8\bar{\nu}_8$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
none 3.8–29.8	95	278 KIM	90 AMY	$\nu_8: e^+e^- \rightarrow$ acoplanar jets
none 9–21.9	95	279 BARTEL	87B JADE	$\nu_8: e^+e^- \rightarrow$ acoplanar jets

277 BARGER 89 used ABE 89B limit for events with large missing transverse momentum. Two-body decay  $\nu_8 \rightarrow \nu g$  is assumed.

278 KIM 90 is at  $E_{\text{cm}} = 50\text{--}60.8$  GeV. The same assumptions as in BARTEL 87B are used.

279 BARTEL 87B is at  $E_{\text{cm}} = 46.3\text{--}46.78$  GeV. The limit assumes the  $\nu_8$  pair production cross section to be eight times larger than that of the corresponding heavy neutrino pair production. This assumption is not valid in general for the weak couplings, and the limit can be sensitive to its  $SU(2)_L \times U(1)_Y$  quantum numbers.

## MASS LIMITS for $W_8$ (Color Octet $W$ Boson)

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
	280 ALBAJAR	89 UA1	$p\bar{p} \rightarrow W_8 X,$ $W_8 \rightarrow W g$

280 ALBAJAR 89 give  $\sigma(W_8 \rightarrow W + \text{jet})/\sigma(W) < 0.019$  (90% CL) for  $m_{W_8} > 220$  GeV.

## REFERENCES FOR Searches for Quark and Lepton Compositeness

ACHARD	02D	PL B531 28	P. Achard <i>et al.</i>	(L3 Collab.)
ADLOFF	02	PL B525 9	C. Adloff <i>et al.</i>	(H1 Collab.)
ACCIARRI	01D	PL B502 37	M. Acciarri <i>et al.</i>	(L3 Collab.)
AFFOLDER	01I	PRL 87 231803	T. Affolder <i>et al.</i>	(CDF Collab.)
BOURILKOV	01	PR D64 071701	D. Bourilkov	
CHEUNG	01B	PL B517 167	K. Cheung	
ABBIENDI	00I	EPJ C14 73	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	00R	EPJ C13 553	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI,G	00D	EPJ C18 253	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBOTT	00E	PR D62 031101	B. Abbott <i>et al.</i>	(D0 Collab.)
ABREU	00A	PL B491 67	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	00S	PL B485 45	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	00E	PL B473 177	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	00G	PL B475 198	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	00P	PL B489 81	M. Acciarri <i>et al.</i>	(L3 Collab.)
ADLOFF	00	PL B479 358	C. Adloff <i>et al.</i>	(H1 Collab.)
ADLOFF	00E	EPJ C17 567	C. Adloff <i>et al.</i>	(H1 Collab.)
AFFOLDER	00I	PR D62 012004	T. Affolder <i>et al.</i>	(CDF Collab.)
BARATE	00I	EPJ C12 183	R. Barate <i>et al.</i>	(ALEPH Collab.)
BOURILKOV	00	PR D62 076005	D. Bourilkov	
BREITWEG	00B	EPJ C14 239	J. Breitweg <i>et al.</i>	(ZEUS Collab.)
ABBIENDI	99	EPJ C6 1	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	99F	EPJ C8 23	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	99P	PL B465 303	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBOTT	99C	PRL 82 2457	B. Abbott <i>et al.</i>	(D0 Collab.)
ABBOTT	99D	PRL 82 4769	B. Abbott <i>et al.</i>	(D0 Collab.)
ABREU	99A	EPJ C11 383	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	99O	EPJ C8 41	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ZARNECKI	99	EPJ C11 539	A.F. Zarnecki	
ABBOTT	98G	PRL 80 666	B. Abbott <i>et al.</i>	(D0 Collab.)



ABREU	98J	PL B433 429	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	98J	PL B433 163	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	98T	PL B439 183	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACKERSTAFF	98	EPJ C1 21	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ACKERSTAFF	98C	EPJ C1 45	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ACKERSTAFF	98V	EPJ C2 441	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ACKER...K...	98B	PL B438 379	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
BARATE	98J	PL B429 201	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	98U	EPJ C4 571	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARGER	98E	PR D57 391	V. Barger <i>et al.</i>	
BERTRAM	98	PL B443 347	I. Bertram, E.H. Simmons	
McFARLAND	98	EPJ C1 509	K.S. McFarland <i>et al.</i>	(CCFR/NuTeV Collab.)
MIURA	98	PR D57 5345	M. Miura <i>et al.</i>	(VENUS Collab.)
ABE	97G	PR D55 R5263	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	97T	PRL 79 2198	F. Abe <i>et al.</i>	(CDF Collab.)
ABREU	97B	PL B393 245	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	97I	ZPHY C74 57	P. Abreu <i>et al.</i>	(DELPHI Collab.)
Also	97L	ZPHY C75 580 erratum	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	97J	ZPHY C74 577	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	97G	PL B401 139	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	97W	PL B413 159	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACKERSTAFF	97	PL B391 197	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ACKERSTAFF	97C	PL B391 221	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ADLOFF	97	NP B483 44	C. Adloff <i>et al.</i>	(H1 Collab.)
ARIMA	97	PR D55 19	T. Arima <i>et al.</i>	(VENUS Collab.)
BREITWEG	97C	ZPHY C76 631	J. Breitweg <i>et al.</i>	(ZEUS Collab.)
DEANDREA	97	PL B409 277	A. Deandrea	(MARS)
ABE	96	PRL 77 438	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	96S	PRL 77 5336	F. Abe <i>et al.</i>	(CDF Collab.)
ABREU	96K	PL B380 480	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	96D	PL B370 211	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	96L	PL B384 323	M. Acciarri <i>et al.</i>	(L3 Collab.)
ALEXANDER	96K	PL B377 222	G. Alexander <i>et al.</i>	(OPAL Collab.)
ALEXANDER	96Q	PL B386 463	G. Alexander <i>et al.</i>	(OPAL Collab.)
BUSKULIC	96W	PL B385 445	D. Buskalic <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	96Z	PL B384 333	D. Buskalic <i>et al.</i>	(ALEPH Collab.)
ABE	95N	PRL 74 3538	F. Abe <i>et al.</i>	(CDF Collab.)
ACCIARRI	95G	PL B353 136	M. Acciarri <i>et al.</i>	(L3 Collab.)
AID	95	PL B353 578	S. Aid <i>et al.</i>	(H1 Collab.)
DERRICK	95B	ZPHY C65 627	M. Derrick <i>et al.</i>	(ZEUS Collab.)
ABE	94	PRL 72 3004	F. Abe <i>et al.</i>	(CDF Collab.)
DIAZCRUZ	94	PR D49 R2149	J.L. Diaz Cruz, O.A. Sampayo	(CINV)
VELISSARIS	94	PL B331 227	C. Velissaris <i>et al.</i>	(AMY Collab.)
ABE	93G	PRL 71 2542	F. Abe <i>et al.</i>	(CDF Collab.)
ABT	93	NP B396 3	I. Abt <i>et al.</i>	(H1 Collab.)
ADRIANI	93M	PRPL 236 1	O. Adriani <i>et al.</i>	(L3 Collab.)
ALITTI	93	NP B400 3	J. Alitti <i>et al.</i>	(UA2 Collab.)
BUSKULIC	93Q	ZPHY C59 215	D. Buskalic <i>et al.</i>	(ALEPH Collab.)
DERRICK	93B	PL B316 207	M. Derrick <i>et al.</i>	(ZEUS Collab.)
ABE	92B	PRL 68 1463	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	92D	PRL 68 1104	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	92M	PRL 69 2896	F. Abe <i>et al.</i>	(CDF Collab.)
ABREU	92C	ZPHY C53 41	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	92D	ZPHY C53 555	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ADRIANI	92B	PL B288 404	O. Adriani <i>et al.</i>	(L3 Collab.)
ADRIANI	92F	PL B292 472	O. Adriani <i>et al.</i>	(L3 Collab.)
BARDADIN-...	92	ZPHY C55 163	M. Bardadin-Otwinowska	(CLER)
DECAMP	92	PRPL 216 253	D. Decamp <i>et al.</i>	(ALEPH Collab.)
HOWELL	92	PL B291 206	B. Howell <i>et al.</i>	(TOPAZ Collab.)
KROHA	92	PR D46 58	H. Kroha	(ROCH)
PDG	92	PR D45, 1 June, Part II	K. Hikasa <i>et al.</i>	(KEK, LBL, BOST+)
SHIMOZAWA	92	PL B284 144	K. Shimozawa <i>et al.</i>	(TOPAZ Collab.)
ABE	91D	PRL 67 2418	F. Abe <i>et al.</i>	(CDF Collab.)
ABREU	91E	PL B268 296	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	91F	NP B367 511	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ADACHI	91	PL B255 613	I. Adachi <i>et al.</i>	(TOPAZ Collab.)
AKRAWY	91F	PL B257 531	M.Z. Akrawy <i>et al.</i>	(OPAL Collab.)
ALITTI	91B	PL B257 232	J. Alitti <i>et al.</i>	(UA2 Collab.)
BEHREND	91B	ZPHY C51 143	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
BEHREND	91C	ZPHY C51 149	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
Also	91B	ZPHY C51 143	H.J. Behrend <i>et al.</i>	(CELLO Collab.)

ABE	90I	ZPHY C48 13	K. Abe <i>et al.</i>	(VENUS Collab.)
ADEVA	90F	PL B247 177	B. Adeva <i>et al.</i>	(L3 Collab.)
ADEVA	90K	PL B250 199	B. Adeva <i>et al.</i>	(L3 Collab.)
ADEVA	90L	PL B250 205	B. Adeva <i>et al.</i>	(L3 Collab.)
ADEVA	90O	PL B252 525	B. Adeva <i>et al.</i>	(L3 Collab.)
AKRAWY	90F	PL B241 133	M.Z. Akrawy <i>et al.</i>	(OPAL Collab.)
AKRAWY	90I	PL B244 135	M.Z. Akrawy <i>et al.</i>	(OPAL Collab.)
AKRAWY	90J	PL B246 285	M.Z. Akrawy <i>et al.</i>	(OPAL Collab.)
DECAMP	90G	PL B236 501	D. Decamp <i>et al.</i>	(ALEPH Collab.)
DECAMP	90O	PL B250 172	D. Decamp <i>et al.</i>	(ALEPH Collab.)
KIM	90	PL B240 243	G.N. Kim <i>et al.</i>	(AMY Collab.)
ABE	89	PRL 62 613	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	89B	PRL 62 1825	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	89D	PRL 63 1447	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	89H	PRL 62 3020	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	89J	ZPHY C45 175	K. Abe <i>et al.</i>	(VENUS Collab.)
ABE	89L	PL B232 425	K. Abe <i>et al.</i>	(VENUS Collab.)
ADACHI	89B	PL B228 553	I. Adachi <i>et al.</i>	(TOPAZ Collab.)
ALBAJAR	89	ZPHY C44 15	C. Albajar <i>et al.</i>	(UA1 Collab.)
BARGER	89	PL B220 464	V. Barger <i>et al.</i>	(WISC, KEK)
BEHREND	89B	PL B222 163	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
BRAUNSCH...	89C	ZPHY C43 549	W. Braunschweig <i>et al.</i>	(TASSO Collab.)
DORENBOS...	89	ZPHY C41 567	J. Dorenbosch <i>et al.</i>	(CHARM Collab.)
HAGIWARA	89	PL B219 369	K. Hagiwara, M. Sakuda, N. Terunuma	(KEK, DURH+)
KIM	89	PL B223 476	S.K. Kim <i>et al.</i>	(AMY Collab.)
ABE	88B	PL B213 400	K. Abe <i>et al.</i>	(VENUS Collab.)
BARINGER	88	PL B206 551	P. Baringer <i>et al.</i>	(HRS Collab.)
BRAUNSCH...	88	ZPHY C37 171	W. Braunschweig <i>et al.</i>	(TASSO Collab.)
BRAUNSCH...	88D	ZPHY C40 163	W. Braunschweig <i>et al.</i>	(TASSO Collab.)
ANSARI	87D	PL B195 613	R. Ansari <i>et al.</i>	(UA2 Collab.)
BARTEL	87B	ZPHY C36 15	W. Bartel <i>et al.</i>	(JADE Collab.)
BEHREND	87C	PL B191 209	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
FERNANDEZ	87B	PR D35 10	E. Fernandez <i>et al.</i>	(MAC Collab.)
ARNISON	86C	PL B172 461	G.T.J. Arnison <i>et al.</i>	(UA1 Collab.)
ARNISON	86D	PL B177 244	G.T.J. Arnison <i>et al.</i>	(UA1 Collab.)
BARTEL	86	ZPHY C31 359	W. Bartel <i>et al.</i>	(JADE Collab.)
BARTEL	86C	ZPHY C30 371	W. Bartel <i>et al.</i>	(JADE Collab.)
BEHREND	86	PL 168B 420	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
BEHREND	86C	PL B181 178	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
DERRICK	86	PL 166B 463	M. Derrick <i>et al.</i>	(HRS Collab.)
Also	86B	PR D34 3286	M. Derrick <i>et al.</i>	(HRS Collab.)
DERRICK	86B	PR D34 3286	M. Derrick <i>et al.</i>	(HRS Collab.)
GRIFOLS	86	PL 168B 264	J.A. Grifols, S. Peris	(BARC)
JODIDIO	86	PR D34 1967	A. Jodidio <i>et al.</i>	(LBL, NWES, TRIU)
Also	88	PR D37 237 erratum	A. Jodidio <i>et al.</i>	(LBL, NWES, TRIU)
APPEL	85	PL 160B 349	J.A. Appel <i>et al.</i>	(UA2 Collab.)
BARTEL	85K	PL 160B 337	W. Bartel <i>et al.</i>	(JADE Collab.)
BERGER	85	ZPHY C28 1	C. Berger <i>et al.</i>	(PLUTO Collab.)
BERGER	85B	ZPHY C27 341	C. Berger <i>et al.</i>	(PLUTO Collab.)
BAGNAIA	84C	PL 138B 430	P. Bagnaia <i>et al.</i>	(UA2 Collab.)
BARTEL	84D	PL 146B 437	W. Bartel <i>et al.</i>	(JADE Collab.)
BARTEL	84E	PL 146B 121	W. Bartel <i>et al.</i>	(JADE Collab.)
EICHTEN	84	RMP 56 579	E. Eichten <i>et al.</i>	(FNAL, LBL, OSU)
ALTHOFF	83C	PL 126B 493	M. Althoff <i>et al.</i>	(TASSO Collab.)
RENARD	82	PL 116B 264	F.M. Renard	(CERN)