### b-baryon ADMIXTURE ( $\Lambda_b$ , $\Xi_b$ , $\Sigma_b$ , $\Omega_b$ )

#### b-baryon ADMIXTURE MEAN LIFE

Each measurement of the *b*-baryon mean life is an average over an admixture of various *b* baryons which decay weakly. Different techniques emphasize different admixtures of produced particles, which could result in a different *b*-baryon mean life. More *b*-baryon flavor specific channels are not included in the measurement.

"OUR EVALUATION" is an average of the data listed below performed by the LEP B Lifetimes Working Group as described in our review "Production and Decay of b-flavored Hadrons" in the  $B^{\pm}$  Section of these Listings. The averaging procedure takes into account correlations between the measurements and asymmetric lifetime errors.

$VALUE (10^{-12})$	s) <u>I</u>	EVTS	DOCUMENT ID		TECN	COMMENT
$1.208 \pm 0.051$	OUR EVAL	UATION				
$1.16 \pm 0.20$	$\pm 0.08$		<sup>1</sup> ABREU	99W	DLPH	$e^+e^-  ightarrow Z$
$1.19 \pm 0.14$	$\pm 0.07$		<sup>2</sup> ABREU	99W	DLPH	$e^+e^- \rightarrow Z$
$1.20\ \pm0.08$	$\pm 0.06$		<sup>3</sup> BARATE	<b>98</b> D	ALEP	$e^+e^- \rightarrow Z$
$1.10 \begin{array}{l} +0.19 \\ -0.17 \end{array}$	$\pm 0.09$		ABREU	<b>96</b> D	DLPH	Excess $\Lambda\mu^-$ impact parameters
1.16 ±0.11	$\pm 0.06$		AKERS	96	OPAL	Excess $\Lambda \ell^-$ , decay lengths and impact parameters
• • • We do	not use the	following	data for averages	, fits,	limits,	•
$1.14 \pm 0.08$	$\pm 0.04$		<sup>4</sup> ABREU	99W	DLPH	$e^+e^- \rightarrow Z$
$1.46 \begin{array}{l} +0.22 \\ -0.21 \end{array}$	$^{+0.07}_{-0.09}$		ABREU	<b>96</b> D	DLPH	Repl. by ABREU 99W
$1.27 \begin{array}{l} +0.35 \\ -0.29 \end{array}$	$\pm 0.09$		ABREU	<b>95</b> S	DLPH	Repl. by ABREU 99W
$1.05 \begin{array}{l} +0.12 \\ -0.11 \end{array}$	$\pm 0.09$	290	BUSKULIC	95L	ALEP	Repl. by BARATE 98D
$1.04 \   ^{+  0.48}_{-  0.38}$	$\pm 0.10$	11	<sup>5</sup> ABREU	93F	DLPH	Excess $\Lambda\mu^-$ , decay lengths
$1.05 \begin{array}{l} +0.23 \\ -0.20 \end{array}$	$\pm 0.08$	157	<sup>6</sup> AKERS	93	OPAL	Excess $\Lambda \ell^-$ , decay lengths
$1.12 \begin{array}{c} +0.32 \\ -0.29 \end{array}$	$\pm 0.16$	101	<sup>7</sup> BUSKULIC	921	ALEP	Excess $\Lambda \ell^-$ , impact parameters
4						

<sup>&</sup>lt;sup>1</sup> Measured using  $\Lambda \ell^-$  decay length.

<sup>&</sup>lt;sup>2</sup> Measured using  $p\ell^-$  decay length.

<sup>&</sup>lt;sup>3</sup> Measured using the excess of  $\Lambda \ell^-$ , lepton impact parameter.

<sup>&</sup>lt;sup>4</sup> This ABREU 99W result is the combined result of the  $\Lambda\ell^-$ ,  $p\ell^-$ , and excess  $\Lambda\mu^-$  impact parameter measurements.

<sup>&</sup>lt;sup>5</sup> ABREU 93F superseded by ABREU 96D.

<sup>&</sup>lt;sup>6</sup> AKERS 93 superseded by AKERS 96.

<sup>&</sup>lt;sup>7</sup> BUSKULIC 921 superseded by BUSKULIC 95L.

#### b-baryon ADMIXTURE $(\Lambda_b, \Xi_b, \Sigma_b, \Omega_b)$

These branching fractions are actually an average over weakly decaying b-baryons weighted by their production rates in Z decay (or high-energy  $p\overline{p}$ ), branching ratios, and detection efficiencies. They scale with the LEP b-baryon production fraction B( $b \rightarrow b$ -baryon) and are evaluated for our value B( $b \rightarrow b$ -baryon) = (11.8  $\pm$  2.0)%.

The branching fractions B(b-baryon  $\to \Lambda \ell^- \overline{\nu}_\ell$  anything) and B( $\Lambda_b^0 \to \Lambda_c^+ \ell^- \overline{\nu}_\ell$  anything) are not pure measurements because the underlying measured products of these with B( $b \to b$ -baryon) were used to determine B( $b \to b$ -baryon), as described in the note "Production and Decay of b-Flavored Hadrons."

	Mode	Fraction $(\Gamma_i/\Gamma)$
$\overline{\Gamma_1}$	$p\mu^-\overline{ u}$ anything	( 4.2 + 1.7 ) %
$\Gamma_2$	$ ho  \ell  \overline{ u}_\ell$ anything	( 4.0± 1.0) %
$\Gamma_3$	<i>p</i> anything	(50 $\pm$ 17 )%
$\Gamma_4$	$arLambda \ell^- \overline{ u}_\ell$ anything	( 2.7± 0.5) %
$\Gamma_5$	$arLambda\ell^+ u_\ell$ anything	
•	$\Lambda$ anything	
$\Gamma_7$	$\Lambda_c^+\ell^-\overline{ u}_\ell$ anything	
Γ <sub>8</sub>	$\Lambda/\overline{\Lambda}$ anything	$(28 \pm 6)\%$
$\Gamma_9$	$ar{arXi}^-\ell^-\overline{ u}_\ell$ anything	$(4.7\pm 1.3) \times 10^{-3}$

#### b-baryon ADMIXTURE ( $\Lambda_b$ , $\Xi_b$ , $\Sigma_b$ , $\Omega_b$ ) BRANCHING RATIOS

#### 

# $\Gamma(\rho\ell\overline{\nu}_{\ell} \text{ anything})/\Gamma_{\text{total}}$ VALUE $0.040\pm0.007\pm0.007$ PACHIC OF TOTAL OF THE OF TH

#### $\Gamma(p\ell\overline{\nu}_{\ell}\text{ anything})/\Gamma(p\text{anything})$

 $0.080 \pm 0.012 \pm 0.014$ 

DOCUMENT ID TECN COMMENT

BARATE 98V ALEP  $e^+e^- \rightarrow Z$ 

 $\Gamma_2/\Gamma_3$ 

HTTP://PDG.LBL.GOV Page 2 Created: 6/12/2002 17:27

<sup>&</sup>lt;sup>8</sup> ABREU 95S reports [B(*b*-baryon  $\rightarrow p \mu^- \overline{\nu}$  anything)  $\times$  B( $\overline{b} \rightarrow b$ -baryon)] = 0.0049  $\pm$  0.0011 $^+$ 0.0015. We divide by our best value B( $\overline{b} \rightarrow b$ -baryon) = (11.8  $\pm$  2.0)  $\times$  10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>&</sup>lt;sup>9</sup> BARATE 98V reports [B(b-baryon  $\to p\ell\overline{\nu}_\ell$  anything)  $\times$  B( $\overline{b}\to b$ -baryon)] = (4.72  $\pm$  0.66  $\pm$  0.44)  $\times$  10<sup>-3</sup>. We divide by our best value B( $\overline{b}\to b$ -baryon) = (11.8  $\pm$  2.0)  $\times$  10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

#### $\Gamma(\Lambda \ell^- \overline{\nu}_\ell \text{ anything}) / \Gamma_{\text{total}}$

 $\Gamma_4/\Gamma$ 

The values and averages in this section serve only to show what values result if one assumes our  $B(b \to b\text{-baryon})$ . They cannot be thought of as measurements since the underlying product branching fractions were also used to determinine  $B(b \to b\text{-baryon})$  as described in the note on "Production and Decay of b-Flavored Hadrons."

VALUE	<b>EVTS</b>	DOCUMENT ID		TECN	COMMENT	
0.027±0.005 OUR AVERAGE						
$0.028\!\pm\!0.004\!\pm\!0.005$		<sup>10</sup> BARATE	<b>98</b> D	ALEP	$e^+e^- \rightarrow Z$	
$0.025 \pm 0.003 \pm 0.004$		<sup>11</sup> AKERS	96	OPAL	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$	
$0.025\!\pm\!0.006\!\pm\!0.004$	262	<sup>12</sup> ABREU	<b>95</b> S	DLPH	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$	
$0.052\!\pm\!0.010\!\pm\!0.009$	290	<sup>13</sup> BUSKULIC	95L	ALEP	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$	
• • • We do not use the	e followir	ng data for averages	, fits,	, limits,	etc. • • •	
seen	157	<sup>14</sup> AKERS	93	OPAL	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$	
$0.059\!\pm\!0.017\!\pm\!0.010$	101	<sup>15</sup> BUSKULIC	921	ALEP	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$	
<sup>10</sup> BARATE 98D reports [B( $b$ -baryon $\rightarrow \Lambda \ell^- \overline{\nu}_{\ell}$ anything) $\times$ B( $\overline{b} \rightarrow b$ -baryon)] =						
$0.00326 \pm 0.00016 \pm 0.00039$ . We divide by our best value B( $\overline{b} \rightarrow b$ -baryon) =						
$(11.8 \pm 2.0) \times 10^{-2}$	<sup>2</sup> . Our 1	irst error is their e	xperii	ment's e	error and our second error	
is the systematic error from using our best value. Measured using the excess of $\Lambda\ell^-$ ,						
lepton impact param						
<sup>11</sup> AKERS 96 reports [B	(b-baryo	on $ ightarrow$ $arLambda \ell^- \overline{ u}_\ell$ anyth	ing)	$\times$ B( $\overline{b}$ –	$\rightarrow b$ -baryon)] = 0.00291 $\pm$	

<sup>11</sup> AKERS 96 reports [B(*b*-baryon  $\to \Lambda \ell^- \overline{\nu}_\ell$  anything)  $\times$  B( $\overline{b} \to b$ -baryon)] = 0.00291  $\pm$  0.00023  $\pm$  0.00025. We divide by our best value B( $\overline{b} \to b$ -baryon) = (11.8  $\pm$  2.0)  $\times$  10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>12</sup> ABREU 95s reports [B(b-baryon  $\to \Lambda \ell^- \overline{\nu}_\ell$  anything)  $\times$  B( $\overline{b} \to b$ -baryon)] = 0.0030  $\pm$  0.0006  $\pm$  0.0004. We divide by our best value B( $\overline{b} \to b$ -baryon) = (11.8  $\pm$  2.0)  $\times$  10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>13</sup> BUSKULIC 95L reports [B(b-baryon  $\rightarrow \Lambda \ell^- \overline{\nu}_\ell$  anything)  $\times$  B( $\overline{b} \rightarrow b$ -baryon)] = 0.0061  $\pm$  0.0006  $\pm$  0.0010. We divide by our best value B( $\overline{b} \rightarrow b$ -baryon) = (11.8  $\pm$  2.0)  $\times$  10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>14</sup> AKERS 93 superseded by AKERS 96.

<sup>15</sup> BUSKULIC 92I reports [B(b-baryon  $\rightarrow \Lambda \ell^- \overline{\nu}_\ell$  anything)  $\times$  B( $\overline{b} \rightarrow b$ -baryon)] = 0.0070  $\pm$  0.0010  $\pm$  0.0018. We divide by our best value B( $\overline{b} \rightarrow b$ -baryon) = (11.8  $\pm$  2.0)  $\times$  10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value. Superseded by BUSKULIC 95L.

#### $\Gamma(\Lambda \ell^+ \nu_\ell \text{ anything})/\Gamma(\Lambda \text{ anything})$

 $\Gamma_5/\Gamma_6$ 

 $\Gamma_8/\Gamma$ 

VALUE	DOCUMENT ID		TECN	COMMENT		
$0.080 \pm 0.012 \pm 0.008$	ABBIENDI	99L	OPAL	$e^+e^-  ightarrow Z$		
• • • We do not use the following data for averages, fits, limits, etc. • •						
$0.070 \pm 0.012 \pm 0.007$	ACKERSTAFF	97N	OPAL	Repl. by ABBIENDI 99L		

## Γ(Λ/Λanything)/Γ<sub>total</sub>

VALUE	<u>DOCUMENT ID</u>	<u>I ECN</u>	COMMENT
0.28±0.06 OUR AVERAGE			
$0.30\!\pm\!0.04\!\pm\!0.05$	<sup>16</sup> ABBIENDI	99L OPAL	$e^+e^-  ightarrow Z$
$0.19^{+0.11}_{-0.07}\pm0.03$	<sup>17</sup> ABREU	95c DLPH	$e^+e^- ightarrow~Z$

• • • We do not use the following data for averages, fits, limits, etc. • • •

HTTP://PDG.LBL.GOV

Page 3

Created: 6/12/2002 17:27

- <sup>16</sup> ABBIENDI 99L reports [B(*b*-baryon  $\rightarrow \Lambda/\overline{\Lambda}$  anything)  $\times$  B( $\overline{b} \rightarrow b$ -baryon)] = 0.035  $\pm$  0.0032  $\pm$  0.0035. We divide by our best value B( $\overline{b} \rightarrow b$ -baryon) = (11.8  $\pm$  2.0)  $\times$  10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- $^{17}$  ABREU 95C reports  $0.28^{+0.17}_{-0.12}$  for B( $\overline{b} \rightarrow b$ -baryon) =  $0.08 \pm 0.02$ . We rescale to our best value B( $\overline{b} \rightarrow b$ -baryon) =  $(11.8 \pm 2.0) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.
- <sup>18</sup> ACKERSTAFF 97N reports [B(b-baryon  $\to \Lambda/\overline{\Lambda}$  anything)  $\times$  B( $\overline{b} \to b$ -baryon)] = 0.0393  $\pm$  0.0046  $\pm$  0.0037. We divide by our best value B( $\overline{b} \to b$ -baryon) = (11.8  $\pm$  2.0)  $\times$  10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

#### $\Gamma(\Xi^-\ell^-\overline{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}$

 $\Gamma_0/\Gamma$ 

<u>VALUE</u>	DOCUMENT ID		TECN	COMMENT
0.0047±0.0013 OUR AVERAGE				
$0.0046\!\pm\!0.0012\!\pm\!0.0008$	<sup>19</sup> BUSKULIC	96T	ALEP	Excess $\underline{\Xi}^-\ell^-$ over $\underline{\Xi}^-\ell^+$
$0.0050 \pm 0.0020 \pm 0.0008$	<sup>20</sup> ABREU	95∨	DLPH	$\Xi$ Excess $\Xi^-\ell^-$ over $\Xi^-\ell^+$

- <sup>19</sup> BUSKULIC 96T reports [B(b-baryon  $\to \Xi^-\ell^-\overline{\nu}_\ell$  anything)  $\times$  B( $\overline{b} \to b$ -baryon)] = 0.00054  $\pm$  0.00011  $\pm$  0.00008. We divide by our best value B( $\overline{b} \to b$ -baryon) = (11.8  $\pm$  2.0)  $\times$  10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- <sup>20</sup> ABREU 95V reports [B(b-baryon  $\to \bar{\Xi}^-\ell^-\bar{\nu}_\ell$  anything)  $\times$  B( $\bar{b} \to b$ -baryon)] = 0.00059  $\pm$  0.00021  $\pm$  0.0001. We divide by our best value B( $\bar{b} \to b$ -baryon) = (11.8  $\pm$  2.0)  $\times$  10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

#### b-baryon ADMIXTURE $(\Lambda_b, \Xi_b, \Sigma_b, \Omega_b)$ REFERENCES

ABBIENDI	99L	EPJ C9 1	G. Abbiendi et al.	(OPAL Collab.)
ABREU	99W	EPJ C10 185	P. Abreu <i>et al.</i>	(DELPHI Collab.)
BARATE	98D	EPJ C2 197	R. Barate et al.	(ALEPH Collab.)
BARATE	98V	EPJ C5 205	R. Barate et al.	(ALEPH Collab.)
ACKERSTAFF	97N	ZPHY C74 423	K. Ackerstaff et al.	(OPAL Collab.)
ABREU	96D	ZPHY C71 199	P. Abreu et al.	(DELPHI Collab.)
AKERS	96	ZPHY C69 195	R. Akers et al.	(OPAL Collab.)
BUSKULIC	96T	PL B384 449	D. Buskulic et al.	(ALEPH Collab.)
ABREU	95C	PL B347 447	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	95S	ZPHY C68 375	P. Abreu et al.	(DELPHI Collab.)
ABREU	95V	ZPHY C68 541	P. Abreu et al.	(DELPHI Collab.)
BUSKULIC	95L	PL B357 685	D. Buskulic et al.	(ALEPH Collab.)
ABREU	93F	PL B311 379	P. Abreu <i>et al.</i>	(ĎELPHI Collab.)
AKERS	93	PL B316 435	R. Akers et al.	(OPAL Collab.)
BUSKULIC	92I	PL B297 449	D. Buskulic et al.	(ALEPH Collab.)
				•

Created: 6/12/2002 17:27