

$\Lambda(1670)$ S_{01} $I(J^P) = 0(\frac{1}{2}^-)$ Status: ***

The measurements of the mass, width, and elasticity published before 1974 are now obsolete and have been omitted. They were last listed in our 1982 edition Physics Letters **111B** (1982).

 $\Lambda(1670)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1660 to 1680 (≈ 1670) OUR ESTIMATE			
1673 ± 2	MANLEY	02	DPWA $\bar{K}N$ multichannel
1670.8 ± 1.7	KOISO	85	DPWA $K^- p \rightarrow \Sigma \pi$
1667 ± 5	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$
1671 ± 3	ALSTON-...	78	DPWA $\bar{K}N \rightarrow \bar{K}N$
1670 ± 5	GOPAL	77	DPWA $\bar{K}N$ multichannel
1675 ± 2	HEPP	76B	DPWA $K^- N \rightarrow \Sigma \pi$
1679 ± 1	KANE	74	DPWA $K^- p \rightarrow \Sigma \pi$
1665 ± 5	PREVOST	74	DPWA $K^- N \rightarrow \Sigma(1385)\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1668.9 ± 2.0	ABAEV	96	DPWA $K^- p \rightarrow \Lambda\eta$
1664	¹ MARTIN	77	DPWA $\bar{K}N$ multichannel

 $\Lambda(1670)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
25 to 50 (≈ 35) OUR ESTIMATE			
23 ± 6	MANLEY	02	DPWA $\bar{K}N$ multichannel
34.1 ± 3.7	KOISO	85	DPWA $K^- p \rightarrow \Sigma \pi$
29 ± 5	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$
29 ± 5	ALSTON-...	78	DPWA $\bar{K}N \rightarrow \bar{K}N$
45 ± 10	GOPAL	77	DPWA $\bar{K}N$ multichannel
46 ± 5	HEPP	76B	DPWA $K^- N \rightarrow \Sigma \pi$
40 ± 3	KANE	74	DPWA $K^- p \rightarrow \Sigma \pi$
19 ± 5	PREVOST	74	DPWA $K^- N \rightarrow \Sigma(1385)\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
21.1 ± 3.6	ABAEV	96	DPWA $K^- p \rightarrow \Lambda\eta$
12	¹ MARTIN	77	DPWA $\bar{K}N$ multichannel

 $\Lambda(1670)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\bar{K}$	20–30 %
Γ_2 $\Sigma\pi$	25–55 %
Γ_3 $\Lambda\eta$	10–25 %
Γ_4 $\Sigma(1385)\pi$	

The above branching fractions are our estimates, not fits or averages.

$\Lambda(1670)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.20 to 0.30 OUR ESTIMATE				
0.37 \pm 0.07	MANLEY 02	DPWA	$\bar{K}N$ multichannel	
0.18 \pm 0.03	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$	
0.17 \pm 0.03	ALSTON-... 78	DPWA	$\bar{K}N \rightarrow \bar{K}N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.20 \pm 0.03	GOPAL 77	DPWA	See GOPAL 80	
0.15	¹ MARTIN 77	DPWA	$\bar{K}N$ multichannel	

$\Gamma(\Lambda\eta)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.30 \pm 0.08	ABAEV 96	DPWA	$K^- p \rightarrow \Lambda\eta$	

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(1670) \rightarrow \Sigma\pi$

VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.38 \pm 0.03	MANLEY 02	DPWA	$\bar{K}N$ multichannel	
-0.26 \pm 0.02	KOISO 85	DPWA	$K^- p \rightarrow \Sigma\pi$	
-0.31 \pm 0.03	GOPAL 77	DPWA	$\bar{K}N$ multichannel	
-0.29 \pm 0.03	HEPP 76B	DPWA	$K^- N \rightarrow \Sigma\pi$	
-0.23 \pm 0.03	LONDON 75	HLBC	$K^- p \rightarrow \Sigma^0\pi^0$	
-0.27 \pm 0.02	KANE 74	DPWA	$K^- p \rightarrow \Sigma\pi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.13	¹ MARTIN 77	DPWA	$\bar{K}N$ multichannel	

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(1670) \rightarrow \Lambda\eta$

VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_3)^{1/2}/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
+0.24 \pm 0.04	MANLEY 02	DPWA	$\bar{K}N$ multichannel	
+0.20 \pm 0.05	BAXTER 73	DPWA	$K^- p \rightarrow$ neutrals	
0.24	KIM 71	DPWA	K-matrix analysis	
0.26	ARMENTEROS69C	HBC		
0.20 or 0.23	BERLEY 65	HBC		

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(1670) \rightarrow \Sigma(1385)\pi$

VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_4)^{1/2}/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.17 \pm 0.06	MANLEY 02	DPWA	$\bar{K}N$ multichannel	
-0.18 \pm 0.05	PREVOST 74	DPWA	$K^- N \rightarrow \Sigma(1385)\pi$	

$\Lambda(1670)$ FOOTNOTES

¹MARTIN 77 obtains identical resonance parameters from a T-matrix pole and from a Breit-Wigner fit.

$\Lambda(1670)$ REFERENCES

MANLEY	02	PRL 88 012002	D.M. Manley <i>et al.</i>	(BNL Crystal Ball Collab.)
ABAEV	96	PR C53 385	V.V. Abaev, B.M.K. Nefkens	(UCLA)
KOISO	85	NP A433 619	H. Koiso <i>et al.</i>	(TOKY, MASA)
PDG	82	PL 111B	M. Roos <i>et al.</i>	(HELS, CIT, CERN)
GOPAL	80	Toronto Conf. 159	G.P. Gopal	(RHEL) IJP
ALSTON-...	78	PR D18 182	M. Alston-Garnjost <i>et al.</i>	(LBL, MTTHO+) IJP
Also	77	PRL 38 1007	M. Alston-Garnjost <i>et al.</i>	(LBL, MTTHO+) IJP
GOPAL	77	NP B119 362	G.P. Gopal <i>et al.</i>	(LOIC, RHEL) IJP
MARTIN	77	NP B127 349	B.R. Martin, M.K. Pidcock, R.G. Moorhouse	(LOUC+) IJP
Also	77B	NP B126 266	B.R. Martin, M.K. Pidcock	(LOUC)
Also	77C	NP B126 285	B.R. Martin, M.K. Pidcock	(LOUC) IJP
HEPP	76B	PL 65B 487	V. Hepp <i>et al.</i>	(CERN, HEIDH, MPIM) IJP
LONDON	75	NP B85 289	G.W. London <i>et al.</i>	(BNL, CERN, EPOL+)
KANE	74	LBL-2452	D.F. Kane	(LBL) IJP
PREVOST	74	NP B69 246	J. Prevost <i>et al.</i>	(SACL, CERN, HEID)
BAXTER	73	NP B67 125	D.F. Baxter <i>et al.</i>	(OXF) IJP
KIM	71	PRL 27 356	J.K. Kim	(HARV) IJP
Also	70	Duke Conf. 161	J.K. Kim	(HARV) IJP
Hyperon Resonances, 1970				
ARMENTEROS	69C	Lund Paper 229	R. Armenteros <i>et al.</i>	(CERN, HEID, SACL) IJP
Values are quoted in LEVI-SETTI 69.				
BERLEY	65	PRL 15 641	D. Berley <i>et al.</i>	(BNL) IJP