

$\Xi(1530) P_{13}$

$$I(J^P) = \frac{1}{2}(3/2^+) \text{ Status: } ****$$


This is the only Ξ resonance whose properties are all reasonably well known. Spin-parity $3/2^+$ is favored by the data.

We use only those determinations of the mass and width that are accompanied by some discussion of systematics and resolution.

$\Xi(1530)$ MASSES

$\Xi(1530)^0$ MASS

<i>VALUE (MeV)</i>	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
1531.80 ± 0.32 OUR FIT				Error includes scale factor of 1.3.
1531.78 ± 0.34 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.
1532.2 ± 0.7		DEBELLEFON 75B	HBC	$K^- p \rightarrow \Xi^- \bar{K} \pi$
1533 ± 1		ROSS 73B	HBC	$K^- p \rightarrow \Xi^- \bar{K} \pi(\pi)$
1531.4 ± 0.8	59	BADIER 72	HBC	$K^- p$ 3.95 GeV/c
1532.0 ± 0.4	1262	BALTAY 72	HBC	$K^- p$ 1.75 GeV/c
1531.3 ± 0.6	324	BORENSTEIN 72	HBC	$K^- p$ 2.2 GeV/c
1532.3 ± 0.7	286	KIRSCH 72	HBC	$K^- p$ 2.87 GeV/c
1528.7 ± 1.1	76	LONDON 66	HBC	$K^- p$ 2.24 GeV/c
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1532.1 ± 0.4	1244	ASTON 85B	LASS	$K^- p$ 11 GeV/c
1532.1 ± 0.6	2700	¹ BAUBILLIER 81B	HBC	$K^- p$ 8.25 GeV/c
1530 ± 1	450	BIAGI 81	SPEC	SPS hyperon beam
1527 ± 6	80	SIXEL 79	HBC	$K^- p$ 10 GeV/c
1535 ± 4	100	SIXEL 79	HBC	$K^- p$ 16 GeV/c
1533.6 ± 1.4	97	BERTHON 74	HBC	Quasi-2-body σ



$\Xi(1530)^0$ mass (MeV)

$\Xi(1530)^-$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1535.0±0.6 OUR FIT				
1535.2±0.8 OUR AVERAGE				
1534.5±1.2		DEBELLEFON 75B	HBC	$K^- p \rightarrow \Xi^- \bar{K} \pi$
1535.3±2.0		ROSS 73B	HBC	$K^- p \rightarrow \Xi^- \bar{K} \pi (\pi)$
1536.2±1.6	185	KIRSCH 72	HBC	$K^- p$ 2.87 GeV/c
1535.7±3.2	38	LONDON 66	HBC	$K^- p$ 2.24 GeV/c
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1540 ±3	48	BERTHON 74	HBC	Quasi-2-body σ
1534.7±1.1	334	BALTAY 72	HBC	$K^- p$ 1.75 GeV/c

$m_{\Xi(1530)^-} - m_{\Xi(1530)}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
3.2±0.6 OUR FIT			
2.9±0.9 OUR AVERAGE			
2.7±1.0	BALTAY 72	HBC	$K^- p$ 1.75 GeV/c
2.0±3.2	MERRILL 66	HBC	$K^- p$ 1.7–2.7 GeV/c
5.7±3.0	PJERROU 65B	HBC	$K^- p$ 1.8–1.95 GeV/c
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
3.9±1.8	² KIRSCH 72	HBC	$K^- p$ 2.87 GeV/c
7 ±4	² LONDON 66	HBC	$K^- p$ 2.24 GeV/c

$\Xi(1530)$ WIDTHS **$\Xi(1530)^0$ WIDTH**

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
9.1±0.5 OUR AVERAGE				
9.5±1.2		DEBELLEFON	75B HBC	$K^- p \rightarrow \Xi^- \bar{K} \pi$
9.1±2.4		ROSS	73B HBC	$K^- p \rightarrow \Xi^- \bar{K} \pi (\pi)$
11 ±2		BADIER	72 HBC	$K^- p$ 3.95 GeV/c
9.0±0.7		BALTAY	72 HBC	$K^- p$ 1.75 GeV/c
8.4±1.4		BORENSTEIN	72 HBC	$\Xi^- \pi^+$
11.0±1.8		KIRSCH	72 HBC	$\Xi^- \pi^+$
7 ±7		BERGE	66 HBC	$K^- p$ 1.5–1.7 GeV/c
8.5±3.5		LONDON	66 HBC	$K^- p$ 2.24 GeV/c
7 ±2		SCHLEIN	63B HBC	$K^- p$ 1.8, 1.95 GeV/c
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
12.8±1.0	2700	¹ BAUBILLIER	81B HBC	$K^- p$ 8.25 GeV/c
19 ±6	80	³ SIXEL	79 HBC	$K^- p$ 10 GeV/c
14 ±5	100	³ SIXEL	79 HBC	$K^- p$ 16 GeV/c

 $\Xi(1530)^-$ WIDTH

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
9.9^{+1.7}_{-1.9} OUR AVERAGE			
9.6±2.8	DEBELLEFON	75B HBC	$K^- p \rightarrow \Xi^- \bar{K} \pi$
8.3±3.6	ROSS	73B HBC	$K^- p \rightarrow \Xi^- \bar{K} \pi (\pi)$
7.8 ^{+3.5} _{-7.8}	BALTAY	72 HBC	$K^- p$ 1.75 GeV/c
16.2±4.6	KIRSCH	72 HBC	$\Xi^- \pi^0, \Xi^0 \pi^-$

 $\Xi(1530)$ POLE POSITIONS **$\Xi(1530)^0$ REAL PART**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
1531.6±0.4	LICHTENBERG74	Using HABIBI 73

 $\Xi(1530)^0$ IMAGINARY PART

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
4.45±0.35	LICHTENBERG74	Using HABIBI 73

 $\Xi(1530)^-$ REAL PART

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
1534.4±1.1	LICHTENBERG74	Using HABIBI 73

 $\Xi(1530)^-$ IMAGINARY PART

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
3.9 ^{+1.75} _{-3.9}	LICHTENBERG74	Using HABIBI 73

$\Xi(1530)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 \quad \Xi \pi$	100 %	
$\Gamma_2 \quad \Xi \gamma$	<4 %	90%

 $\Xi(1530)$ BRANCHING RATIOS

$\Gamma(\Xi \gamma)/\Gamma_{\text{total}}$	Γ_2/Γ			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.04	90	KALBFLEISCH 75	HBC	$K^- p$ 2.18 GeV/c

 $\Xi(1530)$ FOOTNOTES

- ¹ BAUBILLIER 81B is a fit to the inclusive spectrum. The resolution (5 MeV) is not unfolded.
² Redundant with data in the mass Listings.
³ SIXEL 79 doesn't unfold the experimental resolution of 15 MeV.

 $\Xi(1530)$ REFERENCES

ASTON	85B	PR D32 2270	D. Aston <i>et al.</i>	(SLAC, CARL, CNRC, CINC)
BAUBILLIER	81B	NP B192 1	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)
BIAGI	81	ZPHY C9 305	S.F. Biagi <i>et al.</i>	(BRIS, CAVE, GEVA+)
SIXEL	79	NP B159 125	P. Sixel <i>et al.</i>	(AACH3, BERL, CERN, LOIC+)
DEBELLEFON	75B	NC 28A 289	A. de Bellefon <i>et al.</i>	(CDEF, SACL)
KALBFLEISCH	75	PR D11 987	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)
BERTHON	74	NC 21A 146	A. Berthon <i>et al.</i>	(CDEF, RHEL, SACL+)
LICHTENBERG	74	PR D10 3865	D.B. Lichtenberg	(IND)
Also	74B	Private Comm.	D.B. Lichtenberg	(IND)
HABIBI	73	Thesis Nevis 199	M. Habibi	(COLU)
ROSS	73B	Purdue Conf. 355	R.T. Ross, J.L. Lloyd, D. Radojicic	(OXF)
BADIER	72	NP B37 429	J. Badier <i>et al.</i>	(EPOL)
BALTAY	72	PL 42B 129	C. Baltay <i>et al.</i>	(COLU, BING)
BORENSTEIN	72	PR D5 1559	S.R. Borenstein <i>et al.</i>	(BNL, MICH) I
KIRSCH	72	NP B40 349	L.E. Kirsch <i>et al.</i>	(BRAN, UMD, SYRA+) I
BERGE	66	PR 147 945	J.P. Berge <i>et al.</i>	(LRL) I
LONDON	66	PR 143 1034	G.W. London <i>et al.</i>	(BNL, SYRA) IJ
MERRILL	66	Thesis UCRL 16455	D.W. Merrill	(LRL) JP
PJERROU	65B	PRL 14 275	G.M. Pjerrou <i>et al.</i>	(UCLA)
SCHLEIN	63B	PRL 11 167	P.E. Schlein <i>et al.</i>	(UCLA) IJP

OTHER RELATED PAPERS

MAZZUCATO	81	NP B178 1	M. Mazzucato <i>et al.</i>	(AMST, CERN, NIJM+)
BRIEFEL	77	PR D16 2706	E. Briefel <i>et al.</i>	(BRAN, UMD, SYRA+)
BRIEFEL	75	PR D12 1859	E. Briefel <i>et al.</i>	(BRAN, UMD, SYRA+)
HUNGERBU...	74	PR D10 2051	V. Hungerbuhler <i>et al.</i>	(YALE, FNAL, BNL+)
BUTTON-...	66	PR 142 883	J. Button-Shafer <i>et al.</i>	(LRL) JP