$\Sigma(1560)$  Bumps

 $I(J^P) = 1(?^?)$  Status: \*\*

#### OMITTED FROM SUMMARY TABLE

This entry lists peaks reported in mass spectra around 1560 MeV without implying that they are necessarily related.

DIONISI 78B observes a 6 standard-deviation enhancement at 1553 MeV in the charged  $\Lambda/\Sigma\pi$  mass spectra from  $K^-p\to (\Lambda/\Sigma)\pi\,K\overline{K}$  at 4.2 GeV/c. In a CERN ISR experiment, LOCK-MAN 78 reports a narrow 6 standard-deviation enhancement at 1572 MeV in  $\Lambda\pi^\pm$  from the reaction  $pp\to\Lambda\pi^+\pi^-X$ . These enhancements are unlikely to be associated with the  $\Sigma(1580)$  (which has not been confirmed by several recent experiments – see the next entry in the Listings).

CARROLL 76 observes a bump at 1550 MeV (as well as one at 1580 MeV) in the isospin-1  $\overline{K}$  N total cross section, but uncertainties in cross section measurements outside the mass range of the experiment preclude estimating its significance.

See also MEADOWS 80 for a review of this state.

# $\Sigma(1560)$ MASS (PRODUCTION EXPERIMENTS)

EVTS	DOCUMENT ID	TECN	CHG	COMMENT
MATE				
121	DIONISI	78B HBC	$\pm$	$K^-p \rightarrow$
				$(Y\pi)K\overline{K}$
40	LOCKMAN	78 SPEC	$\pm$	$pp \rightarrow \Lambda \pi^+ \pi^- X$
	121	121 DIONISI	121 DIONISI 78B HBC	MATE 121 DIONISI 78B HBC $\pm$

# $\Sigma$ (1560) WIDTH (PRODUCTION EXPERIMENTS)

VALUE (MeV)	EVTS	DOCUMENT ID		TECN	CHG	COMMENT
$79 \pm 30$	121	DIONISI	<b>78</b> B	HBC	$\pm$	$K^-p \rightarrow$
15   6	40	1.00000000	70	CDEC		$(Y\pi)K\overline{K}$ $pp \to \Lambda\pi^+\pi^-X$
$15\pm$ 6	40	LOCKMAN	78	SPEC	土	$pp \rightarrow \Lambda \pi^+ \pi^- X$

# $\Sigma$ (1560) DECAY MODES (PRODUCTION EXPERIMENTS)

	Mode	Fraction $(\Gamma_i/\Gamma)$
Γ <sub>1</sub> Γ <sub>2</sub>	$\Lambda\pi$ $\Sigma\pi$	seen

Created: 5/30/2003 10:57

### $\Sigma$ (1560) BRANCHING RATIOS (PRODUCTION EXPERIMENTS)

 $\Gamma(\Sigma\pi)/[\Gamma(\Lambda\pi)+\Gamma(\Sigma\pi)]$  $\Gamma_2/(\Gamma_1+\Gamma_2)$ **COMMENT** DOCUMENT ID TECN CHG  $0.35 \!\pm\! 0.12$ **78B HBC** DIONISI  $K^-p \rightarrow$  $(Y\pi)K\overline{K}$  $\Gamma(\Lambda\pi)/\Gamma_{\text{total}}$  $\Gamma_1/\Gamma$ DOCUMENT ID **COMMENT** TECN CHG  $pp \rightarrow \Lambda \pi^+ \pi^- X$ **LOCKMAN** 78 SPEC seen

## $\Sigma$ (1560) FOOTNOTES (PRODUCTION EXPERIMENTS)

# $\Sigma$ (1560) REFERENCES (PRODUCTION EXPERIMENTS)

 MEADOWS
 80
 Toronto
 Conf.
 283

 DIONISI
 78B
 PL 78B
 154

 LOCKMAN
 78
 Saclay DPHPE
 78-01

 CARROLL
 76
 PRL 37
 806

B.T. Meadows C. Dionisi, R. Armenteros, J. Diaz W. Lockman *et al.* A.S. Carroll *et al.*  (CINC) (CERN, AMST+) I (UCLA, SACL) (BNL) I

Created: 5/30/2003 10:57

 $<sup>^{</sup>m 1}$  The width observed by LOCKMAN 78 is consistent with experimental resolution.