

$\rho_3(2250)$

$$I^G(J^{PC}) = 1^+(3^{--})$$

OMITTED FROM SUMMARY TABLE

Contains results mostly from formation experiments. For further production experiments see the Further States entry. See also $\rho(2150)$, $f_2(2150)$, $f_4(2300)$, $\rho_5(2350)$.

 $\rho_3(2250)$ MASS **$\bar{p}p \rightarrow \pi\pi$ or $K\bar{K}$**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
~ 2232	HASAN	94	RVUE	$\bar{p}p \rightarrow \pi\pi$
~ 2007	HASAN	94	RVUE	$\bar{p}p \rightarrow \pi\pi$
~ 2090	¹ OAKDEN	94	RVUE	0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~ 2250	² MARTIN	80B	RVUE	
~ 2300	² MARTIN	80C	RVUE	
~ 2140	³ CARTER	78B	CNTR 0	0.7–2.4 $\bar{p}p \rightarrow K^-K^+$
~ 2150	⁴ CARTER	77	CNTR 0	0.7–2.4 $\bar{p}p \rightarrow \pi\pi$

¹ See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

² $I(J^P) = 1(3^-)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.

³ $I = 0, 1$. $J^P = 3^-$ from Barrelet-zero analysis.

⁴ $I(J^P) = 1(3^-)$ from amplitude analysis.

S-CHANNEL $\bar{N}N$

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2260 ± 20	¹⁶ ANISOVICH	02	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
~ 2190	⁵ CUTTS	78B	CNTR	0.97–3 $\bar{p}p \rightarrow \bar{N}N$
2155 ± 15	^{5,6} COUPLAND	77	CNTR 0	0.7–2.4 $\bar{p}p \rightarrow \bar{p}p$
2193 ± 2	^{5,7} ALSPECTOR	73	CNTR	$\bar{p}p$ S channel
2190 ± 10	⁸ ABRAMS	70	CNTR	S channel $\bar{p}N$

⁵ Isospins 0 and 1 not separated.

⁶ From a fit to the total elastic cross section.

⁷ Referred to as T or T region by ALSPECTOR 73.

⁸ Seen as bump in $I = 1$ state. See also COOPER 68. PEASLEE 75 confirm $\bar{p}p$ results of ABRAMS 70, no narrow structure.

 $\pi^-p \rightarrow \eta\pi\pi$

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$2290 \pm 20 \pm 30$	AMELIN	00	VES	37 $\pi^-p \rightarrow \eta\pi^+\pi^-n$

$\rho_3(2250)$ WIDTH **$\bar{p}p \rightarrow \pi\pi$ or $K\bar{K}$**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
~ 220	HASAN	94	RVUE	$\bar{p}p \rightarrow \pi\pi$
~ 287	HASAN	94	RVUE	$\bar{p}p \rightarrow \pi\pi$
~ 60	⁹ OAKDEN	94	RVUE	0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~ 250	¹⁰ MARTIN	80B	RVUE	
~ 200	¹⁰ MARTIN	80C	RVUE	
~ 150	¹¹ CARTER	78B	CNTR 0	0.7–2.4 $\bar{p}p \rightarrow K^-K^+$
~ 200	¹² CARTER	77	CNTR 0	0.7–2.4 $\bar{p}p \rightarrow \pi\pi$

⁹See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

¹⁰ $I(J^P) = 1(3^-)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.

¹¹ $I = 0, 1$. $J^P = 3^-$ from Barrelet-zero analysis.

¹² $I(J^P) = 1(3^-)$ from amplitude analysis.

S-CHANNEL $\bar{N}N$

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
160±25	¹⁷ ANISOVICH	02	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
135±75	^{13,14} COUPLAND	77	CNTR 0	0.7–2.4 $\bar{p}p \rightarrow \bar{p}p$
98±8	¹⁴ ALSPECTOR	73	CNTR	$\bar{p}p$ S channel
~ 85	¹⁵ ABRAMS	70	CNTR	S channel $\bar{p}N$

¹³From a fit to the total elastic cross section.

¹⁴Isospins 0 and 1 not separated.

¹⁵Seen as bump in $I = 1$ state. See also COOPER 68. PEASLEE 75 confirm $\bar{p}p$ results of ABRAMS 70, no narrow structure.

 $\pi^-p \rightarrow \eta\pi\pi$

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
230±50±80	AMELIN	00	VES 37	$\pi^-p \rightarrow \eta\pi^+\pi^-n$

¹⁶From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

¹⁷From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

$\rho_3(2250)$ REFERENCES

ANISOVICH	02	PL B542 8	A.V. Anisovich <i>et al.</i>	
ANISOVICH	01D	PL B508 6	A.V. Anisovich <i>et al.</i>	
ANISOVICH	01E	PL B513 281	A.V. Anisovich <i>et al.</i>	
AMELIN	00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)
MARTIN	80B	NP B176 355	B.R. Martin, D. Morgan	(LOUC, RHEL) JP
MARTIN	80C	NP B169 216	A.D. Martin, M.R. Pennington	(DURH) JP
CARTER	78B	NP B141 467	A.A. Carter	(LOQM)
CUTTS	78B	PR D17 16	D. Cutts <i>et al.</i>	(STON, WISC)
CARTER	77	PL 67B 117	A.A. Carter <i>et al.</i>	(LOQM, RHEL) JP
COUPLAND	77	PL 71B 460	M. Coupland <i>et al.</i>	(LOQM, RHEL)
PEASLEE	75	PL 57B 189	D.C. Peaslee <i>et al.</i>	(CANB, BARI, BROW+)
ALSPECTOR	73	PRL 30 511	J. Alspector <i>et al.</i>	(RUTG, UPNJ)
ABRAMS	70	PR D1 1917	R.J. Abrams <i>et al.</i>	(BNL)
COOPER	68	PRL 20 1059	W.A. Cooper <i>et al.</i>	(ANL)

OTHER RELATED PAPERS

MARTIN	79B	PL 86B 93	A.D. Martin, M.R. Pennington	(DURH)
CARTER	78	NP B132 176	A.A. Carter	(LOQM) JP
CARTER	77B	PL 67B 122	A.A. Carter	(LOQM) JP
CARTER	77C	NP B127 202	A.A. Carter <i>et al.</i>	(LOQM, DARE, RHEL)
ZEMANY	76	NP B103 537	P.D. Zemany <i>et al.</i>	(MSU)
EISENHAND...	75	NP B96 109	E. Eisenhandler <i>et al.</i>	(LOQM, LIVP, DARE+)
BERTANZA	74	NC 23A 209	L. Bertanza <i>et al.</i>	(PISA, PADO, TORI)
BETTINI	73	NC 15A 563	A. Bettini <i>et al.</i>	(PADO, LBL, PISA+)
DONNACHIE	73	LNC 7 285	A. Donnachie, P.R. Thomas	(MCHS)
NICHOLSON	73	PR D7 2572	H. Nicholson <i>et al.</i>	(CIT, ROCH, BNL)
FIELDS	71	PRL 27 1749	T. Fields <i>et al.</i>	(ANL, OXF)
YOH	71	PRL 26 922	J.K. Yoh <i>et al.</i>	(CIT, BNL, ROCH)
ABRAMS	67C	PRL 18 1209	R.J. Abrams <i>et al.</i>	(BNL)