τ

$$J = \frac{1}{2}$$

τ discovery paper was PERL 75. $e^+e^- \to \tau^+\tau^-$ cross-section threshold behavior and magnitude are consistent with pointlike spin-1/2 Dirac particle. BRANDELIK 78 ruled out pointlike spin-0 or spin-1 particle. FELDMAN 78 ruled out $J = 3/2$. KIRKBY 79 also ruled out J=integer, $J = 3/2$.

τ MASS

VALUE (MeV)	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
$1776.99^{+0.29}_{-0.26}$ OUR AVERAGE					
$1775.1 \pm 1.6 \pm 1.0$	13.3k	[1] ABBIENDI	00A	OPAL	1990–1995 LEP runs
$1778.2 \pm 0.8 \pm 1.2$		ANASTASSOV	97	CLEO	$E^{ee}_{\rm cm}$= 10.6 GeV
$1776.96^{+0.18}_{-0.21}{}^{+0.25}_{-0.17}$	65	[2] BAI	96	BES	$E^{ee}_{\rm cm}$= 3.54–3.57 GeV
$1776.3 \pm 2.4 \pm 1.4$	11k	[3] ALBRECHT	92M	ARG	$E^{ee}_{\rm cm}$= 9.4–10.6 GeV
$1783\ ^{+3}_{-4}$	692	[4] BACINO	78B	DLCO	$E^{ee}_{\rm cm}$= 3.1–7.4 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$1777.8 \pm 0.7 \pm 1.7$	35k	[5] BALEST	93	CLEO	Repl. by ANASTASSOV 97
$1776.9\ ^{+0.4}_{-0.5} \pm 0.2$	14	[6] BAI	92	BES	Repl. by BAI 96

[1] ABBIENDI 00A fit τ pseudomass spectrum in $\tau \to \pi^\pm \le 2\pi^0 \nu_\tau$ and $\tau \to \pi^\pm\pi^+\pi^- \le 1\pi^0 \nu_\tau$ decays. Result assumes m_{ν_τ}=0.

[2] BAI 96 fit $\sigma(e^+e^- \to \tau^+\tau^-)$ at different energies near threshold.

[3] ALBRECHT 92M fit τ pseudomass spectrum in $\tau^- \to 2\pi^-\pi^+\nu_\tau$ decays. Result assumes m_{ν_τ}=0.

[4] BACINO 78B value comes from $e^\pm X^\mp$ threshold. Published mass 1782 MeV increased by 1 MeV using the high precision $\psi(2S)$ mass measurement of ZHOLENTZ 80 to eliminate the absolute SPEAR energy calibration uncertainty.

[5] BALEST 93 fit spectra of minimum kinematically allowed τ mass in events of the type $e^+e^- \to \tau^+\tau^- \to (\pi^+ n\pi^0\nu_\tau)(\pi^- m\pi^0\nu_\tau)$ $n \le 2$, $m \le 2$, $1 \le n+m \le 3$. If $m_{\nu_\tau} \neq 0$, result increases by $(m^2_{\nu_\tau}/1100$ MeV).

[6] BAI 92 fit $\sigma(e^+e^- \to \tau^+\tau^-)$ near threshold using $e\mu$ events.

$(m_{\tau^+} - m_{\tau^-})/m_{\rm average}$

A test of *CPT* invariance.

VALUE	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
$< 3.0 \times 10^{-3}$	90	ABBIENDI	00A	OPAL	1990–1995 LEP runs

τ MEAN LIFE

VALUE (10^{-15} s)	EVTS	DOCUMENT ID		TECN	COMMENT
290.6± 1.1 OUR AVERAGE					
293.2± 2.0± 1.5		ACCIARRI	00B	L3	1991–1995 LEP runs
290.1± 1.5± 1.1		BARATE	97R	ALEP	1989–1994 LEP runs
291.4± 3.0		ABREU	96B	DLPH	1991–1993 LEP runs
289.2± 1.7± 1.2		ALEXANDER	96E	OPAL	1990–1994 LEP runs
289.0± 2.8± 4.0	57.4k	BALEST	96	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
291.2± 2.0± 1.2		BARATE	97I	ALEP	Repl. by BARATE 97R
290.1± 4.0	34k	ACCIARRI	96K	L3	Repl. by ACCIARRI 00B
297 ± 9 ± 5	1671	ABE	95Y	SLD	1992–1993 SLC runs
304 ± 14 ± 7	4100	BATTLE	92	CLEO	E_{cm}^{ee}= 10.6 GeV
301 ± 29	3780	KLEINWORT	89	JADE	E_{cm}^{ee}= 35–46 GeV
288 ± 16 ±17	807	AMIDEI	88	MRK2	E_{cm}^{ee}= 29 GeV
306 ± 20 ±14	695	BRAUNSCH...	88C	TASS	E_{cm}^{ee}= 36 GeV
299 ± 15 ±10	1311	ABACHI	87C	HRS	E_{cm}^{ee}= 29 GeV
295 ± 14 ±11	5696	ALBRECHT	87P	ARG	E_{cm}^{ee}= 9.3–10.6 GeV
309 ± 17 ± 7	3788	BAND	87B	MAC	E_{cm}^{ee}= 29 GeV
325 ± 14 ±18	8470	BEBEK	87C	CLEO	E_{cm}^{ee}= 10.5 GeV
460 ±190	102	FELDMAN	82	MRK2	E_{cm}^{ee}= 29 GeV

τ MAGNETIC MOMENT ANOMALY

The q^2 dependence is expected to be small providing no thresholds are nearby.

$\mu_\tau/(e\hbar/2m_\tau)-1 = (g_\tau-2)/2$

For a theoretical calculation [$(g_\tau-2)/2 = 11773(3) \times 10^{-7}$], see SAMUEL 91B.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
> −0.052 and < 0.058 (CL = 95%) OUR LIMIT					
> −0.052 and < 0.058	95	ACCIARRI	98E	L3	1991–1995 LEP runs
• • • We do not use the following data for averages, fits, limits, etc. • • •					
> −0.007 and < 0.005	95	[7] GONZALEZ-S..00		RVUE	$e^+e^- \to \tau^+\tau^-$ and $W \to \tau\nu_\tau$
> −0.068 and < 0.065	95	[8] ACKERSTAFF	98N	OPAL	1990–1995 LEP runs
> −0.004 and < 0.006	95	[9] ESCRIBANO	97	RVUE	$Z \to \tau^+\tau^-$ at LEP
<0.01	95	[10] ESCRIBANO	93	RVUE	$Z \to \tau^+\tau^-$ at LEP
<0.12	90	GRIFOLS	91	RVUE	$Z \to \tau\tau\gamma$ at LEP
<0.023	95	[11] SILVERMAN	83	RVUE	$e^+e^- \to \tau^+\tau^-$ at PETRA

[7] GONZALEZ-SPRINBERG 00 use data on tau lepton production at LEP1, SLC, and LEP2, and data from colliders and LEP2 to determine limits. Assume imaginary component is zero.

[8] ACKERSTAFF 98N use $Z \to \tau^+\tau^-\gamma$ events. The limit applies to an average of the form factor for off-shell τ's having p^2 ranging from m_τ^2 to $(M_Z-m_\tau)^2$.

[9] ESCRIBANO 97 use preliminary experimental results.

[10] ESCRIBANO 93 limit derived from $\Gamma(Z \to \tau^+\tau^-)$, and is on the absolute value of the magnetic moment anomaly.
[11] SILVERMAN 83 limit is derived from $e^+e^- \to \tau^+\tau^-$ total cross-section measurements for q^2 up to (37 GeV)^2.

τ ELECTRIC DIPOLE MOMENT (d_τ)

A nonzero value is forbidden by both T invariance and P invariance.

The q^2 dependence is expected to be small providing no thresholds are nearby.

Re(d_τ)

VALUE (10^{-16} e cm)	CL%	DOCUMENT ID		TECN	COMMENT
(> −0.22 and < 0.45)	95	[12] INAMI	03	BELL	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<4.6	95	[13] ALBRECHT	00	ARG	E_{cm}^{ee}= 10.4 GeV
> −3.1 and < 3.1	95	ACCIARRI	98E	L3	1991–1995 LEP runs
> −3.8 and < 3.6	95	[14] ACKERSTAFF	98N	OPAL	1990–1995 LEP runs
<0.11	95	[15,16] ESCRIBANO	97	RVUE	$Z \to \tau^+\tau^-$ at LEP
<0.5	95	[17] ESCRIBANO	93	RVUE	$Z \to \tau^+\tau^-$ at LEP
<7	90	GRIFOLS	91	RVUE	$Z \to \tau\tau\gamma$ at LEP
<1.6	90	DELAGUILA	90	RVUE	$e^+e^- \to \tau^+\tau^-$ E_{cm}^{ee}= 35 GeV

[12] INAMI 03 use $e^+e^- \to \tau^+\tau^-$ events.
[13] ALBRECHT 00 use $e^+e^- \to \tau^+\tau^-$ events. Limit is on the absolute value of Re(d_τ).
[14] ACKERSTAFF 98N use $Z \to \tau^+\tau^-\gamma$ events. The limit applies to an average of the form factor for off-shell τ's having p^2 ranging from m_τ^2 to $(M_Z - m_\tau)^2$.
[15] ESCRIBANO 97 derive the relationship $|d_\tau| = \cot\theta_W \, |d_\tau^W|$ using effective Lagrangian methods, and use a conference result $|d_\tau^W| < 5.8 \times 10^{-18}$ e cm at 95% CL (L. Silvestris, ICHEP96) to obtain this result.
[16] ESCRIBANO 97 use preliminary experimental results.
[17] ESCRIBANO 93 limit derived from $\Gamma(Z \to \tau^+\tau^-)$, and is on the absolute value of the electric dipole moment.

Im(d_τ)

VALUE (10^{-16} e cm)	CL%	DOCUMENT ID		TECN	COMMENT
(> −0.25 and < 0.008)	95	[18] INAMI	03	BELL	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.8	95	[19] ALBRECHT	00	ARG	E_{cm}^{ee}= 10.4 GeV

[18] INAMI 03 use $e^+e^- \to \tau^+\tau^-$ events.
[19] ALBRECHT 00 use $e^+e^- \to \tau^+\tau^-$ events. Limit is on the absolute value of Im(d_τ).

τ WEAK DIPOLE MOMENT (d_τ^W)

A nonzero value is forbidden by *CP* invariance.

The q^2 dependence is expected to be small providing no thresholds are nearby.

Re(d_τ^W)

VALUE (10^{-17} *e* cm)	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
<0.56	95	ACKERSTAFF	97L	OPAL	1991–1995 LEP runs
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<3.0	90	[20] ACCIARRI	98C	L3	1991–1995 LEP runs
<0.78	95	[21] AKERS	95F	OPAL	Repl. by ACKER-STAFF 97L
<1.5	95	[21] BUSKULIC	95C	ALEP	1990–1992 LEP runs
<7.0	95	[21] ACTON	92F	OPAL	$Z \to \tau^+\tau^-$ at LEP
<3.7	95	[21] BUSKULIC	92J	ALEP	Repl. by BUSKULIC 95C

[20] ACCIARRI 98C limit is on the absolute value of the real part of the weak dipole moment.
[21] Limit is on the absolute value of the real part of the weak dipole moment, and applies for $q^2 = m_Z^2$.

Im(d_τ^W)

VALUE (10^{-17} *e* cm)	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
<1.5	95	ACKERSTAFF	97L	OPAL	1991–1995 LEP runs
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<4.5	95	[22] AKERS	95F	OPAL	Repl. by ACKER-STAFF 97L

[22] Limit is on the absolute value of the imaginary part of the weak dipole moment, and applies for $q^2 = m_Z^2$.

τ WEAK ANOMALOUS MAGNETIC DIPOLE MOMENT (a_τ^W)

Electroweak radiative corrections are expected to contribute at the 10^{-6} level. See BERNABEU 95.

The q^2 dependence is expected to be small providing no thresholds are nearby.

Re(a_τ^W)

VALUE		*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
$<4.5 \times 10^{-3}$		90	[23] ACCIARRI	98C	L3	1991–1995 LEP runs
• • • We do not use the following data for averages, fits, limits, etc. • • •						
> -0.0024 and < 0.0025		95	[24] GONZALEZ-S..00		RVUE	$e^+e^- \to \tau^+\tau^-$ and $W \to \tau\nu_\tau$

[23] ACCIARRI 98C limit is on the absolute value of the real part of the weak anomalous magnetic dipole moment.
[24] GONZALEZ-SPRINBERG 00 use data on tau lepton production at LEP1, SLC, and LEP2, and data from colliders and LEP2 to determine limits. Assume imaginary component is zero.

$\mathrm{Im}(a_\tau^w)$

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<9.9\times10^{-3}$	90	[25] ACCIARRI	98C	L3	1991–1995 LEP runs

[25] ACCIARRI 98C limit is on the absolute value of the imaginary part of the weak anomalous magnetic dipole moment.

τ^- DECAY MODES

τ^+ modes are charge conjugates of the modes below. "$h^\pm$" stands for $\pi^\pm$ or $K^\pm$. "ℓ" stands for e or μ. "Neutrals" stands for γ's and/or π^0's.

	Mode		Fraction (Γ_i/Γ)	Scale factor/ Confidence level
	Modes with one charged particle			
Γ_1	particle$^-$ ≥ 0 neutrals $\geq 0K^0\nu_\tau$ ("1-prong")		(85.35 ± 0.07) %	S=1.1
Γ_2	particle$^-$ ≥ 0 neutrals $\geq 0K_L^0\nu_\tau$		(84.71 ± 0.07) %	S=1.1
Γ_3	$\mu^-\overline{\nu}_\mu\nu_\tau$	[a]	(17.36 ± 0.05) %	
Γ_4	$\mu^-\overline{\nu}_\mu\nu_\tau\gamma$	[b]	(3.6 ± 0.4) × 10^{-3}	
Γ_5	$e^-\overline{\nu}_e\nu_\tau$	[a]	(17.84 ± 0.06) %	
Γ_6	$e^-\overline{\nu}_e\nu_\tau\gamma$	[b]	(1.75 ± 0.18) %	
Γ_7	$h^-\geq 0K_L^0\nu_\tau$		(12.30 ± 0.10) %	S=1.4
Γ_8	$h^-\nu_\tau$		(11.75 ± 0.10) %	S=1.4
Γ_9	$\pi^-\nu_\tau$	[a]	(11.06 ± 0.11) %	S=1.4
Γ_{10}	$K^-\nu_\tau$	[a]	(6.86 ± 0.23) × 10^{-3}	
Γ_{11}	$h^-\geq 1$ neutrals ν_τ		(36.92 ± 0.14) %	S=1.1
Γ_{12}	$h^-\pi^0\nu_\tau$		(25.86 ± 0.13) %	S=1.1
Γ_{13}	$\pi^-\pi^0\nu_\tau$	[a]	(25.41 ± 0.14) %	S=1.1
Γ_{14}	$\pi^-\pi^0$ non-$\rho(770)\nu_\tau$		(3.0 ± 3.2) × 10^{-3}	
Γ_{15}	$K^-\pi^0\nu_\tau$	[a]	(4.50 ± 0.30) × 10^{-3}	
Γ_{16}	$h^-\geq 2\pi^0\nu_\tau$		(10.76 ± 0.15) %	S=1.1
Γ_{17}	$h^-2\pi^0\nu_\tau$		(9.39 ± 0.14) %	S=1.1
Γ_{18}	$h^-2\pi^0\nu_\tau$(ex.K^0)		(9.23 ± 0.14) %	S=1.1
Γ_{19}	$\pi^-2\pi^0\nu_\tau$(ex.K^0)	[a]	(9.17 ± 0.14) %	S=1.1
Γ_{20}	$\pi^-2\pi^0\nu_\tau$(ex.K^0), scalar		< 9 × 10^{-3}	CL=95%
Γ_{21}	$\pi^-2\pi^0\nu_\tau$(ex.K^0), vector		< 7 × 10^{-3}	CL=95%
Γ_{22}	$K^-2\pi^0\nu_\tau$(ex.K^0)	[a]	(5.8 ± 2.3) × 10^{-4}	
Γ_{23}	$h^-\geq 3\pi^0\nu_\tau$		(1.37 ± 0.11) %	S=1.1
Γ_{24}	$h^-3\pi^0\nu_\tau$		(1.21 ± 0.10) %	
Γ_{25}	$\pi^-3\pi^0\nu_\tau$(ex.K^0)	[a]	(1.08 ± 0.10) %	
Γ_{26}	$K^-3\pi^0\nu_\tau$(ex.K^0, η)	[a]	(3.7 $^{+2.2}_{-2.0}$) × 10^{-4}	
Γ_{27}	$h^-4\pi^0\nu_\tau$(ex.K^0)		(1.6 ± 0.6) × 10^{-3}	

Γ_{28}	$h^- 4\pi^0 \nu_\tau$ (ex. K^0,η)	[a]	(1.0 $^{+0.6}_{-0.5}$) $\times 10^{-3}$	
Γ_{29}	$K^- \geq 0\pi^0 \geq 0K^0 \geq 0\gamma \ \nu_\tau$		(1.56 $\pm$ 0.04) %	
Γ_{30}	$K^- \geq 1$ (π^0 or K^0 or γ) ν_τ		(8.74 $\pm$ 0.35) $\times 10^{-3}$	

Modes with K^0's

Γ_{31}	K^0_S (particles)$^- \nu_\tau$		(9.2 $\pm$ 0.4) $\times 10^{-3}$	S=1.1
Γ_{32}	$h^- \overline{K}^0 \nu_\tau$		(1.05 $\pm$ 0.04) %	S=1.1
Γ_{33}	$\pi^- \overline{K}^0 \nu_\tau$	[a]	(8.9 $\pm$ 0.4) $\times 10^{-3}$	S=1.1
Γ_{34}	$\pi^- \overline{K}^0$ (non-$K^*(892)^-$) ν_τ		$<$ 1.7 $\times 10^{-3}$	CL=95%
Γ_{35}	$K^- K^0 \nu_\tau$	[a]	(1.54 $\pm$ 0.16) $\times 10^{-3}$	
Γ_{36}	$K^- K^0 \geq 0\pi^0 \nu_\tau$		(3.09 $\pm$ 0.24) $\times 10^{-3}$	
Γ_{37}	$h^- \overline{K}^0 \pi^0 \nu_\tau$		(5.2 $\pm$ 0.4) $\times 10^{-3}$	
Γ_{38}	$\pi^- \overline{K}^0 \pi^0 \nu_\tau$	[a]	(3.7 $\pm$ 0.4) $\times 10^{-3}$	
Γ_{39}	$\overline{K}^0 \rho^- \nu_\tau$		(2.2 $\pm$ 0.5) $\times 10^{-3}$	
Γ_{40}	$K^- K^0 \pi^0 \nu_\tau$	[a]	(1.55 $\pm$ 0.20) $\times 10^{-3}$	
Γ_{41}	$\pi^- \overline{K}^0 \geq 1\pi^0 \nu_\tau$		(3.2 $\pm$ 1.0) $\times 10^{-3}$	
Γ_{42}	$\pi^- \overline{K}^0 \pi^0 \pi^0 \nu_\tau$		(2.6 $\pm$ 2.4) $\times 10^{-4}$	
Γ_{43}	$K^- K^0 \pi^0 \pi^0 \nu_\tau$		$<$ 1.6 $\times 10^{-4}$	CL=95%
Γ_{44}	$\pi^- K^0 \overline{K}^0 \nu_\tau$		(1.59 $\pm$ 0.29) $\times 10^{-3}$	S=1.1
Γ_{45}	$\pi^- K^0_S K^0_S \nu_\tau$	[a]	(2.4 $\pm$ 0.5) $\times 10^{-4}$	
Γ_{46}	$\pi^- K^0_S K^0_L \nu_\tau$	[a]	(1.10 $\pm$ 0.28) $\times 10^{-3}$	S=1.1
Γ_{47}	$\pi^- K^0 \overline{K}^0 \pi^0 \nu_\tau$		(3.1 $\pm$ 2.3) $\times 10^{-4}$	
Γ_{48}	$\pi^- K^0_S K^0_S \pi^0 \nu_\tau$		$<$ 2.0 $\times 10^{-4}$	CL=95%
Γ_{49}	$\pi^- K^0_S K^0_L \pi^0 \nu_\tau$		(3.1 $\pm$ 1.2) $\times 10^{-4}$	
Γ_{50}	$K^0 h^+ h^- h^- \geq 0$ neutrals ν_τ		$<$ 1.7 $\times 10^{-3}$	CL=95%
Γ_{51}	$K^0 h^+ h^- h^- \nu_\tau$		(2.3 $\pm$ 2.0) $\times 10^{-4}$	

Modes with three charged particles

Γ_{52}	$h^- h^- h^+ \geq 0$ neutrals $\geq 0K^0_L \nu_\tau$		(15.20 $\pm$ 0.07) %	S=1.1
Γ_{53}	$h^- h^- h^+ \geq 0$ neutrals ν_τ (ex. $K^0_S \to \pi^+ \pi^-$) ("3-prong")		(14.57 $\pm$ 0.07) %	S=1.1
Γ_{54}	$h^- h^- h^+ \nu_\tau$		(10.01 $\pm$ 0.09) %	S=1.2
Γ_{55}	$h^- h^- h^+ \nu_\tau$ (ex. K^0)		(9.65 $\pm$ 0.09) %	S=1.2
Γ_{56}	$h^- h^- h^+ \nu_\tau$ (ex. K^0,ω)		(9.61 $\pm$ 0.09) %	S=1.2
Γ_{57}	$\pi^- \pi^+ \pi^- \nu_\tau$		(9.52 $\pm$ 0.10) %	S=1.2
Γ_{58}	$\pi^- \pi^+ \pi^- \nu_\tau$ (ex. K^0)		(9.22 $\pm$ 0.10) %	S=1.2
Γ_{59}	$\pi^- \pi^+ \pi^- \nu_\tau$ (ex. K^0), non-axial vector		$<$ 2.4 %	CL=95%
Γ_{60}	$\pi^- \pi^+ \pi^- \nu_\tau$ (ex. K^0,ω)	[a]	(9.18 $\pm$ 0.10) %	S=1.2
Γ_{61}	$h^- h^- h^+ \geq 1$ neutrals ν_τ		(5.19 $\pm$ 0.10) %	S=1.3
Γ_{62}	$h^- h^- h^+ \geq 1$ neutrals ν_τ (ex. $K^0_S \to \pi^+ \pi^-$)		(4.92 $\pm$ 0.10) %	S=1.3

Γ_{63}	$h^-\,h^-\,h^+\,\pi^0\,\nu_\tau$		(4.53 ± 0.09) %		S=1.3
Γ_{64}	$h^-\,h^-\,h^+\,\pi^0\,\nu_\tau$(ex.$K^0$)		(4.35 ± 0.09) %		S=1.3
Γ_{65}	$h^-\,h^-\,h^+\,\pi^0\,\nu_\tau$(ex. K^0, ω)		(2.62 ± 0.09) %		S=1.2
Γ_{66}	$\pi^-\,\pi^+\,\pi^-\,\pi^0\,\nu_\tau$		(4.37 ± 0.10) %		S=1.3
Γ_{67}	$\pi^-\,\pi^+\,\pi^-\,\pi^0\,\nu_\tau$(ex.$K^0$)		(4.24 ± 0.10) %		S=1.3
Γ_{68}	$\pi^-\,\pi^+\,\pi^-\,\pi^0\,\nu_\tau$(ex.$K^0$,$\omega$)	[a]	(2.51 ± 0.09) %		S=1.2
Γ_{69}	$h^-\,\rho\,\pi^0\,\nu_\tau$				
Γ_{70}	$h^-\,\rho^+\,h^-\,\nu_\tau$				
Γ_{71}	$h^-\,\rho^-\,h^+\,\nu_\tau$				
Γ_{72}	$h^-\,h^-\,h^+\,2\pi^0\,\nu_\tau$		(5.5 ± 0.4) × 10^{-3}		
Γ_{73}	$h^-\,h^-\,h^+\,2\pi^0\,\nu_\tau$(ex.$K^0$)		(5.4 ± 0.4) × 10^{-3}		
Γ_{74}	$h^-\,h^-\,h^+\,2\pi^0\,\nu_\tau$(ex.$K^0$,$\omega$,$\eta$)	[a]	(1.1 ± 0.4) × 10^{-3}		
Γ_{75}	$h^-\,h^-\,h^+\,3\pi^0\,\nu_\tau$	[a]	(2.3 ± 0.8) × 10^{-4}		S=1.6
Γ_{76}	$K^-\,h^+\,h^-\,\geq 0$ neutrals ν_τ		(6.5 ± 0.5) × 10^{-3}		S=1.4
Γ_{77}	$K^-\,h^+\,\pi^-\,\nu_\tau$(ex.$K^0$)		(4.4 ± 0.5) × 10^{-3}		S=1.5
Γ_{78}	$K^-\,h^+\,\pi^-\,\pi^0\,\nu_\tau$(ex.$K^0$)		(1.10 ± 0.22) × 10^{-3}		
Γ_{79}	$K^-\,\pi^+\,\pi^-\,\geq 0$ neutrals ν_τ		(4.5 ± 0.5) × 10^{-3}		S=1.4
Γ_{80}	$K^-\,\pi^+\,\pi^-\,\geq 0\pi^0\,\nu_\tau$(ex.$K^0$)		(3.5 ± 0.5) × 10^{-3}		S=1.4
Γ_{81}	$K^-\,\pi^+\,\pi^-\,\nu_\tau$		(3.3 ± 0.5) × 10^{-3}		S=1.5
Γ_{82}	$K^-\,\pi^+\,\pi^-\,\nu_\tau$(ex.$K^0$)	[a]	(2.8 ± 0.5) × 10^{-3}		S=1.5
Γ_{83}	$K^-\,\rho^0\,\nu_\tau \rightarrow$ $K^-\,\pi^+\,\pi^-\,\nu_\tau$		(1.3 ± 0.5) × 10^{-3}		
Γ_{84}	$K^-\,\pi^+\,\pi^-\,\pi^0\,\nu_\tau$		(1.24 ± 0.25) × 10^{-3}		
Γ_{85}	$K^-\,\pi^+\,\pi^-\,\pi^0\,\nu_\tau$(ex.$K^0$)		(7.0 ± 2.4) × 10^{-4}		
Γ_{86}	$K^-\,\pi^+\,\pi^-\,\pi^0\,\nu_\tau$(ex.$K^0$,$\eta$)	[a]	(6.4 ± 2.4) × 10^{-4}		
Γ_{87}	$K^-\,\pi^+\,K^-\,\geq 0$ neut. ν_τ		< 9	× 10^{-4}	CL=95%
Γ_{88}	$K^-\,K^+\,\pi^-\,\geq 0$ neut. ν_τ		(2.00 ± 0.23) × 10^{-3}		
Γ_{89}	$K^-\,K^+\,\pi^-\,\nu_\tau$	[a]	(1.60 ± 0.19) × 10^{-3}		
Γ_{90}	$K^-\,K^+\,\pi^-\,\pi^0\,\nu_\tau$	[a]	(4.0 ± 1.6) × 10^{-4}		
Γ_{91}	$K^-\,K^+\,K^-\,\geq 0$ neut. ν_τ		< 2.1	× 10^{-3}	CL=95%
Γ_{92}	$K^-\,K^+\,K^-\,\nu_\tau$		< 1.9	× 10^{-4}	CL=90%
Γ_{93}	$\pi^-\,K^+\,\pi^-\,\geq 0$ neut. ν_τ		< 2.5	× 10^{-3}	CL=95%
Γ_{94}	$e^-\,e^-\,e^+\,\overline{\nu}_e\,\nu_\tau$		(2.8 ± 1.5) × 10^{-5}		
Γ_{95}	$\mu^-\,e^-\,e^+\,\overline{\nu}_\mu\,\nu_\tau$		< 3.6	× 10^{-5}	CL=90%

Modes with five charged particles

Γ_{96}	$3h^-2h^+\,\geq 0$ neutrals ν_τ (ex. $K^0_S \rightarrow \pi^-\pi^+$) ("5-prong")		(1.00 ± 0.06) × 10^{-3}		
Γ_{97}	$3h^-2h^+\,\nu_\tau$(ex.K^0)	[a]	(8.2 ± 0.6) × 10^{-4}		
Γ_{98}	$3h^-2h^+\,\pi^0\,\nu_\tau$(ex.$K^0$)	[a]	(1.81 ± 0.27) × 10^{-4}		
Γ_{99}	$3h^-2h^+\,2\pi^0\,\nu_\tau$		< 1.1	× 10^{-4}	CL=90%

Miscellaneous other allowed modes

Γ_{100}	$(5\pi)^-\nu_\tau$		$(8.0 \pm 0.7) \times 10^{-3}$		
Γ_{101}	$4h^-3h^+ \geq 0$ neutrals ν_τ ("7-prong")		$< 2.4 \times 10^{-6}$		CL=90%
Γ_{102}	$X^-(S=-1)\nu_\tau$		(2.86 ± 0.09) %		S=1.1
Γ_{103}	$K^*(892)^- \geq 0$ neutrals $\geq 0K^0_L\nu_\tau$		(1.42 ± 0.18) %		S=1.4
Γ_{104}	$K^*(892)^-\nu_\tau$		(1.29 ± 0.05) %		
Γ_{105}	$K^*(892)^0K^- \geq 0$ neutrals ν_τ		$(3.2 \pm 1.4) \times 10^{-3}$		
Γ_{106}	$K^*(892)^0K^-\nu_\tau$		$(2.1 \pm 0.4) \times 10^{-3}$		
Γ_{107}	$\overline{K}^*(892)^0\pi^- \geq 0$ neutrals ν_τ		$(3.8 \pm 1.7) \times 10^{-3}$		
Γ_{108}	$\overline{K}^*(892)^0\pi^-\nu_\tau$		$(2.2 \pm 0.5) \times 10^{-3}$		
Γ_{109}	$(\overline{K}^*(892)\pi)^-\nu_\tau \to \pi^-\overline{K}^0\pi^0\nu_\tau$		$(1.0 \pm 0.4) \times 10^{-3}$		
Γ_{110}	$K_1(1270)^-\nu_\tau$		$(4.7 \pm 1.1) \times 10^{-3}$		
Γ_{111}	$K_1(1400)^-\nu_\tau$		$(1.7 \pm 2.6) \times 10^{-3}$		S=1.7
Γ_{112}	$K^*(1410)^-\nu_\tau$		$(1.5 \, {}^{+1.4}_{-1.0}) \times 10^{-3}$		
Γ_{113}	$K^*_0(1430)^-\nu_\tau$		$< 5 \times 10^{-4}$		CL=95%
Γ_{114}	$K^*_2(1430)^-\nu_\tau$		$< 3 \times 10^{-3}$		CL=95%
Γ_{115}	$a_0(980)^- \geq 0$ neutrals ν_τ				
Γ_{116}	$\eta\pi^-\nu_\tau$		$< 1.4 \times 10^{-4}$		CL=95%
Γ_{117}	$\eta\pi^-\pi^0\nu_\tau$	[a]	$(1.74 \pm 0.24) \times 10^{-3}$		
Γ_{118}	$\eta\pi^-\pi^0\pi^0\nu_\tau$		$(1.5 \pm 0.5) \times 10^{-4}$		
Γ_{119}	$\eta K^-\nu_\tau$	[a]	$(2.7 \pm 0.6) \times 10^{-4}$		
Γ_{120}	$\eta K^*(892)^-\nu_\tau$		$(2.9 \pm 0.9) \times 10^{-4}$		
Γ_{121}	$\eta K^-\pi^0\nu_\tau$		$(1.8 \pm 0.9) \times 10^{-4}$		
Γ_{122}	$\eta\overline{K}^0\pi^-\nu_\tau$		$(2.2 \pm 0.7) \times 10^{-4}$		
Γ_{123}	$\eta\pi^+\pi^-\pi^- \geq 0$ neutrals ν_τ		$< 3 \times 10^{-3}$		CL=90%
Γ_{124}	$\eta\pi^-\pi^+\pi^-\nu_\tau$		$(2.3 \pm 0.5) \times 10^{-4}$		
Γ_{125}	$\eta a_1(1260)^-\nu_\tau \to \eta\pi^-\rho^0\nu_\tau$		$< 3.9 \times 10^{-4}$		CL=90%
Γ_{126}	$\eta\eta\pi^-\nu_\tau$		$< 1.1 \times 10^{-4}$		CL=95%
Γ_{127}	$\eta\eta\pi^-\pi^0\nu_\tau$		$< 2.0 \times 10^{-4}$		CL=95%
Γ_{128}	$\eta'(958)\pi^-\nu_\tau$		$< 7.4 \times 10^{-5}$		CL=90%
Γ_{129}	$\eta'(958)\pi^-\pi^0\nu_\tau$		$< 8.0 \times 10^{-5}$		CL=90%
Γ_{130}	$\phi\pi^-\nu_\tau$		$< 2.0 \times 10^{-4}$		CL=90%
Γ_{131}	$\phi K^-\nu_\tau$		$< 6.7 \times 10^{-5}$		CL=90%
Γ_{132}	$f_1(1285)\pi^-\nu_\tau$		$(5.8 \pm 2.3) \times 10^{-4}$		
Γ_{133}	$f_1(1285)\pi^-\nu_\tau \to \eta\pi^-\pi^+\pi^-\nu_\tau$		$(1.3 \pm 0.4) \times 10^{-4}$		
Γ_{134}	$\pi(1300)^-\nu_\tau \to (\rho\pi)^-\nu_\tau \to (3\pi)^-\nu_\tau$		$< 1.0 \times 10^{-4}$		CL=90%
Γ_{135}	$\pi(1300)^-\nu_\tau \to ((\pi\pi)_{S-\text{wave}}\,\pi)^-\nu_\tau \to (3\pi)^-\nu_\tau$		$< 1.9 \times 10^{-4}$		CL=90%

Γ_{136}	$h^- \omega \geq 0$ neutrals ν_τ		(2.37 ± 0.08) %	
Γ_{137}	$h^- \omega \nu_\tau$	[a]	(1.94 ± 0.07) %	
Γ_{138}	$h^- \omega \pi^0 \nu_\tau$	[a]	(4.3 ± 0.5) $\times 10^{-3}$	
Γ_{139}	$h^- \omega 2\pi^0 \nu_\tau$		(1.4 ± 0.5) $\times 10^{-4}$	
Γ_{140}	$2h^- h^+ \omega \nu_\tau$		(1.20 ± 0.22) $\times 10^{-4}$	

Lepton Family number (LF), Lepton number (L), or Baryon number (B) violating modes

L means lepton number violation (*e.g.* $\tau^- \to e^+ \pi^- \pi^-$). Following common usage, LF means lepton family violation *and not* lepton number violation (*e.g.* $\tau^- \to e^- \pi^+ \pi^-$). B means baryon number violation.

Γ_{141}	$e^- \gamma$	LF	< 2.7	$\times 10^{-6}$	CL=90%
Γ_{142}	$\mu^- \gamma$	LF	< 1.1	$\times 10^{-6}$	CL=90%
Γ_{143}	$e^- \pi^0$	LF	< 3.7	$\times 10^{-6}$	CL=90%
Γ_{144}	$\mu^- \pi^0$	LF	< 4.0	$\times 10^{-6}$	CL=90%
Γ_{145}	$e^- K^0_S$	LF	< 9.1	$\times 10^{-7}$	CL=90%
Γ_{146}	$\mu^- K^0_S$	LF	< 9.5	$\times 10^{-7}$	CL=90%
Γ_{147}	$e^- \eta$	LF	< 8.2	$\times 10^{-6}$	CL=90%
Γ_{148}	$\mu^- \eta$	LF	< 9.6	$\times 10^{-6}$	CL=90%
Γ_{149}	$e^- \rho^0$	LF	< 2.0	$\times 10^{-6}$	CL=90%
Γ_{150}	$\mu^- \rho^0$	LF	< 6.3	$\times 10^{-6}$	CL=90%
Γ_{151}	$e^- K^*(892)^0$	LF	< 5.1	$\times 10^{-6}$	CL=90%
Γ_{152}	$\mu^- K^*(892)^0$	LF	< 7.5	$\times 10^{-6}$	CL=90%
Γ_{153}	$e^- \overline{K}^*(892)^0$	LF	< 7.4	$\times 10^{-6}$	CL=90%
Γ_{154}	$\mu^- \overline{K}^*(892)^0$	LF	< 7.5	$\times 10^{-6}$	CL=90%
Γ_{155}	$e^- \phi$	LF	< 6.9	$\times 10^{-6}$	CL=90%
Γ_{156}	$\mu^- \phi$	LF	< 7.0	$\times 10^{-6}$	CL=90%
Γ_{157}	$e^- e^+ e^-$	LF	< 2.9	$\times 10^{-6}$	CL=90%
Γ_{158}	$e^- \mu^+ \mu^-$	LF	< 1.8	$\times 10^{-6}$	CL=90%
Γ_{159}	$e^+ \mu^- \mu^-$	LF	< 1.5	$\times 10^{-6}$	CL=90%
Γ_{160}	$\mu^- e^+ e^-$	LF	< 1.7	$\times 10^{-6}$	CL=90%
Γ_{161}	$\mu^+ e^- e^-$	LF	< 1.5	$\times 10^{-6}$	CL=90%
Γ_{162}	$\mu^- \mu^+ \mu^-$	LF	< 1.9	$\times 10^{-6}$	CL=90%
Γ_{163}	$e^- \pi^+ \pi^-$	LF	< 2.2	$\times 10^{-6}$	CL=90%
Γ_{164}	$e^+ \pi^- \pi^-$	L	< 1.9	$\times 10^{-6}$	CL=90%
Γ_{165}	$\mu^- \pi^+ \pi^-$	LF	< 8.2	$\times 10^{-6}$	CL=90%
Γ_{166}	$\mu^+ \pi^- \pi^-$	L	< 3.4	$\times 10^{-6}$	CL=90%
Γ_{167}	$e^- \pi^+ K^-$	LF	< 6.4	$\times 10^{-6}$	CL=90%
Γ_{168}	$e^- \pi^- K^+$	LF	< 3.8	$\times 10^{-6}$	CL=90%
Γ_{169}	$e^+ \pi^- K^-$	L	< 2.1	$\times 10^{-6}$	CL=90%
Γ_{170}	$e^- K^0_S K^0_S$	LF	< 2.2	$\times 10^{-6}$	CL=90%
Γ_{171}	$e^- K^+ K^-$	LF	< 6.0	$\times 10^{-6}$	CL=90%
Γ_{172}	$e^+ K^- K^-$	L	< 3.8	$\times 10^{-6}$	CL=90%
Γ_{173}	$\mu^- \pi^+ K^-$	LF	< 7.5	$\times 10^{-6}$	CL=90%

Γ_{174}	$\mu^- \pi^- K^+$	*LF*	$<$	7.4	$\times 10^{-6}$	CL=90%
Γ_{175}	$\mu^+ \pi^- K^-$	*L*	$<$	7.0	$\times 10^{-6}$	CL=90%
Γ_{176}	$\mu^- K^0_S K^0_S$	*LF*	$<$	3.4	$\times 10^{-6}$	CL=90%
Γ_{177}	$\mu^- K^+ K^-$	*LF*	$<$	1.5	$\times 10^{-5}$	CL=90%
Γ_{178}	$\mu^+ K^- K^-$	*L*	$<$	6.0	$\times 10^{-6}$	CL=90%
Γ_{179}	$e^- \pi^0 \pi^0$	*LF*	$<$	6.5	$\times 10^{-6}$	CL=90%
Γ_{180}	$\mu^- \pi^0 \pi^0$	*LF*	$<$	1.4	$\times 10^{-5}$	CL=90%
Γ_{181}	$e^- \eta \eta$	*LF*	$<$	3.5	$\times 10^{-5}$	CL=90%
Γ_{182}	$\mu^- \eta \eta$	*LF*	$<$	6.0	$\times 10^{-5}$	CL=90%
Γ_{183}	$e^- \pi^0 \eta$	*LF*	$<$	2.4	$\times 10^{-5}$	CL=90%
Γ_{184}	$\mu^- \pi^0 \eta$	*LF*	$<$	2.2	$\times 10^{-5}$	CL=90%
Γ_{185}	$\overline{p}\gamma$	*L,B*	$<$	3.5	$\times 10^{-6}$	CL=90%
Γ_{186}	$\overline{p}\pi^0$	*L,B*	$<$	1.5	$\times 10^{-5}$	CL=90%
Γ_{187}	$\overline{p}2\pi^0$	*L,B*	$<$	3.3	$\times 10^{-5}$	CL=90%
Γ_{188}	$\overline{p}\eta$	*L,B*	$<$	8.9	$\times 10^{-6}$	CL=90%
Γ_{189}	$\overline{p}\pi^0 \eta$	*L,B*	$<$	2.7	$\times 10^{-5}$	CL=90%
Γ_{190}	e^- light boson	*LF*	$<$	2.7	$\times 10^{-3}$	CL=95%
Γ_{191}	μ^- light boson	*LF*	$<$	5	$\times 10^{-3}$	CL=95%

[*a*] Basis mode for the τ.

[*b*] See the Particle Listings below for the energy limits used in this measurement.

CONSTRAINED FIT INFORMATION

An overall fit to 64 branching ratios uses 126 measurements and one constraint to determine 31 parameters. The overall fit has a $\chi^2 = 57.3$ for 96 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\rm total}$. The fit constrains the x_i whose labels appear in this array to sum to one.

	x_3	x_5	x_9	x_{10}	x_{13}	x_{15}	x_{19}	x_{22}	x_{25}	x_{26}
x_5	1									
x_9	−1	−1								
x_{10}	0	0	−21							
x_{13}	−14	−14	−25	1						
x_{15}	0	0	1	−3	−22					
x_{19}	−14	−15	−25	0	−36	4				
x_{22}	0	0	1	−2	5	−21	−16			
x_{25}	−8	−8	−15	0	−18	5	−24	3		
x_{26}	0	0	1	−2	5	−19	3	−13	−22	
x_{28}	−4	−4	−7	0	−10	0	−12	0	−7	0
x_{33}	−2	−2	−12	0	−3	1	−8	1	−2	1
x_{35}	−1	−1	−4	−1	2	−12	−1	−8	2	−8
x_{38}	−2	−2	−3	0	−5	3	−5	2	−9	2
x_{40}	−1	−1	0	−1	2	−16	1	−11	−1	−10
x_{45}	0	0	−2	0	0	0	0	0	0	0
x_{46}	−2	−2	−2	0	−4	1	−4	1	−2	1
x_{60}	−1	−1	−2	0	−3	0	−3	0	−2	0
x_{68}	−1	−1	−3	0	−3	0	−4	0	−2	0
x_{74}	0	0	0	0	0	0	0	0	0	0
x_{75}	0	0	0	0	0	0	0	0	0	0
x_{82}	0	0	0	0	0	0	0	0	0	0
x_{86}	0	0	0	0	0	0	0	0	0	0
x_{89}	0	0	0	0	0	0	0	0	0	0
x_{90}	0	0	0	0	0	0	0	0	0	0
x_{97}	0	0	−1	0	−1	0	−1	0	0	0
x_{98}	0	0	0	0	0	0	0	0	0	0
x_{117}	−1	−1	−1	0	−2	0	−2	0	−1	0
x_{119}	0	0	0	0	0	−3	0	−2	−2	−2
x_{137}	−1	−1	−2	0	−3	0	−3	0	−2	0
x_{138}	−1	−1	−1	0	−2	0	−2	0	−1	0

x_{33}	−1									
x_{35}	0	−5								
x_{38}	−1	−7	0							
x_{40}	0	−2	−15	−19						
x_{45}	0	−2	−1	−2	0					
x_{46}	−1	−12	−4	−10	−3	−3				
x_{60}	−1	−7	−3	3	2	0	0			
x_{68}	−1	4	2	−5	−2	0	0	−45		
x_{74}	2	1	0	1	0	0	0	−7	−7	
x_{75}	0	0	0	0	0	0	0	−3	−3	0
x_{82}	0	0	0	0	0	0	0	−43	2	0
x_{86}	0	0	0	0	0	0	0	5	−15	0
x_{89}	0	0	0	0	0	0	0	−8	−2	0
x_{90}	0	0	0	0	0	0	0	−6	−1	0
x_{97}	0	0	0	0	0	0	0	−1	−1	0
x_{98}	0	0	0	0	0	0	0	0	0	0
x_{117}	−14	0	0	0	0	0	0	−1	−1	−14
x_{119}	0	0	−1	0	−2	0	0	0	0	0
x_{137}	−1	2	1	−2	−1	0	0	−22	−28	−3
x_{138}	−1	1	0	1	0	0	0	−8	−10	−44
	x_{28}	x_{33}	x_{35}	x_{38}	x_{40}	x_{45}	x_{46}	x_{60}	x_{68}	x_{74}

x_{82}	0									
x_{86}	0	−19								
x_{89}	0	−14	8							
x_{90}	0	10	−46	−14						
x_{97}	0	0	0	0	0					
x_{98}	0	0	0	0	0	−19				
x_{117}	0	0	0	0	0	0	0			
x_{119}	0	0	−6	0	0	0	0	0		
x_{137}	−1	−1	2	−1	2	0	0	0	0	
x_{138}	−1	0	0	0	−1	0	0	0	0	−4
	x_{75}	x_{82}	x_{86}	x_{89}	x_{90}	x_{97}	x_{98}	x_{117}	x_{119}	x_{137}

A REVIEW GOES HERE – Check our WWW List of Reviews

τ^- BRANCHING RATIOS

$\Gamma(\text{particle}^- \geq 0\ \text{neutrals} \geq 0 K^0 \nu_\tau(\text{"1-prong"}))/\Gamma_{\text{total}}$ Γ_1/Γ

$\Gamma_1/\Gamma = (\Gamma_3+\Gamma_5+\Gamma_9+\Gamma_{10}+\Gamma_{13}+\Gamma_{15}+\Gamma_{19}+\Gamma_{22}+\Gamma_{25}+\Gamma_{26}+\Gamma_{28}+\Gamma_{33}+\Gamma_{35}+\Gamma_{38}+\Gamma_{40}+2\Gamma_{45}+\Gamma_{46}+0.708\Gamma_{117}+0.715\Gamma_{119}+0.09\Gamma_{137}+0.09\Gamma_{138})/\Gamma$

The charged particle here can be e, μ, or hadron. In many analyses, the sum of the topological branching fractions (1, 3, and 5 prongs) is constrained to be unity. Since the 5-prong fraction is very small, the measured 1-prong and 3-prong fractions are highly correlated and cannot be treated as independent quantities in our overall fit. We arbitrarily choose to use the 3-prong fraction in our fit, and leave the 1-prong fraction out. We do, however, use these 1-prong measurements in our average below. The measurements used only for the average are marked "avg," whereas "f&a" marks a result used for the fit and the average.

VALUE (%)		*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
85.35 ±0.07 OUR FIT Error includes scale factor of 1.1.						
85.26 ±0.13 OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.						
85.316±0.093±0.049	avg	78k	[26] ABREU	01M	DLPH	1992–1995 LEP runs
85.274±0.105±0.073	avg		[27] ACHARD	01D	L3	1992–1995 LEP runs
84.48 ±0.27 ±0.23	avg		ACTON	92H	OPAL	1990–1991 LEP runs
85.45 $^{+0.69}_{-0.73}$ ±0.65	f&a		DECAMP	92C	ALEP	1989–1990 LEP runs

[26] The correlation coefficients between this meausrement and the ABREU 01M measurements of B($\tau \to$ 3-prong) and B($\tau \to$ 5-prong) are -0.98 and -0.08 respectively.

[27] The correlation coefficients between this measurement and the ACHARD 01D measurements of B($\tau \to$ "3-prong") and B($\tau \to$ "5-prong") are -0.978 and -0.082 respectively.



$\Gamma\left(\text{particle}^- \geq 0 \text{ neutrals } \geq 0 K^0 \nu_\tau (\text{“1-prong”})\right)/\Gamma_{\text{total}}$

$\Gamma(\text{particle}^- \geq 0 \text{ neutrals } \geq 0 K_L^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_2/Γ

$\Gamma_2/\Gamma = (\Gamma_3+\Gamma_5+\Gamma_9+\Gamma_{10}+\Gamma_{13}+\Gamma_{15}+\Gamma_{19}+\Gamma_{22}+\Gamma_{25}+\Gamma_{26}+\Gamma_{28}+0.6569\Gamma_{33}+0.6569\Gamma_{35}+0.6569\Gamma_{38}+0.6569\Gamma_{40}+1.0985\Gamma_{45}+0.3139\Gamma_{46}+0.708\Gamma_{117}+0.715\Gamma_{119}+0.09\Gamma_{137}+0.09\Gamma_{138})/\Gamma$

VALUE (%)		EVTS	DOCUMENT ID		TECN	COMMENT
84.71±0.07 OUR FIT	Error includes scale factor of 1.1.					
85.1 ±0.4 OUR AVERAGE						
85.6 ±0.6 ±0.3	avg	3300	[28] ADEVA	91F	L3	E_{cm}^{ee}= 88.3–94.3 GeV
84.9 ±0.4 ±0.3	avg		BEHREND	89B	CELL	E_{cm}^{ee}= 14–47 GeV
84.7 ±0.8 ±0.6	avg		[29] AIHARA	87B	TPC	E_{cm}^{ee}= 29 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •						
86.4 ±0.3 ±0.3			ABACHI	89B	HRS	E_{cm}^{ee}= 29 GeV
87.1 ±1.0 ±0.7			[30] BURCHAT	87	MRK2	E_{cm}^{ee}= 29 GeV
87.2 ±0.5 ±0.8			SCHMIDKE	86	MRK2	E_{cm}^{ee}= 29 GeV
84.7 ±1.1 $^{+1.6}_{-1.3}$		169	[31] ALTHOFF	85	TASS	E_{cm}^{ee}= 34.5 GeV
86.1 ±0.5 ±0.9			BARTEL	85F	JADE	E_{cm}^{ee}= 34.6 GeV
87.8 ±1.3 ±3.9			[32] BERGER	85	PLUT	E_{cm}^{ee}= 34.6 GeV
86.7 ±0.3 ±0.6			FERNANDEZ	85	MAC	E_{cm}^{ee}= 29 GeV

[28] Not independent of ADEVA 91F $\Gamma(h^-h^-h^+ \geq 0 \text{ neutrals } \geq 0 K_L^0 \nu_\tau)/\Gamma_{\text{total}}$ value.

[29] Not independent of AIHARA 87B $\Gamma(\mu^- \overline{\nu}_\mu \nu_\tau)/\Gamma_{\text{total}}$, $\Gamma(e^- \overline{\nu}_e \nu_\tau)/\Gamma_{\text{total}}$, and $/\Gamma_{\text{total}}$ values.

[30] Not independent of SCHMIDKE 86 value (also not independent of BURCHAT 87 value for $\Gamma(h^-h^-h^+ \geq 0 \text{ neutrals } \geq 0 K_L^0 \nu_\tau)/\Gamma_{\text{total}}$.

[31] Not independent of ALTHOFF 85 $\Gamma(\mu^-\overline{\nu}_\mu\nu_\tau)/\Gamma_{\rm total}$, $\Gamma(e^-\overline{\nu}_e\nu_\tau)/\Gamma_{\rm total}$, $/\Gamma_{\rm total}$, and $\Gamma(h^-h^-h^+ \geq 0$ neutrals $\geq 0K^0_L\nu_\tau)/\Gamma_{\rm total}$ values.

[32] Not independent of (1-prong + $0\pi^0$) and (1-prong + $\geq 1\pi^0$) values.

$\Gamma(\mu^-\overline{\nu}_\mu\nu_\tau)/\Gamma_{\rm total}$ Γ_3/Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

To minimize the effect of experiments with large systematic errors, we exclude experiments which together would contribute 5% of the weight in the average.

VALUE (%)				*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
17.36	**±0.05**	**OUR FIT**						
17.33	**±0.06**	**OUR AVERAGE**						
17.34	±0.09	±0.06	f&a	31.4k	ABBIENDI	03	OPAL	1990–1995 LEP runs
17.342	±0.110	±0.067	f&a	21.5k	[33] ACCIARRI	01F	L3	1991–1995 LEP runs
17.325	±0.095	±0.077	f&a	27.7k	ABREU	99X	DLPH	1991–1995 LEP runs
17.37	±0.08	±0.18	avg		[34] ANASTASSOV	97	CLEO	$E^{ee}_{\rm cm}$= 10.6 GeV
17.31	±0.11	±0.05	f&a	20.7k	BUSKULIC	96C	ALEP	1991–1993 LEP runs
17.35	±0.41	±0.37	f&a		DECAMP	92C	ALEP	1989–1990 LEP runs
• • • We do not use the following data for averages, fits, limits, etc. • • •								
17.02	±0.19	±0.24		6586	ABREU	95T	DLPH	Repl.. by ABREU 99X
17.36	±0.27			7941	AKERS	95I	OPAL	Repl. by ABBIENDI 03
17.6	±0.4	±0.4		2148	ADRIANI	93M	L3	Repl. by ACCIARRI 01F
17.4	±0.3	±0.5			[35] ALBRECHT	93G	ARG	$E^{ee}_{\rm cm}$= 9.4–10.6 GeV
17.7	±0.8	±0.4		568	BEHREND	90	CELL	$E^{ee}_{\rm cm}$= 35 GeV
17.4	±1.0			2197	ADEVA	88	MRKJ	$E^{ee}_{\rm cm}$= 14–16 GeV
17.7	±1.2	±0.7			AIHARA	87B	TPC	$E^{ee}_{\rm cm}$= 29 GeV
18.3	±0.9	±0.8			BURCHAT	87	MRK2	$E^{ee}_{\rm cm}$= 29 GeV
18.6	±0.8	±0.7		558	[36] BARTEL	86D	JADE	$E^{ee}_{\rm cm}$= 34.6 GeV
12.9	±1.7	$^{+0.7}_{-0.5}$			ALTHOFF	85	TASS	$E^{ee}_{\rm cm}$= 34.5 GeV
18.0	±0.9	±0.5		473	[36] ASH	85B	MAC	$E^{ee}_{\rm cm}$= 29 GeV
18.0	±1.0	±0.6			[37] BALTRUSAIT...	85	MRK3	$E^{ee}_{\rm cm}$= 3.77 GeV
19.4	±1.6	±1.7		153	BERGER	85	PLUT	$E^{ee}_{\rm cm}$= 34.6 GeV
17.6	±2.6	±2.1		47	BEHREND	83C	CELL	$E^{ee}_{\rm cm}$= 34 GeV
17.8	±2.0	±1.8			BERGER	81B	PLUT	$E^{ee}_{\rm cm}$= 9–32 GeV

[33] The correlation coefficient between this measurement and the ACCIARRI 01F measurement of $B(\tau^- \to e^-\overline{\nu}_e\nu_\tau)$ is 0.08.

[34] The correlation coefficients between this measurement and the ANASTASSOV 97 measurements of $B(e\overline{\nu}_e\nu_\tau)$, $B(\mu\overline{\nu}_\mu\nu_\tau)/B(e\overline{\nu}_e\nu_\tau)$, $B(h^-\nu_\tau)$, and $B(h^-\nu_\tau)/B(e\overline{\nu}_e\nu_\tau)$ are 0.50, 0.58, 0.50, and 0.08 respectively.

[35] Not independent of ALBRECHT 92D $\Gamma(\mu^-\overline{\nu}_\mu\nu_\tau)/\Gamma(e^-\overline{\nu}_e\nu_\tau)$ and ALBRECHT 93G $\Gamma(\mu^-\overline{\nu}_\mu\nu_\tau)\times\Gamma(e^-\overline{\nu}_e\nu_\tau)/\Gamma^2_{\rm total}$ values.

[36] Modified using B($e^-\overline{\nu}_e\nu_\tau$)/B("1 prong") and B("1 prong") ,= 0.855.

[37] Error correlated with BALTRUSAITIS 85 $e\nu\overline{\nu}$ value.

$\Gamma(\mu^-\overline{\nu}_\mu\nu_\tau\gamma)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
0.361 ± 0.016 ± 0.035		[38] BERGFELD	00	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.30 ± 0.04 ± 0.05	116	[39] ALEXANDER	96S	OPAL	1991–1994 LEP runs
0.23 ± 0.10	10	[40] WU	90	MRK2	E_{cm}^{ee}= 29 GeV

[38] BERGFELD 00 impose requirements on detected γ's corresponding to a τ-rest-frame energy cutoff $E_\gamma^* > 10$ MeV. For $E_\gamma^* > 20$ MeV, they quote $(3.04 \pm 0.14 \pm 0.30) \times 10^{-3}$.

[39] ALEXANDER 96S impose requirements on detected γ's corresponding to a τ-rest-frame energy cutoff E_γ >20 MeV.

[40] WU 90 reports $\Gamma(\mu^-\overline{\nu}_\mu\nu_\tau\gamma)/\Gamma(\mu^-\overline{\nu}_\mu\nu_\tau) = 0.013 \pm 0.006$, which is converted to $\Gamma(\mu^-\overline{\nu}_\mu\nu_\tau\gamma)/\Gamma_{\text{total}}$ using $\Gamma(\mu^-\overline{\nu}_\mu\nu_\tau\gamma)/\Gamma_{\text{total}} = 17.35\%$. Requirements on detected γ's correspond to a τ rest frame energy cutoff $E_\gamma > 37$ MeV.

$\Gamma(e^-\overline{\nu}_e\nu_\tau)/\Gamma_{\text{total}}$ Γ_5/Γ

To minimize the effect of experiments with large systematic errors, we exclude experiments which together would contribute 5% of the weight in the average.

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
17.84 ± 0.06 OUR FIT					
17.81 ± 0.06 OUR AVERAGE					
17.806 ± 0.104 ± 0.076	24.7k	[41] ACCIARRI	01F	L3	1991–1995 LEP runs
17.81 ± 0.09 ± 0.06	33.1k	ABBIENDI	99H	OPAL	1991–1995 LEP runs
17.877 ± 0.109 ± 0.110	23.3k	ABREU	99X	DLPH	1991–1995 LEP runs
17.76 ± 0.06 ± 0.17		[42] ANASTASSOV	97	CLEO	E_{cm}^{ee}= 10.6 GeV
17.79 ± 0.12 ± 0.06	20.6k	BUSKULIC	96C	ALEP	1991–1993 LEP runs
18.09 ± 0.45 ± 0.45		DECAMP	92C	ALEP	1989–1990 LEP runs
• • • We do not use the following data for averages, fits, limits, etc. • • •					
17.78 ± 0.10 ± 0.09	25.3k	ALEXANDER	96D	OPAL	Repl. by ABBIENDI 99H
17.51 ± 0.23 ± 0.31	5059	ABREU	95T	DLPH	Repl.. by ABREU 99X
17.9 ± 0.4 ± 0.4	2892	ADRIANI	93M	L3	Repl. by ACCIARRI 01F
17.5 ± 0.3 ± 0.5		[43] ALBRECHT	93G	ARG	E_{cm}^{ee}= 9.4–10.6 GeV
17.97 ± 0.14 ± 0.23	3970	AKERIB	92	CLEO	Repl. by ANASTASSOV 97
19.1 ± 0.4 ± 0.6	2960	[44] AMMAR	92	CLEO	E_{cm}^{ee}= 10.5–10.9 GeV
17.0 ± 0.5 ± 0.6	1.7k	ABACHI	90	HRS	E_{cm}^{ee}= 29 GeV
18.4 ± 0.8 ± 0.4	644	BEHREND	90	CELL	E_{cm}^{ee}= 35 GeV
16.3 ± 0.3 ± 3.2		JANSSEN	89	CBAL	E_{cm}^{ee}= 9.4–10.6 GeV
18.4 ± 1.2 ± 1.0		AIHARA	87B	TPC	E_{cm}^{ee}= 29 GeV
19.1 ± 0.8 ± 1.1		BURCHAT	87	MRK2	E_{cm}^{ee}= 29 GeV
16.8 ± 0.7 ± 0.9	515	[44] BARTEL	86D	JADE	E_{cm}^{ee}= 34.6 GeV
20.4 ± 3.0 $^{+1.4}_{-0.9}$		ALTHOFF	85	TASS	E_{cm}^{ee}= 34.5 GeV

17.8	±0.9	±0.6	390	[44] ASH	85B	MAC	E_{cm}^{ee}= 29 GeV
18.2	±0.7	±0.5		[45] BALTRUSAIT..	85	MRK3	E_{cm}^{ee}= 3.77 GeV
13.0	±1.9	±2.9		BERGER	85	PLUT	E_{cm}^{ee}= 34.6 GeV
18.3	±2.4	±1.9	60	BEHREND	83C	CELL	E_{cm}^{ee}= 34 GeV
16.0	±1.3		459	[46] BACINO	78B	DLCO	E_{cm}^{ee}= 3.1–7.4 GeV

[41] The correlation coefficient between this measurement and the ACCIARRI 01F measurement of B($\tau^- \to \mu^- \overline{\nu}_\mu \nu_\tau$) is 0.08.

[42] The correlation coefficients between this measurement and the ANASTASSOV 97 measurements of B($\mu\overline{\nu}_\mu\nu_\tau$), B($\mu\overline{\nu}_\mu\nu_\tau$)/B($e\overline{\nu}_e\nu_\tau$), B($h^-\nu_\tau$), and B($h^-\nu_\tau$)/B($e\overline{\nu}_e\nu_\tau$) are 0.50, −0.42, 0.48, and −0.39 respectively.

[43] Not independent of ALBRECHT 92D $\Gamma(\mu^-\overline{\nu}_\mu\nu_\tau)/\Gamma(e^-\overline{\nu}_e\nu_\tau)$ and ALBRECHT 93G $\Gamma(\mu^-\overline{\nu}_\mu\nu_\tau)\times\Gamma(e^-\overline{\nu}_e\nu_\tau)/\Gamma^2_{\text{total}}$ values.

[44] Modified using B($e^-\overline{\nu}_e\nu_\tau$)/B("1 prong") and B("1 prong") ,= 0.855.

[45] Error correlated with BALTRUSAITIS 85 $\Gamma(\mu^-\overline{\nu}_\mu\nu_\tau)/\Gamma_{\text{total}}$.

[46] BACINO 78B value comes from fit to events with $e^\pm$ and one other nonelectron charged prong.

$\Gamma(e^-\overline{\nu}_e\nu_\tau\gamma)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (%)	DOCUMENT ID		TECN	COMMENT
1.75±0.06±0.17	[47] BERGFELD	00	CLEO	E_{cm}^{ee}= 10.6 GeV

[47] BERGFELD 00 impose requirements on detected γ's corresponding to a τ-rest-frame energy cutoff $E^*_\gamma > 10$ MeV.

$\Gamma(\mu^-\overline{\nu}_\mu\nu_\tau)/\Gamma(e^-\overline{\nu}_e\nu_\tau)$ Γ_3/Γ_5

Standard Model prediction including mass effects is 0.9726.

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE		DOCUMENT ID		TECN	COMMENT
0.973 ±0.004 OUR FIT					
0.978 ±0.011 OUR AVERAGE					
0.9777±0.0063±0.0087	f&a	[48] ANASTASSOV 97		CLEO	E_{cm}^{ee}= 10.6 GeV
0.997 ±0.035 ±0.040	f&a	ALBRECHT	92D	ARG	E_{cm}^{ee}= 9.4–10.6 GeV

[48] The correlation coefficients between this measurement and the ANASTASSOV 97 measurements of B($\mu\overline{\nu}_\mu\nu_\tau$), B($e\overline{\nu}_e\nu_\tau$), B($h^-\nu_\tau$), and B($h^-\nu_\tau$)/B($e\overline{\nu}_e\nu_\tau$) are 0.58, −0.42, 0.07, and 0.45 respectively.

$\Gamma(h^- \geq 0K^0_L\,\nu_\tau)/\Gamma_{\text{total}}$ Γ_7/Γ

$\Gamma_7/\Gamma = (\Gamma_9+\Gamma_{10}+\frac{1}{2}\Gamma_{33}+\frac{1}{2}\Gamma_{35}+\Gamma_{45})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID		TECN	COMMENT
12.30±0.10 OUR FIT Error includes scale factor of 1.4.						
12.44±0.14 OUR AVERAGE						
12.44±0.11±0.11	f&a	15k	[49] BUSKULIC	96	ALEP	1991–1993 LEP run
12.47±0.26±0.43	f&a	2967	[50] ACCIARRI	95	L3	1992 LEP run
12.4 ±0.7 ±0.7	f&a	283	[51] ABREU	92N	DLPH	1990 LEP run

12.98 ± 0.44 ± 0.33	f&a		[52] DECAMP	92C	ALEP	1989–1990 LEP runs
12.1 ± 0.7 ± 0.5	f&a	309	ALEXANDER	91D	OPAL	1990 LEP run
11.3 ± 0.5 ± 0.8	avg	798	[53] FORD	87	MAC	E_{cm}^{ee} = 29 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

11.7 ± 0.6 ± 0.8		[54] ALBRECHT	92D	ARG	E_{cm}^{ee} = 9.4–10.6 GeV
12.3 ± 0.9 ± 0.5	1338	BEHREND	90	CELL	E_{cm}^{ee} = 35 GeV
11.1 ± 1.1 ± 1.4		[55] BURCHAT	87	MRK2	E_{cm}^{ee} = 29 GeV
12.3 ± 0.6 ± 1.1	328	[56] BARTEL	86D	JADE	E_{cm}^{ee} = 34.6 GeV
13.0 ± 2.0 ± 4.0		BERGER	85	PLUT	E_{cm}^{ee} = 34.6 GeV
11.2 ± 1.7 ± 1.2	34	[57] BEHREND	83C	CELL	E_{cm}^{ee} = 34 GeV

[49] BUSKULIC 96 quote 11.78 ± 0.11 ± 0.13 We add 0.66 to undo their correction for unseen K_L^0 and modify the systematic error accordingly.

[50] ACCIARRI 95 with 0.65% added to remove their correction for $\pi^- K_L^0$ backgrounds.

[51] ABREU 92N with 0.5% added to remove their correction for $K^*(892)^-$ backgrounds.

[52] DECAMP 92C quote $B(h^- \geq 0 K_L^0 \geq 0 (K_S^0 \to \pi^+ \pi^-) \nu_\tau) = 13.32 \pm 0.44 \pm 0.33$. We subtract 0.35 to correct for their inclusion of the K_S^0 decays.

[53] FORD 87 result for $B(\pi^- \nu_\tau)$ with 0.67% added to remove their K^- correction and adjusted for 1992 B("1 prong").

[54] Not independent of ALBRECHT 92D $\Gamma(\mu^- \overline{\nu}_\mu \nu_\tau)/\Gamma(e^- \overline{\nu}_e \nu_\tau)$, $\Gamma(\mu^- \overline{\nu}_\mu \nu_\tau) \times \Gamma(e^- \overline{\nu}_e \nu_\tau)$, and $\Gamma(h^- \geq 0 K_L^0 \nu_\tau)/\Gamma(e^- \overline{\nu}_e \nu_\tau)$ values.

[55] BURCHAT 87 with 1.1% added to remove their correction for K^- and $K^*(892)^-$ backgrounds.

[56] BARTEL 86D result for $B(\pi^- \nu_\tau)$ with 0.59% added to remove their K^- correction and adjusted for 1992 B("1 prong").

[57] BEHREND 83C quote $B(\pi^- \nu_\tau) = 9.9 \pm 1.7 \pm 1.3$ after subtracting 1.3 ± 0.5 to correct for $B(K^- \nu_\tau)$.

$\Gamma(h^- \nu_\tau)/\Gamma_{total}$ $\Gamma_8/\Gamma = (\Gamma_9+\Gamma_{10})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		*DOCUMENT ID*	*TECN*	*COMMENT*
11.75 ± 0.10 OUR FIT Error includes scale factor of 1.4.				
11.65 ± 0.21 OUR AVERAGE Error includes scale factor of 1.9.				
11.98 ± 0.13 ± 0.16	f&a	ACKERSTAFF 98M	OPAL	1991–1995 LEP runs
11.52 ± 0.05 ± 0.12	f&a	[58] ANASTASSOV 97	CLEO	E_{cm}^{ee} = 10.6 GeV

[58] The correlation coefficients between this measurement and the ANASTASSOV 97 measurements of $B(\mu \overline{\nu}_\mu \nu_\tau)$, $B(e \overline{\nu}_e \nu_\tau)$, $B(\mu \overline{\nu}_\mu \nu_\tau)/B(e \overline{\nu}_e \nu_\tau)$, and $B(h^- \nu_\tau)/B(e \overline{\nu}_e \nu_\tau)$ are 0.50, 0.48, 0.07, and 0.63 respectively.

$\Gamma(h^- \nu_\tau)/\Gamma(e^- \overline{\nu}_e \nu_\tau)$ $\Gamma_8/\Gamma_5 = (\Gamma_9+\Gamma_{10})/\Gamma_5$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE		*DOCUMENT ID*	*TECN*	*COMMENT*
0.659 ± 0.006 OUR FIT Error includes scale factor of 1.3.				
0.6484 ± 0.0041 ± 0.0060 avg		[59] ANASTASSOV 97	CLEO	E_{cm}^{ee} = 10.6 GeV

[59] The correlation coefficients between this measurement and the ANASTASSOV 97 measurements of $B(\mu\overline{\nu}_\mu\nu_\tau)$, $B(e\overline{\nu}_e\nu_\tau)$, $B(\mu\overline{\nu}_\mu\nu_\tau)/B(e\overline{\nu}_e\nu_\tau)$, and $B(h^-\nu_\tau)$ are 0.08, −0.39, 0.45, and 0.63 respectively.

$\Gamma(\pi^-\nu_\tau)/\Gamma_{total}$ Γ_9/Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID		TECN	COMMENT
11.06±0.11 OUR FIT Error includes scale factor of 1.4.						
11.07±0.18 OUR AVERAGE						
11.06±0.11±0.14	avg		[60] BUSKULIC	96	ALEP	LEP 1991–1993 data
11.7 ±0.4 ±1.8	f&a	1138	BLOCKER	82D	MRK2	E_{cm}^{ee}= 3.5–6.7 GeV

[60] Not independent of BUSKULIC 96 $B(h^-\nu_\tau)$ and $B(K^-\nu_\tau)$ values.

$\Gamma(K^-\nu_\tau)/\Gamma_{total}$ Γ_{10}/Γ

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
0.686±0.023 OUR FIT					
0.685±0.023 OUR AVERAGE					
0.658±0.027±0.029		[61] ABBIENDI	01J	OPAL	1990–1995 LEP runs
0.696±0.025±0.014	2032	BARATE	99K	ALEP	1991–1995 LEP runs
0.85 ±0.18	27	ABREU	94K	DLPH	LEP 1992 Z data
0.66 ±0.07 ±0.09	99	BATTLE	94	CLEO	$E_{cm}^{ee} \approx$ 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.72 ±0.04 ±0.04	728	BUSKULIC	96	ALEP	Repl. by BARATE 99K
0.59 ±0.18	16	MILLS	84	DLCO	E_{cm}^{ee}= 29 GeV
1.3 ±0.5	15	BLOCKER	82B	MRK2	E_{cm}^{ee}= 3.9–6.7 GeV

[61] The correlation coefficient between this measurement and the ABBIENDI 01J $B(\tau^- \to K^- \geq 0\pi^0 \geq 0K^0 \geq 0\gamma\ \nu_\tau)$ is 0.60.

$\Gamma(h^- \geq 1 \text{ neutrals}\nu_\tau)/\Gamma_{total}$ Γ_{11}/Γ

$\Gamma_{11}/\Gamma = (\Gamma_{13}+\Gamma_{15}+\Gamma_{19}+\Gamma_{22}+\Gamma_{25}+\Gamma_{26}+\Gamma_{28}+0.157\Gamma_{33}+0.157\Gamma_{35}+0.157\Gamma_{38}+0.157\Gamma_{40}+0.0985\Gamma_{45}+0.708\Gamma_{117}+0.715\Gamma_{119}+0.09\Gamma_{137}+0.09\Gamma_{138})/\Gamma$

VALUE (%)	DOCUMENT ID		TECN	COMMENT
36.92±0.14 OUR FIT Error includes scale factor of 1.1.				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
36.14±0.33±0.58	[62] AKERS	94E	OPAL	1991–1992 LEP runs
38.4 ±1.2 ±1.0	[63] BURCHAT	87	MRK2	E_{cm}^{ee}= 29 GeV
42.7 ±2.0 ±2.9	BERGER	85	PLUT	E_{cm}^{ee}= 34.6 GeV

[62] Not independent of ACKERSTAFF 98M $B(h^-\pi^0\nu_\tau)$ and $B(h^- \geq 2\pi^0\nu_\tau)$ values.

[63] BURCHAT 87 quote for $B(\pi^\pm \geq 1 \text{ neutral}\nu_\tau) = 0.378 \pm 0.012 \pm 0.010$. We add 0.006 to account for contribution from $(K^{*-}\nu_\tau)$ which they fixed at BR = 0.013.

$\Gamma(h^-\pi^0\nu_\tau)/\Gamma_{total}$ $\Gamma_{12}/\Gamma=(\Gamma_{13}+\Gamma_{15})/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
25.86±0.13 OUR FIT Error includes scale factor of 1.1.					
25.76±0.15 OUR AVERAGE					
25.89±0.17±0.29		ACKERSTAFF	98M	OPAL	1991–1995 LEP runs
25.76±0.15±0.13	31k	BUSKULIC	96	ALEP	LEP 1991–1993 data
25.05±0.35±0.50	6613	ACCIARRI	95	L3	1992 LEP run
25.87±0.12±0.42	51k	[64] ARTUSO	94	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
25.98±0.36±0.52		[65] AKERS	94E	OPAL	Repl. by ACKERSTAFF 98M
22.9 ±0.8 ±1.3	283	[66] ABREU	92N	DLPH	E_{cm}^{ee}= 88.2–94.2 GeV
23.1 ±0.4 ±0.9	1249	[67] ALBRECHT	92Q	ARG	E_{cm}^{ee}= 10 GeV
25.02±0.64±0.88	1849	DECAMP	92C	ALEP	1989–1990 LEP runs
22.0 ±0.8 ±1.9	779	ANTREASYAN	91	CBAL	E_{cm}^{ee}= 9.4–10.6 GeV
22.6 ±1.5 ±0.7	1101	BEHREND	90	CELL	E_{cm}^{ee}= 35 GeV
23.1 ±1.9 ±1.6		BEHREND	84	CELL	E_{cm}^{ee}= 14,22 GeV

[64] ARTUSO 94 reports the combined result from three independent methods, one of which (23% of the $\tau^-\to h^-\pi^0\nu_\tau$) is normalized to the inclusive one-prong branching fraction, taken as 0.854 ± 0.004. Renormalization to the present value causes negligible change.

[65] AKERS 94E quote $(26.25\pm0.36\pm0.52)\times10^{-2}$; we subtract 0.27% from their number to correct for $\tau^-\to h^-K_L^0\nu_\tau$.

[66] ABREU 92N with 0.5% added to remove their correction for $K^*(892)^-$ backgrounds.

[67] ALBRECHT 92Q with 0.5% added to remove their correction for $\tau^-\to K^*(892)^-\nu_\tau$ background.

$\Gamma(\pi^-\pi^0\nu_\tau)/\Gamma_{total}$ Γ_{13}/Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID		TECN	COMMENT
25.41±0.14 OUR FIT Error includes scale factor of 1.1.						
25.31±0.18 OUR AVERAGE						
25.30±0.15±0.13	avg		[68] BUSKULIC	96	ALEP	LEP 1991–1993 data
25.36±0.44	avg		[69] ARTUSO	94	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •						
21.5 ±0.4 ±1.9		4400	[70,71] ALBRECHT	88L	ARG	E_{cm}^{ee}= 10 GeV
23.0 ±1.3 ±1.7		582	ADLER	87B	MRK3	E_{cm}^{ee}= 3.77 GeV
25.8 ±1.7 ±2.5			[72] BURCHAT	87	MRK2	E_{cm}^{ee}= 29 GeV
22.3 ±0.6 ±1.4		629	[71] YELTON	86	MRK2	E_{cm}^{ee}= 29 GeV

[68] Not independent of BUSKULIC 96 B($h^-\pi^0\nu_\tau$) and B($K^-\pi^0\nu_\tau$) values.

[69] Not independent of ARTUSO 94 B($h^-\pi^0\nu_\tau$) and BATTLE 94 B($K^-\pi^0\nu_\tau$) values.

[70] The authors divide by ($\Gamma_3+\Gamma_5+\Gamma_9+\Gamma_{10}$)/Γ = 0.467 to obtain this result.

[71] Experiment had no hadron identification. Kaon corrections were made, but insufficient information is given to permit their removal.

[72] BURCHAT 87 value is not independent of YELTON 86 value. Nonresonant decays included.

$\Gamma(\pi^-\pi^0 \text{ non-}\rho(770)\nu_\tau)/\Gamma_{total}$ Γ_{14}/Γ

VALUE (%)	DOCUMENT ID		TECN	COMMENT
0.3 ±0.1 ±0.3	[73] BEHREND	84	CELL	E_{cm}^{ee}= 14,22 GeV

[73] BEHREND 84 assume a flat nonresonant mass distribution down to the $\rho(770)$ mass, using events with mass above 1300 to set the level.

$\Gamma(K^-\pi^0\nu_\tau)/\Gamma_{total}$ Γ_{15}/Γ

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
0.450±0.030 OUR FIT					
0.449±0.034 OUR AVERAGE					
0.444±0.026±0.024	923	BARATE	99K	ALEP	1991–1995 LEP runs
0.51 ±0.10 ±0.07	37	BATTLE	94	CLEO	$E_{cm}^{ee} \approx$ 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.52 ±0.04 ±0.05	395	BUSKULIC	96	ALEP	Repl. by BARATE 99K

$\Gamma(h^- \geq 2\pi^0\nu_\tau)/\Gamma_{total}$ Γ_{16}/Γ

$\Gamma_{16}/\Gamma = (\Gamma_{19}+\Gamma_{22}+\Gamma_{25}+\Gamma_{26}+\Gamma_{28}+0.157\Gamma_{33}+0.157\Gamma_{35}+0.157\Gamma_{38}+0.157\Gamma_{40}+0.0985\Gamma_{45}+0.319\Gamma_{117}+0.322\Gamma_{119})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID		TECN	COMMENT
10.76±0.15 OUR FIT Error includes scale factor of 1.1.						
10.0 ±0.4 OUR AVERAGE						
9.91±0.31±0.27	f&a		ACKERSTAFF	98M	OPAL	1991–1995 LEP runs
12.0 ±1.4 ±2.5	f&a		[74] BURCHAT	87	MRK2	E_{cm}^{ee}= 29 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •						
9.89±0.34±0.55			[75] AKERS	94E	OPAL	Repl. by ACKERSTAFF 98M
14.0 ±1.2 ±0.6		938	[76] BEHREND	90	CELL	E_{cm}^{ee}= 35 GeV
13.9 ±2.0 $^{+1.9}_{-2.2}$			[77] AIHARA	86E	TPC	E_{cm}^{ee}= 29 GeV

[74] Error correlated with BURCHAT 87 $\Gamma(\rho^-\nu_e)/\Gamma$(total) value.

[75] AKERS 94E not independent of AKERS 94E B($h^- \geq 1\pi^0\nu_\tau$) and B($h^-\pi^0\nu_\tau$) measurements.

[76] No independent of BEHREND 90 $\Gamma(h^-2\pi^0\nu_\tau$(exp. K^0)) and $\Gamma(h^- \geq 3\pi^0\nu_\tau)$.

[77] AIHARA 86E (TPC) quote B($2\pi^0\pi^-\nu_\tau$) + 1.6B($3\pi^0\pi^-\nu_\tau$) + 1.1B($\pi^0\eta\pi^-\nu_\tau$).

$\Gamma(h^-2\pi^0\nu_\tau)/\Gamma_{total}$ Γ_{17}/Γ

$\Gamma_{17}/\Gamma = (\Gamma_{19}+\Gamma_{22}+0.157\Gamma_{33}+0.157\Gamma_{35})/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
9.39±0.14 OUR FIT Error includes scale factor of 1.1.					
9.48±0.13±0.10	12k	[78] BUSKULIC	96	ALEP	LEP 1991–1993 data

[78] BUSKULIC 96 quote 9.29 ± 0.13 ± 0.10. We add 0.19 to undo their correction for $\tau^- \to h^- K^0\nu_\tau$.

$\Gamma(h^- 2\pi^0 \nu_\tau (\text{ex.} K^0))/\Gamma_{\text{total}}$ Γ_{18}/Γ

$\Gamma_{18}/\Gamma = (\Gamma_{19}+\Gamma_{22})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. f&a marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID		TECN	COMMENT
9.23±0.14 OUR FIT	Error includes scale factor of 1.1.					
9.08±0.34 OUR AVERAGE						
8.88±0.37±0.42	f&a	1060	ACCIARRI	95	L3	1992 LEP run
8.96±0.16±0.44	avg		[79] PROCARIO	93	CLEO	$E_{\text{cm}}^{ee} \approx$ 10.6 GeV
10.38±0.66±0.82	f&a	809	[80] DECAMP	92C	ALEP	1989–1990 LEP runs
• • • We do not use the following data for averages, fits, limits, etc. • • •						
5.7 ±0.5 $^{+1.7}_{-1.0}$		133	[81] ANTREASYAN	91	CBAL	E_{cm}^{ee}= 9.4–10.6 GeV
10.0 ±1.5 ±1.1		333	[82] BEHREND	90	CELL	E_{cm}^{ee}= 35 GeV
8.7 ±0.4 ±1.1		815	[83] BAND	87	MAC	E_{cm}^{ee}= 29 GeV
6.2 ±0.6 ±1.2			[84] GAN	87	MRK2	E_{cm}^{ee}= 29 GeV
6.0 ±3.0 ±1.8			BEHREND	84	CELL	E_{cm}^{ee}= 14,22 GeV

[79] PROCARIO 93 entry is obtained from $B(h^- 2\pi^0 \nu_\tau)/B(h^- \pi^0 \nu_\tau)$ using ARTUSO 94 result for $B(h^- \pi^0 \nu_\tau)$.

[80] We subtract 0.0015 to account for $\tau^- \to K^*(892)^- \nu_\tau$ contribution.

[81] ANTREASYAN 91 subtract 0.001 to account for the $\tau^- \to K^*(892)^- \nu_\tau$ contribution.

[82] BEHREND 90 subtract 0.002 to account for the $\tau^- \to K^*(892)^- \nu_\tau$ contribution.

[83] BAND 87 assume $B(\pi^- 3\pi^0 \nu_\tau) = 0.01$ and $B(\pi^- \pi^0 \eta \nu_\tau) = 0.005$.

[84] GAN 87 analysis use photon multiplicity distribution.

$\Gamma(h^- 2\pi^0 \nu_\tau (\text{ex.} K^0))/\Gamma(h^- \pi^0 \nu_\tau)$ Γ_{18}/Γ_{12}

$\Gamma_{18}/\Gamma_{12} = (\Gamma_{19}+\Gamma_{22})/(\Gamma_{13}+\Gamma_{15})$

VALUE	DOCUMENT ID		TECN	COMMENT
0.357±0.006 OUR FIT	Error includes scale factor of 1.1.			
0.342±0.006±0.016	[85] PROCARIO	93	CLEO	$E_{\text{cm}}^{ee} \approx$ 10.6 GeV

[85] PROCARIO 93 quote 0.345 ± 0.006 ± 0.016 after correction for 2 kaon backgrounds assuming $B(K^{*-} \nu_\tau)$=1.42 ± 0.18% and $B(h^- K^0 \pi^0 \nu_\tau)$=0.48 ± 0.48%. We multiply by 0.990 ± 0.010 to remove these corrections to $B(h^- \pi^0 \nu_\tau)$.

$\Gamma(\pi^- 2\pi^0 \nu_\tau (\text{ex.} K^0))/\Gamma_{\text{total}}$ Γ_{19}/Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		DOCUMENT ID		TECN	COMMENT
9.17±0.14 OUR FIT	Error includes scale factor of 1.1.				
9.21±0.13±0.11	**avg**	[86] BUSKULIC	96	ALEP	LEP 1991–1993 data

[86] Not independent of BUSKULIC 96 $B(h^- 2\pi^0 \nu_\tau (\text{ex. } K^0))$ and $B(K^- 2\pi^0 \nu_\tau (\text{ex. } K^0))$ values.

$\Gamma(\pi^- 2\pi^0 \nu_\tau (\text{ex.} K^0), \text{scalar})/\Gamma(\pi^- 2\pi^0 \nu_\tau (\text{ex.} K^0))$ Γ_{20}/Γ_{19}

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.094	95	[87] BROWDER	00	CLEO	4.7 fb^{-1} E_{cm}^{ee}= 10.6 GeV

[87] Model-independent limit from structure function analysis on contribution to B($\tau^- \to \pi^- 2\pi^0 \nu_\tau$(ex. K^0)) from scalars.

$\Gamma(\pi^- 2\pi^0 \nu_\tau (\text{ex.} K^0), \text{vector})/\Gamma(\pi^- 2\pi^0 \nu_\tau (\text{ex.} K^0))$ Γ_{21}/Γ_{19}

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.073	95	[88] BROWDER	00	CLEO	4.7 fb^{-1} E_{cm}^{ee}= 10.6 GeV

[88] Model-independent limit from structure function analysis on contribution to B($\tau^- \to \pi^- 2\pi^0 \nu_\tau$(ex. K^0)) from vectors.

$\Gamma(K^- 2\pi^0 \nu_\tau (\text{ex.} K^0))/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
0.058±0.023 OUR FIT					
0.058±0.024 OUR AVERAGE					
0.056±0.020±0.015	131	BARATE	99K	ALEP	1991–1995 LEP runs
0.09 ±0.10 ±0.03	3	[89] BATTLE	94	CLEO	$E_{cm}^{ee} \approx$ 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.08 ±0.02 ±0.02	59	BUSKULIC	96	ALEP	Repl. by BARATE 99K

[89] BATTLE 94 quote 0.14 ± 0.10 ± 0.03 or < 0.3% at 90% CL. We subtract (0.05 ± 0.02)% to account for $\tau^- \to K^-(K^0 \to \pi^0\pi^0)\nu_\tau$ background.

$\Gamma(h^- \geq 3\pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{23}/Γ

$\Gamma_{23}/\Gamma = (\Gamma_{25}+\Gamma_{26}+\Gamma_{28}+0.157\Gamma_{38}+0.157\Gamma_{40}+0.0985\Gamma_{45}+0.319\Gamma_{117}+0.322\Gamma_{119})/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
1.37±0.11 OUR FIT Error includes scale factor of 1.1.					
1.53±0.40±0.46	186	DECAMP	92C	ALEP	1989–1990 LEP runs
• • • We do not use the following data for averages, fits, limits, etc. • • •					
3.2 ±1.0 ±1.0		BEHREND	90	CELL	E_{cm}^{ee}= 35 GeV

$\Gamma(h^- 3\pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{24}/Γ

$\Gamma_{24}/\Gamma = (\Gamma_{25}+\Gamma_{26}+0.157\Gamma_{38}+0.157\Gamma_{40}+0.322\Gamma_{119})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID		TECN	COMMENT
1.21±0.10 OUR FIT						
1.22±0.10 OUR AVERAGE						
1.24±0.09±0.11	f&a	2.3k	[90] BUSKULIC	96	ALEP	LEP 1991–1993 data
1.70±0.24±0.38	f&a	293	ACCIARRI	95	L3	1992 LEP run
1.15±0.08±0.13	avg		[91] PROCARIO	93	CLEO	$E_{cm}^{ee} \approx$ 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$0.0\ ^{+1.4}_{-0.1}$		$^{+1.1}_{-0.1}$	[92] GAN	87	MRK2	E_{cm}^{ee}= 29 GeV

[90] BUSKULIC 96 quote $B(h^-3\pi^0\nu_\tau(\text{ex. } K^0)) = 1.17 \pm 0.09 \pm 0.11$. We add 0.07 to remove their correction for K^0 backgrounds.

[91] PROCARIO 93 entry is obtained from $B(h^-3\pi^0\nu_\tau)/B(h^-\pi^0\nu_\tau)$ using ARTUSO 94 result for $B(h^-\pi^0\nu_\tau)$.

[92] Highly correlated with GAN 87 $\Gamma(\eta\pi^-\pi^0\nu_\tau)/\Gamma_{\text{total}}$ value. Authors quote $B(\pi^\pm 3\pi^0\nu_\tau) + 0.67B(\pi^\pm\eta\pi^0\nu_\tau) = 0.047 \pm 0.010 \pm 0.011$.

$\Gamma(h^-3\pi^0\nu_\tau)/\Gamma(h^-\pi^0\nu_\tau)$ Γ_{24}/Γ_{12}

$\Gamma_{24}/\Gamma_{12} = (\Gamma_{25}+\Gamma_{26}+0.157\Gamma_{38}+0.157\Gamma_{40}+0.322\Gamma_{119})/(\Gamma_{13}+\Gamma_{15})$

VALUE	DOCUMENT ID	TECN	COMMENT
0.047±0.004 OUR FIT	Error includes scale factor of 1.1.		
0.044±0.003±0.005	[93] PROCARIO 93	CLEO	$E_{\text{cm}}^{ee} \approx$ 10.6 GeV

[93] PROCARIO 93 quote 0.041 ± 0.003 ± 0.005 after correction for 2 kaon backgrounds assuming $B(K^{*-}\nu_\tau)$=1.42 ± 0.18% and $B(h^-K^0\pi^0\nu_\tau)$=0.48 ± 0.48%. We add 0.003 ± 0.003 and multiply the sum by 0.990 ± 0.010 to remove these corrections.

$\Gamma(\pi^-3\pi^0\nu_\tau(\text{ex.}K^0))/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (%)	DOCUMENT ID
1.08±0.10 OUR FIT	

$\Gamma(K^-3\pi^0\nu_\tau(\text{ex.}K^0,\eta))/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$0.037^{+0.022}_{-0.020}$ OUR FIT				
0.037±0.021±0.011	22	BARATE 99K	ALEP	1991–1995 LEP runs
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.05 ±0.13		[94] BUSKULIC 94E	ALEP	Repl. by BARATE 99K

[94] BUSKULIC 94E quote $B(K^- \geq 0\pi^0 \geq 0K^0\nu_\tau) - [B(K^-\nu_\tau) + B(K^-\pi^0\nu_\tau) + B(K^-K^0\nu_\tau) + B(K^-\pi^0\pi^0\nu_\tau) + B(K^-\pi^0K^0\nu_\tau)] = 0.05 \pm 0.13\%$ accounting for common systematic errors in BUSKULIC 94E and BUSKULIC 94F measurements of these modes. We assume $B(K^- \geq 2K^0\nu_\tau)$ and $B(K^- \geq 4\pi^0\nu_\tau)$ are negligible.

$\Gamma(h^-4\pi^0\nu_\tau(\text{ex.}K^0))/\Gamma_{\text{total}}$ Γ_{27}/Γ

$\Gamma_{27}/\Gamma = (\Gamma_{28}+0.319\Gamma_{117})/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.16±0.06 OUR FIT				
0.16±0.06 OUR AVERAGE				
0.16±0.04±0.09	232	[95] BUSKULIC 96	ALEP	LEP 1991–1993 data
0.16±0.05±0.05		[96] PROCARIO 93	CLEO	$E_{\text{cm}}^{ee} \approx$ 10.6 GeV

[95] BUSKULIC 96 quote result for $\tau^- \to h^- \geq 4\pi^0\nu_\tau$. We assume $B(h^- \geq 5\pi^0\nu_\tau)$ is negligible.

[96] PROCARIO 93 quotes $B(h^-4\pi^0\nu_\tau)/B(h^-\pi^0\nu_\tau) = 0.006 \pm 0.002 \pm 0.002$. We multiply by the ARTUSO 94 result for $B(h^-\pi^0\nu_\tau)$ to obtain $B(h^-4\pi^0\nu_\tau)$. PROCARIO 93 assume $B(h^- \geq 5\ \pi^0\nu_\tau)$ is small and do not correct for it.

$\Gamma(h^- 4\pi^0 \nu_\tau (\text{ex.} K^0, \eta))/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (%)	DOCUMENT ID
$0.10^{+0.06}_{-0.05}$ **OUR FIT**	

$\Gamma(K^- \geq 0\pi^0 \geq 0K^0 \geq 0\gamma\, \nu_\tau)/\Gamma_{\text{total}}$ Γ_{29}/Γ

$\Gamma_{29}/\Gamma = (\Gamma_{10}+\Gamma_{15}+\Gamma_{22}+\Gamma_{26}+\Gamma_{35}+\Gamma_{40}+0.715\Gamma_{119})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)			EVTS	DOCUMENT ID		TECN	COMMENT
1.56 ±0.04 OUR FIT							
1.53 ±0.04 OUR AVERAGE							
1.528±0.039±0.040		f&a		[97] ABBIENDI	01J	OPAL	1990–1995 LEP runs
1.520±0.040±0.041		avg	4006	[98] BARATE	99K	ALEP	1991–1995 LEP runs
1.54 ±0.24		f&a		ABREU	94K	DLPH	LEP 1992 Z data
1.70 ±0.12	±0.19	f&a	202	[99] BATTLE	94	CLEO	$E_{\text{cm}}^{ee} \approx$ 10.6 GeV
1.6 ±0.4	±0.2	f&a	35	AIHARA	87B	TPC	E_{cm}^{ee}= 29 GeV
1.71 ±0.29		f&a	53	MILLS	84	DLCO	E_{cm}^{ee}= 29 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •							
1.70 ±0.05	±0.06		1610	[100] BUSKULIC	96	ALEP	Repl. by BARATE 99K

[97] The correlation coefficient between this measurement and the ABBIENDI 01J B($\tau^- \to K^- \nu_\tau$) is 0.60.

[98] Not independent of BARATE 99K B($K^- \nu_\tau$), B($K^- \pi^0 \nu_\tau$), B($K^- 2\pi^0 \nu_\tau$(ex. K^0)), B($K^- 3\pi^0 \nu_\tau$(ex. K^0)), B($K^- K^0 \nu_\tau$), and B($K^- K^0 \pi^0 \nu_\tau$) values.

[99] BATTLE 94 quote 1.60 ± 0.12 ± 0.19. We add 0.10 ± 0.02 to correct for their rejection of $K_S^0 \to \pi^+ \pi^-$ decays.

[100] Not independent of BUSKULIC 96 B($K^- \nu_\tau$), B($K^- \pi^0 \nu_\tau$), B($K^- 2\pi^0 \nu_\tau$), B($K^- K^0 \nu_\tau$), and B($K^- K^0 \pi^0 \nu_\tau$) values.

$\Gamma(K^- \geq 1(\pi^0 \text{ or } K^0 \text{ or } \gamma)\, \nu_\tau)/\Gamma_{\text{total}}$ Γ_{30}/Γ

$\Gamma_{30}/\Gamma = (\Gamma_{15}+\Gamma_{22}+\Gamma_{26}+\Gamma_{35}+\Gamma_{40}+0.715\Gamma_{119})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)			EVTS	DOCUMENT ID		TECN	COMMENT
0.874±0.035 OUR FIT							
0.86 ±0.05 OUR AVERAGE							
0.869±0.031±0.034		avg		[101] ABBIENDI	01J	OPAL	1990–1995 LEP runs
0.69 ±0.25		avg		[102] ABREU	94K	DLPH	LEP 1992 Z data
1.2 ±0.5	$^{+0.2}_{-0.4}$	f&a	9	AIHARA	87B	TPC	E_{cm}^{ee}= 29 GeV

[101] Not independent of ABBIENDI 01J B($\tau^- \to K^- \nu_\tau$) and B($\tau^- \to K^- \geq 0\pi^0 \geq 0K^0 \geq 0\gamma\, \nu_\tau$) values.

[102] Not independent of ABREU 94K B($K^- \nu_\tau$) and B($K^- \geq 0$ neutrals ν_τ) measurements.

$\Gamma(K^0_S(\text{particles})^- \nu_\tau)/\Gamma_{\text{total}}$ Γ_{31}/Γ

$\Gamma_{31}/\Gamma = (\frac{1}{2}\Gamma_{33}+\frac{1}{2}\Gamma_{35}+\frac{1}{2}\Gamma_{38}+\frac{1}{2}\Gamma_{40}+\Gamma_{45}+\Gamma_{46})/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.92 ±0.04 OUR FIT	Error includes scale factor of 1.1.			
0.97 ±0.07 OUR AVERAGE				
0.970±0.058±0.062	929	BARATE	98E ALEP	1991–1995 LEP runs
0.97 ±0.09 ±0.06	141	AKERS	94G OPAL	E^{ee}_{cm}= 88–94 GeV

$\Gamma(h^- \overline{K}^0 \nu_\tau)/\Gamma_{\text{total}}$ $\Gamma_{32}/\Gamma = (\Gamma_{33}+\Gamma_{35})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID		TECN	COMMENT
1.05 ±0.04 OUR FIT	Error includes scale factor of 1.1.					
0.90 ±0.07 OUR AVERAGE						
1.01 ±0.11 ±0.07	avg	555	[103] BARATE	98E	ALEP	1991–1995 LEP runs
0.855±0.036±0.073	f&a	1242	COAN	96	CLEO	$E^{ee}_{\text{cm}} \approx$ 10.6 GeV

[103] Not independent of BARATE 98E B($\tau^- \to \pi^- \overline{K}^0 \nu_\tau$) and B($\tau^- \to K^- K^0 \nu_\tau$) values.

$\Gamma(\pi^- \overline{K}^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{33}/Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID		TECN	COMMENT
0.89 ±0.04 OUR FIT	Error includes scale factor of 1.1.					
0.88 ±0.05 OUR AVERAGE	Error includes scale factor of 1.2.					
0.933±0.068±0.049	f&a	377	ABBIENDI	00C	OPAL	1991–1995 LEP runs
0.928±0.045±0.034	f&a	937	[104] BARATE	99K	ALEP	1991–1995 LEP runs
0.855±0.117±0.066	avg	509	[105] BARATE	98E	ALEP	1991–1995 LEP runs
0.704±0.041±0.072	avg		[106] COAN	96	CLEO	$E^{ee}_{\text{cm}} \approx$ 10.6 GeV
0.95 ±0.15 ±0.06	f&a		[107] ACCIARRI	95F	L3	1991–1993 LEP runs
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.79 ±0.10 ±0.09		98	[108] BUSKULIC	96	ALEP	Repl. by BARATE 99K

[104] BARATE 99K measure K^0's by detecting K^0_L's in their hadron calorimeter.

[105] BARATE 98E reconstruct K^0's using $K^0_S \to \pi^+ \pi^-$ decays. Not independent of BARATE 98E B(K^0 particles$^- \nu_\tau$) value.

[106] Not independent of COAN 96 B($h^- K^0 \nu_\tau$) and B($K^- K^0 \nu_\tau$) measurements.

[107] ACCIARRI 95F do not identify π^-/K^- and assume B($K^- K^0 \nu_\tau$) = (0.29 ± 0.12)%.

[108] BUSKULIC 96 measure K^0's by detecting K^0_L's in their hadron calorimeter.

$\Gamma(\pi^- \overline{K}^0 (\text{non-}K^*(892)^-)\nu_\tau)/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<0.17	95	ACCIARRI	95F L3	1991–1993 LEP runs

$\Gamma(K^- K^0 \nu_\tau)/\Gamma_{total}$ Γ_{35}/Γ

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
0.154±0.016 OUR FIT					
0.158±0.017 OUR AVERAGE					
0.162±0.021±0.011	150	[109] BARATE	99K	ALEP	1991–1995 LEP runs
0.158±0.042±0.017	46	[110] BARATE	98E	ALEP	1991–1995 LEP runs
0.151±0.021±0.022	111	COAN	96	CLEO	$E_{cm}^{ee} \approx$ 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.26 ±0.09 ±0.02	13	[111] BUSKULIC	96	ALEP	Repl. by BARATE 99K

[109] BARATE 99K measure K^0's by detecting K_L^0's in their hadron calorimeter.

[110] BARATE 98E reconstruct K^0's using $K_S^0 \to \pi^+ \pi^-$ decays.

[111] BUSKULIC 96 measure K^0's by detecting K_L^0's in their hadron calorimeter.

$\Gamma(K^- K^0 \geq 0\pi^0 \nu_\tau)/\Gamma_{total}$ $\Gamma_{36}/\Gamma = (\Gamma_{35}+\Gamma_{40})/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
0.309±0.024 OUR FIT					
0.330±0.055±0.039	124	ABBIENDI	00C	OPAL	1991–1995 LEP runs

$\Gamma(h^- \overline{K}^0 \pi^0 \nu_\tau)/\Gamma_{total}$ $\Gamma_{37}/\Gamma = (\Gamma_{38}+\Gamma_{40})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID		TECN	COMMENT
0.52 ±0.04 OUR FIT						
0.50 ±0.06 OUR AVERAGE Error includes scale factor of 1.2.						
0.446±0.052±0.046	avg	157	[112] BARATE	98E	ALEP	1991–1995 LEP runs
0.562±0.050±0.048	f&a	264	COAN	96	CLEO	$E_{cm}^{ee} \approx$ 10.6 GeV

[112] Not independent of BARATE 98E B($\tau^- \to \pi^- \overline{K}^0 \pi^0 \tau$) and B($\tau^- \to K^- K^0 \pi^0 \nu_\tau$) values.

$\Gamma(\pi^- \overline{K}^0 \pi^0 \nu_\tau)/\Gamma_{total}$ Γ_{38}/Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID		TECN	COMMENT
0.37 ±0.04 OUR FIT						
0.36 ±0.04 OUR AVERAGE						
0.347±0.053±0.037	f&a	299	[113] BARATE	99K	ALEP	1991–1995 LEP runs
0.294±0.073±0.037	f&a	142	[114] BARATE	98E	ALEP	1991–1995 LEP runs
0.417±0.058±0.044	avg		[115] COAN	96	CLEO	$E_{cm}^{ee} \approx$ 10.6 GeV
0.41 ±0.12 ±0.03	f&a		[116] ACCIARRI	95F	L3	1991–1993 LEP runs
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.32 ±0.11 ±0.05		23	[117] BUSKULIC	96	ALEP	Repl. by BARATE 99K

[113] BARATE 99K measure K^0's by detecting K_L^0's in their hadron calorimeter.

[114] BARATE 98E reconstruct K^0's using $K_S^0 \to \pi^+ \pi^-$ decays.

[115] Not independent of COAN 96 B($h^- K^0 \pi^0 \nu_\tau$) and B($K^- K^0 \pi^0 \nu_\tau$) measurements.

[116] ACCIARRI 95F do not identify π^-/K^- and assume B($K^- K^0 \pi^0 \nu_\tau$) = (0.05 ± 0.05)%.

[117] BUSKULIC 96 measure K^0's by detecting K_L^0's in their hadron calorimeter.

$\Gamma(\overline{K}^0\rho^-\nu_\tau)/\Gamma_{total}$ Γ_{39}/Γ

VALUE (%)	DOCUMENT ID		TECN	COMMENT
0.22 ±0.05 OUR AVERAGE				
0.250±0.057±0.044	[118] BARATE	99K	ALEP	1991–1995 LEP runs
0.188±0.054±0.038	[119] BARATE	98E	ALEP	1991–1995 LEP runs

[118] BARATE 99K measure K^0's by detecting K^0_L's in hadron calorimeter. They determine the $\overline{K}^0\rho^-$ fraction in $\tau^- \to \pi^-\overline{K}^0\pi^0\nu_\tau$ decays to be (0.72 ± 0.12 ± 0.10) and multiply their B($\pi^-\overline{K}^0\pi^0\nu_\tau$) measurement by this fraction to obtain the quoted result.

[119] BARATE 98E reconstruct K^0's using $K^0_S \to \pi^+\pi^-$ decays. They determine the $\overline{K}^0\rho^-$ fraction in $\tau^- \to \pi^-\overline{K}^0\pi^0\nu_\tau$ decays to be (0.64 ± 0.09 ± 0.10) and multiply their B($\pi^-\overline{K}^0\pi^0\nu_\tau$) measurement by this fraction to obtain the quoted result.

$\Gamma(K^-K^0\pi^0\nu_\tau)/\Gamma_{total}$ Γ_{40}/Γ

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
0.155±0.020 OUR FIT					
0.144±0.023 OUR AVERAGE					
0.143±0.025±0.015	78	[120] BARATE	99K	ALEP	1991–1995 LEP runs
0.152±0.076±0.021	15	[121] BARATE	98E	ALEP	1991–1995 LEP runs
0.145±0.036±0.020	32	COAN	96	CLEO	$E^{ee}_{cm} \approx$ 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.10 ±0.05 ±0.03	5	[122] BUSKULIC	96	ALEP	Repl. by BARATE 99K

[120] BARATE 99K measure K^0's by detecting K^0_L's in their hadron calorimeter.

[121] BARATE 98E reconstruct K^0's using $K^0_S \to \pi^+\pi^-$ decays.

[122] BUSKULIC 96 measure K^0's by detecting K^0_L's in their hadron calorimeter.

$\Gamma(\pi^-\overline{K}^0 \geq 1\pi^0\nu_\tau)/\Gamma_{total}$ $\Gamma_{41}/\Gamma = (\Gamma_{38}+\Gamma_{42})/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
0.324±0.074±0.066	148	ABBIENDI	00C	OPAL	1991–1995 LEP runs

$\Gamma(\pi^-\overline{K}^0\pi^0\pi^0\nu_\tau)/\Gamma_{total}$ Γ_{42}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
0.26±0.24			[123] BARATE	99R	ALEP	1991–1995 LEP runs
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.66	95	17	[124] BARATE	99K	ALEP	1991–1995 LEP runs
0.58±0.33±0.14		5	[125] BARATE	98E	ALEP	1991–1995 LEP runs

[123] BARATE 99R combine the BARATE 98E and BARATE 99K measurements to obtain this value.

[124] BARATE 99K measure K^0's by detecting K^0_L's in their hadron calorimeter.

[125] BARATE 98E reconstruct K^0's using $K^0_S \to \pi^+\pi^-$ decays.

$\Gamma(K^- K^0 \pi^0 \pi^0 \nu_\tau)/\Gamma_{total}$ Γ_{43}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<0.16 \times 10^{-3}$	95	[126] BARATE	99R	ALEP	1991–1995 LEP runs

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.18 \times 10^{-3}$	95	[127] BARATE	99K	ALEP	1991–1995 LEP runs
$<0.39 \times 10^{-3}$	95	[128] BARATE	98E	ALEP	1991–1995 LEP runs

[126] BARATE 99R combine the BARATE 98E and BARATE 99K bounds to obtain this value.
[127] BARATE 99K measure K^0's by detecting K^0_L's in hadron calorimeter.
[128] BARATE 98E reconstruct K^0's by using $K^0_S \to \pi^+ \pi^-$ decays.

$\Gamma(\pi^- K^0 \overline{K^0} \nu_\tau)/\Gamma_{total}$ $\Gamma_{44}/\Gamma = (2\Gamma_{45}+\Gamma_{46})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID		TECN	COMMENT
0.159 ± 0.029 OUR FIT	Error includes scale factor of 1.1.					
$0.153 \pm 0.030 \pm 0.016$	**avg**	74	[129] BARATE	98E	ALEP	1991–1995 LEP runs

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.31 \pm 0.12 \pm 0.04$		[130] ACCIARRI	95F	L3	1991–1993 LEP runs

[129] BARATE 98E obtain this value by adding twice their $B(\pi^- K^0_S K^0_S \nu_\tau)$ value to their $B(\pi^- K^0_S K^0_L \nu_\tau)$ value.
[130] ACCIARRI 95F assume $B(\pi^- K^0_S K^0_S \nu) = B(\pi^- K^0_S K^0_L \nu) = 1/2 B(\pi^- K^0_S K^0_L \nu)$.

$\Gamma(\pi^- K^0_S K^0_S \nu_\tau)/\Gamma_{total}$ Γ_{45}/Γ

Bose-Einstein correlations might make the mixing fraction different than 1/4.

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
0.024 ± 0.005 OUR FIT					
0.024 ± 0.005 OUR AVERAGE					
$0.026 \pm 0.010 \pm 0.005$	6	BARATE	98E	ALEP	1991–1995 LEP runs
$0.023 \pm 0.005 \pm 0.003$	42	COAN	96	CLEO	$E^{ee}_{cm} \approx$ 10.6 GeV

$\Gamma(\pi^- K^0_S K^0_L \nu_\tau)/\Gamma_{total}$ Γ_{46}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.110 ± 0.028 OUR FIT	Error includes scale factor of 1.1.			
$0.101 \pm 0.023 \pm 0.013$	68	BARATE	98E ALEP	1991–1995 LEP runs

$\Gamma(\pi^- K^0 \overline{K^0} \pi^0 \nu_\tau)/\Gamma_{total}$ Γ_{47}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
$(0.31 \pm 0.23) \times 10^{-3}$	[131] BARATE	99R	ALEP	1991–1995 LEP runs

[131] BARATE 99R combine BARATE 98E $\Gamma(\pi^- K^0_S K^0_S \pi^0 \nu_\tau)/\Gamma_{total}$ and $\Gamma(\pi^- K^0_S K^0_L \pi^0 \nu_\tau)/\Gamma_{total}$ measurements to obtain this value.

$\Gamma(\pi^- K^0_S K^0_S \pi^0 \nu_\tau)/\Gamma_{total}$ Γ_{48}/Γ

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<0.020	95	BARATE	98E ALEP	1991–1995 LEP runs

$\Gamma(\pi^- K_S^0 K_L^0 \pi^0 \nu_\tau)/\Gamma_{total}$ Γ_{49}/Γ

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
0.031 ± 0.011 ± 0.005	11	BARATE	98E	ALEP	1991–1995 LEP runs

$\Gamma(K^0 h^+ h^- h^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{total}$ Γ_{50}/Γ

VALUE (%)	CL%	DOCUMENT ID		TECN	COMMENT
<0.17	95	TSCHIRHART	88	HRS	E_{cm}^{ee} = 29 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.27	90	BELTRAMI	85	HRS	E_{cm}^{ee} = 29 GeV

$\Gamma(K^0 h^+ h^- h^- \nu_\tau)/\Gamma_{total}$ Γ_{51}/Γ

VALUE (%)	EVTS		DOCUMENT ID		TECN	COMMENT
0.023 ± 0.019 ± 0.007	6	[132]	BARATE	98E	ALEP	1991–1995 LEP runs

[132] BARATE 98E reconstruct K^0's using $K_S^0 \to \pi^+ \pi^-$ decays.

$\Gamma(h^- h^- h^+ \geq 0 \text{ neutrals } \geq 0 K_L^0 \nu_\tau)/\Gamma_{total}$ Γ_{52}/Γ

$\Gamma_{52}/\Gamma = (0.3431\Gamma_{33}+0.3431\Gamma_{35}+0.3431\Gamma_{38}+0.3431\Gamma_{40}+0.4307\Gamma_{45}+0.6861\Gamma_{46}+\Gamma_{60}+\Gamma_{68}+\Gamma_{74}+\Gamma_{75}+\Gamma_{82}+\Gamma_{86}+\Gamma_{89}+\Gamma_{90}+0.285\Gamma_{117}+0.285\Gamma_{119}+0.9101\Gamma_{137}+0.9101\Gamma_{138})/\Gamma$

VALUE (%)	EVTS		DOCUMENT ID		TECN	COMMENT
15.20 ± 0.07 OUR FIT Error includes scale factor of 1.1.						
14.8 ± 0.4 OUR AVERAGE						
14.4 ± 0.6 ± 0.3			ADEVA	91F	L3	E_{cm}^{ee} = 88.3–94.3 GeV
15.0 ± 0.4 ± 0.3			BEHREND	89B	CELL	E_{cm}^{ee} = 14–47 GeV
15.1 ± 0.8 ± 0.6			AIHARA	87B	TPC	E_{cm}^{ee} = 29 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •						
13.5 ± 0.3 ± 0.3			ABACHI	89B	HRS	E_{cm}^{ee} = 29 GeV
12.8 ± 1.0 ± 0.7		[133]	BURCHAT	87	MRK2	E_{cm}^{ee} = 29 GeV
12.1 ± 0.5 ± 1.2			RUCKSTUHL	86	DLCO	E_{cm}^{ee} = 29 GeV
12.8 ± 0.5 ± 0.8	1420		SCHMIDKE	86	MRK2	E_{cm}^{ee} = 29 GeV
15.3 ± 1.1 $^{+1.3}_{-1.6}$	367		ALTHOFF	85	TASS	E_{cm}^{ee} = 34.5 GeV
13.6 ± 0.5 ± 0.8			BARTEL	85F	JADE	E_{cm}^{ee} = 34.6 GeV
12.2 ± 1.3 ± 3.9		[134]	BERGER	85	PLUT	E_{cm}^{ee} = 34.6 GeV
13.3 ± 0.3 ± 0.6			FERNANDEZ	85	MAC	E_{cm}^{ee} = 29 GeV
24 ± 6	35		BRANDELIK	80	TASS	E_{cm}^{ee} = 30 GeV
32 ± 5	692	[135]	BACINO	78B	DLCO	E_{cm}^{ee} = 3.1–7.4 GeV
35 ± 11		[135]	BRANDELIK	78	DASP	Assumes $V-A$ decay
18 ± 6.5	33	[135]	JAROS	78	MRK1	$E_{cm}^{ee} > 6$ GeV

[133] BURCHAT 87 value is not independent of SCHMIDKE 86 value.

[134] Not independent of BERGER 85 $\Gamma(\mu^- \overline{\nu}_\mu \nu_\tau)/\Gamma_{total}$, $\Gamma(e^- \overline{\nu}_e \nu_\tau)/\Gamma_{total}$, $\Gamma(h^- \geq 1 \text{ neutrals} \nu_\tau)/\Gamma_{total}$, and $\Gamma(h^- \geq 0 K_L^0 \nu_\tau)/\Gamma_{total}$, and therefore not used in the fit.

[135] Low energy experiments are not in average or fit because the systematic errors in background subtraction are judged to be large.

$\Gamma(h^-h^-h^+ \geq 0$ neutrals ν_τ(ex. $K^0_S \to \pi^+\pi^-$)("3-prong"))/Γ_{total} Γ_{53}/Γ

$\Gamma_{53}/\Gamma = (\Gamma_{60}+\Gamma_{68}+\Gamma_{74}+\Gamma_{75}+\Gamma_{82}+\Gamma_{86}+\Gamma_{89}+\Gamma_{90}+0.285\Gamma_{117}+0.285\Gamma_{119}+0.9101\Gamma_{137}+0.9101\Gamma_{138})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)			EVTS	DOCUMENT ID		TECN	COMMENT
14.57 ±0.07 OUR FIT Error includes scale factor of 1.1.							
14.59 ±0.08 OUR AVERAGE Error includes scale factor of 1.1.							
14.569 ±0.093	±0.048	f&a	23k	[136] ABREU	01M	DLPH	1992–1995 LEP runs
14.556 ±0.105	±0.076	f&a		[137] ACHARD	01D	L3	1992–1995 LEP runs
14.96 ±0.09	±0.22	f&a	10.4k	AKERS	95Y	OPAL	1991–1994 LEP runs
14.22 ±0.10	±0.37	avg		[138] BALEST	95C	CLEO	$E^{ee}_{cm} \approx$ 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •							
15.26 ±0.26	±0.22			ACTON	92H	OPAL	Repl. by AKERS 95Y
13.3 ±0.3	±0.8			[139] ALBRECHT	92D	ARG	E^{ee}_{cm}= 9.4–10.6 GeV
14.35 $^{+0.40}_{-0.45}$	±0.24			DECAMP	92C	ALEP	1989–1990 LEP runs

[136] The correlation coefficients between this meausrement and the ABREU 01M measurements of B($\tau \to$ 1-prong) and B($\tau \to$ 5-prong) are −0.98 and −0.08 respectively.

[137] The correlation coefficients between this measurement and the ACHARD 01D measurements of B($\tau \to$ "1-prong") and B($\tau \to$ "5-prong") are −0.978 and −0.19 respectively.

[138] Not independent of BALEST 95C B($h^-h^-h^+\nu_\tau$) and B($h^-h^-h^+\pi^0\nu_\tau$) values, and BORTOLETTO 93 B($h^-h^-h^+2\pi^0\nu_\tau$)/B($h^-h^-h^+ \geq 0$ neutrals ν_τ) value.

[139] This ALBRECHT 92D value is not independent of their $\Gamma(\mu^-\overline{\nu}_\mu\nu_\tau)\Gamma(e^-\overline{\nu}_e\nu_\tau)/\Gamma^2_{total}$ value.

$\Gamma(h^-h^-h^+\nu_\tau)/\Gamma_{total}$ Γ_{54}/Γ

$\Gamma_{54}/\Gamma = (0.3431\Gamma_{33}+0.3431\Gamma_{35}+\Gamma_{60}+\Gamma_{82}+\Gamma_{89}+0.0221\Gamma_{137})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)			EVTS	DOCUMENT ID		TECN	COMMENT
10.01±0.09 OUR FIT Error includes scale factor of 1.2.							
9.8 ±0.6 OUR AVERAGE Error includes scale factor of 4.4. See the ideogram below.							
7.6 ±0.1	±0.5	avg	7.5k	[140] ALBRECHT	96E	ARG	E^{ee}_{cm}= 9.4–10.6 GeV
9.92±0.10	±0.09	f&a	11.2k	[141] BUSKULIC	96	ALEP	LEP 1991–1993 data
9.49±0.36	±0.63	f&a		DECAMP	92C	ALEP	1989–1990 LEP runs
• • • We do not use the following data for averages, fits, limits, etc. • • •							
8.7 ±0.7	±0.3		694	[142] BEHREND	90	CELL	E^{ee}_{cm}= 35 GeV
7.0 ±0.3	±0.7		1566	[143] BAND	87	MAC	E^{ee}_{cm}= 29 GeV
6.7 ±0.8	±0.9			[144] BURCHAT	87	MRK2	E^{ee}_{cm}= 29 GeV
6.4 ±0.4	±0.9			[145] RUCKSTUHL	86	DLCO	E^{ee}_{cm}= 29 GeV
7.8 ±0.5	±0.8		890	SCHMIDKE	86	MRK2	E^{ee}_{cm}= 29 GeV
8.4 ±0.4	±0.7		1255	[145] FERNANDEZ	85	MAC	E^{ee}_{cm}= 29 GeV
9.7 ±2.0	±1.3			BEHREND	84	CELL	E^{ee}_{cm}= 14,22 GeV

[140] ALBRECHT 96E not independent of ALBRECHT 93C $\Gamma(h^-h^-h^+\nu_\tau(\text{ex. } K^0) \times \Gamma(\text{particle}^- \geq 0 \text{ neutrals } \geq 0 K^0_L \nu_\tau)/\Gamma^2_{\text{total}}$ value.

[141] BUSKULIC 96 quote $B(h^-h^-h^+\nu_\tau(\text{ex. } K^0)) = 9.50 \pm 0.10 \pm 0.11$. We add 0.42 to remove their K^0 correction and reduce the systematic error accordingly.

[142] BEHREND 90 subtract 0.3% to account for the $\tau^- \to K^*(892)^-\nu_\tau$ contribution to measured events.

[143] BAND 87 subtract for charged kaon modes; not independent of FERNANDEZ 85 value.

[144] BURCHAT 87 value is not independent of SCHMIDKE 86 value.

[145] Value obtained by multiplying paper's $R = B(h^-h^-h^+\nu_\tau)/B(\text{3-prong})$ by B(3-prong) = 0.143 and subtracting 0.3% for $K^*(892)$ background.



$\Gamma\left(h^-h^-h^+\nu_\tau\right)/\Gamma_{\text{total}}$ (%)

$\Gamma(h^-h^-h^+\nu_\tau(\text{ex.}K^0))/\Gamma_{\text{total}}$ Γ_{55}/Γ

$\Gamma_{55}/\Gamma = (\Gamma_{60}+\Gamma_{82}+\Gamma_{89}+0.0221\Gamma_{137})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
9.65±0.09 OUR FIT Error includes scale factor of 1.2.						
9.57±0.11 OUR AVERAGE						
9.50±0.10±0.11	avg	11.2k	[146] BUSKULIC	96	ALEP	LEP 1991–1993 data
9.87±0.10±0.24	avg		[147] AKERS	95Y	OPAL	1991–1994 LEP runs
9.51±0.07±0.20	f&a	37.7k	BALEST	95C	CLEO	$E^{ee}_{\text{cm}} \approx$ 10.6 GeV

[146] Not independent of BUSKULIC 96 $B(h^-h^-h^+\nu_\tau)$ value.

[147] Not independent of AKERS 95Y $B(h^-h^-h^+ \geq 0 \text{ neutrals}\nu_\tau(\text{ex. } K^0_S \to \pi^+\pi^-))$ and $B(h^-h^-h^+\nu_\tau(\text{ex. } K^0))/B(h^-h^-h^+ \geq 0 \text{ neutrals}\nu_\tau(\text{ex. } K^0_S \to \pi^+\pi^-))$ values.

$\Gamma(h^-h^-h^+\nu_\tau(\text{ex.}K^0))/\Gamma(h^-h^-h^+ \geq 0 \text{ neutrals } \nu_\tau(\text{ex. } K^0_S \to \pi^+\pi^-)$ ("3-prong")) Γ_{55}/Γ_{53}

$\Gamma_{55}/\Gamma_{53} = (\Gamma_{60}+\Gamma_{82}+\Gamma_{89}+0.0221\Gamma_{137})/(\Gamma_{60}+\Gamma_{68}+\Gamma_{74}+\Gamma_{75}+\Gamma_{82}+\Gamma_{86}+\Gamma_{89}+\Gamma_{90}+0.285\Gamma_{117}+0.285\Gamma_{119}+0.9101\Gamma_{137}+0.9101\Gamma_{138})$

VALUE	DOCUMENT ID	TECN	COMMENT
0.662±0.006 OUR FIT Error includes scale factor of 1.3.			
0.660±0.004±0.014	AKERS	95Y OPAL	1991–1994 LEP runs

$\Gamma(h^-h^-h^+\nu_\tau(\text{ex.}K^0,\omega))/\Gamma_{\text{total}}$ $\Gamma_{56}/\Gamma = (\Gamma_{60}+\Gamma_{82}+\Gamma_{89})/\Gamma$

VALUE (%)	DOCUMENT ID
9.61±0.09 OUR FIT Error includes scale factor of 1.2.	

$\Gamma(\pi^-\pi^+\pi^-\nu_\tau)/\Gamma_{\text{total}}$ $\Gamma_{57}/\Gamma = (0.3431\Gamma_{33}+\Gamma_{60}+0.0221\Gamma_{137})/\Gamma$

VALUE (%)	DOCUMENT ID
9.52±0.10 OUR FIT Error includes scale factor of 1.2.	

$\Gamma(\pi^-\pi^+\pi^-\nu_\tau(\text{ex.}K^0))/\Gamma_{\text{total}}$ $\Gamma_{58}/\Gamma = (\Gamma_{60}+0.0221\Gamma_{137})/\Gamma$

VALUE (%)	DOCUMENT ID
9.22±0.10 OUR FIT Error includes scale factor of 1.2.	

$\Gamma(\pi^-\pi^+\pi^-\nu_\tau(\text{ex.}K^0)\text{, non-axial vector})/\Gamma(\pi^-\pi^+\pi^-\nu_\tau(\text{ex.}K^0))$ $\Gamma_{59}/\Gamma_{58} = \Gamma_{59}/(\Gamma_{60}+0.0221\Gamma_{137})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.261	95	[148] ACKERSTAFF 97R	OPAL	1992–1994 LEP runs

[148] Model-independent limit from structure function analysis on contribution to B($\tau^- \to \pi^-\pi^+\pi^-\nu_\tau$(ex. K^0)) from non-axial vectors.

$\Gamma(\pi^-\pi^+\pi^-\nu_\tau(\text{ex.}K^0,\omega))/\Gamma_{\text{total}}$ Γ_{60}/Γ

VALUE (%)	DOCUMENT ID
9.18±0.10 OUR FIT Error includes scale factor of 1.2.	

$\Gamma(h^-h^-h^+ \geq 1 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$ Γ_{61}/Γ

$\Gamma_{61}/\Gamma = (0.3431\Gamma_{38}+0.3431\Gamma_{40}+0.4307\Gamma_{45}+0.6861\Gamma_{46}+\Gamma_{68}+\Gamma_{74}+\Gamma_{75}+\Gamma_{86}+\Gamma_{90}+0.285\Gamma_{117}+0.285\Gamma_{119}+0.888\Gamma_{137}+0.9101\Gamma_{138})/\Gamma$

VALUE (%)		EVTS	DOCUMENT ID		TECN	COMMENT
5.19±0.10 OUR FIT Error includes scale factor of 1.3.						
• • • We do not use the following data for averages, fits, limits, etc. • • •						
5.6 ±0.7 ±0.3		352	[149] BEHREND	90	CELL	E^{ee}_{cm}= 35 GeV
4.2 ±0.5 ±0.9		203	[150] ALBRECHT	87L	ARG	E^{ee}_{cm}= 10 GeV
6.1 ±0.8 ±0.9			[151] BURCHAT	87	MRK2	E^{ee}_{cm}= 29 GeV
7.6 ±0.4 ±0.9			[152,153] RUCKSTUHL	86	DLCO	E^{ee}_{cm}= 29 GeV
4.7 ±0.5 ±0.8		530	[154] SCHMIDKE	86	MRK2	E^{ee}_{cm}= 29 GeV
5.6 ±0.4 ±0.7			[153] FERNANDEZ	85	MAC	E^{ee}_{cm}= 29 GeV
6.2 ±2.3 ±1.7			BEHREND	84	CELL	E^{ee}_{cm}= 14,22 GeV

[149] BEHREND 90 value is not independent of BEHREND 90 B($3h\nu_\tau \geq 1$ neutrals) + B(5-prong).

[150] ALBRECHT 87L measure the product of branching ratios B($3\pi^\pm\pi^0\nu_\tau$) B(($e\overline{\nu}$ or $\mu\overline{\nu}$ or π or K or ρ)ν_τ) = 0.029 and use the PDG 86 values for the second branching ratio which sum to 0.69 ± 0.03 to get the quoted value.

[151] BURCHAT 87 value is not independent of SCHMIDKE 86 value.

[152] Contributions from kaons and from $>1\pi^0$ are subtracted. Not independent of (3-prong + $0\pi^0$) and (3-prong + $\geq 0\pi^0$) values.

[153] Value obtained using paper's R = B($h^-h^-h^+\nu_\tau$)/B(3-prong) and current B(3-prong) = 0.143.

[154] Not independent of SCHMIDKE 86 $h^-h^-h^+\nu_\tau$ and $h^-h^-h^+(\geq 0\pi^0)\nu_\tau$ values.

$\Gamma(h^-h^-h^+ \geq 1$ neutrals ν_τ(ex. $K^0_S \to \pi^+\pi^-))/\Gamma_{total}$ Γ_{62}/Γ

$\Gamma_{62}/\Gamma = (\Gamma_{68}+\Gamma_{74}+\Gamma_{75}+\Gamma_{86}+\Gamma_{90}+0.285\Gamma_{117}+0.285\Gamma_{119}+0.888\Gamma_{137}+0.9101\Gamma_{138})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID		TECN	COMMENT
4.92±0.10 OUR FIT	Error includes scale factor of 1.3.					
5.07±0.24 OUR AVERAGE						
5.09±0.10±0.23	avg		[155] AKERS	95Y	OPAL	1991–1994 LEP runs
4.95±0.29±0.65	f&a	570	DECAMP	92C	ALEP	1989–1990 LEP runs

[155] Not independent of AKERS 95Y B($h^-h^-h^+ \geq 0$ neutralsν_τ(ex. $K^0_S \to \pi^+\pi^-$)) and B($h^-h^-h^+ \geq 0$ neutralsν_τ(ex. K^0))/B($h^-h^-h^+ \geq 0$ neutralsν_τ(ex. $K^0_S \to \pi^+\pi^-$)) values.

$\Gamma(h^-h^-h^+\pi^0\nu_\tau)/\Gamma_{total}$ Γ_{63}/Γ

$\Gamma_{63}/\Gamma = (0.3431\Gamma_{38}+0.3431\Gamma_{40}+\Gamma_{68}+\Gamma_{86}+\Gamma_{90}+0.231\Gamma_{119}+0.888\Gamma_{137}+0.0221\Gamma_{138})/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
4.53±0.09 OUR FIT	Error includes scale factor of 1.3.				
4.45±0.09±0.07	6.1k	[156] BUSKULIC	96	ALEP	LEP 1991–1993 data

[156] BUSKULIC 96 quote B($h^-h^-h^+\pi^0\nu_\tau$(ex. K^0)) = 4.30 ± 0.09 ± 0.09. We add 0.15 to remove their K^0 correction and reduce the systematic error accordingly.

$\Gamma(h^-h^-h^+\pi^0\nu_\tau$(ex.$K^0$))$/\Gamma_{total}$ Γ_{64}/Γ

$\Gamma_{64}/\Gamma = (\Gamma_{68}+\Gamma_{86}+\Gamma_{90}+0.231\Gamma_{119}+0.888\Gamma_{137}+0.0221\Gamma_{138})/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
4.35±0.09 OUR FIT	Error includes scale factor of 1.3.			
4.23±0.06±0.22	7.2k	BALEST	95C CLEO	$E^{ee}_{cm} \approx$ 10.6 GeV

$\Gamma(h^-h^-h^+\pi^0\nu_\tau$(ex. K^0,ω))$/\Gamma_{total}$ $\Gamma_{65}/\Gamma = (\Gamma_{68}+\Gamma_{86}+\Gamma_{90}+0.231\Gamma_{119})/\Gamma$

VALUE (%)	DOCUMENT ID
2.62±0.09 OUR FIT	Error includes scale factor of 1.2.

$\Gamma(\pi^-\pi^+\pi^-\pi^0\nu_\tau)/\Gamma_{\rm total}$
$\Gamma_{66}/\Gamma = (0.3431\Gamma_{38}+\Gamma_{68}+0.888\Gamma_{137}+0.0221\Gamma_{138})/\Gamma$

VALUE (%)	DOCUMENT ID
4.37±0.10 OUR FIT Error includes scale factor of 1.3.	

$\Gamma(\pi^-\pi^+\pi^-\pi^0\nu_\tau(\text{ex.}K^0))/\Gamma_{\rm total}$ $\Gamma_{67}/\Gamma = (\Gamma_{68}+0.888\Gamma_{137}+0.0221\Gamma_{138})/\Gamma$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
4.24±0.10 OUR FIT Error includes scale factor of 1.3.			
4.19±0.10±0.21	[157] EDWARDS	00A CLEO	4.7 fb^{-1} $E_{\rm cm}^{ee}$= 10.6 GeV

[157] EDWARDS 00A quote $(4.19 \pm 0.10) \times 10^{-2}$ with a 5% systematic error.

$\Gamma(\pi^-\pi^+\pi^-\pi^0\nu_\tau(\text{ex.}K^0,\omega))/\Gamma_{\rm total}$ Γ_{68}/Γ

VALUE (%)	DOCUMENT ID
2.51±0.09 OUR FIT Error includes scale factor of 1.2.	

$\Gamma(h^-\rho\pi^0\nu_\tau)/\Gamma(h^-h^-h^+\pi^0\nu_\tau)$ Γ_{69}/Γ_{63}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.30±0.04±0.02	393	ALBRECHT	91D ARG	$E_{\rm cm}^{ee}$= 9.4–10.6 GeV

$\Gamma(h^-\rho^+h^-\nu_\tau)/\Gamma(h^-h^-h^+\pi^0\nu_\tau)$ Γ_{70}/Γ_{63}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.10±0.03±0.04	142	ALBRECHT	91D ARG	$E_{\rm cm}^{ee}$= 9.4–10.6 GeV

$\Gamma(h^-\rho^-h^+\nu_\tau)/\Gamma(h^-h^-h^+\pi^0\nu_\tau)$ Γ_{71}/Γ_{63}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.26±0.05±0.01	370	ALBRECHT	91D ARG	$E_{\rm cm}^{ee}$= 9.4–10.6 GeV

$\Gamma(h^-h^-h^+2\pi^0\nu_\tau)/\Gamma_{\rm total}$ Γ_{72}/Γ
$\Gamma_{72}/\Gamma = (0.4307\Gamma_{45}+\Gamma_{74}+0.236\Gamma_{117}+0.888\Gamma_{138})/\Gamma$

VALUE (%)	DOCUMENT ID
0.55±0.04 OUR FIT	

$\Gamma(h^-h^-h^+2\pi^0\nu_\tau(\text{ex.}K^0))/\Gamma_{\rm total}$ Γ_{73}/Γ
$\Gamma_{73}/\Gamma = (\Gamma_{74}+0.236\Gamma_{117}+0.888\Gamma_{138})/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.54±0.04 OUR FIT				
0.50±0.07±0.07	1.8k	BUSKULIC	96 ALEP	LEP 1991–1993 data

$\Gamma(h^-h^-h^+2\pi^0\nu_\tau(\text{ex.}K^0))/\Gamma(h^-h^-h^+ \geq 0\,\text{neutrals}\, \geq 0K_L^0\nu_\tau)$ Γ_{73}/Γ_{52}
$\Gamma_{73}/\Gamma_{52} = (\Gamma_{74}+0.236\Gamma_{117}+0.888\Gamma_{138})/(0.3431\Gamma_{33}+0.3431\Gamma_{35}+0.3431\Gamma_{38}+0.3431\Gamma_{40}+0.4307\Gamma_{45}+0.6861\Gamma_{46}+\Gamma_{60}+\Gamma_{68}+\Gamma_{74}+\Gamma_{75}+\Gamma_{82}+\Gamma_{86}+\Gamma_{89}+\Gamma_{90}+0.285\Gamma_{117}+0.285\Gamma_{119}+0.9101\Gamma_{137}+0.9101\Gamma_{138})$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0354±0.0028 OUR FIT				
0.034 ±0.002 ±0.003	668	BORTOLETTO93	CLEO	$E_{\rm cm}^{ee} \approx$ 10.6 GeV

$\Gamma(h^- h^- h^+ 2\pi^0 \nu_\tau (\text{ex.}K^0,\omega,\eta))/\Gamma_{\text{total}}$ Γ_{74}/Γ

VALUE (%)	DOCUMENT ID
0.11±0.04 OUR FIT	

$\Gamma(h^- h^- h^+ 3\pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{75}/Γ

VALUE (%)		EVTS	DOCUMENT ID		TECN	COMMENT
0.023 ±0.008 OUR FIT	Error includes scale factor of 1.6.					
0.023 ±0.005 OUR AVERAGE						
0.022 ±0.003 ±0.004		139	ANASTASSOV	01	CLEO	E_{cm}^{ee}= 10.6 GeV
0.11 ±0.04 ±0.05		440	[158] BUSKULIC	96	ALEP	LEP 1991–1993 data
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.0285±0.0056±0.0051		57	ANDERSON	97	CLEO	Repl. by ANASTASSOV 01

[158] BUSKULIC 96 state their measurement is for $B(h^- h^+ \geq 3\pi^0 \nu_\tau)$. We assume that $B(h^- h^- h^+ \geq 4\pi^0 \nu_\tau)$ is very small.

$\Gamma(K^- h^+ h^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$
$\Gamma_{76}/\Gamma = (0.3431\Gamma_{35}+0.3431\Gamma_{40}+\Gamma_{82}+\Gamma_{86}+\Gamma_{89}+\Gamma_{90}+0.285\Gamma_{119})/\Gamma$

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
0.65±0.05 OUR FIT	Error includes scale factor of 1.4.			
<0.6	90	AIHARA	84C TPC	E_{cm}^{ee}= 29 GeV

$\Gamma(K^- h^+ \pi^- \nu_\tau (\text{ex.}K^0))/\Gamma_{\text{total}}$ $\Gamma_{77}/\Gamma = (\Gamma_{82}+\Gamma_{89})/\Gamma$

VALUE (%)	DOCUMENT ID
0.44±0.05 OUR FIT	Error includes scale factor of 1.5.

$\Gamma(K^- h^+ \pi^- \nu_\tau (\text{ex.}K^0))/\Gamma(\pi^- \pi^+ \pi^- \nu_\tau (\text{ex.}K^0))$
$\Gamma_{77}/\Gamma_{58} = (\Gamma_{82}+\Gamma_{89})/(\Gamma_{60}+0.0221\Gamma_{137})$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
4.7 ±0.6 OUR FIT	Error includes scale factor of 1.5.			
5.44±0.21±0.53	7.9k	RICHICHI	99 CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(K^- h^+ \pi^- \pi^0 \nu_\tau (\text{ex.}K^0))/\Gamma_{\text{total}}$ $\Gamma_{78}/\Gamma = (\Gamma_{86}+\Gamma_{90}+0.231\Gamma_{119})/\Gamma$

VALUE (%)	DOCUMENT ID
0.110±0.022 OUR FIT	

$\Gamma(K^- h^+ \pi^- \pi^0 \nu_\tau (\text{ex.}K^0))/\Gamma(\pi^- \pi^+ \pi^- \pi^0 \nu_\tau (\text{ex.}K^0))$
$\Gamma_{78}/\Gamma_{67} = (\Gamma_{86}+\Gamma_{90}+0.231\Gamma_{119})/(\Gamma_{68}+0.888\Gamma_{137}+0.0221\Gamma_{138})$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
2.6 ±0.5 OUR FIT				
2.61±0.45±0.42	719	RICHICHI	99 CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(K^- \pi^+ \pi^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$
$\Gamma_{79}/\Gamma = (0.3431\Gamma_{35}+0.3431\Gamma_{40}+\Gamma_{82}+\Gamma_{86}+0.285\Gamma_{119})/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.45±0.05 OUR FIT	Error includes scale factor of 1.4.			
$\mathbf{0.58^{+0.15}_{-0.13}\pm0.12}$	20	[159] BAUER	94 TPC	E_{cm}^{ee}= 29 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.22^{+0.16}_{-0.13}\pm0.05$	9	[160] MILLS	85 DLCO	E_{cm}^{ee}= 29 GeV

[159] We multiply 0.58% by 0.20, the relative systematic error quoted by BAUER 94, to obtain the systematic error.

[160] Error correlated with MILLS 85 ($K K \pi \nu$) value. We multiply 0.22% by 0.23, the relative systematic error quoted by MILLS 85, to obtain obtain the systematic error.

$\Gamma(K^-\pi^+\pi^- \geq 0\pi^0\nu_\tau(\text{ex.}K^0))/\Gamma_{\text{total}}$ $\Gamma_{80}/\Gamma = (\Gamma_{82}+\Gamma_{86}+0.231\Gamma_{119})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		*DOCUMENT ID*		*TECN*	*COMMENT*
0.35 ±0.05 OUR FIT Error includes scale factor of 1.4.					
0.30 ±0.05 OUR AVERAGE					
0.343±0.073±0.031	f&a	ABBIENDI	00D	OPAL	1990–1995 LEP runs
0.275±0.064	avg	[161] BARATE	98	ALEP	1991–1995 LEP runs

[161] Not independent of BARATE 98 $\Gamma(\tau^- \to K^-\pi^+\pi^-\nu_\tau)/\Gamma_{\text{total}}$ and $\Gamma(\tau^- \to K^-\pi^+\pi^-\pi^0\nu_\tau)/\Gamma_{\text{total}}$ values.

$\Gamma(K^-\pi^+\pi^-\nu_\tau)/\Gamma_{\text{total}}$ $\Gamma_{81}/\Gamma = (0.3431\Gamma_{35}+\Gamma_{82})/\Gamma$

VALUE (%)	*DOCUMENT ID*
0.33±0.05 OUR FIT Error includes scale factor of 1.5.	

$\Gamma(K^-\pi^+\pi^-\nu_\tau(\text{ex.}K^0))/\Gamma_{\text{total}}$ Γ_{82}/Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
0.28 ±0.05 OUR FIT Error includes scale factor of 1.5.					
0.28 ±0.05 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.					
0.360±0.082±0.048 avg		ABBIENDI	00D	OPAL	1990–1995 LEP runs
0.346±0.023±0.056 avg	158	[162] RICHICHI	99	CLEO	E_{cm}^{ee}= 10.6 GeV
0.214±0.037±0.029 f&a		BARATE	98	ALEP	1991–1995 LEP runs

[162] Not independent of RICHICHI 99 $\Gamma(\tau^- \to K^-h^+\pi^-\nu_\tau(\text{ex.}K^0))/\Gamma(\tau^- \to \pi^-\pi^+\pi^-\nu_\tau(\text{ex.}K^0))$, $\Gamma(\tau^- \to K^-K^+\pi^-\nu_\tau)/\Gamma(\tau^- \to \pi^-\pi^+\pi^-\nu_\tau(\text{ex.}K^0))$ and BALEST 95C $\Gamma(\tau^- \to h^-h^-h^+\nu_\tau(\text{ex.}K^0))/\Gamma_{\text{total}}$ values.



$\Gamma\left(K^-\pi^+\pi^-\nu_\tau(\text{ex.}K^0)\right)/\Gamma_{\text{total}}$ (%)

$\Gamma\left(K^-\rho^0\nu_\tau \to K^-\pi^+\pi^-\nu_\tau\right)/\Gamma\left(K^-\pi^+\pi^-\nu_\tau(\text{ex.}K^0)\right)$ **Γ_{83}/Γ_{82}**

VALUE	*DOCUMENT ID*		*TECN*	*COMMENT*
0.48±0.14±0.10	[163] ASNER	00B	CLEO	E_{cm}^{ee}= 10.6 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.39±0.14	[164] BARATE	99R	ALEP	1991–1995 LEP runs

[163] ASNER 00B assume $\tau^- \to K^-\pi^+\pi^-\nu_\tau$ (ex. K^0) decays proceed only through $K\rho$ and $K^*\pi$ intermediate states. They assume the resonance structure of $\tau^- \to K^-\pi^+\pi^-\nu_\tau$ (ex. K^0) decays is dominated by $K_1(1270)^-$ and $K_1(1400)^-$ resonances, and assume $B(K_1(1270) \to K^*(892)\pi) = (16 \pm 5)\%$, $B(K_1(1270) \to K\rho) = (42 \pm 6)\%$, and $B(K_1(1400) \to K\rho) = 0$.

[164] BARATE 99R assume $\tau^- \to K^-\pi^+\pi^-\nu_\tau$ (ex. K^0) decays proceed only through $K\rho$ and $K^*\pi$ intermediate states. The quoted error is statistical only.

$\Gamma\left(K^-\pi^+\pi^-\pi^0\nu_\tau\right)/\Gamma_{\text{total}}$ **$\Gamma_{84}/\Gamma = (0.3431\Gamma_{40}+\Gamma_{86}+0.231\Gamma_{119})/\Gamma$**

VALUE (units 10^{-4})	*DOCUMENT ID*
12.4±2.5 OUR FIT	

$\Gamma\left(K^-\pi^+\pi^-\pi^0\nu_\tau(\text{ex.}K^0)\right)/\Gamma_{\text{total}}$ **$\Gamma_{85}/\Gamma = (\Gamma_{86}+0.231\Gamma_{119})/\Gamma$**

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (units 10^{-4})		CL%	DOCUMENT ID		TECN	COMMENT
7.0±2.4 OUR FIT						
7.0±2.5 OUR AVERAGE						
7.5±2.6±1.8	avg		[165] RICHICHI	99	CLEO	E_{cm}^{ee}= 10.6 GeV
6.1±3.9±1.8	f&a		BARATE	98	ALEP	1991–1995 LEP runs
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<17		95	ABBIENDI	00D	OPAL	1990–1995 LEP runs

[165] Not independent of RICHICHI 99 $\Gamma(\tau^- \to K^- h^+ \pi^- \nu_\tau(\text{ex.}K^0))/\Gamma(\tau^- \to \pi^- \pi^+ \pi^- \nu_\tau(\text{ex.}K^0))$, $\Gamma(\tau^- \to K^- K^+ \pi^- \nu_\tau)/\Gamma(\tau^- \to \pi^- \pi^+ \pi^- \nu_\tau(\text{ex.}K^0))$ and BALEST 95C $\Gamma(\tau^- \to h^- h^- h^+ \nu_\tau(\text{ex.}K^0))/\Gamma_{\text{total}}$ values.

$\Gamma(K^-\pi^+\pi^-\pi^0\nu_\tau(\text{ex.}K^0,\eta))/\Gamma_{\text{total}}$ Γ_{86}/Γ

Test of lepton family number conservation.

VALUE (units 10^{-4})	DOCUMENT ID
6.4±2.4 OUR FIT	

$\Gamma(K^-\pi^+K^- \geq 0 \text{ neut. } \nu_\tau)/\Gamma_{\text{total}}$ Γ_{87}/Γ

VALUE (%)	CL%	DOCUMENT ID		TECN	COMMENT
<0.09	95	BAUER	94	TPC	E_{cm}^{ee}= 29 GeV

$\Gamma(K^-K^+\pi^- \geq 0 \text{ neut. } \nu_\tau)/\Gamma_{\text{total}}$ $\Gamma_{88}/\Gamma=(\Gamma_{89}+\Gamma_{90})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID		TECN	COMMENT
0.200±0.023 OUR FIT						
0.203±0.031 OUR AVERAGE						
0.159±0.053±0.020	f&a		ABBIENDI	00D	OPAL	1990–1995 LEP runs
0.238±0.042	avg		[166] BARATE	98	ALEP	1991–1995 LEP runs
$0.15\ ^{+0.09}_{-0.07}\ \pm0.03$	f&a	4	[167] BAUER	94	TPC	E_{cm}^{ee}= 29 GeV

[166] Not independent of BARATE 98 $\Gamma(\tau^- \to K^- K^+ \pi^- \nu_\tau)/\Gamma_{\text{total}}$ and $\Gamma(\tau^- \to K^- K^+ \pi^- \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ values.

[167] We multiply 0.15% by 0.20, the relative systematic error quoted by BAUER 94, to obtain the systematic error.

$\Gamma(K^-K^+\pi^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_{89}/Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID		TECN	COMMENT
0.160±0.019 OUR FIT						
0.151±0.019 OUR AVERAGE						
0.087±0.056±0.040	avg		ABBIENDI	00D	OPAL	1990–1995 LEP runs
0.145±0.013±0.028	avg	2.3k	[168] RICHICHI	99	CLEO	E_{cm}^{ee}= 10.6 GeV
0.163±0.021±0.017	f&a		BARATE	98	ALEP	1991–1995 LEP runs
$0.22\ ^{+0.17}_{-0.11}\ \pm0.05$	f&a	9	[169] MILLS	85	DLCO	E_{cm}^{ee}= 29 GeV

[168] Not independent of RICHICHI 99 $\Gamma(\tau^- \to K^- K^+ \pi^- \nu_\tau)/\Gamma(\tau^- \to \pi^- \pi^+ \pi^- \nu_\tau(\text{ex.}K^0))$ and BALEST 95C $\Gamma(\tau^- \to h^- h^- h^+ \nu_\tau(\text{ex.}K^0))/\Gamma_{\text{total}}$ values.

[169] Error correlated with MILLS 85 ($K\pi\pi\pi^0\nu$) value. We multiply 0.22% by 0.23, the relative systematic error quoted by MILLS 85, to obtain obtain the systematic error.

$\Gamma(K^- K^+ \pi^- \nu_\tau)/\Gamma(\pi^- \pi^+ \pi^- \nu_\tau(\text{ex.}K^0))$ $\Gamma_{89}/\Gamma_{58} = \Gamma_{89}/(\Gamma_{60}+0.0221\Gamma_{137})$

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
1.74±0.20 OUR FIT					
1.60±0.15±0.30	2.3k	RICHICHI	99	CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(K^- K^+ \pi^- \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{90}/Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (units 10^{-4})		CL%	EVTS	DOCUMENT ID		TECN	COMMENT
4.0±1.6 OUR FIT							
4.4±1.8 OUR AVERAGE	Error includes scale factor of 1.1.						
3.3±1.8±0.7	avg		158 [170]	RICHICHI	99	CLEO	E_{cm}^{ee}= 10.6 GeV
7.5±2.9±1.5	f&a			BARATE	98	ALEP	1991–1995 LEP runs
• • • We do not use the following data for averages, fits, limits, etc. • • •							
<27		95		ABBIENDI	00D	OPAL	1990–1995 LEP runs

[170] Not independent of RICHICHI 99 $\Gamma(\tau^- \to K^- K^+ \pi^- \nu_\tau)/\Gamma(\tau^- \to \pi^- \pi^+ \pi^- \nu_\tau(\text{ex.}K^0))$ and BALEST 95C $\Gamma(\tau^- \to h^- h^- h^+ \nu_\tau(\text{ex.}K^0))/\Gamma_{\text{total}}$ values.

$\Gamma(K^- K^+ \pi^- \pi^0 \nu_\tau)/\Gamma(\pi^- \pi^+ \pi^- \pi^0 \nu_\tau(\text{ex.}K^0))$ $\Gamma_{90}/\Gamma_{67} = \Gamma_{90}/(\Gamma_{68}+0.888\Gamma_{137}+0.0221\Gamma_{138})$

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
0.9 ±0.4 OUR FIT					
0.79±0.44±0.16	158	[171] RICHICHI	99	CLEO	E_{cm}^{ee}= 10.6 GeV

[171] RICHICHI 99 also quote a 95%CL upper limit of 0.0157 for this measurement.

$\Gamma(K^- K^+ K^- \geq 0 \text{ neut. } \nu_\tau)/\Gamma_{\text{total}}$ Γ_{91}/Γ

VALUE (%)	CL%	DOCUMENT ID		TECN	COMMENT
<0.21	95	BAUER	94	TPC	E_{cm}^{ee}= 29 GeV

$\Gamma(K^- K^+ K^- \nu_\tau)/\Gamma_{\text{total}}$ Γ_{92}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.9 \times 10^{-4}$	90	BARATE	98	ALEP	1991–1995 LEP runs

$\Gamma(\pi^- K^+ \pi^- \geq 0 \text{ neut. } \nu_\tau)/\Gamma_{\text{total}}$ Γ_{93}/Γ

VALUE (%)	CL%	DOCUMENT ID		TECN	COMMENT
<0.25	95	BAUER	94	TPC	E_{cm}^{ee}= 29 GeV

$\Gamma(e^- e^- e^+ \overline{\nu}_e \nu_\tau)/\Gamma_{total}$ Γ_{94}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID		TECN	COMMENT
2.8±1.4±0.4	5	ALAM	96	CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(\mu^- e^- e^+ \overline{\nu}_\mu \nu_\tau)/\Gamma_{total}$ Γ_{95}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN	COMMENT
<3.6	90	ALAM	96	CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(3h^-2h^+ \geq 0$ neutrals ν_τ(ex. $K_S^0 \to \pi^-\pi^+$)("5-prong"))$/\Gamma_{total}$ Γ_{96}/Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average. $\Gamma_{96}/\Gamma = (\Gamma_{97}+\Gamma_{98})/\Gamma$

VALUE (%)		EVTS	DOCUMENT ID		TECN	COMMENT
0.100±0.006 OUR FIT						
0.111±0.008 OUR AVERAGE	Error includes scale factor of 1.1.					
0.115±0.013±0.006	f&a	112	[172] ABREU	01M	DLPH	1992–1995 LEP runs
0.170±0.022±0.026	f&a		[173] ACHARD	01D	L3	1992–1995 LEP runs
0.119±0.013±0.008	avg	119	[174] ACKERSTAFF	99E	OPAL	1991–1995 LEP runs
0.097±0.005±0.011	f&a	419	GIBAUT	94B	CLEO	E_{cm}^{ee}= 10.6 GeV
0.102±0.029	f&a	13	BYLSMA	87	HRS	E_{cm}^{ee}= 29 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.26 ±0.06 ±0.05			ACTON	92H	OPAL	E_{cm}^{ee}= 88.2–94.2 GeV
0.10 $^{+0.05}_{-0.04}$ ±0.03			DECAMP	92C	ALEP	1989–1990 LEP runs
0.16 ±0.13 ±0.04			BEHREND	89B	CELL	E_{cm}^{ee}= 14–47 GeV
0.3 ±0.1 ±0.2			BARTEL	85F	JADE	E_{cm}^{ee}= 34.6 GeV
0.13 ±0.04		10	BELTRAMI	85	HRS	Repl. by BYLSMA 87
0.16 ±0.08 ±0.04		4	BURCHAT	85	MRK2	E_{cm}^{ee}= 29 GeV
1.0 ±0.4		10	BEHREND	82	CELL	Repl. by BEHREND 89B

[172] The correlation coefficients between this meausrement and the ABREU 01M measurements of B($\tau \to$ 1-prong) and B($\tau \to$ 3-prong) are −0.08 and −0.08 respectively.

[173] The correlation coefficients between this measurement and the ACHARD 01D measurements of B($\tau \to$ "1-prong") and B($\tau \to$ "3-prong") are −0.082 and −0.19 respectively.

[174] Not independent of ACKERSTAFF 99E B($\tau^- \to 3h^-2h^+\nu_\tau$(ex. K^0)) and B($\tau^- \to 3h^-2h^+\pi^0\nu_\tau$(ex. K^0)) measurements.

$\Gamma(3h^-2h^+\nu_\tau$(ex.K^0))$/\Gamma_{total}$ Γ_{97}/Γ

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
0.082±0.006 OUR FIT					
0.076±0.007 OUR AVERAGE					
0.091±0.014±0.006	97	ACKERSTAFF	99E	OPAL	1991–1995 LEP runs
0.080±0.011±0.013	58	BUSKULIC	96	ALEP	LEP 1991–1993 data
0.077±0.005±0.009	295	GIBAUT	94B	CLEO	E_{cm}^{ee}= 10.6 GeV
0.064±0.023±0.01	12	ALBRECHT	88B	ARG	E_{cm}^{ee}= 10 GeV
0.051±0.020	7	BYLSMA	87	HRS	E_{cm}^{ee}= 29 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.067±0.030	5	[175] BELTRAMI	85	HRS	Repl. by BYLSMA 87

[175] The error quoted is statistical only.

$\Gamma(3h^-2h^+\pi^0\nu_\tau(\text{ex.}K^0))/\Gamma_{\text{total}}$ Γ_{98}/Γ

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
0.0181±0.0027 OUR FIT					
0.0172±0.0027 OUR AVERAGE					
0.017 ±0.002 ±0.002	231	ANASTASSOV	01	CLEO	E_{cm}^{ee}= 10.6 GeV
0.027 ±0.018 ±0.009	23	ACKERSTAFF	99E	OPAL	1991–1995 LEP runs
0.018 ±0.007 ±0.012	18	BUSKULIC	96	ALEP	LEP 1991–1993 data
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.019 ±0.004 ±0.004	31	GIBAUT	94B	CLEO	Repl. by ANASTASSOV 01
0.051 ±0.022	6	BYLSMA	87	HRS	E_{cm}^{ee}= 29 GeV
0.067 ±0.030	5	[176] BELTRAMI	85	HRS	Repl. by BYLSMA 87

[176] The error quoted is statistical only.

$\Gamma(3h^-2h^+2\pi^0\nu_\tau)/\Gamma_{\text{total}}$ Γ_{99}/Γ

VALUE (%)	CL%	DOCUMENT ID		TECN	COMMENT
<0.011	90	GIBAUT	94B	CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma((5\pi)^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_{100}/Γ

$\Gamma_{100}/\Gamma = (\Gamma_{28}+\Gamma_{45}+\Gamma_{74}+\Gamma_{97}+0.553\Gamma_{117}+0.888\Gamma_{138})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		DOCUMENT ID		TECN	COMMENT
0.80±0.07 OUR FIT					
0.61±0.06±0.08	**avg**	[177] GIBAUT	94B	CLEO	E_{cm}^{ee}= 10.6 GeV

[177] Not independent of GIBAUT 94B B($3h^-2h^+\nu_\tau$), PROCARIO 93 B($h^-4\pi^0\nu_\tau$), and BORTOLETTO 93 B($2h^-h^+2\pi^0\nu_\tau$)/B("3prong") measurements. Result is corrected for η contributions.

$\Gamma(4h^-3h^+\geq 0\text{ neutrals }\nu_\tau(\text{"7-prong"}))/\Gamma_{\text{total}}$ Γ_{101}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<2.4\times10^{-6}$	90	EDWARDS	97B	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<1.8\times10^{-5}$	95	ACKERSTAFF	97J	OPAL	1990–1995 LEP runs
$<2.9\times10^{-4}$	90	BYLSMA	87	HRS	E_{cm}^{ee}= 29 GeV

$\Gamma(X^-(S=-1)\nu_\tau)/\Gamma_{\text{total}}$

$\Gamma_{102}/\Gamma = (\Gamma_{10}+\Gamma_{15}+\Gamma_{22}+\Gamma_{26}+\Gamma_{33}+\Gamma_{38}+\Gamma_{82}+\Gamma_{86}+\Gamma_{119})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		DOCUMENT ID		TECN	COMMENT
2.86±0.09 OUR FIT	Error includes scale factor of 1.1.				
2.87±0.12	**avg**	[178] BARATE	99R	ALEP	1991–1995 LEP runs

[178] BARATE 99R perform a combined analysis of all ALEPH LEP 1 data on τ branching fraction measurements for decay modes having total strangeness equal to -1.

$\Gamma(K^*(892)^- \geq 0 \text{ neutrals } \geq 0 K_L^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{103}/Γ

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
1.42±0.18 OUR AVERAGE	Error includes scale factor of 1.4. See the ideogram below.				
$1.19 \pm 0.15 \,^{+0.13}_{-0.18}$	104	ALBRECHT	95H	ARG	E_{cm}^{ee}= 9.4–10.6 GeV
1.94±0.27±0.15	74	[179] AKERS	94G	OPAL	E_{cm}^{ee}= 88–94 GeV
1.43±0.11±0.13	475	[180] GOLDBERG	90	CLEO	E_{cm}^{ee}= 9.4–10.9 GeV

[179] AKERS 94G reject events in which a K_S^0 accompanies the $K^*(892)^-$. We do not correct for them.

[180] GOLDBERG 90 estimates that 10% of observed $K^*(892)$ are accompanied by a π^0.



$\Gamma(K^*(892)^- \geq 0 \text{ neutrals } \geq 0 K_L^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{103}/Γ

$\Gamma(K^*(892)^- \nu_\tau)/\Gamma_{\text{total}}$ Γ_{104}/Γ

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
1.29 ±0.05 OUR AVERAGE					
1.326±0.063		BARATE	99R	ALEP	1991–1995 LEP runs
1.11 ±0.12		[181] COAN	96	CLEO	$E_{\text{cm}}^{ee} \approx$ 10.6 GeV
1.42 ±0.22 ±0.09		[182] ACCIARRI	95F	L3	1991–1993 LEP runs
$1.23 \pm 0.21 \,^{+0.11}_{-0.21}$	54	[183] ALBRECHT	88L	ARG	E_{cm}^{ee}= 10 GeV
1.9 ±0.3 ±0.4	44	[184] TSCHIRHART	88	HRS	E_{cm}^{ee}= 29 GeV
1.5 ±0.4 ±0.4	15	[185] AIHARA	87C	TPC	E_{cm}^{ee}= 29 GeV
1.3 ±0.3 ±0.3	31	YELTON	86	MRK2	E_{cm}^{ee}= 29 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.39 ±0.09 ±0.10		[186] BUSKULIC	96	ALEP	Repl. by BARATE 99R
1.45 ±0.13 ±0.11	273	[187] BUSKULIC	94F	ALEP	Repl. by BUSKULIC 96
1.7 ±0.7	11	DORFAN	81	MRK2	E_{cm}^{ee}= 4.2–6.7 GeV

[181] Not independent of COAN 96 $B(\pi^- \overline{K}^0 \nu_\tau)$ and BATTLE 94 $B(K^- \pi^0 \nu_\tau)$ measurements. $K\pi$ final states are consistent with and assumed to originate from $K^*(892)^-$ production.

[182] This result is obtained from their $B(\pi^- \overline{K}^0 \nu_\tau)$ assuming all those decays originate in $K^*(892)^-$ decays.

[183] The authors divide by $\Gamma_2/\Gamma = 0.865$ to obtain this result.

[184] Not independent of TSCHIRHART 88 $\Gamma(\tau^- \to h^- \overline{K}^0 \geq 0$ neutrals $\geq 0 K^0_L \nu_\tau)/\Gamma$(total).

[185] Decay π^- identified in this experiment, is assumed in the others.

[186] Not independent of BUSKULIC 96 $B(\pi^- \overline{K}^0 \nu_\tau)$ and $B(K^- \pi^0 \nu_\tau)$ measurements.

[187] BUSKULIC 94F obtain this result from BUSKULIC 94F $B(\overline{K}^0 \pi^- \nu_\tau)$ and BUSKULIC 94E $B(K^- \pi^0 \nu_\tau)$ assuming all of those decays originate in $K^*(892)^-$ decays.

$\Gamma(K^*(892)^- \nu_\tau)/\Gamma(\pi^- \pi^0 \nu_\tau)$ Γ_{104}/Γ_{13}

VALUE	DOCUMENT ID		TECN	COMMENT
0.075±0.027	[188] ABREU	94K	DLPH	LEP 1992 Z data

[188] ABREU 94K quote $B(\tau^- \to K^*(892)^- \nu_\tau) B(K^*(892)^- \to K^- \pi^0)/B(\tau^- \to \rho^- \nu_\tau) = 0.025 \pm 0.009$. We divide by $B(K^*(892)^- \to K^- \pi^0) = 0.333$ to obtain this result.

$\Gamma(K^*(892)^0 K^- \geq 0$ neutrals $\nu_\tau)/\Gamma_{total}$ Γ_{105}/Γ

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
0.32±0.08±0.12	119	GOLDBERG	90	CLEO	E^{ee}_{cm}= 9.4–10.9 GeV

$\Gamma(K^*(892)^0 K^- \nu_\tau)/\Gamma_{total}$ Γ_{106}/Γ

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
0.21 ±0.04 OUR AVERAGE					
0.213±0.048		[189] BARATE	98	ALEP	1991–1995 LEP runs
0.20 ±0.05 ±0.04	47	ALBRECHT	95H	ARG	E^{ee}_{cm}= 9.4–10.6 GeV

[189] BARATE 98 measure the K^- ($\rho^0 \to \pi^+ \pi^-$) fraction in $\tau^- \to K^- \pi^+ \pi^- \nu_\tau$ decays to be (35 ± 11)% and derive this result from their measurement of $\Gamma(\tau^- \to K^- \pi^+ \pi^- \nu_\tau)/\Gamma_{total}$ assuming the intermediate states are all $K^- \rho$ and $K^- K^*(892)^0$.

$\Gamma(\overline{K}^*(892)^0 \pi^- \geq 0$ neutrals $\nu_\tau)/\Gamma_{total}$ Γ_{107}/Γ

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
0.38±0.11±0.13	105	GOLDBERG	90	CLEO	E^{ee}_{cm}= 9.4–10.9 GeV

$\Gamma(\overline{K}^*(892)^0 \pi^- \nu_\tau)/\Gamma_{total}$ Γ_{108}/Γ

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
0.22 ±0.05 OUR AVERAGE					
0.209±0.058		[190] BARATE	98	ALEP	1991–1995 LEP runs
0.25 ±0.10 ±0.05	27	ALBRECHT	95H	ARG	E^{ee}_{cm}= 9.4–10.6 GeV

[190] BARATE 98 measure the $K^- K^*(892)^0$ fraction in $\tau^- \to K^- K^+ \pi^- \nu_\tau$ decays to be (87 ± 13)% and derive this result from their measurement of $\Gamma(\tau^- \to K^- K^+ \pi^- \nu_\tau)/\Gamma_{total}$.

$\Gamma((\overline{K}^*(892)\pi)^-\nu_\tau \to \pi^-\overline{K}^0\pi^0\nu_\tau)/\Gamma_{total}$ Γ_{109}/Γ

VALUE (%)	DOCUMENT ID		TECN	COMMENT
0.10 ±0.04 OUR AVERAGE				
0.097±0.044±0.036	[191] BARATE	99K	ALEP	1991–1995 LEP runs
0.106±0.037±0.032	[192] BARATE	98E	ALEP	1991–1995 LEP runs

[191] BARATE 99K measure K^0's by detecting K^0_L's in their hadron calorimeter. They determine the $\overline{K}^0\rho^-$ fraction in $\tau^- \to \pi^-\overline{K}^0\pi^0\nu_\tau$ decays to be (0.72 ± 0.12 ± 0.10) and multiply their B($\pi^-\overline{K}^0\pi^0\nu_\tau$) measurement by one minus this fraction to obtain the quoted result.

[192] BARATE 98E reconstruct K^0's using $K^0_S \to \pi^+\pi^-$ decays. They determine the $\overline{K}^0\rho^-$ fraction in $\tau^- \to \pi^-\overline{K}^0\pi^0\nu_\tau$ decays to be (0.64 ± 0.09 ± 0.10) and multiply their B($\pi^-\overline{K}^0\pi^0\nu_\tau$) measurement by one minus this fraction to obtain the quoted result.

$\Gamma(K_1(1270)^-\nu_\tau)/\Gamma_{total}$ Γ_{110}/Γ

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
0.47±0.11 OUR AVERAGE					
0.48±0.11		BARATE	99R	ALEP	1991–1995 LEP runs
$0.41^{+0.41}_{-0.35}\pm0.10$	5	[193] BAUER	94	TPC	E^{ee}_{cm}= 29 GeV

[193] We multiply 0.41% by 0.25, the relative systematic error quoted by BAUER 94, to obtain the systematic error.

$\Gamma(K_1(1400)^-\nu_\tau)/\Gamma_{total}$ Γ_{111}/Γ

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
0.17±0.26 OUR AVERAGE Error includes scale factor of 1.7.					
0.05±0.17		BARATE	99R	ALEP	1991–1995 LEP runs
$0.76^{+0.40}_{-0.33}\pm0.20$	11	[194] BAUER	94	TPC	E^{ee}_{cm}= 29 GeV

[194] We multiply 0.76% by 0.25, the relative systematic error quoted by BAUER 94, to obtain the systematic error.

$[\Gamma(K_1(1270)^-\nu_\tau)+\Gamma(K_1(1400)^-\nu_\tau)]/\Gamma_{total}$ $(\Gamma_{110}+\Gamma_{111})/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
$\mathbf{1.17^{+0.41}_{-0.37}\pm0.29}$	16	[195] BAUER	94	TPC	E^{ee}_{cm}= 29 GeV

[195] We multiply 1.17% by 0.25, the relative systematic error quoted by BAUER 94, to obtain the systematic error. Not independent of BAUER 94 B($K_1(1270)^-\nu_\tau$) and BAUER 94 B($K_1(1400)^-\nu_\tau$) measurements.

$\Gamma(K_1(1270)^-\nu_\tau)/[\Gamma(K_1(1270)^-\nu_\tau)+\Gamma(K_1(1400)^-\nu_\tau)]$ $\Gamma_{110}/(\Gamma_{110}+\Gamma_{111})$

VALUE	DOCUMENT ID		TECN	COMMENT
0.69±0.15 OUR AVERAGE				
0.71±0.16±0.11	[196] ABBIENDI	00D	OPAL	1990–1995 LEP runs
0.66±0.19±0.13	[197] ASNER	00B	CLEO	E^{ee}_{cm}= 10.6 GeV

[196] ABBIENDI 00D assume the resonance structure of $\tau^- \to K^-\pi^+\pi^-\nu_\tau$ decays is dominated by the $K_1(1270)^-$ and $K_1(1400)^-$ resonances.

[197] ASNER 00B assume the resonance structure of $\tau^- \to K^-\pi^+\pi^-\nu_\tau$ (ex. K^0) decays is dominated by $K_1(1270)^-$ and $K_1(1400)^-$ resonances.

$\Gamma(K^*(1410)^-\nu_\tau)/\Gamma_{total}$ Γ_{112}/Γ

VALUE (units 10^{-3})	DOCUMENT ID		TECN	COMMENT
$1.5^{+1.4}_{-1.0}$	BARATE	99R	ALEP	1991–1995 LEP runs

$\Gamma(K_0^*(1430)^-\nu_\tau)/\Gamma_{total}$ Γ_{113}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID		TECN	COMMENT
<0.5	95	BARATE	99R	ALEP	1991–1995 LEP runs

$\Gamma(K_2^*(1430)^-\nu_\tau)/\Gamma_{total}$ Γ_{114}/Γ

VALUE (%)	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
<0.3	95		TSCHIRHART	88	HRS	E_{cm}^{ee}= 29 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.33	95		[198] ACCIARRI	95F	L3	1991–1993 LEP runs
<0.9	95	0	DORFAN	81	MRK2	E_{cm}^{ee}= 4.2–6.7 GeV

[198] ACCIARRI 95F quote $B(\tau^- \to K^*(1430)^- \to \pi^- \overline{K}^0 \nu_\tau) < 0.11\%$. We divide by $B(K^*(1430)^- \to \pi^- \overline{K}^0) = 0.33$ to obtain the limit shown.

$\Gamma(a_0(980)^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{total} \times B(a_0(980) \to K^0 K^-)$ $\Gamma_{115}/\Gamma \times B$

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<2.8	90	GOLDBERG	90	CLEO	E_{cm}^{ee}= 9.4–10.9 GeV

$\Gamma(\eta\pi^-\nu_\tau)/\Gamma_{total}$ Γ_{116}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
< 1.4	95	0	BARTELT	96	CLEO	$E_{cm}^{ee} \approx$ 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 6.2	95		BUSKULIC	97C	ALEP	1991–1994 LEP runs
< 3.4	95		ARTUSO	92	CLEO	$E_{cm}^{ee} \approx$ 10.6 GeV
< 90	95		ALBRECHT	88M	ARG	$E_{cm}^{ee} \approx$ 10 GeV
<140	90		BEHREND	88	CELL	E_{cm}^{ee}= 14–46.8 GeV
<180	95		BARINGER	87	CLEO	E_{cm}^{ee}= 10.5 GeV
<250	90	0	COFFMAN	87	MRK3	E_{cm}^{ee}= 3.77 GeV
510 ±100±120		65	DERRICK	87	HRS	E_{cm}^{ee}= 29 GeV
<100	95		GAN	87B	MRK2	E_{cm}^{ee}= 29 GeV

$\Gamma(\eta\pi^-\pi^0\nu_\tau)/\Gamma_{total}$ Γ_{117}/Γ

VALUE (%)	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
0.174±0.024 OUR FIT						
0.173±0.024 OUR AVERAGE						
0.18 ±0.04 ±0.02			BUSKULIC	97C	ALEP	1991–1994 LEP runs
0.17 ±0.02 ±0.02		125	ARTUSO	92	CLEO	$E_{cm}^{ee} \approx$ 10.6 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.10		95		ALBRECHT	88M	ARG	$E_{cm}^{ee} \approx 10$ GeV
<2.10		95		BARINGER	87	CLEO	E_{cm}^{ee}= 10.5 GeV
4.20	$^{+0.70}_{-1.20}$ ±1.60		[199]	GAN	87	MRK2	E_{cm}^{ee}= 29 GeV

[199] Highly correlated with GAN 87 $\Gamma(\pi^- 3\pi^0 \nu_\tau)/\Gamma$(total) value.

$\Gamma(\eta \pi^- \pi^0 \pi^0 \nu_\tau)/\Gamma_{total}$ Γ_{118}/Γ

VALUE (units 10^{-4})	CL%	EVTS		DOCUMENT ID		TECN	COMMENT
1.5±0.5		30	[200]	ANASTASSOV	01	CLEO	E_{cm}^{ee}= 10.6 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.4±0.6±0.3	15	[201]	BERGFELD	97	CLEO	Repl. by ANASTASSOV 01
< 4.3	95		ARTUSO	92	CLEO	$E_{cm}^{ee} \approx$ 10.6 GeV
<120	95		ALBRECHT	88M	ARG	$E_{cm}^{ee} \approx$ 10 GeV

[200] Weighted average of BERGFELD 97 and ANASTASSOV 01 value of $(1.5 \pm 0.6 \pm 0.3) \times 10^{-4}$ obtained using η's reconstructed from $\eta \to \pi^+ \pi^- \pi^0$ decays.

[201] BERGFELD 97 reconstruct η's using $\eta \to \gamma\gamma$ decays.

$\Gamma(\eta K^- \nu_\tau)/\Gamma_{total}$ Γ_{119}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
2.7±0.6 OUR FIT						
2.7±0.6 OUR AVERAGE						
$2.9^{+1.3}_{-1.2}$±0.7			BUSKULIC	97C	ALEP	1991–1994 LEP runs
2.6±0.5±0.5	85		BARTELT	96	CLEO	$E_{cm}^{ee} \approx$ 10.6 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4.7	95	ARTUSO	92	CLEO	$E_{cm}^{ee} \approx$ 10.6 GeV

$\Gamma(\eta K^*(892)^- \nu_\tau)/\Gamma_{total}$ Γ_{120}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
2.90±0.80±0.42	25	BISHAI	99	CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(\eta K^- \pi^0 \nu_\tau)/\Gamma_{total}$ Γ_{121}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
1.77±0.56±0.71	36	BISHAI	99	CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(\eta \overline{K}^0 \pi^- \nu_\tau)/\Gamma_{total}$ Γ_{122}/Γ

VALUE (units 10^{-4})	EVTS		DOCUMENT ID		TECN	COMMENT
2.20±0.70±0.22	15	[202]	BISHAI	99	CLEO	E_{cm}^{ee}= 10.6 GeV

[202] We multiply the BISHAI 99 measurement $B(\tau^- \to \eta K_S^0 \pi^- \nu_\tau) = (1.10 \pm 0.35 \pm 0.11) \times 10^{-4}$ by 2 to obtain the listed value.

$\Gamma(\eta \pi^+ \pi^- \pi^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{total}$ Γ_{123}/Γ

VALUE (%)	CL%	DOCUMENT ID		TECN	COMMENT
<0.3	90	ABACHI	87B	HRS	E_{cm}^{ee}= 29 GeV

$\Gamma(\eta\pi^-\pi^+\pi^-\nu_\tau)/\Gamma_{total}$ Γ_{124}/Γ

VALUE (units 10^{-4})	EVTS		DOCUMENT ID		TECN	COMMENT
2.3 ± 0.5	170	[203]	ANASTASSOV	01	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$3.4^{+0.6}_{-0.5}\pm0.6$	89	[204]	BERGFELD	97	CLEO	Repl. by ANASTASSOV 01

[203] Weighted average of BERGFELD 97 and ANASTASSOV 01 measurements using η's reconstructed from $\eta \to \pi^+\pi^-\pi^0$ and $\eta \to 3\pi^0$ decays.
[204] BERGFELD 97 reconstruct η's using $\eta \to \gamma\gamma$ and $\eta \to 3\pi^0$ decays.

$\Gamma(\eta a_1(1260)^-\nu_\tau \to \eta\pi^-\rho^0\nu_\tau)/\Gamma_{total}$ Γ_{125}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<3.9\times10^{-4}$	90	BERGFELD	97	CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(\eta\eta\pi^-\nu_\tau)/\Gamma_{total}$ Γ_{126}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
< 1.1	95	ARTUSO	92	CLEO	$E_{cm}^{ee}\approx$ 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<83	95	ALBRECHT	88M	ARG	$E_{cm}^{ee}\approx$ 10 GeV

$\Gamma(\eta\eta\pi^-\pi^0\nu_\tau)/\Gamma_{total}$ Γ_{127}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
< 2.0	95	ARTUSO	92	CLEO	$E_{cm}^{ee}\approx$ 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<90	95	ALBRECHT	88M	ARG	$E_{cm}^{ee}\approx$ 10 GeV

$\Gamma(\eta'(958)\pi^-\nu_\tau)/\Gamma_{total}$ Γ_{128}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<7.4\times10^{-5}$	90	BERGFELD	97	CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(\eta'(958)\pi^-\pi^0\nu_\tau)/\Gamma_{total}$ Γ_{129}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<8.0\times10^{-5}$	90	BERGFELD	97	CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(\phi\pi^-\nu_\tau)/\Gamma_{total}$ Γ_{130}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<2.0\times10^{-4}$	90	[205] AVERY	97	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<3.5\times10^{-4}$	90	ALBRECHT	95H	ARG	E_{cm}^{ee}= 9.4–10.6 GeV

[205] AVERY 97 limit varies from (1.2–2.0) $\times10^{-4}$ depending on decay model assumptions.

$\Gamma(\phi K^-\nu_\tau)/\Gamma_{total}$ Γ_{131}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<6.7\times10^{-5}$	90	[206] AVERY	97	CLEO	E_{cm}^{ee}= 10.6 GeV

[206] AVERY 97 limit varies from (5.4–6.7) $\times10^{-5}$ depending on decay model assumptions.

$\Gamma(f_1(1285)\pi^-\nu_\tau)/\Gamma_{total}$ Γ_{132}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
$5.8^{+1.4}_{-1.3}\pm1.8$	54	BERGFELD	97	CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(f_1(1285)\pi^-\nu_\tau \to \eta\pi^-\pi^+\pi^-\nu_\tau)/\Gamma(\eta\pi^-\pi^+\pi^-\nu_\tau)$ $\Gamma_{133}/\Gamma_{124}$

VALUE	DOCUMENT ID		TECN	COMMENT
0.55±0.14	BERGFELD	97	CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(\pi(1300)^-\nu_\tau \to (\rho\pi)^-\nu_\tau \to (3\pi)^-\nu_\tau)/\Gamma_{total}$ Γ_{134}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.0\times10^{-4}$	90	ASNER	00	CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(\pi(1300)^-\nu_\tau \to ((\pi\pi)_{S\text{-wave}}\pi)^-\nu_\tau \to (3\pi)^-\nu_\tau)/\Gamma_{total}$ Γ_{135}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.9\times10^{-4}$	90	ASNER	00	CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(h^-\omega \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{total}$ Γ_{136}/Γ

$\Gamma_{136}/\Gamma = (\Gamma_{137}+\Gamma_{138})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID		TECN	COMMENT
2.37±0.08 OUR FIT						
1.65±0.3 ±0.2	**avg**	1513	ALBRECHT	88M	ARG	$E_{cm}^{ee}\approx$ 10 GeV

$\Gamma(h^-\omega\nu_\tau)/\Gamma_{total}$ Γ_{137}/Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID		TECN	COMMENT
1.94±0.07 OUR FIT						
1.92±0.07 OUR AVERAGE						
1.91±0.07±0.06	f&a	5803	BUSKULIC	97C	ALEP	1991–1994 LEP
1.95±0.07±0.11	avg	2223	[207] BALEST	95C	CLEO	E_{cm}^{ee} (runs) ≈ 10.6 GeV
1.60±0.27±0.41	f&a	139	BARINGER	87	CLEO	E_{cm}^{ee}= 10.5 GeV

[207] Not independent of BALEST 95C $B(\tau^-\to h^-\omega\nu_\tau)/B(\tau^-\to h^-h^-h^+\pi^0\nu_\tau)$ value.

$\Gamma(h^-\omega\nu_\tau)/\Gamma(h^-h^-h^+\pi^0\nu_\tau(\text{ex.}K^0))$ Γ_{137}/Γ_{64}

$\Gamma_{137}/\Gamma_{64} = \Gamma_{137}/(\Gamma_{68}+\Gamma_{86}+\Gamma_{90}+0.231\Gamma_{119}+0.888\Gamma_{137}+0.0221\Gamma_{138})$

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.446±0.015 OUR FIT					
0.453±0.019 OUR AVERAGE					
0.431±0.033	2350	[208] BUSKULIC	96	ALEP	LEP 1991–1993 data
0.464±0.016±0.017	2223	[209] BALEST	95C	CLEO	$E_{cm}^{ee}\approx$ 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.37 ±0.05 ±0.02	458	[210] ALBRECHT	91D	ARG	E_{cm}^{ee}= 9.4–10.6 GeV

[208] BUSKULIC 96 quote the fraction of $\tau \to h^- h^- h^+ \pi^0 \nu_\tau$(ex. K^0) decays which originate in a $h^- \omega$ final state = 0.383 ± 0.029. We divide this by the $\omega(782) \to \pi^+ \pi^- \pi^0$ branching fraction (0.888).

[209] BALEST 95C quote the fraction of $\tau^- \to h^- h^- h^+ \pi^0 \nu_\tau$ (ex. K^0) decays which originate in a $h^- \omega$ final state equals 0.412 ± 0.014 ± 0.015. We divide this by the $\omega(782) \to \pi^+ \pi^- \pi^0$ branching fraction (0.888).

[210] ALBRECHT 91D quote the fraction of $\tau^- \to h^- h^- h^+ \pi^0 \nu_\tau$ decays which originate in a $\pi^- \omega$ final state equals $0.33 \pm 0.04 \pm 0.02$. We divide this by the $\omega(782) \to \pi^+ \pi^- \pi^0$ branching fraction (0.888).

$\Gamma(h^- \omega \pi^0 \nu_\tau)/\Gamma_{total}$ Γ_{138}/Γ

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
0.43±0.05 OUR FIT					
0.43±0.06±0.05	7283	BUSKULIC	97C	ALEP	1991–1994 LEP runs

$\Gamma(h^- \omega 2\pi^0 \nu_\tau)/\Gamma_{total}$ Γ_{139}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
1.4 ±0.4 ±0.3	53	ANASTASSOV	01	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$1.89^{+0.74}_{-0.67} \pm 0.40$	19	ANDERSON	97	CLEO	Repl. by ANASTASSOV 01

$\Gamma(h^- \omega \pi^0 \nu_\tau)/\Gamma(h^- h^- h^+ \geq 0 \text{ neutrals } \geq 0 K_L^0 \nu_\tau)$ Γ_{138}/Γ_{52}

$\Gamma_{138}/\Gamma_{52} = \Gamma_{138}/(0.3431\Gamma_{33}+0.3431\Gamma_{35}+0.3431\Gamma_{38}+0.3431\Gamma_{40}+0.4307\Gamma_{45}+0.6861\Gamma_{46}+\Gamma_{60}+\Gamma_{68}+\Gamma_{74}+\Gamma_{75}+\Gamma_{82}+\Gamma_{86}+\Gamma_{89}+\Gamma_{90}+0.285\Gamma_{117}+0.285\Gamma_{119}+0.9101\Gamma_{137}+0.9101\Gamma_{138})$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE		EVTS	DOCUMENT ID		TECN	COMMENT
0.0286±0.0031 OUR FIT						
0.028 ±0.003 ±0.003	**avg**	430	[211] BORTOLETTO	93	CLEO	$E_{cm}^{ee} \approx$ 10.6 GeV

[211] Not independent of BORTOLETTO 93 $\Gamma(\tau^- \to h^- \omega \pi^0 \nu_\tau)/\Gamma(\tau^- \to h^- h^- h^+ 2\pi^0 \nu_\tau$(ex.$K^0$)) value.

$\Gamma(h^- \omega \pi^0 \nu_\tau)/\Gamma(h^- h^- h^+ 2\pi^0 \nu_\tau(\text{ex.}K^0))$ Γ_{138}/Γ_{73}

$\Gamma_{138}/\Gamma_{73} = \Gamma_{138}/(\Gamma_{74}+0.236\Gamma_{117}+0.888\Gamma_{138})$

VALUE	DOCUMENT ID		TECN	COMMENT
0.81±0.08 OUR FIT				
0.81±0.06±0.06	BORTOLETTO	93	CLEO	$E_{cm}^{ee} \approx$ 10.6 GeV

$\Gamma(2h^- h^+ \omega \nu_\tau)/\Gamma_{total}$ Γ_{140}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
1.2 ±0.2 ±0.1	110	ANASTASSOV	01	CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(e^-\gamma)/\Gamma_{total}$ Γ_{141}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<2.7\times10^{-6}$	90	EDWARDS	97	CLEO	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<1.1\times10^{-4}$	90	ABREU	95U	DLPH	1990–1993 LEP runs
$<1.2\times10^{-4}$	90	ALBRECHT	92K	ARG	E_{cm}^{ee}= 10 GeV
$<2.0\times10^{-4}$	90	KEH	88	CBAL	E_{cm}^{ee}= 10 GeV
$<6.4\times10^{-4}$	90	HAYES	82	MRK2	E_{cm}^{ee}= 3.8–6.8 GeV

$\Gamma(\mu^-\gamma)/\Gamma_{total}$ Γ_{142}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.1\times10^{-6}$	90	AHMED	00	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<3.0\times10^{-6}$	90	EDWARDS	97	CLEO	
$<6.2\times10^{-5}$	90	ABREU	95U	DLPH	1990–1993 LEP runs
$<0.42\times10^{-5}$	90	BEAN	93	CLEO	E_{cm}^{ee}= 10.6 GeV
$<3.4\times10^{-5}$	90	ALBRECHT	92K	ARG	E_{cm}^{ee}= 10 GeV
$<55\times10^{-5}$	90	HAYES	82	MRK2	E_{cm}^{ee}= 3.8–6.8 GeV

$\Gamma(e^-\pi^0)/\Gamma_{total}$ Γ_{143}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<3.7\times10^{-6}$	90	BONVICINI	97	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<17\times10^{-5}$	90	ALBRECHT	92K	ARG	E_{cm}^{ee}= 10 GeV
$<14\times10^{-5}$	90	KEH	88	CBAL	E_{cm}^{ee}= 10 GeV
$<210\times10^{-5}$	90	HAYES	82	MRK2	E_{cm}^{ee}= 3.8–6.8 GeV

$\Gamma(\mu^-\pi^0)/\Gamma_{total}$ Γ_{144}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<4.0\times10^{-6}$	90	BONVICINI	97	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<4.4\times10^{-5}$	90	ALBRECHT	92K	ARG	E_{cm}^{ee}= 10 GeV
$<82\times10^{-5}$	90	HAYES	82	MRK2	E_{cm}^{ee}= 3.8–6.8 GeV

$\Gamma(e^-K_S^0)/\Gamma_{total}$ Γ_{145}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<9.1\times10^{-7}$	90	CHEN	02C	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<1.3\times10^{-3}$	90	HAYES	82	MRK2	E_{cm}^{ee}= 3.8–6.8 GeV

$\Gamma(\mu^- K^0_S)/\Gamma_{total}$ Γ_{146}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<9.5 \times 10^{-7}$	90	CHEN	02C	CLEO	E^{ee}_{cm}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<1.0 \times 10^{-3}$	90	HAYES	82	MRK2	E^{ee}_{cm}= 3.8–6.8 GeV

$\Gamma(e^- \eta)/\Gamma_{total}$ Γ_{147}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$< 8.2 \times 10^{-6}$	90	BONVICINI	97	CLEO	E^{ee}_{cm}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$< 6.3 \times 10^{-5}$	90	ALBRECHT	92K	ARG	E^{ee}_{cm}= 10 GeV
$<24 \times 10^{-5}$	90	KEH	88	CBAL	E^{ee}_{cm}= 10 GeV

$\Gamma(\mu^- \eta)/\Gamma_{total}$ Γ_{148}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<9.6 \times 10^{-6}$	90	BONVICINI	97	CLEO	E^{ee}_{cm}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<7.3 \times 10^{-5}$	90	ALBRECHT	92K	ARG	E^{ee}_{cm}= 10 GeV

$\Gamma(e^- \rho^0)/\Gamma_{total}$ Γ_{149}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$< 2.0 \times 10^{-6}$	90	BLISS	98	CLEO	E^{ee}_{cm}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$< 0.42 \times 10^{-5}$	90	[212] BARTELT	94	CLEO	Repl. by BLISS 98
$< 1.9 \times 10^{-5}$	90	ALBRECHT	92K	ARG	E^{ee}_{cm}= 10 GeV
$<37 \times 10^{-5}$	90	HAYES	82	MRK2	E^{ee}_{cm}= 3.8–6.8 GeV

[212] BARTELT 94 assume phase space decays.

$\Gamma(\mu^- \rho^0)/\Gamma_{total}$ Γ_{150}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$< 6.3 \times 10^{-6}$	90	BLISS	98	CLEO	E^{ee}_{cm}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$< 0.57 \times 10^{-5}$	90	[213] BARTELT	94	CLEO	Repl. by BLISS 98
$< 2.9 \times 10^{-5}$	90	ALBRECHT	92K	ARG	E^{ee}_{cm}= 10 GeV
$<44 \times 10^{-5}$	90	HAYES	82	MRK2	E^{ee}_{cm}= 3.8–6.8 GeV

[213] BARTELT 94 assume phase space decays.

$\Gamma(e^- K^*(892)^0)/\Gamma_{total}$ Γ_{151}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<5.1 $\times 10^{-6}$	90	BLISS	98	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<0.63 \times 10^{-5}$	90	[214] BARTELT	94	CLEO	Repl. by BLISS 98
$<3.8 \times 10^{-5}$	90	ALBRECHT	92K	ARG	E_{cm}^{ee}= 10 GeV

[214] BARTELT 94 assume phase space decays.

$\Gamma(\mu^- K^*(892)^0)/\Gamma_{total}$ Γ_{152}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<7.5 $\times 10^{-6}$	90	BLISS	98	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<0.94 \times 10^{-5}$	90	[215] BARTELT	94	CLEO	Repl. by BLISS 98
$<4.5 \times 10^{-5}$	90	ALBRECHT	92K	ARG	E_{cm}^{ee}= 10 GeV

[215] BARTELT 94 assume phase space decays.

$\Gamma(e^- \overline{K}^*(892)^0)/\Gamma_{total}$ Γ_{153}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<7.4 $\times 10^{-6}$	90	BLISS	98	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<1.1 \times 10^{-5}$	90	[216] BARTELT	94	CLEO	Repl. by BLISS 98

[216] BARTELT 94 assume phase space decays.

$\Gamma(\mu^- \overline{K}^*(892)^0)/\Gamma_{total}$ Γ_{154}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<7.5 $\times 10^{-6}$	90	BLISS	98	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<0.87 \times 10^{-5}$	90	[217] BARTELT	94	CLEO	Repl. by BLISS 98

[217] BARTELT 94 assume phase space decays.

$\Gamma(e^- \phi)/\Gamma_{total}$ Γ_{155}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<6.9 $\times 10^{-6}$	90	BLISS	98	CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(\mu^- \phi)/\Gamma_{total}$ Γ_{156}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<7.0 $\times 10^{-6}$	90	BLISS	98	CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(e^- e^+ e^-)/\Gamma_{total}$ Γ_{157}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$< 2.9 \times 10^{-6}$	90	BLISS	98	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$< 0.33 \times 10^{-5}$	90	[218] BARTELT	94	CLEO	Repl. by BLISS 98
$< 1.3 \times 10^{-5}$	90	ALBRECHT	92K	ARG	E_{cm}^{ee}= 10 GeV
$< 2.7 \times 10^{-5}$	90	BOWCOCK	90	CLEO	E_{cm}^{ee}= 10.4–10.9
$<40 \times 10^{-5}$	90	HAYES	82	MRK2	E_{cm}^{ee}= 3.8–6.8 GeV

[218] BARTELT 94 assume phase space decays.

$\Gamma(e^- \mu^+ \mu^-)/\Gamma_{total}$ Γ_{158}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$< 1.8 \times 10^{-6}$	90	BLISS	98	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$< 0.36 \times 10^{-5}$	90	[219] BARTELT	94	CLEO	Repl. by BLISS 98
$< 1.9 \times 10^{-5}$	90	ALBRECHT	92K	ARG	E_{cm}^{ee}= 10 GeV
$< 2.7 \times 10^{-5}$	90	BOWCOCK	90	CLEO	E_{cm}^{ee}= 10.4–10.9
$<33 \times 10^{-5}$	90	HAYES	82	MRK2	E_{cm}^{ee}= 3.8–6.8 GeV

[219] BARTELT 94 assume phase space decays.

$\Gamma(e^+ \mu^- \mu^-)/\Gamma_{total}$ Γ_{159}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.5 \times 10^{-6}$	90	BLISS	98	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<0.35 \times 10^{-5}$	90	[220] BARTELT	94	CLEO	Repl. by BLISS 98
$<1.8 \times 10^{-5}$	90	ALBRECHT	92K	ARG	E_{cm}^{ee}= 10 GeV
$<1.6 \times 10^{-5}$	90	BOWCOCK	90	CLEO	E_{cm}^{ee}= 10.4–10.9

[220] BARTELT 94 assume phase space decays.

$\Gamma(\mu^- e^+ e^-)/\Gamma_{total}$ Γ_{160}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$< 1.7 \times 10^{-6}$	90	BLISS	98	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$< 0.34 \times 10^{-5}$	90	[221] BARTELT	94	CLEO	Repl. by BLISS 98
$< 1.4 \times 10^{-5}$	90	ALBRECHT	92K	ARG	E_{cm}^{ee}= 10 GeV
$< 2.7 \times 10^{-5}$	90	BOWCOCK	90	CLEO	E_{cm}^{ee}= 10.4–10.9
$<44 \times 10^{-5}$	90	HAYES	82	MRK2	E_{cm}^{ee}= 3.8–6.8 GeV

[221] BARTELT 94 assume phase space decays.

$\Gamma(\mu^+ e^- e^-)/\Gamma_{total}$ Γ_{161}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<1.5 $\times 10^{-6}$	90	BLISS	98	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<0.34 \times 10^{-5}$	90	[222] BARTELT	94	CLEO	Repl. by BLISS 98
$<1.4 \times 10^{-5}$	90	ALBRECHT	92K	ARG	E_{cm}^{ee}= 10 GeV
$<1.6 \times 10^{-5}$	90	BOWCOCK	90	CLEO	E_{cm}^{ee}= 10.4–10.9

[222] BARTELT 94 assume phase space decays.

$\Gamma(\mu^- \mu^+ \mu^-)/\Gamma_{total}$ Γ_{162}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
< 1.9 $\times 10^{-6}$	90	BLISS	98	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$< 0.43 \times 10^{-5}$	90	[223] BARTELT	94	CLEO	Repl. by BLISS 98
$< 1.9 \times 10^{-5}$	90	ALBRECHT	92K	ARG	E_{cm}^{ee}= 10 GeV
$< 1.7 \times 10^{-5}$	90	BOWCOCK	90	CLEO	E_{cm}^{ee}= 10.4–10.9
$<49 \times 10^{-5}$	90	HAYES	82	MRK2	E_{cm}^{ee}= 3.8–6.8 GeV

[223] BARTELT 94 assume phase space decays.

$\Gamma(e^- \pi^+ \pi^-)/\Gamma_{total}$ Γ_{163}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<2.2 $\times 10^{-6}$	90	BLISS	98	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<0.44 \times 10^{-5}$	90	[224] BARTELT	94	CLEO	Repl. by BLISS 98
$<2.7 \times 10^{-5}$	90	ALBRECHT	92K	ARG	E_{cm}^{ee}= 10 GeV
$<6.0 \times 10^{-5}$	90	BOWCOCK	90	CLEO	E_{cm}^{ee}= 10.4–10.9

[224] BARTELT 94 assume phase space decays.

$\Gamma(e^+ \pi^- \pi^-)/\Gamma_{total}$ Γ_{164}/Γ

Test of lepton number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<1.9 $\times 10^{-6}$	90	BLISS	98	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<0.44 \times 10^{-5}$	90	[225] BARTELT	94	CLEO	Repl. by BLISS 98
$<1.8 \times 10^{-5}$	90	ALBRECHT	92K	ARG	E_{cm}^{ee}= 10 GeV
$<1.7 \times 10^{-5}$	90	BOWCOCK	90	CLEO	E_{cm}^{ee}= 10.4–10.9

[225] BARTELT 94 assume phase space decays.

$\Gamma(\mu^- \pi^+ \pi^-)/\Gamma_{total}$ Γ_{165}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<8.2 $\times 10^{-6}$	90	BLISS	98	CLEO	E_{cm}^{ee}= 10.6 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.74 \times 10^{-5}$	90	[226]	BARTELT	94	CLEO	Repl. by BLISS 98
$<3.6 \times 10^{-5}$	90		ALBRECHT	92K	ARG	E_{cm}^{ee}= 10 GeV
$<3.9 \times 10^{-5}$	90		BOWCOCK	90	CLEO	E_{cm}^{ee}= 10.4–10.9

[226] BARTELT 94 assume phase space decays.

$\Gamma(\mu^+ \pi^- \pi^-)/\Gamma_{total}$ Γ_{166}/Γ

Test of lepton number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<3.4 \times 10^{-6}$	90	BLISS	98	CLEO	E_{cm}^{ee}= 10.6 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.69 \times 10^{-5}$	90	[227]	BARTELT	94	CLEO	Repl. by BLISS 98
$<6.3 \times 10^{-5}$	90		ALBRECHT	92K	ARG	E_{cm}^{ee}= 10 GeV
$<3.9 \times 10^{-5}$	90		BOWCOCK	90	CLEO	E_{cm}^{ee}= 10.4–10.9

[227] BARTELT 94 assume phase space decays.

$\Gamma(e^- \pi^+ K^-)/\Gamma_{total}$ Γ_{167}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<6.4 \times 10^{-6}$	90	BLISS	98	CLEO	E_{cm}^{ee}= 10.6 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.77 \times 10^{-5}$	90	[228]	BARTELT	94	CLEO	Repl. by BLISS 98
$<2.9 \times 10^{-5}$	90		ALBRECHT	92K	ARG	E_{cm}^{ee}= 10 GeV
$<5.8 \times 10^{-5}$	90		BOWCOCK	90	CLEO	E_{cm}^{ee}= 10.4–10.9

[228] BARTELT 94 assume phase space decays.

$\Gamma(e^- \pi^- K^+)/\Gamma_{total}$ Γ_{168}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<3.8 \times 10^{-6}$	90	BLISS	98	CLEO	E_{cm}^{ee}= 10.6 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.46 \times 10^{-5}$	90	[229]	BARTELT	94	CLEO	Repl. by BLISS 98
$<5.8 \times 10^{-5}$	90		BOWCOCK	90	CLEO	E_{cm}^{ee}= 10.4–10.9

[229] BARTELT 94 assume phase space decays.

$\Gamma(e^+ \pi^- K^-)/\Gamma_{total}$ Γ_{169}/Γ

Test of lepton number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<2.1 \times 10^{-6}$	90	BLISS	98	CLEO	E_{cm}^{ee}= 10.6 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.45 \times 10^{-5}$	90	[230]	BARTELT	94	CLEO	Repl. by BLISS 98
$<2.0 \times 10^{-5}$	90		ALBRECHT	92K	ARG	E_{cm}^{ee}= 10 GeV
$<4.9 \times 10^{-5}$	90		BOWCOCK	90	CLEO	E_{cm}^{ee}= 10.4–10.9

[230] BARTELT 94 assume phase space decays.

$\Gamma(e^- K^0_S K^0_S)/\Gamma_{total}$ Γ_{170}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<2.2 \times 10^{-6}$	90	CHEN	02C	CLEO	E^{ee}_{cm}= 10.6 GeV

$\Gamma(e^- K^+ K^-)/\Gamma_{total}$ Γ_{171}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<6.0 \times 10^{-6}$	90	BLISS	98	CLEO	E^{ee}_{cm}= 10.6 GeV

$\Gamma(e^+ K^- K^-)/\Gamma_{total}$ Γ_{172}/Γ

Test of lepton number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<3.8 \times 10^{-6}$	90	BLISS	98	CLEO	E^{ee}_{cm}= 10.6 GeV

$\Gamma(\mu^- \pi^+ K^-)/\Gamma_{total}$ Γ_{173}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$< 7.5 \times 10^{-6}$	90	BLISS	98	CLEO	E^{ee}_{cm}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$< 0.87 \times 10^{-5}$	90	[231] BARTELT	94	CLEO	Repl. by BLISS 98
$<11 \times 10^{-5}$	90	ALBRECHT	92K	ARG	E^{ee}_{cm}= 10 GeV
$< 7.7 \times 10^{-5}$	90	BOWCOCK	90	CLEO	E^{ee}_{cm}= 10.4–10.9

[231] BARTELT 94 assume phase space decays.

$\Gamma(\mu^- \pi^- K^+)/\Gamma_{total}$ Γ_{174}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<7.4 \times 10^{-6}$	90	BLISS	98	CLEO	E^{ee}_{cm}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<1.5 \times 10^{-5}$	90	[232] BARTELT	94	CLEO	Repl. by BLISS 98
$<7.7 \times 10^{-5}$	90	BOWCOCK	90	CLEO	E^{ee}_{cm}= 10.4–10.9

[232] BARTELT 94 assume phase space decays.

$\Gamma(\mu^+ \pi^- K^-)/\Gamma_{total}$ Γ_{175}/Γ

Test of lepton number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<7.0 \times 10^{-6}$	90	BLISS	98	CLEO	E^{ee}_{cm}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<2.0 \times 10^{-5}$	90	[233] BARTELT	94	CLEO	Repl. by BLISS 98
$<5.8 \times 10^{-5}$	90	ALBRECHT	92K	ARG	E^{ee}_{cm}= 10 GeV
$<4.0 \times 10^{-5}$	90	BOWCOCK	90	CLEO	E^{ee}_{cm}= 10.4–10.9

[233] BARTELT 94 assume phase space decays.

$\Gamma(\mu^- K^0_S K^0_S)/\Gamma_{total}$ Γ_{176}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<3.4 \times 10^{-6}$	90	CHEN	02C	CLEO	E^{ee}_{cm}= 10.6 GeV

$\Gamma(\mu^- K^+ K^-)/\Gamma_{total}$ Γ_{177}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<15 \times 10^{-6}$	90	BLISS	98	CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(\mu^+ K^- K^-)/\Gamma_{total}$ Γ_{178}/Γ

Test of lepton number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<6.0 \times 10^{-6}$	90	BLISS	98	CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(e^- \pi^0 \pi^0)/\Gamma_{total}$ Γ_{179}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<6.5 \times 10^{-6}$	90	BONVICINI	97	CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(\mu^- \pi^0 \pi^0)/\Gamma_{total}$ Γ_{180}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<14 \times 10^{-6}$	90	BONVICINI	97	CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(e^- \eta \eta)/\Gamma_{total}$ Γ_{181}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<35 \times 10^{-6}$	90	BONVICINI	97	CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(\mu^- \eta \eta)/\Gamma_{total}$ Γ_{182}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<60 \times 10^{-6}$	90	BONVICINI	97	CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(e^- \pi^0 \eta)/\Gamma_{total}$ Γ_{183}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<24 \times 10^{-6}$	90	BONVICINI	97	CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(\mu^- \pi^0 \eta)/\Gamma_{total}$ Γ_{184}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<22 \times 10^{-6}$	90	BONVICINI	97	CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(\overline{p}\gamma)/\Gamma_{total}$ Γ_{185}/Γ

Test of lepton number and baryon number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$< 3.5 \times 10^{-6}$	90	GODANG	99	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<29 \times 10^{-5}$	90	ALBRECHT	92K	ARG	E_{cm}^{ee}= 10 GeV

$\Gamma(\overline{p}\pi^0)/\Gamma_{total}$ Γ_{186}/Γ

Test of lepton number and baryon number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<15\times10^{-6}$	90	GODANG	99	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<66\times10^{-5}$	90	ALBRECHT	92K	ARG	E_{cm}^{ee}= 10 GeV

$\Gamma(\overline{p}2\pi^0)/\Gamma_{total}$ Γ_{187}/Γ

Test of lepton number and baryon number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<33\times10^{-6}$	90	GODANG	99	CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(\overline{p}\eta)/\Gamma_{total}$ Γ_{188}/Γ

Test of lepton number and baryon number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<8.9\times10^{-6}$	90	GODANG	99	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<130\times10^{-5}$	90	ALBRECHT	92K	ARG	E_{cm}^{ee}= 10 GeV

$\Gamma(\overline{p}\pi^0\eta)/\Gamma_{total}$ Γ_{189}/Γ

Test of lepton number and baryon number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<27\times10^{-6}$	90	GODANG	99	CLEO	E_{cm}^{ee}= 10.6 GeV

$\Gamma(e^-\text{ light boson})/\Gamma(e^-\overline{\nu}_e\nu_\tau)$ Γ_{190}/Γ_5

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.015	95	[234] ALBRECHT	95G	ARG	E_{cm}^{ee}= 9.4–10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.018	95	[235] ALBRECHT	90E	ARG	E_{cm}^{ee}= 9.4–10.6 GeV
<0.040	95	[236] BALTRUSAIT...85		MRK3	E_{cm}^{ee}= 3.77 GeV

[234] ALBRECHT 95G limit holds for bosons with mass < 0.4 GeV. The limit rises to 0.036 for a mass of 1.0 GeV, then falls to 0.006 at the upper mass limit of 1.6 GeV.
[235] ALBRECHT 90E limit applies for spinless boson with mass < 100 MeV, and rises to 0.050 for mass = 500 MeV.
[236] BALTRUSAITIS 85 limit applies for spinless boson with mass < 100 MeV.

$\Gamma(\mu^-\text{ light boson})/\Gamma(e^-\overline{\nu}_e\nu_\tau)$ Γ_{191}/Γ_5

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.026	95	[237] ALBRECHT	95G	ARG	E_{cm}^{ee}= 9.4–10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.033	95	[238] ALBRECHT	90E	ARG	E_{cm}^{ee}= 9.4–10.6 GeV
<0.125	95	[239] BALTRUSAIT...85		MRK3	E_{cm}^{ee}= 3.77 GeV

[237] ALBRECHT 95G limit holds for bosons with mass < 1.3 GeV. The limit rises to 0.034 for a mass of 1.4 GeV, then falls to 0.003 at the upper mass limit of 1.6 GeV.
[238] ALBRECHT 90E limit applies for spinless boson with mass < 100 MeV, and rises to 0.071 for mass = 500 MeV.
[239] BALTRUSAITIS 85 limit applies for spinless boson with mass < 100 MeV.

τ-DECAY PARAMETERS

A REVIEW GOES HERE – Check our WWW List of Reviews

$\rho^\tau(e \text{ or } \mu)$ PARAMETER

($V-A$) theory predicts $\rho = 0.75$.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.745±0.008 OUR FIT					
0.749±0.008 OUR AVERAGE					
0.742±0.014±0.006	81k	HEISTER	01E	ALEP	1991–1995 LEP runs
0.775±0.023±0.020	36k	ABREU	00L	DLPH	1992–1995 runs
0.781±0.028±0.018	46k	ACKERSTAFF	99D	OPAL	1990–1995 LEP runs
0.762±0.035	54k	ACCIARRI	98R	L3	1991–1995 LEP runs
0.731±0.031		[240] ALBRECHT	98	ARG	E_{cm}^{ee}= 9.5–10.6 GeV
0.72 ±0.09 ±0.03		[241] ABE	97O	SLD	1993–1995 SLC runs
0.747±0.010±0.006	55k	ALEXANDER	97F	CLEO	E_{cm}^{ee}= 10.6 GeV
0.79 ±0.10 ±0.10	3732	FORD	87B	MAC	E_{cm}^{ee}= 29 GeV
0.71 ±0.09 ±0.03	1426	BEHRENDS	85	CLEO	e^+e^- near $\Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.735±0.013±0.008	31k	AMMAR	97B	CLEO	Repl. by ALEXANDER 97F
0.794±0.039±0.031	18k	ACCIARRI	96H	L3	Repl. by ACCIARRI 98R
0.732±0.034±0.020	8.2k	[242] ALBRECHT	95	ARG	E_{cm}^{ee}= 9.5–10.6 GeV
0.738±0.038		[243] ALBRECHT	95C	ARG	Repl. by ALBRECHT 98
0.751±0.039±0.022		BUSKULIC	95D	ALEP	Repl. by HEISTER 01E
0.742±0.035±0.020	8000	ALBRECHT	90E	ARG	E_{cm}^{ee}= 9.4–10.6 GeV

[240] Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 98, ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E. ALBRECHT 98 use tau pair events of the type $\tau^-\tau^+ \to (\ell^-\overline{\nu}_\ell\nu_\tau)(\pi^+\pi^0\overline{\nu}_\tau)$, and their charged conjugates.

[241] ABE 97O assume $\eta^\tau = 0$ in their fit. Letting η^τ vary in the fit gives a ρ^τ value of 0.69 ± 0.13 ± 0.05.

[242] Value is from a simultaneous fit for the ρ^τ and η^τ decay parameters to the lepton energy spectrum. Not independent of ALBRECHT 90E $\rho^\tau(e \text{ or } \mu)$ value which assumes η^τ=0. Result is strongly correlated with ALBRECHT 95C.

[243] Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E.

$\rho^\tau(e)$ PARAMETER

($V-A$) theory predicts $\rho = 0.75$.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.747±0.010 OUR FIT					
0.744±0.010 OUR AVERAGE					
0.747±0.019±0.014	44k	HEISTER	01E	ALEP	1991–1995 LEP runs
0.744±0.036±0.037	17k	ABREU	00L	DLPH	1992–1995 runs
0.779±0.047±0.029	25k	ACKERSTAFF	99D	OPAL	1990–1995 LEP runs
0.68 ±0.04 ±0.07		[244] ALBRECHT	98	ARG	E_{cm}^{ee}= 9.5–10.6 GeV
0.71 ±0.14 ±0.05		ABE	97O	SLD	1993–1995 SLC runs
0.747±0.012±0.004	34k	ALEXANDER	97F	CLEO	E_{cm}^{ee}= 10.6 GeV

0.735±0.036±0.020	4.7k	[245]	ALBRECHT	95	ARG	E_{cm}^{ee}= 9.5–10.6 GeV
0.79 ±0.08 ±0.06	3230	[246]	ALBRECHT	93G	ARG	E_{cm}^{ee}= 9.4–10.6 GeV
0.64 ±0.06 ±0.07	2753		JANSSEN	89	CBAL	E_{cm}^{ee}= 9.4–10.6 GeV
0.62 ±0.17 ±0.14	1823		FORD	87B	MAC	E_{cm}^{ee}= 29 GeV
0.60 ±0.13	699		BEHRENDS	85	CLEO	e^+e^- near $\Upsilon(4S)$
0.72 ±0.10 ±0.11	594		BACINO	79B	DLCO	E_{cm}^{ee}= 3.5–7.4 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.732±0.014±0.009	19k		AMMAR	97B	CLEO	Repl. by ALEXANDER 97F
0.793±0.050±0.025			BUSKULIC	95D	ALEP	Repl. by HEISTER 01E
0.747±0.045±0.028	5106		ALBRECHT	90E	ARG	Repl. by ALBRECHT 95

[244] ALBRECHT 98 use tau pair events of the type $\tau^-\tau^+ \to (\ell^-\overline{\nu}_\ell\nu_\tau)(\pi^+\pi^0\overline{\nu}_\tau)$, and their charged conjugates.

[245] ALBRECHT 95 use tau pair events of the type $\tau^-\tau^+ \to (\ell^-\overline{\nu}_\ell\nu_\tau)(h^+h^-h^+(\pi^0)\overline{\nu}_\tau)$ and their charged conjugates.

[246] ALBRECHT 93G use tau pair events of the type $\tau^-\tau^+ \to (\mu^-\overline{\nu}_\mu\nu_\tau)(e^+\nu_e\overline{\nu}_\tau)$ and their charged conjugates.

$\rho^\tau(\mu)$ PARAMETER

($V-A$) theory predicts $\rho = 0.75$.

VALUE	EVTS		DOCUMENT ID		TECN	COMMENT
0.763±0.020 OUR FIT						
0.770±0.022 OUR AVERAGE						
0.776±0.045±0.019	46k		HEISTER	01E	ALEP	1991–1995 LEP runs
0.999±0.098±0.045	22k		ABREU	00L	DLPH	1992–1995 runs
0.777±0.044±0.016	27k		ACKERSTAFF	99D	OPAL	1990–1995 LEP runs
0.69 ±0.06 ±0.06		[247]	ALBRECHT	98	ARG	E_{cm}^{ee}= 9.5–10.6 GeV
0.54 ±0.28 ±0.14			ABE	97O	SLD	1993–1995 SLC runs
0.750±0.017±0.045	22k		ALEXANDER	97F	CLEO	E_{cm}^{ee}= 10.6 GeV
0.76 ±0.07 ±0.08	3230		ALBRECHT	93G	ARG	E_{cm}^{ee}= 9.4–10.6 GeV
0.734±0.055±0.027	3041		ALBRECHT	90E	ARG	E_{cm}^{ee}= 9.4–10.6 GeV
0.89 ±0.14 ±0.08	1909		FORD	87B	MAC	E_{cm}^{ee}= 29 GeV
0.81 ±0.13	727		BEHRENDS	85	CLEO	e^+e^- near $\Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.747±0.048±0.044	13k		AMMAR	97B	CLEO	Repl. by ALEXANDER 97F
0.693±0.057±0.028			BUSKULIC	95D	ALEP	Repl. by HEISTER 01E

[247] ALBRECHT 98 use tau pair events of the type $\tau^-\tau^+ \to (\ell^-\overline{\nu}_\ell\nu_\tau)(\pi^+\pi^0\overline{\nu}_\tau)$, and their charged conjugates.

$\xi^\tau(e \text{ or } \mu)$ PARAMETER

($V-A$) theory predicts $\xi = 1$.

VALUE	EVTS		DOCUMENT ID		TECN	COMMENT
0.985±0.030 OUR FIT						
0.981±0.031 OUR AVERAGE						
0.986±0.068±0.031	81k		HEISTER	01E	ALEP	1991–1995 LEP runs
0.929±0.070±0.030	36k		ABREU	00L	DLPH	1992–1995 runs
0.98 ±0.22 ±0.10	46k		ACKERSTAFF	99D	OPAL	1990–1995 LEP runs
0.70 ±0.16	54k		ACCIARRI	98R	L3	1991–1995 LEP runs
1.03 ±0.11		[248]	ALBRECHT	98	ARG	E_{cm}^{ee}= 9.5–10.6 GeV
1.05 ±0.35 ±0.04		[249]	ABE	97O	SLD	1993–1995 SLC runs
1.007±0.040±0.015	55k		ALEXANDER	97F	CLEO	E_{cm}^{ee}= 10.6 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.94 ±0.21 ±0.07	18k		ACCIARRI	96H	L3	Repl. by ACCIARRI 98R
0.97 ±0.14		[250]	ALBRECHT	95C	ARG	Repl. by ALBRECHT 98
1.18 ±0.15 ±0.16			BUSKULIC	95D	ALEP	Repl. by HEISTER 01E
0.90 ±0.15 ±0.10	3230	[251]	ALBRECHT	93G	ARG	E_{cm}^{ee}= 9.4–10.6 GeV

[248] Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 98, ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E. ALBRECHT 98 use tau pair events of the type $\tau^-\tau^+ \to (\ell^-\overline{\nu}_\ell\nu_\tau)(\pi^+\pi^0\overline{\nu}_\tau)$, and their charged conjugates.

[249] ABE 97O assume $\eta^\tau = 0$ in their fit. Letting η^τ vary in the fit gives a ξ^τ value of 1.02 ± 0.36 ± 0.05.

[250] Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E. ALBRECHT 95C uses events of the type $\tau^-\tau^+ \to (\ell^-\overline{\nu}_\ell\nu_\tau)\,(h^+h^-h^+\overline{\nu}_\tau)$ and their charged conjugates.

[251] ALBRECHT 93G measurement determines $|\xi^\tau|$ for the case $\xi^\tau(e) = \xi^\tau(\mu)$, but the authors point out that other LEP experiments determine the sign to be positive.

$\xi^\tau(e)$ PARAMETER

($V-A$) theory predicts $\xi = 1$.

VALUE	EVTS		DOCUMENT ID	TECN	COMMENT
0.994±0.040 OUR FIT					
1.00 ±0.04 OUR AVERAGE					
1.011±0.094±0.038	44k		HEISTER	01E ALEP	1991–1995 LEP runs
1.01 ±0.12 ±0.05	17k		ABREU	00L DLPH	1992–1995 runs
1.13 ±0.39 ±0.14	25k		ACKERSTAFF	99D OPAL	1990–1995 LEP runs
1.11 ±0.20 ±0.08		[252]	ALBRECHT	98 ARG	E_{cm}^{ee}= 9.5–10.6 GeV
1.16 ±0.52 ±0.06			ABE	97O SLD	1993–1995 SLC runs
0.979±0.048±0.016	34k		ALEXANDER	97F CLEO	E_{cm}^{ee}= 10.6 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.03 ±0.23 ±0.09		BUSKULIC	95D	ALEP	Repl. by HEISTER 01E

[252] ALBRECHT 98 use tau pair events of the type $\tau^-\tau^+ \to (\ell^-\overline{\nu}_\ell\nu_\tau)(\pi^+\pi^0\overline{\nu}_\tau)$, and their charged conjugates.

$\xi^\tau(\mu)$ PARAMETER

($V-A$) theory predicts $\xi = 1$.

VALUE	EVTS		DOCUMENT ID	TECN	COMMENT
1.030±0.059 OUR FIT					
1.06 ±0.06 OUR AVERAGE					
1.030±0.120±0.050	46k		HEISTER	01E ALEP	1991–1995 LEP runs
1.16 ±0.19 ±0.06	22k		ABREU	00L DLPH	1992–1995 runs
0.79 ±0.41 ±0.09	27k		ACKERSTAFF	99D OPAL	1990–1995 LEP runs
1.26 ±0.27 ±0.14		[253]	ALBRECHT	98 ARG	E_{cm}^{ee}= 9.5–10.6 GeV
0.75 ±0.50 ±0.14			ABE	97O SLD	1993–1995 SLC runs
1.054±0.069±0.047	22k		ALEXANDER	97F CLEO	E_{cm}^{ee}= 10.6 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.23 ±0.22 ±0.10		BUSKULIC	95D	ALEP	Repl. by HEISTER 01E

[253] ALBRECHT 98 use tau pair events of the type $\tau^-\tau^+ \to (\ell^-\overline{\nu}_\ell\nu_\tau)(\pi^+\pi^0\overline{\nu}_\tau)$, and their charged conjugates.

η^τ(e or μ) PARAMETER

($V-A$) theory predicts $\eta = 0$.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.013±0.020 OUR FIT					
0.015±0.021 OUR AVERAGE					
0.012±0.026±0.004	81k	HEISTER	01E	ALEP	1991–1995 LEP runs
−0.005±0.036±0.037		ABREU	00L	DLPH	1992–1995 runs
0.027±0.055±0.005	46k	ACKERSTAFF	99D	OPAL	1990–1995 LEP runs
0.27 ±0.14	54k	ACCIARRI	98R	L3	1991–1995 LEP runs
−0.13 ±0.47 ±0.15		ABE	97O	SLD	1993–1995 SLC runs
−0.015±0.061±0.062	31k	AMMAR	97B	CLEO	E_{cm}^{ee}= 10.6 GeV
0.03 ±0.18 ±0.12	8.2k	ALBRECHT	95	ARG	E_{cm}^{ee}= 9.5–10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.25 ±0.17 ±0.11	18k	ACCIARRI	96H	L3	Repl. by ACCIARRI 98R
−0.04 ±0.15 ±0.11		BUSKULIC	95D	ALEP	Repl. by HEISTER 01E

$\eta^\tau(\mu)$ PARAMETER

($V-A$) theory predicts $\eta = 0$.

VALUE	EVTS		DOCUMENT ID		TECN	COMMENT
0.094±0.073 OUR FIT						
0.17 ±0.15 OUR AVERAGE Error includes scale factor of 1.2.						
0.160±0.150±0.060	46k		HEISTER	01E	ALEP	1991–1995 LEP runs
0.72 ±0.32 ±0.15			ABREU	00L	DLPH	1992–1995 runs
−0.59 ±0.82 ±0.45		[254]	ABE	97O	SLD	1993–1995 SLC runs
0.010±0.149±0.171	13k	[255]	AMMAR	97B	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.010±0.065±0.001	27k	[256]	ACKERSTAFF	99D	OPAL	1990–1995 LEP runs
−0.24 ±0.23 ±0.18			BUSKULIC	95D	ALEP	Repl. by HEISTER 01E

[254] Highly correlated (corr. = 0.92) with ABE 97O $\rho^\tau(\mu)$ measurement.

[255] Highly correlated (corr. = 0.949) with AMMAR 97B $\rho^\tau(\mu)$ value.

[256] ACKERSTAFF 99D result is dominated by a constraint on η^τ from the OPAL measurements of the τ lifetime and B($\tau^- \to \mu^- \overline{\nu}_\mu \nu_\tau$) assuming lepton universality for the total coupling strength.

$(\delta\xi)^\tau$(e or μ) PARAMETER

($V-A$) theory predicts $(\delta\xi) = 0.75$.

VALUE	EVTS		DOCUMENT ID		TECN	COMMENT
0.746±0.021 OUR FIT						
0.744±0.022 OUR AVERAGE						
0.776±0.045±0.024	81k		HEISTER	01E	ALEP	1991–1995 LEP runs
0.779±0.070±0.028	36k		ABREU	00L	DLPH	1992–1995 runs
0.65 ±0.14 ±0.07	46k		ACKERSTAFF	99D	OPAL	1990–1995 LEP runs
0.70 ±0.11	54k		ACCIARRI	98R	L3	1991–1995 LEP runs
0.63 ±0.09		[257]	ALBRECHT	98	ARG	E_{cm}^{ee}= 9.5–10.6 GeV
0.88 ±0.27 ±0.04		[258]	ABE	97O	SLD	1993–1995 SLC runs
0.745±0.026±0.009	55k		ALEXANDER	97F	CLEO	E_{cm}^{ee}= 10.6 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.81 ±0.14 ±0.06	18k		ACCIARRI	96H L3	Repl. by ACCIARRI 98R
0.65 ±0.12		[259]	ALBRECHT	95C ARG	Repl. by ALBRECHT 98
0.88 ±0.11 ±0.07			BUSKULIC	95D ALEP	Repl. by HEISTER 01E

[257] Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 98, ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E. ALBRECHT 98 use tau pair events of the type $\tau^-\tau^+ \to (\ell^-\overline{\nu}_\ell\nu_\tau)(\pi^+\pi^0\overline{\nu}_\tau)$, and their charged conjugates.

[258] ABE 97O assume $\eta^\tau = 0$ in their fit. Letting η^τ vary in the fit gives a $(\rho\xi)^\tau$ value of 0.87 ± 0.27 ± 0.04.

[259] Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E. ALBRECHT 95C uses events of the type $\tau^-\tau^+ \to (\ell^-\overline{\nu}_\ell\nu_\tau)\,(h^+h^-h^+\overline{\nu}_\tau)$ and their charged conjugates.

$(\delta\xi)^\tau(e)$ PARAMETER

$(V-A)$ theory predicts $(\delta\xi) = 0.75$.

VALUE	EVTS		DOCUMENT ID	TECN	COMMENT
0.734±0.028 OUR FIT					
0.731±0.029 OUR AVERAGE					
0.778±0.066±0.024	44k		HEISTER	01E ALEP	1991–1995 LEP runs
0.85 ±0.12 ±0.04	17k		ABREU	00L DLPH	1992–1995 runs
0.72 ±0.31 ±0.14	25k		ACKERSTAFF	99D OPAL	1990–1995 LEP runs
0.56 ±0.14 ±0.06		[260]	ALBRECHT	98 ARG	$E^{ee}_{\rm cm}$= 9.5–10.6 GeV
0.85 ±0.43 ±0.08			ABE	97O SLD	1993–1995 SLC runs
0.720±0.032±0.010	34k		ALEXANDER	97F CLEO	$E^{ee}_{\rm cm}$= 10.6 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.11 ±0.17 ±0.07			BUSKULIC	95D ALEP	Repl. by HEISTER 01E

[260] ALBRECHT 98 use tau pair events of the type $\tau^-\tau^+ \to (\ell^-\overline{\nu}_\ell\nu_\tau)(\pi^+\pi^0\overline{\nu}_\tau)$, and their charged conjugates.

$(\delta\xi)^\tau(\mu)$ PARAMETER

$(V-A)$ theory predicts $(\delta\xi) = 0.75$.

VALUE	EVTS		DOCUMENT ID	TECN	COMMENT
0.778±0.037 OUR FIT					
0.79 ±0.04 OUR AVERAGE					
0.786±0.066±0.028	46k		HEISTER	01E ALEP	1991–1995 LEP runs
0.86 ±0.13 ±0.04	22k		ABREU	00L DLPH	1992–1995 runs
0.63 ±0.23 ±0.05	27k		ACKERSTAFF	99D OPAL	1990–1995 LEP runs
0.73 ±0.18 ±0.10		[261]	ALBRECHT	98 ARG	$E^{ee}_{\rm cm}$= 9.5–10.6 GeV
0.82 ±0.32 ±0.07			ABE	97O SLD	1993–1995 SLC runs
0.786±0.041±0.032	22k		ALEXANDER	97F CLEO	$E^{ee}_{\rm cm}$= 10.6 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.71 ±0.14 ±0.06			BUSKULIC	95D ALEP	Repl. by HEISTER 01E

[261] ALBRECHT 98 use tau pair events of the type $\tau^-\tau^+ \to (\ell^-\overline{\nu}_\ell\nu_\tau)(\pi^+\pi^0\overline{\nu}_\tau)$, and their charged conjugates.

$\xi^\tau(\pi)$ PARAMETER

($V-A$) theory predicts $\xi^\tau(\pi) = 1$.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.993±0.022 OUR FIT					
0.994±0.023 OUR AVERAGE					
0.994±0.020±0.014	27k	HEISTER	01E	ALEP	1991–1995 LEP runs
0.81 ±0.17 ±0.02		ABE	97O	SLD	1993–1995 SLC runs
1.03 ±0.06 ±0.04	2.0k	COAN	97	CLEO	E_{cm}^{ee}= 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.987±0.057±0.027		BUSKULIC	95D	ALEP	Repl. by HEISTER 01E
0.95 ±0.11 ±0.05		[262] BUSKULIC	94D	ALEP	1990+1991 LEP run

[262] Superseded by BUSKULIC 95D.

$\xi^\tau(\rho)$ PARAMETER

($V-A$) theory predicts $\xi^\tau(\rho) = 1$.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.994±0.008 OUR FIT					
0.994±0.009 OUR AVERAGE					
0.987±0.012±0.011	59k	HEISTER	01E	ALEP	1991–1995 LEP runs
0.99 ±0.12 ±0.04		ABE	97O	SLD	1993–1995 SLC runs
0.995±0.010±0.003	66k	ALEXANDER	97F	CLEO	E_{cm}^{ee}= 10.6 GeV
1.022±0.028±0.030	1.7k	[263] ALBRECHT	94E	ARG	E_{cm}^{ee}= 9.4–10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.045±0.058±0.032		BUSKULIC	95D	ALEP	Repl. by HEISTER 01E
1.03 ±0.11 ±0.05		[264] BUSKULIC	94D	ALEP	1990+1991 LEP run

[263] ALBRECHT 94E measure the square of this quantity and use the sign determined by ALBRECHT 90I to obtain the quoted result.

[264] Superseded by BUSKULIC 95D.

$\xi^\tau(a_1)$ PARAMETER

($V-A$) theory predicts $\xi^\tau(a_1) = 1$.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
1.001±0.027 OUR FIT					
1.002±0.028 OUR AVERAGE					
1.000±0.016±0.024	35k	[265] HEISTER	01E	ALEP	1991–1995 LEP runs
1.02 ±0.13 ±0.03	17.2k	ASNER	00	CLEO	E_{cm}^{ee}= 10.6 GeV
1.29 ±0.26 ±0.11	7.4k	[266] ACKERSTAFF	97R	OPAL	1992–1994 LEP runs
0.85 $^{+0.15}_{-0.17}$ ±0.05		ALBRECHT	95C	ARG	E_{cm}^{ee}= 9.5–10.6 GeV
1.25 ±0.23 $^{+0.15}_{-0.08}$	7.5k	ALBRECHT	93C	ARG	E_{cm}^{ee}= 9.4–10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.08 $^{+0.46}_{-0.41}$ $^{+0.14}_{-0.25}$	2.6k	[267] AKERS	95P	OPAL	Repl. by ACKERSTAFF 97R
0.937±0.116±0.064		BUSKULIC	95D	ALEP	Repl. by HEISTER 01E

[265] HEISTER 01E quote 1.000 ± 0.016 ± 0.013 ± 0.020 where the errors are statistical, systematic, and an uncertainty due to the final state model. We combine the systematic error and model uncertainty.

[266] ACKERSTAFF 97R obtain this result with a model independent fit to the hadronic structure functions. Fitting with the model of Kuhn and Santamaria (ZPHY **C48**, 445 (1990)) gives 0.87 ± 0.16 ± 0.04, and with the model of of Isgur *et al.* (PR **D39**,1357 (1989)) they obtain 1.20 ± 0.21 ± 0.14.

[267] AKERS 95P obtain this result with a model independent fit to the hadronic structure functions. Fitting with the model of Kuhn and Santamaria (ZPHY **C48**, 445 (1990)) gives $0.87 \pm 0.27^{+0.05}_{-0.06}$, and with the model of of Isgur *et al.* (PR **D39**,1357 (1989)) they obtain $1.10 \pm 0.31^{+0.13}_{-0.14}$.

ξ^τ(all hadronic modes) PARAMETER

($V-A$) theory predicts $\xi^\tau = 1$.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.995±0.007 OUR FIT					
0.997±0.007 OUR AVERAGE					
0.992±0.007±0.008	102k	[268] HEISTER	01E	ALEP	1991–1995 LEP runs
0.997±0.027±0.011	39k	[269] ABREU	00L	DLPH	1992–1995 runs
1.02 ±0.13 ±0.03	17.2k	[270] ASNER	00	CLEO	$E^{ee}_{\rm cm}$= 10.6 GeV
1.032±0.031	37k	[271] ACCIARRI	98R	L3	1991–1995 LEP runs
0.93 ±0.10 ±0.04		ABE	97O	SLD	1993–1995 SLC runs
1.29 ±0.26 ±0.11	7.4k	[272] ACKERSTAFF	97R	OPAL	1992–1994 LEP runs
0.995±0.010±0.003	66k	[273] ALEXANDER	97F	CLEO	$E^{ee}_{\rm cm}$= 10.6 GeV
1.03 ±0.06 ±0.04	2.0k	[274] COAN	97	CLEO	$E^{ee}_{\rm cm}$= 10.6 GeV
1.017±0.039		[275] ALBRECHT	95C	ARG	$E^{ee}_{\rm cm}$= 9.5–10.6 GeV
$1.25 \pm 0.23\ ^{+0.15}_{-0.08}$	7.5k	[276] ALBRECHT	93C	ARG	$E^{ee}_{\rm cm}$= 9.4–10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.970±0.053±0.011	14k	[277] ACCIARRI	96H	L3	Repl. by ACCIARRI 98R
$1.08\ ^{+0.46}_{-0.41}\ ^{+0.14}_{-0.25}$	2.6k	[278] AKERS	95P	OPAL	Repl. by ACKERSTAFF 97R
1.006±0.032±0.019		[279] BUSKULIC	95D	ALEP	Repl. by HEISTER 01E
1.022±0.028±0.030	1.7k	[280] ALBRECHT	94E	ARG	$E^{ee}_{\rm cm}$= 9.4–10.6 GeV
0.99 ±0.07 ±0.04		[281] BUSKULIC	94D	ALEP	1990+1991 LEP run

[268] HEISTER 01E quote 0.992 ± 0.007 ± 0.006 ± 0.005 where the errors are statistical, systematic, and an uncertainty due to the final state model. We combine the systematic error and model uncertainty. They use $\tau \to \pi\nu_\tau$, $\tau \to K\nu_\tau$, $\tau \to \rho\nu_\tau$, and $\tau \to a_1\nu_\tau$ decays.

[269] ABREU 00L use $\tau^- \to h^- \geq 0\pi^0\,\nu_\tau$ decays.

[270] ASNER 00 use $\tau^- \to \pi^- 2\pi^0\,\nu_\tau$ decays.

[271] ACCIARRI 98R use $\tau \to \pi\nu_\tau$, $\tau \to K\nu_\tau$, and $\tau \to \rho\nu_\tau$ decays.

[272] ACKERSTAFF 97R use $\tau \to a_1\nu_\tau$ decays.

[273] ALEXANDER 97F use $\tau \to \rho\nu_\tau$ decays.

[274] COAN 97 use h^+h^- energy correlations.

[275] Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E.

[276] Uses $\tau \to a_1\nu_\tau$ decays. Replaced by ALBRECHT 95C.

[277] ACCIARRI 96H use $\tau \to \pi\nu_\tau$, $\tau \to K\nu_\tau$, and $\tau \to \rho\nu_\tau$ decays.

[278] AKERS 95P use $\tau \to a_1\nu_\tau$ decays.

[279] BUSKULIC 95D use $\tau \to \pi \nu_\tau$, $\tau \to \rho \nu_\tau$, and $\tau \to a_1 \nu_\tau$ decays.

[280] ALBRECHT 94E measure the square of this quantity and use the sign determined by ALBRECHT 90I to obtain the quoted result. Uses $\tau \to a_1 \nu_\tau$ decays. Replaced by ALBRECHT 95C.

[281] BUSKULIC 94D use $\tau \to \pi \nu_\tau$ and $\tau \to \rho \nu_\tau$ decays. Superseded by BUSKULIC 95D.

τ REFERENCES

ABBIENDI	03	PL B551 35	G. Abbiendi *et al.*	(OPAL Collab.)
INAMI	03	PL B551 16	K. Inami *et al.*	(BELLE Collab.)
CHEN	02C	PR D66 071101R	S. Chen *et al.*	(CLEO Collab.)
ABBIENDI	01J	EPJ C19 653	G. Abbiendi *et al.*	(OPAL Collab.)
ABREU	01M	EPJ C20 617	P. Abreu *et al.*	(DELPHI Collab.)
ACCIARRI	01F	PL B507 47	M. Acciarri *et al.*	(L3 Collab.)
ACHARD	01D	PL B519 189	P. Achard *et al.*	(L3 Collab.)
ANASTASSOV	01	PRL 86 4467	A. Anastassov *et al.*	(CLEO Collab.)
HEISTER	01E	EPJ C22 217	A. Heister *et al.*	(ALEPH Collab.)
ABBIENDI	00A	PL B492 23	G. Abbiendi *et al.*	(OPAL Collab.)
ABBIENDI	00C	EPJ C13 213	G. Abbiendi *et al.*	(OPAL Collab.)
ABBIENDI	00D	EPJ C13 197	G. Abbiendi *et al.*	(OPAL Collab.)
ABREU	00L	EPJ C16 229	P. Abreu *et al.*	(DELPHI Collab.)
ACCIARRI	00B	PL B479 67	M. Acciarri *et al.*	(L3 Collab.)
AHMED	00	PR D61 071101R	S. Ahmed *et al.*	(CLEO Collab.)
ALBRECHT	00	PL B485 37	H. Albrecht *et al.*	(ARGUS Collab.)
ASNER	00	PR D61 012002	D.M. Asner *et al.*	(CLEO Collab.)
ASNER	00B	PR D62 072006	D.M. Asner *et al.*	(CLEO Collab.)
BERGFELD	00	PRL 84 830	T. Bergfeld *et al.*	(CLEO Collab.)
BROWDER	00	PR D61 052004	T.E. Browder *et al.*	(CLEO Collab.)
EDWARDS	00A	PR D61 072003	K.W. Edwards *et al.*	(CLEO Collab.)
GONZALEZ-S...	00	NP B582 3	G.A. Gonzalez-Sprinberg *et al.*	
ABBIENDI	99H	PL B447 134	G. Abbiendi *et al.*	(OPAL Collab.)
ABREU	99X	EPJ C10 201	P. Abreu *et al.*	(DELPHI Collab.)
ACKERSTAFF	99D	EPJ C8 3	K. Ackerstaff *et al.*	(OPAL Collab.)
ACKERSTAFF	99E	EPJ C8 183	K. Ackerstaff *et al.*	(OPAL Collab.)
BARATE	99K	EPJ C10 1	R. Barate *et al.*	(ALEPH Collab.)
BARATE	99R	EPJ C11 599	R. Barate *et al.*	(ALEPH Collab.)
BISHAI	99	PRL 82 281	M. Bishai *et al.*	(CLEO Collab.)
GODANG	99	PR D59 091303	R. Godang *et al.*	(CLEO Collab.)
RICHICHI	99	PR D60 112002	S.J. Richichi *et al.*	(CLEO Collab.)
ACCIARRI	98C	PL B426 207	M. Acciarri *et al.*	(L3 Collab.)
ACCIARRI	98E	PL B434 169	M. Aciarri *et al.*	(L3 Collab.)
ACCIARRI	98R	PL B438 405	M. Acciarri *et al.*	(L3 Collab.)
ACKERSTAFF	98M	EPJ C4 193	K. Ackerstaff *et al.*	(OPAL Collab.)
ACKERSTAFF	98N	PL B431 188	K. Ackerstaff *et al.*	(OPAL Collab.)
ALBRECHT	98	PL B431 179	H. Albrecht *et al.*	(ARGUS Collab.)
BARATE	98	EPJ C1 65	R. Barate *et al.*	(ALEPH Collab.)
BARATE	98E	EPJ C4 29	R. Barate *et al.*	(ALEPH Collab.)
BLISS	98	PR D57 5903	D.W. Bliss *et al.*	(CLEO Collab.)
ABE	97O	PRL 78 4691	K. Abe *et al.*	(SLD Collab.)
ACKERSTAFF	97J	PL B404 213	K. Ackerstaff *et al.*	(OPAL Collab.)
ACKERSTAFF	97L	ZPHY C74 403	K. Ackerstaff *et al.*	(OPAL Collab.)
ACKERSTAFF	97R	ZPHY C75 593	K. Ackerstaff *et al.*	(OPAL Collab.)
ALEXANDER	97F	PR D56 5320	J.P. Alexander *et al.*	(CLEO Collab.)
AMMAR	97B	PRL 78 4686	R. Ammar *et al.*	(CLEO Collab.)
ANASTASSOV	97	PR D55 2559	A. Anastassov *et al.*	(CLEO Collab.)
Also	98B	PR D58 119903 (erratum)	A. Anastassov *et al.*	(CLEO Collab.)
ANDERSON	97	PRL 79 3814	S. Anderson *et al.*	(CLEO Collab.)
AVERY	97	PR D55 R1119	P. Avery *et al.*	(CLEO Collab.)
BARATE	97I	ZPHY C74 387	R. Barate *et al.*	(ALEPH Collab.)
BARATE	97R	PL B414 362	R. Barate *et al.*	(ALEPH Collab.)
BERGFELD	97	PRL 79 2406	T. Bergfeld *et al.*	(CLEO Collab.)
BONVICINI	97	PRL 79 1221	G. Bonvicini *et al.*	(CLEO Collab.)
BUSKULIC	97C	ZPHY C74 263	D. Buskulic *et al.*	(ALEPH Collab.)
COAN	97	PR D55 7291	T.E. Coan *et al.*	(CLEO Collab.)
EDWARDS	97	PR D55 R3919	K.W. Edwards *et al.*	(CLEO Collab.)
EDWARDS	97B	PR D56 R5297	K.W. Edwards *et al.*	(CLEO Collab.)
ESCRIBANO	97	PL B395 369	R. Escribano, E. Masso	(BARC, PARIT)
ABREU	96B	PL B365 448	P. Abreu *et al.*	(DELPHI Collab.)

ACCIARRI	96H	PL B377 313	M. Acciarri *et al.*	(L3 Collab.)
ACCIARRI	96K	PL B389 187	M. Acciarri *et al.*	(L3 Collab.)
ALAM	96	PRL 76 2637	M.S. Alam *et al.*	(CLEO Collab.)
ALBRECHT	96E	PRPL 276 223	H. Albrecht *et al.*	(ARGUS Collab.)
ALEXANDER	96D	PL B369 163	G. Alexander *et al.*	(OPAL Collab.)
ALEXANDER	96E	PL B374 341	G. Alexander *et al.*	(OPAL Collab.)
ALEXANDER	96S	PL B388 437	G. Alexander *et al.*	(OPAL Collab.)
BAI	96	PR D53 20	J.Z. Bai *et al.*	(BES Collab.)
BALEST	96	PL B388 402	R. Balest *et al.*	(CLEO Collab.)
BARTELT	96	PRL 76 4119	J.E. Bartelt *et al.*	(CLEO Collab.)
BUSKULIC	96	ZPHY C70 579	D. Buskulic *et al.*	(ALEPH Collab.)
BUSKULIC	96C	ZPHY C70 561	D. Buskulic *et al.*	(ALEPH Collab.)
COAN	96	PR D53 6037	T.E. Coan *et al.*	(CLEO Collab.)
ABE	95Y	PR D52 4828	K. Abe *et al.*	(SLD Collab.)
ABREU	95T	PL B357 715	P. Abreu *et al.*	(DELPHI Collab.)
ABREU	95U	PL B359 411	P. Abreu *et al.*	(DELPHI Collab.)
ACCIARRI	95	PL B345 93	M. Acciarri *et al.*	(L3 Collab.)
ACCIARRI	95F	PL B352 487	M. Acciarri *et al.*	(L3 Collab.)
AKERS	95F	ZPHY C66 31	R. Akers *et al.*	(OPAL Collab.)
AKERS	95I	ZPHY C66 543	R. Akers *et al.*	(OPAL Collab.)
AKERS	95P	ZPHY C67 45	R. Akers *et al.*	(OPAL Collab.)
AKERS	95Y	ZPHY C68 555	R. Akers *et al.*	(OPAL Collab.)
ALBRECHT	95	PL B341 441	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	95C	PL B349 576	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	95G	ZPHY C68 25	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	95H	ZPHY C68 215	H. Albrecht *et al.*	(ARGUS Collab.)
BALEST	95C	PRL 75 3809	R. Balest *et al.*	(CLEO Collab.)
BERNABEU	95	NP B436 474	J. Bernabeu *et al.*	
BUSKULIC	95C	PL B346 371	D. Buskulic *et al.*	(ALEPH Collab.)
BUSKULIC	95D	PL B346 379	D. Buskulic *et al.*	(ALEPH Collab.)
Also	95P	PL B363 265 erratum	D. Buskulic *et al.*	(ALEPH Collab.)
ABREU	94K	PL B334 435	P. Abreu *et al.*	(DELPHI Collab.)
AKERS	94E	PL B328 207	R. Akers *et al.*	(OPAL Collab.)
AKERS	94G	PL B339 278	R. Akers *et al.*	(OPAL Collab.)
ALBRECHT	94E	PL B337 383	H. Albrecht *et al.*	(ARGUS Collab.)
ARTUSO	94	PRL 72 3762	M. Artuso *et al.*	(CLEO Collab.)
BARTELT	94	PRL 73 1890	J.E. Bartelt *et al.*	(CLEO Collab.)
BATTLE	94	PRL 73 1079	M. Battle *et al.*	(CLEO Collab.)
BAUER	94	PR D50 R13	D.A. Bauer *et al.*	(TPC/2gamma Collab.)
BUSKULIC	94D	PL B321 168	D. Buskulic *et al.*	(ALEPH Collab.)
BUSKULIC	94E	PL B332 209	D. Buskulic *et al.*	(ALEPH Collab.)
BUSKULIC	94F	PL B332 219	D. Buskulic *et al.*	(ALEPH Collab.)
GIBAUT	94B	PRL 73 934	D. Gibaut *et al.*	(CLEO Collab.)
ADRIANI	93M	PRPL 236 1	O. Adriani *et al.*	(L3 Collab.)
ALBRECHT	93C	ZPHY C58 61	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	93G	PL B316 608	H. Albrecht *et al.*	(ARGUS Collab.)
BALEST	93	PR D47 R3671	R. Balest *et al.*	(CLEO Collab.)
BEAN	93	PRL 70 138	A. Bean *et al.*	(CLEO Collab.)
BORTOLETTO	93	PRL 71 1791	D. Bortoletto *et al.*	(CLEO Collab.)
ESCRIBANO	93	PL B301 419	R. Escribano, E. Masso	(BARC)
PROCARIO	93	PRL 70 1207	M. Procario *et al.*	(CLEO Collab.)
ABREU	92N	ZPHY C55 555	P. Abreu *et al.*	(DELPHI Collab.)
ACTON	92F	PL B281 405	D.P. Acton *et al.*	(OPAL Collab.)
ACTON	92H	PL B288 373	P.D. Acton *et al.*	(OPAL Collab.)
AKERIB	92	PRL 69 3610	D.S. Akerib *et al.*	(CLEO Collab.)
Also	93B	PRL 71 3395 (erratum)	D.S. Akerib *et al.*	(CLEO Collab.)
ALBRECHT	92D	ZPHY C53 367	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	92K	ZPHY C55 179	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	92M	PL B292 221	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	92Q	ZPHY C56 339	H. Albrecht *et al.*	(ARGUS Collab.)
AMMAR	92	PR D45 3976	R. Ammar *et al.*	(CLEO Collab.)
ARTUSO	92	PRL 69 3278	M. Artuso *et al.*	(CLEO Collab.)
BAI	92	PRL 69 3021	J.Z. Bai *et al.*	(BES Collab.)
BATTLE	92	PL B291 488	M. Battle *et al.*	(CLEO Collab.)
BUSKULIC	92J	PL B297 459	D. Buskulic *et al.*	(ALEPH Collab.)
DECAMP	92C	ZPHY C54 211	D. Decamp *et al.*	(ALEPH Collab.)
ADEVA	91F	PL B265 451	B. Adeva *et al.*	(L3 Collab.)
ALBRECHT	91D	PL B260 259	H. Albrecht *et al.*	(ARGUS Collab.)
ALEXANDER	91D	PL B266 201	G. Alexander *et al.*	(OPAL Collab.)
ANTREASYAN	91	PL B259 216	D. Antreasyan *et al.*	(Crystal Ball Collab.)
GRIFOLS	91	PL B255 611	J.A. Grifols, A. Mendez	(BARC)

SAMUEL	91B	PRL 67 668	M.A. Samuel, G.W. Li, R. Mendel	(OKSU, WONT)
Also	92B	PRL 69 995	M.A. Samuel, G.W. Li, R. Mendel	(OKSU, WONT)
Erratum.				
ABACHI	90	PR D41 1414	S. Abachi *et al.*	(HRS Collab.)
ALBRECHT	90E	PL B246 278	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	90I	PL B250 164	H. Albrecht *et al.*	(ARGUS Collab.)
BEHREND	90	ZPHY C46 537	H.J. Behrend *et al.*	(CELLO Collab.)
BOWCOCK	90	PR D41 805	T.J.V. Bowcock *et al.*	(CLEO Collab.)
DELAGUILA	90	PL B252 116	F. del Aguila, M. Sher	(BARC, WILL)
GOLDBERG	90	PL B251 223	M. Goldberg *et al.*	(CLEO Collab.)
WU	90	PR D41 2339	D.Y. Wu *et al.*	(Mark II Collab.)
ABACHI	89B	PR D40 902	S. Abachi *et al.*	(HRS Collab.)
BEHREND	89B	PL B222 163	H.J. Behrend *et al.*	(CELLO Collab.)
JANSSEN	89	PL B228 273	H. Janssen *et al.*	(Crystal Ball Collab.)
KLEINWORT	89	ZPHY C42 7	C. Kleinwort *et al.*	(JADE Collab.)
ADEVA	88	PR D38 2665	B. Adeva *et al.*	(Mark-J Collab.)
ALBRECHT	88B	PL B202 149	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	88L	ZPHY C41 1	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	88M	ZPHY C41 405	H. Albrecht *et al.*	(ARGUS Collab.)
AMIDEI	88	PR D37 1750	D. Amidei *et al.*	(Mark II Collab.)
BEHREND	88	PL B200 226	H.J. Behrend *et al.*	(CELLO Collab.)
BRAUNSCH...	88C	ZPHY C39 331	W. Braunschweig *et al.*	(TASSO Collab.)
KEH	88	PL B212 123	S. Keh *et al.*	(Crystal Ball Collab.)
TSCHIRHART	88	PL B205 407	R. Tschirhart *et al.*	(HRS Collab.)
ABACHI	87B	PL B197 291	S. Abachi *et al.*	(HRS Collab.)
ABACHI	87C	PRL 59 2519	S. Abachi *et al.*	(HRS Collab.)
ADLER	87B	PRL 59 1527	J. Adler *et al.*	(Mark III Collab.)
AIHARA	87B	PR D35 1553	H. Aihara *et al.*	(TPC Collab.)
AIHARA	87C	PRL 59 751	H. Aihara *et al.*	(TPC Collab.)
ALBRECHT	87L	PL B185 223	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	87P	PL B199 580	H. Albrecht *et al.*	(ARGUS Collab.)
BAND	87	PL B198 297	H.R. Band *et al.*	(MAC Collab.)
BAND	87B	PRL 59 415	H.R. Band *et al.*	(MAC Collab.)
BARINGER	87	PRL 59 1993	P. Baringer *et al.*	(CLEO Collab.)
BEBEK	87C	PR D36 690	C. Bebek *et al.*	(CLEO Collab.)
BURCHAT	87	PR D35 27	P.R. Burchat *et al.*	(Mark II Collab.)
BYLSMA	87	PR D35 2269	B.G. Bylsma *et al.*	(HRS Collab.)
COFFMAN	87	PR D36 2185	D.M. Coffman *et al.*	(Mark III Collab.)
DERRICK	87	PL B189 260	M. Derrick *et al.*	(HRS Collab.)
FORD	87	PR D35 408	W.T. Ford *et al.*	(MAC Collab.)
FORD	87B	PR D36 1971	W.T. Ford *et al.*	(MAC Collab.)
GAN	87	PRL 59 411	K.K. Gan *et al.*	(Mark II Collab.)
GAN	87B	PL B197 561	K.K. Gan *et al.*	(Mark II Collab.)
AIHARA	86E	PRL 57 1836	H. Aihara *et al.*	(TPC Collab.)
BARTEL	86D	PL B182 216	W. Bartel *et al.*	(JADE Collab.)
PDG	86	PL 170B	M. Aguilar-Benitez *et al.*	(CERN, CIT+)
RUCKSTUHL	86	PRL 56 2132	W. Ruckstuhl *et al.*	(DELCO Collab.)
SCHMIDKE	86	PRL 57 527	W.B. Schmidke *et al.*	(Mark II Collab.)
YELTON	86	PRL 56 812	J.M. Yelton *et al.*	(Mark II Collab.)
ALTHOFF	85	ZPHY C26 521	M. Althoff *et al.*	(TASSO Collab.)
ASH	85B	PRL 55 2118	W.W. Ash *et al.*	(MAC Collab.)
BALTRUSAIT...	85	PRL 55 1842	R.M. Baltrusaitis *et al.*	(Mark III Collab.)
BARTEL	85F	PL 161B 188	W. Bartel *et al.*	(JADE Collab.)
BEHRENDS	85	PR D32 2468	S. Behrends *et al.*	(CLEO Collab.)
BELTRAMI	85	PRL 54 1775	I. Beltrami *et al.*	(HRS Collab.)
BERGER	85	ZPHY C28 1	C. Berger *et al.*	(PLUTO Collab.)
BURCHAT	85	PRL 54 2489	P.R. Burchat *et al.*	(Mark II Collab.)
FERNANDEZ	85	PRL 54 1624	E. Fernandez *et al.*	(MAC Collab.)
MILLS	85	PRL 54 624	G.B. Mills *et al.*	(DELCO Collab.)
AIHARA	84C	PR D30 2436	H. Aihara *et al.*	(TPC Collab.)
BEHREND	84	ZPHY C23 103	H.J. Behrend *et al.*	(CELLO Collab.)
MILLS	84	PRL 52 1944	G.B. Mills *et al.*	(DELCO Collab.)
BEHREND	83C	PL 127B 270	H.J. Behrend *et al.*	(CELLO Collab.)
SILVERMAN	83	PR D27 1196	D.J. Silverman, G.L. Shaw	(UCI)
BEHREND	82	PL 114B 282	H.J. Behrend *et al.*	(CELLO Collab.)
BLOCKER	82B	PRL 48 1586	C.A. Blocker *et al.*	(Mark II Collab.)
BLOCKER	82D	PL 109B 119	C.A. Blocker *et al.*	(Mark II Collab.) J
FELDMAN	82	PRL 48 66	G.J. Feldman *et al.*	(Mark II Collab.)
HAYES	82	PR D25 2869	K.G. Hayes *et al.*	(Mark II Collab.)
BERGER	81B	PL 99B 489	C. Berger *et al.*	(PLUTO Collab.)
DORFAN	81	PRL 46 215	J.M. Dorfan *et al.*	(Mark II Collab.)

BRANDELIK	80	PL 92B 199	R. Brandelik *et al.*	(TASSO Collab.)
ZHOLENTZ	80	PL 96B 214	A.A. Zholents *et al.*	(NOVO)
Also	81	SJNP 34 814	A.A. Zholents *et al.*	(NOVO)
		Translated from YAF 34 1471.		
BACINO	79B	PRL 42 749	W.J. Bacino *et al.*	(DELCO Collab.)
KIRKBY	79	SLAC-PUB-2419	J. Kirkby	(SLAC) J
Batavia Lepton Photon Conference.				
BACINO	78B	PRL 41 13	W.J. Bacino *et al.*	(DELCO Collab.) J
Also	78	Tokyo Conf. 249	J. Kirz	(STON)
Also	80	PL 96B 214	A.A. Zholents *et al.*	(NOVO)
BRANDELIK	78	PL 73B 109	R. Brandelik *et al.*	(DASP Collab.) J
FELDMAN	78	Tokyo Conf. 777	G.J. Feldman	(SLAC) J
JAROS	78	PRL 40 1120	J. Jaros *et al.*	(SLAC, LBL, NWES, HAWA)
PERL	75	PRL 35 1489	M.L. Perl *et al.*	(LBL, SLAC)

OTHER RELATED PAPERS

RAHAL-CAL...	98	IJMP A13 695	G. Rahal-Callot	(ETH)
GENTILE	96	PRPL 274 287	S. Gentile, M. Pohl	(ROMAI, ETH)
WEINSTEIN	93	ARNPS 43 457	A.J. Weinstein, R. Stroynowski	(CIT, SMU)
PERL	92	RPP 55 653	M.L. Perl	(SLAC)
PICH	90	MPL A5 1995	A. Pich	(VALE)
BARISH	88	PRPL 157 1	B.C. Barish, R. Stroynowski	(CIT)
GAN	88	IJMP A3 531	K.K. Gan, M.L. Perl	(SLAC)
HAYES	88	PR D38 3351	K.G. Hayes, M.L. Perl	(SLAC)
PERL	80	ARNPS 30 299	M.L. Perl	(SLAC)