

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+) \text{ Status: } ***$$

According to the quark model, the Ξ_c^+ (quark content *usc*) and Ξ_c^0 form an isospin doublet, and the spin-parity ought to be $J^P = 1/2^+$. None of I , J , or P has actually been measured.

Ξ_c^+ MASS


The fit uses the Ξ_c^+ and Ξ_c^0 mass and mass-difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2466.3 ± 1.4 OUR FIT				
2466.4 ± 1.5 OUR AVERAGE				
2465.8 ± 1.9 ± 2.5	90	FRABETTI	98 E687	γ Be, $\bar{E}_\gamma = 220$ GeV
2467.0 ± 1.6 ± 2.0	147	EDWARDS	96 CLE2	$e^+ e^- \approx \gamma(4S)$
2465.1 ± 3.6 ± 1.9	30	ALBRECHT	90F ARG	$e^+ e^-$ at $\gamma(4S)$
2467 ± 3 ± 4	23	ALAM	89 CLEO	$e^+ e^-$ 10.6 GeV
2466.5 ± 2.7 ± 1.2	5	BARLAG	89C ACCM	π^- Cu 230 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2464.4 ± 2.0 ± 1.4	30	FRABETTI	93B E687	See FRABETTI 98
2459 ± 5 ± 30	56	¹ COTEUS	87 SPEC	$nA \simeq 600$ GeV
2460 ± 25	82	BIAGI	83 SPEC	Σ^- Be 135 GeV

¹ Although COTEUS 87 claims to agree well with BIAGI 83 on the mass and width, there appears to be a discrepancy between the two experiments. BIAGI 83 sees a single peak (stated significance about 6 standard deviations) in the $\Lambda K^- \pi^+ \pi^+$ mass spectrum. COTEUS 87 sees two peaks in the same spectrum, one at the Ξ_c^+ mass, the other 75 MeV lower. The latter is attributed to $\Xi_c^+ \rightarrow \Sigma^0 K^- \pi^+ \pi^+ \rightarrow (\Lambda \gamma) K^- \pi^+ \pi^+$, with the γ unseen. The combined significance of the double peak is stated to be 5.5 standard deviations. But the absence of any trace of a lower peak in BIAGI 83 seems to us to throw into question the interpretation of the lower peak of COTEUS 87.

Ξ_c^+ MEAN LIFE

VALUE (10^{-15} s)	EVTS	DOCUMENT ID	TECN	COMMENT
442 ± 26 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			
503 ± 47 ± 18	250	MAHMOOD	02 CLE2	$e^+ e^- \approx \gamma(4S)$
439 ± 22 ± 9	532	LINK	01D FOCS	γ nucleus, $\bar{E}_\gamma \approx 180$ GeV
$340^{+70}_{-50} \pm 20$	56	FRABETTI	98 E687	γ Be, $\bar{E}_\gamma = 220$ GeV
$400^{+180}_{-120} \pm 100$	102	COTEUS	87 SPEC	$nA \simeq 600$ GeV
$480^{+210}_{-150} \pm 200$	53	BIAGI	85C SPEC	Σ^- Be 135 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$410^{+110}_{-80} \pm 20$	30	FRABETTI	93B E687	See FRABETTI 98
200^{+110}_{-60}	6	BARLAG	89C ACCM	π^- (K^-) Cu 230 GeV



Ξ_c^+ mean life

Ξ_c^+ DECAY MODES

No absolute branching fractions have been measured. THE FOLLOWING ARE BRANCHING RATIOS RELATIVE TO $\Xi^- \pi^+ \pi^+$.

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 \Lambda K^- \pi^+ \pi^+$	[a] 0.58 ± 0.18	
$\Gamma_2 \Lambda \bar{K}^*(892)^0 \pi^+$	[a,b] < 0.29	90%
$\Gamma_3 \Sigma(1385)^+ K^- \pi^+$	[a,b] < 0.41	90%
$\Gamma_4 \Sigma^+ K^- \pi^+$	[a] 1.18 ± 0.31	
$\Gamma_5 \Sigma^+ \bar{K}^*(892)^0$	[a,b] 0.92 ± 0.30	
$\Gamma_6 \Sigma^0 K^- \pi^+ \pi^+$	[a] 0.49 ± 0.26	
$\Gamma_7 \Xi^0 \pi^+$	[a] 0.55 ± 0.16	
$\Gamma_8 \Xi^- \pi^+ \pi^+$	[a] defined as 1	
$\Gamma_9 \Xi(1530)^0 \pi^+$	[a,b] < 0.2	90%
$\Gamma_{10} \Xi^0 \pi^+ \pi^0$	[a] 2.34 ± 0.68	
$\Gamma_{11} \Xi^0 \pi^+ \pi^+ \pi^-$	[a] 1.74 ± 0.50	
$\Gamma_{12} \Xi^0 e^+ \nu_e$	[a] $2.3 \begin{array}{l} +0.7 \\ -0.9 \end{array}$	
$\Gamma_{13} p K^- \pi^+$	[a] 0.21 ± 0.03	
$\Gamma_{14} p \bar{K}^*(892)^0$	[a,b] 0.12 ± 0.02	

[a] No absolute branching fractions have been measured. The following are branching *ratios* relative to $\Xi^- \pi^+ \pi^+$.

[b] This branching fraction includes all the decay modes of the final-state resonance.

Ξ_c^+ BRANCHING RATIOS

$\Gamma(\Lambda K^- \pi^+ \pi^+)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
-------	------	-------------	------	---------	-------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	56	COTEUS	87	SPEC	$nA \simeq 600 \text{ GeV}$
seen	82	² BIAGI	83	SPEC	$\Sigma^- \text{ Be } 135 \text{ GeV}$

² BIAGI 85B looks for but does not see the Ξ_c^+ in $pK^- \bar{K}^0 \pi^+$ ($\Gamma(pK^- \bar{K}^0 \pi^+)/\Gamma(\Lambda K^- \pi^+ \pi^+) < 0.08$ with 90% CL), $p2K^- 2\pi^+$ ($\Gamma(p2K^- 2\pi^+)/\Gamma(\Lambda K^- \pi^+ \pi^+) < 0.03$, 90% CL), $\Omega^- K^+ \pi^+$, $\Lambda K^{*0} \pi^+$, and $\Sigma(1385)^+ K^- \pi^+$.

$\Gamma(\Lambda K^- \pi^+ \pi^+)/\Gamma(\Xi^- \pi^+ \pi^+)$

Γ_1/Γ_8

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ_8
0.58 ± 0.16 ± 0.07	61	BERGFELD	96	CLE2	$e^+ e^- \approx \Upsilon(4S)$

$\Gamma(\Lambda \bar{K}^*(892)^0 \pi^+)/\Gamma(\Lambda K^- \pi^+ \pi^+)$

Γ_2/Γ_1

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.5	90	BERGFELD	96	CLE2 $e^+ e^- \approx \Upsilon(4S)$

$\Gamma(\Sigma(1385)^+ K^- \pi^+)/\Gamma(\Lambda K^- \pi^+ \pi^+)$

Γ_3/Γ_1

Unseen decay modes of the $\Sigma(1385)^+$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.7	90	BERGFELD	96	CLE2 $e^+ e^- \approx \Upsilon(4S)$

$\Gamma(\Sigma^+ K^- \pi^+)/\Gamma(\Xi^- \pi^+ \pi^+)$

Γ_4/Γ_8

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.18 ± 0.26 ± 0.17	119	BERGFELD	96	CLE2 $e^+ e^- \approx \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.92 ± 0.20 ± 0.07	3 JUN	00 SELX	Σ^- nucleus, 600 GeV
0.09 ^{+0.13 + 0.03} _{-0.06 - 0.02}	5 BARLAG	89C ACCM	$2 \Sigma^+ K^- \pi^+, 3 \Xi^- \pi^+ \pi^+$

³ This JUN 00 result is redundant with other results given below.

$\Gamma(\Sigma^+ \bar{K}^*(892)^0)/\Gamma(\Xi^- \pi^+ \pi^+)$

Γ_5/Γ_8

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.92 ± 0.27 ± 0.14	61	BERGFELD	96	CLE2 $e^+ e^- \approx \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	59	AVERY	95	CLE2 $e^+ e^- \approx \Upsilon(4S)$
------	----	-------	----	-------------------------------------

$\Gamma(\Sigma^0 K^- \pi^+ \pi^+)/\Gamma(\Lambda K^- \pi^+ \pi^+)$ Γ_6/Γ_1

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.84±0.36	47	4 COTEUS	87	SPEC $nA \simeq 600$ GeV

⁴ See, however, the note on the COTEUS 87 Ξ_c^+ mass measurement.

 $\Gamma(\Xi^0 \pi^+)/\Gamma(\Xi^- \pi^+ \pi^+)$ Γ_7/Γ_8

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.55±0.13±0.09	39	EDWARDS	96	CLE2 $e^+ e^- \approx \gamma(4S)$

 $\Gamma(\Xi^- \pi^+ \pi^+)/\Gamma_{\text{total}}$ Γ_8/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				

seen	131	BERGFELD	96	CLE2 $e^+ e^- \approx \gamma(4S)$
seen	160	AVERY	95	CLE2 $e^+ e^- \approx \gamma(4S)$
seen	30	FRABETTI	93B E687	γ Be, $\bar{E}_\gamma = 220$ GeV
seen	30	ALBRECHT	90F ARG	$e^+ e^-$ at $\gamma(4S)$
seen	23	ALAM	89	CLEO $e^+ e^-$ 10.6 GeV

 $\Gamma(\Xi(1530)^0 \pi^+)/\Gamma(\Xi^- \pi^+ \pi^+)$ Γ_9/Γ_8

Unseen decay modes of the $\Xi(1530)^0$ are included.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.2	90	BERGFELD	96	CLE2 $e^+ e^- \approx \gamma(4S)$

 $\Gamma(\Xi^0 \pi^+ \pi^0)/\Gamma(\Xi^- \pi^+ \pi^+)$ Γ_{10}/Γ_8

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.34±0.57±0.37	81	EDWARDS	96	CLE2 $e^+ e^- \approx \gamma(4S)$

 $\Gamma(\Xi(1530)^0 \pi^+)/\Gamma(\Xi^0 \pi^+ \pi^0)$ Γ_9/Γ_{10}

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.3	90	EDWARDS	96	CLE2 $e^+ e^- \approx \gamma(4S)$

 $\Gamma(\Xi^0 \pi^+ \pi^+ \pi^-)/\Gamma(\Xi^- \pi^+ \pi^+)$ Γ_{11}/Γ_8

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.74±0.42±0.27	57	EDWARDS	96	CLE2 $e^+ e^- \approx \gamma(4S)$

 $\Gamma(\Xi^0 e^+ \nu_e)/\Gamma(\Xi^- \pi^+ \pi^+)$ Γ_{12}/Γ_8

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.3±0.6^{+0.3}_{-0.6}	41	ALEXANDER	95B CLE2	$e^+ e^- \approx \gamma(4S)$

 $\Gamma(p K^- \pi^+)/\Gamma(\Sigma^+ K^- \pi^+)$ Γ_{13}/Γ_4

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.22±0.06±0.03	76	JUN	00 SELX	Σ^- nucleus, 600 GeV

$\Gamma(pK^-\pi^+)/\Gamma(\Xi^-\pi^+\pi^+)$

<u>VALUE</u>	<u>EVTS</u>
0.214±0.034 OUR AVERAGE	
0.234±0.047±0.022	202
0.20 ±0.04 ±0.02	76

 Γ_{13}/Γ_8

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
LINK	01B FOCS	γ nucleus
JUN	00 SELX	Σ^- nucleus, 600 GeV

 $\Gamma(p\bar{K}^*(892)^0)/\Gamma(pK^-\pi^+)$ Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

<u>VALUE</u>
0.54±0.09±0.05

 Γ_{14}/Γ_{13}

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
LINK	01B FOCS	γ nucleus

 Ξ_c^+ REFERENCES

MAHMOOD	02	PR D65 031102	A.H. Mahmood <i>et al.</i>	(CLEO Collab.)
LINK	01B	PL B512 277	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	01D	PL B523 53	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
JUN	00	PRL 84 1857	S.Y. Jun <i>et al.</i>	(FNAL SELEX Collab.)
FRAEBETTI	98	PL B427 211	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
BERGFELD	96	PL B365 431	T. Bergfeld <i>et al.</i>	(CLEO Collab.)
EDWARDS	96	PL B373 261	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
ALEXANDER	95B	PRL 74 3113	J. Alexander <i>et al.</i>	(CLEO Collab.)
Also	95E	PRL 75 4155 (erratum)	J. Alexander <i>et al.</i>	(CLEO Collab.)
AVERY	95	PRL 75 4364	P. Avery <i>et al.</i>	(CLEO Collab.)
FRAEBETTI	93B	PRL 70 1381	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ALBRECHT	90F	PL B247 121	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALAM	89	PL B226 401	M.S. Alam <i>et al.</i>	(CLEO Collab.)
BARLAG	89C	PL B233 522	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
COTEUS	87	PRL 59 1530	P. Coteus <i>et al.</i>	(FNAL E400 Collab.)
BIAGI	85B	ZPHY C28 175	S.F. Biagi <i>et al.</i>	(CERN WA62 Collab.)
BIAGI	85C	PL 150B 230	S.F. Biagi <i>et al.</i>	(CERN WA62 Collab.)
BIAGI	83	PL 122B 455	S.F. Biagi <i>et al.</i>	(CERN WA62 Collab.)