$\Upsilon(1S)$

$I^G(J^{PC}) = 0^-(1^{--})$

$\Upsilon(1S)$ MASS

VALUE (MeV)	*DOCUMENT ID*		*TECN*	*COMMENT*
9460.30±0.26 OUR AVERAGE	Error includes scale factor of 3.3.			
9460.51±0.09±0.05	[1] ARTAMONOV	00	MD1	$e^+e^- \to$ hadrons
9459.97±0.11±0.07	MACKAY	84	REDE	$e^+e^- \to$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •				
9460.60±0.09±0.05	[2,3] BARU	92B	REDE	$e^+e^- \to$ hadrons
9460.59±0.12	BARU	86	REDE	$e^+e^- \to$ hadrons
9460.6 ±0.4	[3,4] ARTAMONOV	84	REDE	$e^+e^- \to$ hadrons

[1] Reanalysis of BARU 92B and ARTAMONOV 84 using new electron mass (COHEN 87).
[2] Superseding BARU 86.
[3] Superseded by ARTAMONOV 00.
[4] Value includes data of ARTAMONOV 82.

$\Upsilon(1S)$ WIDTH

VALUE (keV)	*DOCUMENT ID*
53.0±1.5 OUR EVALUATION	See the Note on Width Determinations of the Υ states

$\Upsilon(1S)$ DECAY MODES

	Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1	$\tau^+\tau^-$	$(2.67^{+0.14}_{-0.16})$ %	
Γ_2	e^+e^-	(2.38±0.11) %	
Γ_3	$\mu^+\mu^-$	(2.48±0.06) %	
	Hadronic decays		
Γ_4	$\eta'(958)$anything	(2.8 ±0.4) %	
Γ_5	$J/\psi(1S)$anything	(1.1 ±0.4) $\times 10^{-3}$	
Γ_6	$\rho\pi$	< 2 $\times 10^{-4}$	90%
Γ_7	$\pi^+\pi^-$	< 5 $\times 10^{-4}$	90%
Γ_8	K^+K^-	< 5 $\times 10^{-4}$	90%
Γ_9	$p\overline{p}$	< 5 $\times 10^{-4}$	90%
Γ_{10}	$\pi^0\pi^+\pi^-$	< 1.84 $\times 10^{-5}$	90%
Γ_{11}	$D^*(2010)^\pm$anything		

Radiative decays

Γ_{12}	$\gamma\pi^+\pi^-$	$(6.3 \pm 1.8) \times 10^{-5}$		
Γ_{13}	$\gamma\pi^0\pi^0$	$(1.7 \pm 0.7) \times 10^{-5}$		
Γ_{14}	$\gamma 2h^+2h^-$	$(7.0 \pm 1.5) \times 10^{-4}$		
Γ_{15}	$\gamma 3h^+3h^-$	$(5.4 \pm 2.0) \times 10^{-4}$		
Γ_{16}	$\gamma 4h^+4h^-$	$(7.4 \pm 3.5) \times 10^{-4}$		
Γ_{17}	$\gamma\pi^+\pi^-K^+K^-$	$(2.9 \pm 0.9) \times 10^{-4}$		
Γ_{18}	$\gamma 2\pi^+2\pi^-$	$(2.5 \pm 0.9) \times 10^{-4}$		
Γ_{19}	$\gamma 3\pi^+3\pi^-$	$(2.5 \pm 1.2) \times 10^{-4}$		
Γ_{20}	$\gamma 2\pi^+2\pi^-K^+K^-$	$(2.4 \pm 1.2) \times 10^{-4}$		
Γ_{21}	$\gamma\pi^+\pi^-p\overline{p}$	$(1.5 \pm 0.6) \times 10^{-4}$		
Γ_{22}	$\gamma 2\pi^+2\pi^-p\overline{p}$	$(4 \pm 6) \times 10^{-5}$		
Γ_{23}	$\gamma 2K^+2K^-$	$(2.0 \pm 2.0) \times 10^{-5}$		
Γ_{24}	$\gamma\eta'(958)$	$< 1.6 \times 10^{-5}$		90%
Γ_{25}	$\gamma\eta$	$< 2.1 \times 10^{-5}$		90%
Γ_{26}	$\gamma f_2'(1525)$	$< 1.4 \times 10^{-4}$		90%
Γ_{27}	$\gamma f_2(1270)$	$(8 \pm 4) \times 10^{-5}$		
Γ_{28}	$\gamma\eta(1405)$	$< 8.2 \times 10^{-5}$		90%
Γ_{29}	$\gamma f_0(1710) \rightarrow \gamma K\overline{K}$	$< 2.6 \times 10^{-4}$		90%
Γ_{30}	$\gamma f_0(2200) \rightarrow \gamma K^+K^-$	$< 2 \times 10^{-4}$		90%
Γ_{31}	$\gamma f_J(2220) \rightarrow \gamma K^+K^-$	$< 1.5 \times 10^{-5}$		90%
Γ_{32}	$\gamma f_J(2220) \rightarrow \gamma\pi^+\pi^-$	$< 1.2 \times 10^{-5}$		90%
Γ_{33}	$\gamma f_J(2220) \rightarrow \gamma p\overline{p}$	$< 1.6 \times 10^{-5}$		90%
Γ_{34}	$\gamma\eta(2225) \rightarrow \gamma\phi\phi$	$< 3 \times 10^{-3}$		90%
Γ_{35}	γX (X = pseudoscalar with $m < 7.2$ GeV)	$< 3 \times 10^{-5}$		90%
Γ_{36}	$\gamma X\overline{X}$ ($X\overline{X}$ = vectors with $m < 3.1$ GeV)	$< 1 \times 10^{-3}$		90%

$\Upsilon(1S)$ $\Gamma(i)\Gamma(e^+e^-)/\Gamma(total)$

$\Gamma(e^+e^-) \times \Gamma(\mu^+\mu^-)/\Gamma_{total}$ — $\Gamma_2\Gamma_3/\Gamma$

VALUE (eV)	*DOCUMENT ID*		*TECN*	*COMMENT*
31.2±1.6±1.7	KOBEL	92	CBAL	$e^+e^- \rightarrow \mu^+\mu^-$

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{total}$ — $\Gamma_0\Gamma_2/\Gamma$

VALUE (keV)	*DOCUMENT ID*		*TECN*	*COMMENT*
1.216±0.027 OUR AVERAGE				
$1.187 \pm 0.023 \pm 0.031$	[5] BARU	92B	MD1	$e^+e^- \rightarrow$ hadrons
$1.23 \pm 0.02 \pm 0.05$	[5] JAKUBOWSKI	88	CBAL	$e^+e^- \rightarrow$ hadrons
$1.37 \pm 0.06 \pm 0.09$	[6] GILES	84B	CLEO	$e^+e^- \rightarrow$ hadrons
$1.23 \pm 0.08 \pm 0.04$	[6] ALBRECHT	82	DASP	$e^+e^- \rightarrow$ hadrons
$1.13 \pm 0.07 \pm 0.11$	[6] NICZYPORUK	82	LENA	$e^+e^- \rightarrow$ hadrons
1.09 ± 0.25	[6] BOCK	80	CNTR	$e^+e^- \rightarrow$ hadrons
1.35 ± 0.14	[7] BERGER	79	PLUT	$e^+e^- \rightarrow$ hadrons

[5] Radiative corrections evaluated following KURAEV 85.
[6] Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.
[7] Radiative corrections reevaluated by ALEXANDER 89 using B($\mu\mu$) = 0.026.

$\Upsilon(1S)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$ Γ_2

VALUE (keV)	*DOCUMENT ID*
1.314±0.029 OUR EVALUATION	

$\Upsilon(1S)$ BRANCHING RATIOS

$\Gamma(\tau^+\tau^-)/\Gamma_{total}$ Γ_1/Γ

VALUE	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
$0.0267^{+0.0014}_{-0.0016}$ OUR AVERAGE					
$0.0261\pm0.0012^{+0.0009}_{-0.0013}$	25k	CINABRO	94B	CLE2	$e^+e^- \to \tau^+\tau^-$
0.027 ±0.004 ±0.002		[8] ALBRECHT	85C	ARG	$\Upsilon(2S) \to \pi^+\pi^-\tau^+\tau^-$
0.034 ±0.004 ±0.004		GILES	83	CLEO	$e^+e^- \to \tau^+\tau^-$

[8] Using B($\Upsilon(1S) \to e\,e$) = B($\Upsilon(1S) \to \mu\mu$) = 0.0256; not used for width evaluations.

$\Gamma(\mu^+\mu^-)/\Gamma_{total}$ Γ_3/Γ

VALUE	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
0.0248±0.0006 OUR AVERAGE					
0.0249±0.0008±0.0013		ALEXANDER	98	CLE2	$\Upsilon(2S) \to \pi^+\pi^-\mu^+\mu^-$
0.0212±0.0020±0.0010		[9] BARU	92	MD1	$e^+e^- \to \mu^+\mu^-$
0.0231±0.0012±0.0010		[9] KOBEL	92	CBAL	$e^+e^- \to \mu^+\mu^-$
0.0252±0.0007±0.0007		CHEN	89B	CLEO	$e^+e^- \to \mu^+\mu^-$
0.0261±0.0009±0.0011		KAARSBERG	89	CSB2	$e^+e^- \to \mu^+\mu^-$
0.0230±0.0025±0.0013	86	ALBRECHT	87	ARG	$\Upsilon(2S) \to \pi^+\pi^-\mu^+\mu^-$
0.029 ±0.003 ±0.002	864	BESSON	84	CLEO	$\Upsilon(2S) \to \pi^+\pi^-\mu^+\mu^-$
0.027 ±0.003 ±0.003		ANDREWS	83	CLEO	$e^+e^- \to \mu^+\mu^-$
0.032 ±0.013 ±0.003		ALBRECHT	82	DASP	$e^+e^- \to \mu^+\mu^-$
0.038 ±0.015 ±0.002		NICZYPORUK	82	LENA	$e^+e^- \to \mu^+\mu^-$
$0.014\ ^{+0.034}_{-0.014}$		BOCK	80	CNTR	$e^+e^- \to \mu^+\mu^-$
0.022 ±0.020		BERGER	79	PLUT	$e^+e^- \to \mu^+\mu^-$

[9] Taking into account interference between the resonance and continuum.

$\Gamma(e^+e^-)/\Gamma_{total}$ Γ_2/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0238±0.0011 OUR AVERAGE					
0.0229±0.0008±0.0011		ALEXANDER	98	CLE2	$\Upsilon(2S) \to \pi^+\pi^- e^+e^-$
0.0242±0.0014±0.0014	307	ALBRECHT	87	ARG	$\Upsilon(2S) \to \pi^+\pi^- e^+e^-$
0.028 ±0.003 ±0.002	826	BESSON	84	CLEO	$\Upsilon(2S) \to \pi^+\pi^- e^+e^-$
0.051 ±0.030		BERGER	80C	PLUT	$e^+e^- \to e^+e^-$

$\Gamma(\eta'(958)\text{anything})/\Gamma_{total}$ Γ_4/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
0.028±0.004±0.002	ARTUSO	03	CLE2	$\Upsilon(1S) \to \eta'$ anything

$\Gamma(J/\psi(1S)\text{anything})/\Gamma_{total}$ Γ_5/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID		TECN	COMMENT
< 0.68	90	ALBRECHT	92J	ARG	$e^+e^- \to e^+e^-$ X, $e^+e^- \to \mu^+\mu^-$ X
1.1 ±0.4±0.2	[10]	FULTON	89	CLEO	$e^+e^- \to \mu^+\mu^-$ X

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.7	90	MASCHMANN	90	CBAL	$e^+e^- \to$ hadrons
<20	90	NICZYPORUK	83	LENA	

[10] Using B($(J/\psi) \to \mu^+\mu^-$) = (6.9 ± 0.9)%.

$\Gamma(\pi^+\pi^-)/\Gamma_{total}$ Γ_7/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<5	90	BARU	92	MD1	$\Upsilon(1S) \to \pi^+\pi^-$

$\Gamma(K^+K^-)/\Gamma_{total}$ Γ_8/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<5	90	BARU	92	MD1	$\Upsilon(1S) \to K^+K^-$

$\Gamma(p\bar{p})/\Gamma_{total}$ Γ_9/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<5	90	[11] BARU	96	MD1	$\Upsilon(1S) \to p\bar{p}$

[11] Supersedes BARU 92 in this node.

$\Gamma(\pi^0\pi^+\pi^-)/\Gamma_{total}$ Γ_{10}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN	COMMENT
<1.84	90	ANASTASSOV	99	CLE2	$e^+e^- \to$ hadrons

$\Gamma(\gamma X)/\Gamma_{total}$ Γ_{35}/Γ

(X = pseudoscalar with $m < 7.2$ GeV)

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN	COMMENT
<3	90	[12] BALEST	95	CLEO	$e^+e^- \to \gamma + X$

[12] For a noninteracting pseudoscalar X with mass < 7.2 GeV.

$\Gamma(\gamma X\overline{X})/\Gamma_{total}$ Γ_{36}/Γ

($X\overline{X}$ = vectors with $m<3.1$ GeV)

VALUE (units 10^{-3})	CL%	DOCUMENT ID		TECN	COMMENT
<1	90	[13] BALEST	95	CLEO	$e^+e^- \to \gamma + X\overline{X}$

[13] For a noninteracting vector X with mass < 3.1 GeV.

$\Gamma(\gamma\pi^+\pi^-)/\Gamma_{total}$ Γ_{12}/Γ

VALUE (units 10^{-5})	DOCUMENT ID		TECN	COMMENT
6.3±1.2±1.3	[14] ANASTASSOV	99	CLE2	$e^+e^- \to$ hadrons

[14] For $m_{\pi\pi}>1$ GeV.

$\Gamma(\gamma\pi^0\pi^0)/\Gamma_{total}$ Γ_{13}/Γ

VALUE (units 10^{-5})	DOCUMENT ID		TECN	COMMENT
1.7±0.6±0.3	[15] ANASTASSOV	99	CLE2	$e^+e^- \to$ hadrons

[15] For $m_{\pi\pi}>1$ GeV.

$\Gamma(\gamma 2\pi^+2\pi^-)/\Gamma_{total}$ Γ_{18}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
2.5±0.7±0.5	26 ± 7	FULTON	90B	CLEO	$e^+e^- \to$ hadrons

$\Gamma(\gamma\pi^+\pi^-K^+K^-)/\Gamma_{total}$ Γ_{17}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
2.9±0.7±0.6	29 ± 8	FULTON	90B	CLEO	$e^+e^- \to$ hadrons

$\Gamma(\gamma\pi^+\pi^-p\overline{p})/\Gamma_{total}$ Γ_{21}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
1.5±0.5±0.3	22 ± 6	FULTON	90B	CLEO	$e^+e^- \to$ hadrons

$\Gamma(\gamma 2K^+2K^-)/\Gamma_{total}$ Γ_{23}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
0.2 ±0.2	2 ± 2	FULTON	90B	CLEO	$e^+e^- \to$ hadrons

$\Gamma(\gamma 3\pi^+3\pi^-)/\Gamma_{total}$ Γ_{19}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
2.5±0.9±0.8	17 ± 5	FULTON	90B	CLEO	$e^+e^- \to$ hadrons

$\Gamma(\gamma 2\pi^+2\pi^-K^+K^-)/\Gamma_{total}$ Γ_{20}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
2.4±0.9±0.8	18 ± 7	FULTON	90B	CLEO	$e^+e^- \to$ hadrons

$\Gamma(\gamma 2\pi^+2\pi^-p\overline{p})/\Gamma_{total}$ Γ_{22}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
0.4±0.4±0.4	7 ± 6	FULTON	90B	CLEO	$e^+e^- \to$ hadrons

$\Gamma(\gamma 2h^+ 2h^-)/\Gamma_{total}$ Γ_{14}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
7.0±1.1±1.0	80 ± 12	FULTON	90B	CLEO	$e^+ e^- \rightarrow$ hadrons

$\Gamma(\gamma 3h^+ 3h^-)/\Gamma_{total}$ Γ_{15}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
5.4±1.5±1.3	39 ± 11	FULTON	90B	CLEO	$e^+ e^- \rightarrow$ hadrons

$\Gamma(\gamma 4h^+ 4h^-)/\Gamma_{total}$ Γ_{16}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
7.4±2.5±2.5	36 ± 12	FULTON	90B	CLEO	$e^+ e^- \rightarrow$ hadrons

$\Gamma(\rho\pi)/\Gamma_{total}$ Γ_6/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
< 2	90	FULTON	90B		$\Upsilon(1S) \rightarrow \rho^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<10	90	BLINOV	90	MD1	$\Upsilon(1S) \rightarrow \rho^0 \pi^0$
<21	90	NICZYPORUK	83	LENA	$\Upsilon(1S) \rightarrow \rho^0 \pi^0$

$\Gamma(D^*(2010)^\pm \text{anything})/\Gamma_{total}$ Γ_{11}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID		TECN	COMMENT
<19	90	[16] ALBRECHT	92J	ARG	$e^+ e^- \rightarrow D^0 \pi^\pm$ X

[16] For $x_p > 0.2$.

$\Gamma(\gamma\eta(1405))/\Gamma_{total}$ Γ_{28}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN	COMMENT
<8.2	90	[17] FULTON	90B	CLEO	$\Upsilon(1S) \rightarrow \gamma K^\pm \pi^\mp K^0_S$

[17] Includes unknown branching ratio of $\eta(1405) \rightarrow K^\pm \pi^\mp K^0_S$.

$\Gamma(\gamma\eta'(958))/\Gamma_{total}$ Γ_{24}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN	COMMENT
< 1.6	90	RICHICHI	01B	CLE2	$\Upsilon(1S) \rightarrow \gamma\eta' \rightarrow \gamma\eta\pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<130	90	SCHMITT	88	CBAL	$\Upsilon(1S) \rightarrow \gamma$ X

$\Gamma(\gamma\eta)/\Gamma_{total}$ Γ_{25}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN	COMMENT
< 2.1	90	MASEK	02	CLEO	$\Upsilon(1S) \rightarrow \gamma\eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<28.2	90	MASEK	02	CLEO	$\eta \rightarrow \gamma\gamma$
< 6.7	90	MASEK	02	CLEO	$\eta \rightarrow \pi^0\pi^0\pi^0$
< 2.6	90	MASEK	02	CLEO	$\eta \rightarrow \pi^+\pi^-\pi^0$
<35	90	SCHMITT	88	CBAL	$\Upsilon(1S) \rightarrow \gamma$ X

$\Gamma(\gamma f_2'(1525))/\Gamma_{total}$ Γ_{26}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN	COMMENT
<14	90	[18] FULTON	90B	CLEO	$\Upsilon(1S) \to \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<19.4	90	[18] ALBRECHT	89	ARG	$\Upsilon(1S) \to \gamma K^+ K^-$

[18] Assuming $B(f_2'(1525) \to K\overline{K}) = 0.71$.

$\Gamma(\gamma f_0(1710) \to \gamma K\overline{K})/\Gamma_{total}$ Γ_{29}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
< 2.6	90	[19] ALBRECHT	89	ARG	$\Upsilon(1S) \to \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 6.3	90	[19] FULTON	90B	CLEO	$\Upsilon(1S) \to \gamma K^+ K^-$
<19	90	[19] FULTON	90B	CLEO	$\Upsilon(1S) \to \gamma K_S^0 K_S^0$
< 8	90	[20] ALBRECHT	89	ARG	$\Upsilon(1S) \to \gamma \pi^+ \pi^-$
<24	90	[21] SCHMITT	88	CBAL	$\Upsilon(1S) \to \gamma X$

[19] Assuming $B(f_0(1710) \to K\overline{K}) = 0.38$.
[20] Assuming $B(f_0(1710) \to \pi\pi) = 0.04$.
[21] Assuming $B(f_0(1710) \to \eta\eta) = 0.18$.

$\Gamma(\gamma f_2(1270))/\Gamma_{total}$ Γ_{27}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{8.1 \pm 2.3^{+2.9}_{-2.7}}$		[22] ANASTASSOV	99	CLE2	$e^+ e^- \to$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<21	90	[22] FULTON	90B	CLEO	$\Upsilon(1S) \to \gamma \pi^+ \pi^-$
<13	90	[22] ALBRECHT	89	ARG	$\Upsilon(1S) \to \gamma \pi^+ \pi^-$
<81	90	SCHMITT	88	CBAL	$\Upsilon(1S) \to \gamma X$

[22] Using $B(f_2(1270) \to \pi\pi) = 0.84$.

$\Gamma(\gamma f_J(2220) \to \gamma K^+ K^-)/\Gamma_{total}$ Γ_{31}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN	COMMENT
< 1.6	90	MASEK	02	CLEO	$\Upsilon(1S) \to \gamma K^+ K^-$
< 1.5	90	[23] FULTON	90B	CLEO	$\Upsilon(1S) \to \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 2.9	90	[23] ALBRECHT	89	ARG	$\Upsilon(1S) \to \gamma K^+ K^-$
<20	90	[23] BARU	89	MD1	$\Upsilon(1S) \to \gamma K^+ K^-$

[23] Including unknown branching ratio of $f_J(2220) \to K^+ K^-$.

$\Gamma(\gamma f_J(2220) \to \gamma \pi^+ \pi^-)/\Gamma_{total}$ Γ_{32}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN	COMMENT
<1.2	90	MASEK	02	CLEO	$\Upsilon(1S) \to \gamma \pi^+ \pi^-$

$\Gamma(\gamma f_J(2220) \to \gamma p\overline{p})/\Gamma_{total}$ Γ_{33}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN	COMMENT
<1.6	90	MASEK	02	CLEO	$\Upsilon(1S) \to \gamma p\overline{p}$

$\Gamma(\gamma\eta(2225) \to \gamma\phi\phi)/\Gamma_{total}$ Γ_{34}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.003	90	[24] BARU	89	MD1	$\Upsilon(1S) \to \gamma K^+ K^- K^+ K^-$

[24] Assuming that the $\eta(2225)$ decays only into $\phi\phi$.

$\Gamma(\gamma f_0(2200) \to \gamma K^+ K^-)/\Gamma_{total}$ Γ_{30}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0002	90	[25] BARU	89	MD1	$\Upsilon(1S) \to \gamma K^+ K^-$

[25] Assuming that the $f_0(2200)$ decays only into $K^+ K^-$.

$\Upsilon(1S)$ REFERENCES

ARTUSO	03	PR D67 052003	M. Artuso *et al.*	(CLEO Collab.)
MASEK	02	PR D65 072002	G. Masek *et al.*	(CLEO Collab.)
RICHICHI	01B	PRL 87 141801	S.J. Richichi *et al.*	(CLEO Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov *et al.*	
ANASTASSOV	99	PRL 82 286	A. Anastassov *et al.*	(CLEO Collab.)
ALEXANDER	98	PR D58 052004	J.P. Alexander *et al.*	(CLEO Collab.)
BARU	96	PRPL 267 71	S.E. Baru *et al.*	(NOVO)
BALEST	95	PR D51 2053	R. Balest *et al.*	(CLEO Collab.)
CINABRO	94B	PL B340 129	D. Cinabro *et al.*	(CLEO Collab.)
ALBRECHT	92J	ZPHY C55 25	H. Albrecht *et al.*	(ARGUS Collab.)
BARU	92	ZPHY C54 229	S.E. Baru *et al.*	(NOVO)
BARU	92B	ZPHY C56 547	S.E. Baru *et al.*	(NOVO)
KOBEL	92	ZPHY C53 193	M. Kobel *et al.*	(Crystal Ball Collab.)
BLINOV	90	PL B245 311	A.E. Blinov *et al.*	(NOVO)
FULTON	90B	PR D41 1401	R. Fulton *et al.*	(CLEO Collab.)
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann *et al.*	(Crystal Ball Collab.)
ALBRECHT	89	ZPHY C42 349	H. Albrecht *et al.*	(ARGUS Collab.)
ALEXANDER	89	NP B320 45	J.P. Alexander *et al.*	(LBL, MICH, SLAC)
BARU	89	ZPHY C42 505	S.E. Baru *et al.*	(NOVO)
CHEN	89B	PR D39 3528	W.Y. Chen *et al.*	(CLEO Collab.)
FULTON	89	PL B224 445	R. Fulton *et al.*	(CLEO Collab.)
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg *et al.*	(CUSB Collab.)
BUCHMUEL...	88	HE e^+e^- Physics 412	W. Buchmueller, S. Cooper	(HANN, DESY, MIT)
		Editors: A. Ali and P. Soeding, World Scientific, Singapore		
JAKUBOWSKI	88	ZPHY C40 49	Z. Jakubowski *et al.*	(Crystal Ball Collab.) IGJPC
SCHMITT	88	ZPHY C40 199	P. Schmitt *et al.*	(Crystal Ball Collab.)
ALBRECHT	87	ZPHY C35 283	H. Albrecht *et al.*	(ARGUS Collab.)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
BARU	86	ZPHY C30 551	S.E. Baru *et al.*	(NOVO)
ALBRECHT	85C	PL 154B 452	H. Albrecht *et al.*	(ARGUS Collab.)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
		Translated from YAF 41 733.		
ARTAMONOV	84	PL 137B 272	A.S. Artamonov *et al.*	(NOVO)
BESSON	84	PR D30 1433	D. Besson *et al.*	(CLEO Collab.)
GILES	84B	PR D29 1285	R. Giles *et al.*	(CLEO Collab.)
MACKAY	84	PR D29 2483	W.W. MacKay *et al.*	(CUSB Collab.)
ANDREWS	83	PRL 50 807	D.E. Andrews *et al.*	(CLEO Collab.)
GILES	83	PRL 50 877	R. Giles *et al.*	(HARV, OSU, ROCH, RUTG+)
NICZYPORUK	83	ZPHY C17 197	B. Niczyporuk *et al.*	(LENA Collab.)
ALBRECHT	82	PL 116B 383	H. Albrecht *et al.*	(DESY, DORT, HEIDH+)
ARTAMONOV	82	PL 118B 225	A.S. Artamonov *et al.*	(NOVO)
NICZYPORUK	82	ZPHY C15 299	B. Niczyporuk *et al.*	(LENA Collab.)
BERGER	80C	PL 93B 497	C. Berger *et al.*	(PLUTO Collab.)
BOCK	80	ZPHY C6 125	P. Bock *et al.*	(HEIDP, MPIM, DESY, HAMB)
BERGER	79	ZPHY C1 343	C. Berger *et al.*	(PLUTO Collab.)

OTHER RELATED PAPERS

KOENIGS...	86	DESY 86/136	K. Koenigsmann	(DESY)
ALBRECHT	84	PL 134B 137	H. Albrecht *et al.*	(ARGUS Collab.)
ARTAMONOV	84	PL 137B 272	A.S. Artamonov *et al.*	(NOVO)
ARTAMONOV	82	PL 118B 225	A.S. Artamonov *et al.*	(NOVO)
BERGER	78	PL 76B 243	C. Berger *et al.*	(PLUTO Collab.)
BIENLEIN	78	PL 78B 360	J.K. Bienlein *et al.*	(DESY, HAMB, HEIDP+)
DARDEN	78	PL 76B 246	C.W. Darden *et al.*	(DESY, DORT, HEIDH+)
GARELICK	78	PR D18 945	D.A. Garelick *et al.*	(NEAS, WASH, TUFTS)
KAPLAN	78	PRL 40 435	D.M. Kaplan *et al.*	(STON, FNAL, COLU)
YOH	78	PRL 41 684	J.K. Yoh *et al.*	(COLU, FNAL, STON)
COBB	77	PL 72B 273	J.H. Cobb *et al.*	(BNL, CERN, SYRA, YALE)
HERB	77	PRL 39 252	S.W. Herb *et al.*	(COLU, FNAL, STON)
INNES	77	PRL 39 1240	W.R. Innes *et al.*	(COLU, FNAL, STON)