

$$I(J^P) = \frac{1}{2}(0^-)$$

Quantum numbers not measured. Values shown are quark-model predictions.

See also the B^\pm/B^0 ADMIXTURE and $B^\pm/B^0/B_s^0/b$ -baryon ADMIXTURE sections.

B^\pm MASS

The fit uses m_{B^+} , $(m_{B^0} - m_{B^+})$, and m_{B^0} to determine m_{B^+} , m_{B^0} , and the mass difference.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
5279.17 ± 0.29 OUR FIT				
5279.1 ± 0.4 OUR AVERAGE				
5279.10 ± 0.41 ± 0.36		¹ ACOSTA	06 CDF	$p\bar{p}$ at 1.96 TeV
5279.1 ± 0.4 ± 0.4	526	² CSORNA	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
5279.1 ± 1.7 ± 1.4	147	ABE	96B CDF	$p\bar{p}$ at 1.8 TeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
5278.8 ± 0.54 ± 2.0	362	ALAM	94 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
5278.3 ± 0.4 ± 2.0		BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
5280.5 ± 1.0 ± 2.0		³ ALBRECHT	90J ARG	$e^+e^- \rightarrow \Upsilon(4S)$
5275.8 ± 1.3 ± 3.0	32	ALBRECHT	87C ARG	$e^+e^- \rightarrow \Upsilon(4S)$
5278.2 ± 1.8 ± 3.0	12	⁴ ALBRECHT	87D ARG	$e^+e^- \rightarrow \Upsilon(4S)$
5278.6 ± 0.8 ± 2.0		BEBEK	87 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Uses exclusively reconstructed final states containing a $J/\psi \rightarrow \mu^+\mu^-$ decays.

² CSORNA 00 uses fully reconstructed $526 B^+ \rightarrow J/\psi(\ell) K^+$ events and invariant masses without beam constraint.

³ ALBRECHT 90J assumes 10580 for $\Upsilon(4S)$ mass. Supersedes ALBRECHT 87C and ALBRECHT 87D.

⁴ Found using fully reconstructed decays with $J/\psi(1S)$. ALBRECHT 87D assume $m_{\Upsilon(4S)} = 10577$ MeV.

B^\pm MEAN LIFE

See $B^\pm/B^0/B_s^0/b$ -baryon ADMIXTURE section for data on B -hadron mean life averaged over species of bottom particles.

“OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements and asymmetric lifetime errors.

VALUE (10^{-12} s)	EVTS	DOCUMENT ID	TECN	COMMENT
1.638 ± 0.011 OUR EVALUATION				
1.635 ± 0.011 ± 0.011		¹ ABE	05B BELL	$e^+e^- \rightarrow \Upsilon(4S)$
1.624 ± 0.014 ± 0.018		² ABDALLAH	04E DLPH	$e^+e^- \rightarrow Z$
1.636 ± 0.058 ± 0.025		³ ACOSTA	02C CDF	$p\bar{p}$ at 1.8 TeV
1.673 ± 0.032 ± 0.023		⁴ AUBERT	01F BABR	$e^+e^- \rightarrow \Upsilon(4S)$
1.648 ± 0.049 ± 0.035		⁵ BARATE	00R ALEP	$e^+e^- \rightarrow Z$

1.643 ± 0.037 ± 0.025		⁶ ABBIENDI	99J	OPAL	$e^+ e^- \rightarrow Z$
1.637 ± 0.058 ^{+0.045} _{-0.043}		⁵ ABE	98Q	CDF	$p\bar{p}$ at 1.8 TeV
1.66 ± 0.06 ± 0.03		⁶ ACCIARRI	98S	L3	$e^+ e^- \rightarrow Z$
1.66 ± 0.06 ± 0.05		⁶ ABE	97J	SLD	$e^+ e^- \rightarrow Z$
1.58 ^{+0.21} ^{+0.04} _{-0.18} _{-0.03}	94	³ BUSKULIC	96J	ALEP	$e^+ e^- \rightarrow Z$
1.61 ± 0.16 ± 0.12		^{5,7} ABREU	95Q	DLPH	$e^+ e^- \rightarrow Z$
1.72 ± 0.08 ± 0.06		⁸ ADAM	95	DLPH	$e^+ e^- \rightarrow Z$
1.52 ± 0.14 ± 0.09		⁵ AKERS	95T	OPAL	$e^+ e^- \rightarrow Z$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
1.695 ± 0.026 ± 0.015		⁴ ABE	02H	BELL	Repl. by ABE 05B
1.68 ± 0.07 ± 0.02		³ ABE	98B	CDF	Repl. by ACOSTA 02C
1.56 ± 0.13 ± 0.06		⁵ ABE	96C	CDF	Repl. by ABE 98Q
1.58 ± 0.09 ± 0.03		⁹ BUSKULIC	96J	ALEP	$e^+ e^- \rightarrow Z$
1.58 ± 0.09 ± 0.04		⁵ BUSKULIC	96J	ALEP	Repl. by BARATE 00R
1.70 ± 0.09		¹⁰ ADAM	95	DLPH	$e^+ e^- \rightarrow Z$
1.61 ± 0.16 ± 0.05	148	³ ABE	94D	CDF	Repl. by ABE 98B
1.30 ^{+0.33} ^{+0.16} _{-0.29}	92	⁵ ABREU	93D	DLPH	Sup. by ABREU 95Q
1.56 ± 0.19 ± 0.13	134	⁸ ABREU	93G	DLPH	Sup. by ADAM 95
1.51 ^{+0.30} ^{+0.12} _{-0.28} _{-0.14}	59	⁵ ACTON	93C	OPAL	Sup. by AKERS 95T
1.47 ^{+0.22} ^{+0.15} _{-0.19} _{-0.14}	77	⁵ BUSKULIC	93D	ALEP	Sup. by BUSKULIC 96J

¹ Measurement performed using a combined fit of CP -violation, mixing and lifetimes.

² Measurement performed using an inclusive reconstruction and B flavor identification technique.

³ Measured mean life using fully reconstructed decays.

⁴ Events are selected in which one B meson is fully reconstructed while the second B meson is reconstructed inclusively.

⁵ Data analyzed using $D/D^* \ell X$ event vertices.

⁶ Data analyzed using charge of secondary vertex.

⁷ ABREU 95Q assumes $B(B^0 \rightarrow D^{*-} \ell^+ \nu_\ell) = 3.2 \pm 1.7\%$.

⁸ Data analyzed using vertex-charge technique to tag B charge.

⁹ Combined result of $D/D^* \ell X$ analysis and fully reconstructed B analysis.

¹⁰ Combined ABREU 95Q and ADAM 95 result.

B^+ DECAY MODES

B^- modes are charge conjugates of the modes below. Modes which do not identify the charge state of the B are listed in the B^\pm/B^0 ADMIXTURE section.

The branching fractions listed below assume 50% $B^0 \bar{B}^0$ and 50% $B^+ B^-$ production at the $\Upsilon(4S)$. We have attempted to bring older measurements up to date by rescaling their assumed $\Upsilon(4S)$ production ratio to 50:50 and their assumed D , D_S , D^* , and ψ branching ratios to current values whenever this would affect our averages and best limits significantly.

Indentation is used to indicate a subchannel of a previous reaction. All resonant subchannels have been corrected for resonance branching fractions to the final state so the sum of the subchannel branching fractions can exceed that of the final state.

For inclusive branching fractions, e.g., $B \rightarrow D^\pm$ anything, the values usually are multiplicities, not branching fractions. They can be greater than one.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	
Semileptonic and leptonic modes			
Γ_1	$\ell^+ \nu_\ell$ anything	[a] (10.99 ± 0.28) %	
Γ_2	$e^+ \nu_e X_c$	(10.8 ± 0.4) %	
Γ_3	$D \ell^+ \nu_\ell$ anything	(9.8 ± 0.7) %	
Γ_4	$\bar{D}^0 \ell^+ \nu_\ell$	[a] (2.24 ± 0.11) %	
Γ_5	$\bar{D}^0 \tau^+ \nu_\tau$	(7 ± 4) × 10 ⁻³	
Γ_6	$\bar{D}^*(2007)^0 \ell^+ \nu_\ell$	[a] (5.68 ± 0.19) %	
Γ_7	$\bar{D}^*(2007)^0 \tau^+ \nu_\tau$	(2.2 ± 0.6) %	
Γ_8	$D^- \pi^+ \ell^+ \nu_\ell$	(4.2 ± 0.5) × 10 ⁻³	
Γ_9	$\bar{D}_0^*(2420)^0 \ell^+ \nu_\ell \times$ $B(\bar{D}_0^{*0} \rightarrow D^+ \pi^-)$	(2.5 ± 0.5) × 10 ⁻³	
Γ_{10}	$\bar{D}_2^*(2460)^0 \ell^+ \nu_\ell \times$ $B(\bar{D}_2^{*0} \rightarrow D^+ \pi^-)$	(1.67 ± 0.30) × 10 ⁻³	S=1.2
Γ_{11}	$D^{(*)} n \pi \ell^+ \nu_\ell (n \geq 1)$	(1.86 ± 0.26) %	
Γ_{12}	$D^{*-} \pi^+ \ell^+ \nu_\ell$	(6.1 ± 0.6) × 10 ⁻³	
Γ_{13}	$\bar{D}_1(2420)^0 \ell^+ \nu_\ell \times B(\bar{D}_1^0 \rightarrow$ $D^{*+} \pi^-)$	(3.2 ± 0.4) × 10 ⁻³	
Γ_{14}	$\bar{D}'_1(2430)^0 \ell^+ \nu_\ell \times B(\bar{D}'_1{}^0 \rightarrow$ $D^{*+} \pi^-)$	(2.7 ± 0.6) × 10 ⁻³	
Γ_{15}	$\bar{D}_2^*(2460)^0 \ell^+ \nu_\ell \times$ $B(\bar{D}_2^{*0} \rightarrow D^{*+} \pi^-)$	(1.8 ± 0.7) × 10 ⁻³	
Γ_{16}	$\pi^0 \ell^+ \nu_\ell$	(7.7 ± 1.2) × 10 ⁻⁵	
Γ_{17}	$\pi^0 e^+ \nu_e$		
Γ_{18}	$\eta \ell^+ \nu_\ell$	(6 ± 4) × 10 ⁻⁵	
Γ_{19}	$\eta' \ell^+ \nu_\ell$	(1.7 ± 2.2) × 10 ⁻⁵	
Γ_{20}	$\omega \ell^+ \nu_\ell$	[a] (1.3 ± 0.6) × 10 ⁻⁴	
Γ_{21}	$\omega \mu^+ \nu_\mu$		
Γ_{22}	$\rho^0 \ell^+ \nu_\ell$	[a] (1.28 ± 0.18) × 10 ⁻⁴	
Γ_{23}	$p \bar{p} e^+ \nu_e$	< 5.2 × 10 ⁻³	CL=90%
Γ_{24}	$e^+ \nu_e$	< 5.2 × 10 ⁻⁶	CL=90%
Γ_{25}	$\mu^+ \nu_\mu$	< 1.7 × 10 ⁻⁶	CL=90%
Γ_{26}	$\tau^+ \nu_\tau$	(1.4 ± 0.4) × 10 ⁻⁴	
Γ_{27}	$e^+ \nu_e \gamma$	< 2.0 × 10 ⁻⁴	CL=90%
Γ_{28}	$\mu^+ \nu_\mu \gamma$	< 5.2 × 10 ⁻⁵	CL=90%

Inclusive modes

Γ_{29}	$D^0 X$	(8.6 \pm 0.7) %
Γ_{30}	$\bar{D}^0 X$	(79 \pm 4) %
Γ_{31}	$D^+ X$	(2.5 \pm 0.5) %
Γ_{32}	$D^- X$	(9.9 \pm 1.2) %
Γ_{33}	$D_s^+ X$	(7.9 $^{+1.4}_{-1.3}$) %
Γ_{34}	$D_s^- X$	(1.10 $^{+0.40}_{-0.32}$) %
Γ_{35}	$\Lambda_c^+ X$	(2.1 $^{+0.9}_{-0.6}$) %
Γ_{36}	$\bar{\Lambda}_c^- X$	(2.8 $^{+1.1}_{-0.9}$) %
Γ_{37}	$\bar{c} X$	(97 \pm 4) %
Γ_{38}	$c X$	(23.4 $^{+2.2}_{-1.8}$) %
Γ_{39}	$\bar{c} c X$	(120 \pm 6) %

D , D^* , or D_s modes

Γ_{40}	$\bar{D}^0 \pi^+$	(4.84 \pm 0.15) $\times 10^{-3}$
Γ_{41}	$D_{CP(+1)} \pi^+$	[b] (2.3 \pm 0.4) $\times 10^{-3}$
Γ_{42}	$D_{CP(-1)} \pi^+$	[b] (2.0 \pm 0.4) $\times 10^{-3}$
Γ_{43}	$\bar{D}^0 \rho^+$	(1.34 \pm 0.18) %
Γ_{44}	$\bar{D}^0 K^+$	(3.68 \pm 0.33) $\times 10^{-4}$
Γ_{45}	$D_{CP(+1)} K^+$	[b] (1.95 \pm 0.27) $\times 10^{-4}$
Γ_{46}	$D_{CP(-1)} K^+$	[b] (1.89 \pm 0.27) $\times 10^{-4}$
Γ_{47}	$[K^- \pi^+]_D K^+$	[c] < 2.8 $\times 10^{-7}$ CL=90%
Γ_{48}	$[K^+ \pi^-]_D K^+$	[c] < 4 $\times 10^{-5}$ CL=90%
Γ_{49}	$[K^- \pi^+ \pi^0]_D K^+$	
Γ_{50}	$[K^+ \pi^- \pi^0]_D K^+$	
Γ_{51}	$[K^- \pi^+]_D K^*(892)^+$	[c]
Γ_{52}	$[K^+ \pi^-]_D K^*(892)^+$	[c]
Γ_{53}	$[K^- \pi^+]_D \pi^+$	[c] (6.3 \pm 1.1) $\times 10^{-7}$
Γ_{54}	$[K^+ \pi^-]_D \pi^+$	(1.9 \pm 0.4) $\times 10^{-4}$
Γ_{55}	$[\pi^+ \pi^- \pi^0]_D K^-$	(4.6 \pm 0.9) $\times 10^{-6}$
Γ_{56}	$\bar{D}^0 K^*(892)^+$	(5.3 \pm 0.4) $\times 10^{-4}$
Γ_{57}	$D_{CP(-1)} K^*(892)^+$	[b] (1.7 \pm 0.7) $\times 10^{-4}$
Γ_{58}	$D_{CP(+1)} K^*(892)^+$	[b] (5.2 \pm 1.2) $\times 10^{-4}$
Γ_{59}	$\bar{D}^0 K^+ \bar{K}^0$	(5.5 \pm 1.6) $\times 10^{-4}$
Γ_{60}	$\bar{D}^0 K^+ \bar{K}^*(892)^0$	(7.5 \pm 1.7) $\times 10^{-4}$
Γ_{61}	$\bar{D}^0 \pi^+ \pi^+ \pi^-$	(1.1 \pm 0.4) %
Γ_{62}	$\bar{D}^0 \pi^+ \pi^+ \pi^-$ nonresonant	(5 \pm 4) $\times 10^{-3}$
Γ_{63}	$\bar{D}^0 \pi^+ \rho^0$	(4.2 \pm 3.0) $\times 10^{-3}$
Γ_{64}	$\bar{D}^0 a_1(1260)^+$	(4 \pm 4) $\times 10^{-3}$
Γ_{65}	$\bar{D}^0 \omega \pi^+$	(4.1 \pm 0.9) $\times 10^{-3}$

Γ_{66}	$D^*(2010)^- \pi^+ \pi^+$	$(1.35 \pm 0.22) \times 10^{-3}$	
Γ_{67}	$D^- \pi^+ \pi^+$	$(1.02 \pm 0.16) \times 10^{-3}$	
Γ_{68}	$D^+ K^0$	$< 5.0 \times 10^{-6}$	CL=90%
Γ_{69}	$\bar{D}^*(2007)^0 \pi^+$	$(5.19 \pm 0.26) \times 10^{-3}$	
Γ_{70}	$\bar{D}_{CP(+1)}^{*0} \pi^+$	[d] $(2.9 \pm 0.6) \times 10^{-3}$	
Γ_{71}	$\bar{D}_{CP(-1)}^{*0} \pi^+$	[d] $(2.5 \pm 1.0) \times 10^{-3}$	
Γ_{72}	$\bar{D}^*(2007)^0 \omega \pi^+$	$(4.5 \pm 1.2) \times 10^{-3}$	
Γ_{73}	$\bar{D}^*(2007)^0 \rho^+$	$(9.8 \pm 1.7) \times 10^{-3}$	
Γ_{74}	$\bar{D}^*(2007)^0 K^+$	$(4.16 \pm 0.33) \times 10^{-4}$	
Γ_{75}	$\bar{D}_{CP(+1)}^{*0} K^+$	[d] $(2.7 \pm 0.4) \times 10^{-4}$	
Γ_{76}	$\bar{D}_{CP(-1)}^{*0} K^+$	[d] $(2.29 \pm 0.32) \times 10^{-4}$	
Γ_{77}	$\bar{D}^*(2007)^0 K^*(892)^+$	$(8.1 \pm 1.4) \times 10^{-4}$	
Γ_{78}	$\bar{D}^*(2007)^0 K^+ \bar{K}^0$	$< 1.06 \times 10^{-3}$	CL=90%
Γ_{79}	$\bar{D}^*(2007)^0 K^+ K^*(892)^0$	$(1.5 \pm 0.4) \times 10^{-3}$	
Γ_{80}	$\bar{D}^*(2007)^0 \pi^+ \pi^+ \pi^-$	$(1.03 \pm 0.12) \%$	
Γ_{81}	$\bar{D}^*(2007)^0 a_1(1260)^+$	$(1.9 \pm 0.5) \%$	
Γ_{82}	$\bar{D}^*(2007)^0 \pi^- \pi^+ \pi^+ \pi^0$	$(1.8 \pm 0.4) \%$	
Γ_{83}	$\bar{D}^{*0} 3\pi^+ 2\pi^-$	$(5.7 \pm 1.2) \times 10^{-3}$	
Γ_{84}	$D^*(2010)^+ \pi^0$	$< 3.6 \times 10^{-6}$	
Γ_{85}	$D^*(2010)^+ K^0$	$< 9.0 \times 10^{-6}$	CL=90%
Γ_{86}	$D^*(2010)^- \pi^+ \pi^+ \pi^0$	$(1.5 \pm 0.7) \%$	
Γ_{87}	$D^*(2010)^- \pi^+ \pi^+ \pi^+ \pi^-$	$(2.6 \pm 0.4) \times 10^{-3}$	
Γ_{88}	$\bar{D}^{**0} \pi^+$	[e] $(5.9 \pm 1.3) \times 10^{-3}$	
Γ_{89}	$\bar{D}_1^*(2420)^0 \pi^+$	$(1.5 \pm 0.6) \times 10^{-3}$	S=1.3
Γ_{90}	$\bar{D}_1(2420)^0 \pi^+ \times B(\bar{D}_1^0 \rightarrow \bar{D}^0 \pi^+ \pi^-)$	$(1.9 \begin{smallmatrix} +0.5 \\ -0.6 \end{smallmatrix}) \times 10^{-4}$	
Γ_{91}	$\bar{D}_2^*(2462)^0 \pi^+ \times B(\bar{D}_2^*(2462)^0 \rightarrow D^- \pi^+)$	$(3.4 \pm 0.8) \times 10^{-4}$	
Γ_{92}	$\bar{D}_0^*(2400)^0 \pi^+ \times B(\bar{D}_0^*(2400)^0 \rightarrow D^- \pi^+)$	$(6.1 \pm 1.9) \times 10^{-4}$	
Γ_{93}	$\bar{D}_1(2421)^0 \pi^+ \times B(\bar{D}_1(2421)^0 \rightarrow D^{*-} \pi^+)$	$(6.8 \pm 1.5) \times 10^{-4}$	
Γ_{94}	$\bar{D}_2^*(2462)^0 \pi^+ \times B(\bar{D}_2^*(2462)^0 \rightarrow D^{*-} \pi^+)$	$(1.8 \pm 0.5) \times 10^{-4}$	
Γ_{95}	$\bar{D}'_1(2427)^0 \pi^+ \times B(\bar{D}'_1(2427)^0 \rightarrow D^{*-} \pi^+)$	$(5.0 \pm 1.2) \times 10^{-4}$	
Γ_{96}	$\bar{D}_1(2420)^0 \pi^+ \times B(\bar{D}_1^0 \rightarrow \bar{D}^{*0} \pi^+ \pi^-)$	$< 6 \times 10^{-6}$	CL=90%
Γ_{97}	$\bar{D}_1^*(2420)^0 \rho^+$	$< 1.4 \times 10^{-3}$	CL=90%
Γ_{98}	$\bar{D}_2^*(2460)^0 \pi^+$	$< 1.3 \times 10^{-3}$	CL=90%

Γ_{99}	$\bar{D}_2^*(2460)^0 \pi^+ \times B(\bar{D}_2^{*0} \rightarrow \bar{D}^{*0} \pi^+ \pi^-)$	< 2.2	$\times 10^{-5}$	CL=90%
Γ_{100}	$\bar{D}_2^*(2460)^0 \rho^+$	< 4.7	$\times 10^{-3}$	CL=90%
Γ_{101}	$\bar{D}^0 D_s^+$	(1.02 \pm 0.17)	%	
Γ_{102}	$D_{s0}(2317)^+ \bar{D}^0 \times B(D_{s0}(2317)^+ \rightarrow D_s^+ \pi^0)$	(7.4 $\begin{smallmatrix} +2.2 \\ -1.7 \end{smallmatrix}$)	$\times 10^{-4}$	
Γ_{103}	$D_{s0}(2317)^+ \bar{D}^0 \times B(D_{s0}(2317)^+ \rightarrow D_s^{*+} \gamma)$	< 7.6	$\times 10^{-4}$	CL=90%
Γ_{104}	$D_{s0}(2317)^+ \bar{D}^*(2007)^0 \times B(D_{s0}(2317)^+ \rightarrow D_s^+ \pi^0)$	(9 \pm 7)	$\times 10^{-4}$	
Γ_{105}	$D_{sJ}(2457)^+ \bar{D}^0$	(3.1 $\begin{smallmatrix} +1.0 \\ -0.9 \end{smallmatrix}$)	$\times 10^{-3}$	
Γ_{106}	$D_{sJ}(2457)^+ \bar{D}^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^+ \gamma)$	(4.7 $\begin{smallmatrix} +1.3 \\ -1.1 \end{smallmatrix}$)	$\times 10^{-4}$	
Γ_{107}	$D_{sJ}(2457)^+ \bar{D}^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^+ \pi^+ \pi^-)$	< 2.2	$\times 10^{-4}$	CL=90%
Γ_{108}	$D_{sJ}(2457)^+ \bar{D}^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^+ \pi^0)$	< 2.7	$\times 10^{-4}$	CL=90%
Γ_{109}	$D_{sJ}(2457)^+ \bar{D}^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^{*+} \gamma)$	< 9.8	$\times 10^{-4}$	CL=90%
Γ_{110}	$D_{sJ}(2457)^+ \bar{D}^*(2007)^0$	(1.20 \pm 0.30)	%	
Γ_{111}	$D_{sJ}(2457)^+ \bar{D}^*(2007)^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^+ \gamma)$	(1.4 $\begin{smallmatrix} +0.7 \\ -0.6 \end{smallmatrix}$)	$\times 10^{-3}$	
Γ_{112}	$\bar{D}^0 D_{s1}(2536)^+ \times B(D_{s1}(2536)^+ \rightarrow D^*(2007)^0 K^+)$	(2.2 \pm 0.7)	$\times 10^{-4}$	
Γ_{113}	$\bar{D}^*(2007)^0 D_{s1}(2536)^+ \times B(D_{s1}(2536)^+ \rightarrow D^*(2007)^0 K^+)$	(5.5 \pm 1.6)	$\times 10^{-4}$	
Γ_{114}	$\bar{D}^0 D_{s1}(2536)^+ \times B(D_{s1}(2536)^+ \rightarrow D^{*+} K^0)$	(2.3 \pm 1.1)	$\times 10^{-4}$	
Γ_{115}	$\bar{D}^0 D_{sJ}(2700)^+ \times B(D_{sJ}(2700)^+ \rightarrow D^0 K^+)$	(1.13 $\begin{smallmatrix} +0.26 \\ -0.40 \end{smallmatrix}$)	$\times 10^{-3}$	
Γ_{116}	$\bar{D}^{*0} D_{s1}(2536)^+ \times B(D_{s1}(2536)^+ \rightarrow D^{*+} K^0)$	(3.9 \pm 2.6)	$\times 10^{-4}$	
Γ_{117}	$\bar{D}^{*0} D_{sJ}(2573)^+ \times B(D_{sJ}(2573)^+ \rightarrow D^0 K^+)$	< 2	$\times 10^{-4}$	CL=90%
Γ_{118}	$\bar{D}^*(2007)^0 D_{sJ}(2573)^+ \times B(D_{sJ}(2573)^+ \rightarrow D^0 K^+)$	< 5	$\times 10^{-4}$	CL=90%
Γ_{119}	$\bar{D}^0 D_s^{*+}$	(7.8 \pm 1.6)	$\times 10^{-3}$	

Γ_{120}	$\bar{D}_s^*(2007)^0 D_s^+$	$(8.4 \pm 1.7) \times 10^{-3}$	
Γ_{121}	$\bar{D}_s^*(2007)^0 D_s^{*+}$	$(1.74 \pm 0.23) \%$	
Γ_{122}	$D_s^{(*)+} \bar{D}^{*0}$	$(2.7 \pm 1.2) \%$	
Γ_{123}	$\bar{D}_s^*(2007)^0 D^*(2010)^+$	$(8.1 \pm 1.7) \times 10^{-4}$	
Γ_{124}	$\bar{D}^0 D^*(2010)^+ + \bar{D}_s^*(2007)^0 D^+$	< 1.30	$\%$ CL=90%
Γ_{125}	$\bar{D}^0 D^*(2010)^+$	$(3.9 \pm 0.5) \times 10^{-4}$	
Γ_{126}	$\bar{D}^0 D^+$	$(3.8 \pm 0.4) \times 10^{-4}$	
Γ_{127}	$\bar{D}^0 D^+ K^0$	< 2.8	$\times 10^{-3}$ CL=90%
Γ_{128}	$D^+ \bar{D}_s^*(2007)^0$	$(6.3 \pm 1.7) \times 10^{-4}$	
Γ_{129}	$\bar{D}_s^*(2007)^0 D^+ K^0$	< 6.1	$\times 10^{-3}$ CL=90%
Γ_{130}	$\bar{D}^0 \bar{D}_s^*(2010)^+ K^0$	$(5.2 \pm 1.2) \times 10^{-3}$	
Γ_{131}	$\bar{D}_s^*(2007)^0 D^*(2010)^+ K^0$	$(7.8 \pm 2.6) \times 10^{-3}$	
Γ_{132}	$\bar{D}^0 D^0 K^+$	$(2.10 \pm 0.26) \times 10^{-3}$	
Γ_{133}	$\bar{D}_s^*(2007)^0 D^0 K^+$	< 3.8	$\times 10^{-3}$ CL=90%
Γ_{134}	$\bar{D}^0 D^*(2007)^0 K^+$	$(4.7 \pm 1.0) \times 10^{-3}$	
Γ_{135}	$\bar{D}_s^*(2007)^0 D^*(2007)^0 K^+$	$(5.3 \pm 1.6) \times 10^{-3}$	
Γ_{136}	$D^- D^+ K^+$	< 4	$\times 10^{-4}$ CL=90%
Γ_{137}	$D^- D^*(2010)^+ K^+$	< 7	$\times 10^{-4}$ CL=90%
Γ_{138}	$D^*(2010)^- D^+ K^+$	$(1.5 \pm 0.4) \times 10^{-3}$	
Γ_{139}	$D^*(2010)^- D^*(2010)^+ K^+$	< 1.8	$\times 10^{-3}$ CL=90%
Γ_{140}	$(\bar{D} + \bar{D}^*)(D + D^*)K$	$(3.5 \pm 0.6) \%$	
Γ_{141}	$D_s^+ \pi^0$	$(1.6 \pm 0.6) \times 10^{-5}$	
Γ_{142}	$D_s^{*+} \pi^0$	< 2.7	$\times 10^{-4}$ CL=90%
Γ_{143}	$D_s^+ \eta$	< 4	$\times 10^{-4}$ CL=90%
Γ_{144}	$D_s^{*+} \eta$	< 6	$\times 10^{-4}$ CL=90%
Γ_{145}	$D_s^+ \rho^0$	< 3.1	$\times 10^{-4}$ CL=90%
Γ_{146}	$D_s^{*+} \rho^0$	< 4	$\times 10^{-4}$ CL=90%
Γ_{147}	$D_s^+ \omega$	< 4	$\times 10^{-4}$ CL=90%
Γ_{148}	$D_s^{*+} \omega$	< 6	$\times 10^{-4}$ CL=90%
Γ_{149}	$D_s^+ a_1(1260)^0$	< 1.8	$\times 10^{-3}$ CL=90%
Γ_{150}	$D_s^{*+} a_1(1260)^0$	< 1.4	$\times 10^{-3}$ CL=90%
Γ_{151}	$D_s^+ \phi$	< 1.9	$\times 10^{-6}$ CL=90%
Γ_{152}	$D_s^{*+} \phi$	< 1.2	$\times 10^{-5}$ CL=90%
Γ_{153}	$D_s^+ \bar{K}^0$	< 9	$\times 10^{-4}$ CL=90%
Γ_{154}	$D_s^{*+} \bar{K}^0$	< 9	$\times 10^{-4}$ CL=90%
Γ_{155}	$D_s^+ \bar{K}^*(892)^0$	< 4	$\times 10^{-4}$ CL=90%
Γ_{156}	$D_s^{*+} \bar{K}^*(892)^0$	< 4	$\times 10^{-4}$ CL=90%
Γ_{157}	$D_s^- \pi^+ K^+$	$(2.0 \pm 0.4) \times 10^{-4}$	
Γ_{158}	$D_s^{*-} \pi^+ K^+$	$(1.7 \pm 0.4) \times 10^{-4}$	
Γ_{159}	$D_s^- \pi^+ K^*(892)^+$	< 5	$\times 10^{-3}$ CL=90%

Γ_{160}	$D_s^{*-} \pi^+ K^*(892)^+$	< 7	$\times 10^{-3}$	CL=90%
Γ_{161}	$D_s^- K^+ K^+$	(1.1 ± 0.4)	$\times 10^{-5}$	
Γ_{162}	$D_s^{*-} K^+ K^+$	< 1.5	$\times 10^{-5}$	CL=90%

Charmonium modes

Γ_{163}	$\eta_c K^+$	(9.1 ± 1.3)	$\times 10^{-4}$	
Γ_{164}	$\eta_c K^*(892)^+$	$(1.2 \begin{smallmatrix} +0.7 \\ -0.6 \end{smallmatrix})$	$\times 10^{-3}$	
Γ_{165}	$\eta_c(2S) K^+$	(3.4 ± 1.8)	$\times 10^{-4}$	
Γ_{166}	$J/\psi(1S) K^+$	(1.007 ± 0.035)	$\times 10^{-3}$	
Γ_{167}	$J/\psi(1S) K^+ \pi^+ \pi^-$	(1.07 ± 0.19)	$\times 10^{-3}$	S=1.9
Γ_{168}	$h_c(1P) K^+ \times B(h_c(1P) \rightarrow J/\psi \pi^+ \pi^-)$	< 3.4	$\times 10^{-6}$	CL=90%
Γ_{169}	$X(3872) K^+$	< 3.2	$\times 10^{-4}$	CL=90%
Γ_{170}	$X(3872) K^+ \times B(X \rightarrow J/\psi \pi^+ \pi^-)$	(9.5 ± 1.9)	$\times 10^{-6}$	S=1.3
Γ_{171}	$X(3872) K^+ \times B(X \rightarrow J/\psi \gamma)$	(3.3 ± 1.0)	$\times 10^{-6}$	
Γ_{172}	$X(3872) K^+ \times B(X \rightarrow D^0 \bar{D}^0)$	< 6.0	$\times 10^{-5}$	CL=90%
Γ_{173}	$X(3872) K^+ \times B(X \rightarrow D^+ D^-)$	< 4.0	$\times 10^{-5}$	CL=90%
Γ_{174}	$X(3872) K^+ \times B(X \rightarrow D^0 \bar{D}^0 \pi^0)$	(1.0 ± 0.4)	$\times 10^{-4}$	
Γ_{175}	$X(3872) K^+ \times B(X \rightarrow \bar{D}^{*0} D^0)$	(1.7 ± 0.6)	$\times 10^{-4}$	
Γ_{176}	$X(3872) K^+ \times B(X(3872) \rightarrow J/\psi(1S) \eta)$	< 7.7	$\times 10^{-6}$	CL=90%
Γ_{177}	$X(3872)^+ K^0 \times B(X(3872)^+ \rightarrow J/\psi(1S) \pi^+ \pi^0)$ [f]	< 2.2	$\times 10^{-5}$	CL=90%
Γ_{178}	$X(4260)^0 K^+ \times B(X^0 \rightarrow J/\psi \pi^+ \pi^-)$	< 2.9	$\times 10^{-5}$	CL=95%
Γ_{179}	$X(3945)^0 K^+ \times B(X^0 \rightarrow J/\psi \gamma)$	< 1.4	$\times 10^{-5}$	CL=90%
Γ_{180}	$Z(3930)^0 K^+ \times B(Z^0 \rightarrow J/\psi \gamma)$	< 2.5	$\times 10^{-6}$	CL=90%
Γ_{181}	$X(3945) K^+$			
Γ_{182}	$J/\psi(1S) K^*(892)^+$	(1.43 ± 0.08)	$\times 10^{-3}$	
Γ_{183}	$J/\psi(1S) K(1270)^+$	(1.8 ± 0.5)	$\times 10^{-3}$	
Γ_{184}	$J/\psi(1S) K(1400)^+$	< 5	$\times 10^{-4}$	CL=90%
Γ_{185}	$J/\psi(1S) \eta K^+$	(1.08 ± 0.33)	$\times 10^{-4}$	
Γ_{186}	$J/\psi(1S) \eta' K^+$	< 8.8	$\times 10^{-5}$	CL=90%
Γ_{187}	$J/\psi(1S) \phi K^+$	(5.2 ± 1.7)	$\times 10^{-5}$	S=1.2
Γ_{188}	$J/\psi(1S) \omega K^+$ nonresonant	(3.5 ± 0.4)	$\times 10^{-4}$	
Γ_{189}	$J/\psi(1S) \pi^+$	(4.9 ± 0.6)	$\times 10^{-5}$	S=1.5
Γ_{190}	$J/\psi(1S) \rho^+$	(5.0 ± 0.8)	$\times 10^{-5}$	
Γ_{191}	$J/\psi(1S) \pi^+ \pi^0$ nonresonant	< 7.3	$\times 10^{-6}$	CL=90%
Γ_{192}	$J/\psi(1S) a_1(1260)^+$	< 1.2	$\times 10^{-3}$	CL=90%
Γ_{193}	$J/\psi(1S) p\bar{\Lambda}$	(1.18 ± 0.31)	$\times 10^{-5}$	

Γ_{194}	$J/\psi(1S)\bar{\Sigma}^0 p$	< 1.1	$\times 10^{-5}$	CL=90%
Γ_{195}	$J/\psi(1S)D^+$	< 1.2	$\times 10^{-4}$	CL=90%
Γ_{196}	$J/\psi(1S)\bar{D}^0 \pi^+$	< 2.5	$\times 10^{-5}$	CL=90%
Γ_{197}	$\psi(2S)\pi^+$	(2.44 ± 0.30)	$\times 10^{-5}$	
Γ_{198}	$\psi(2S)K^+$	(6.48 ± 0.35)	$\times 10^{-4}$	
Γ_{199}	$\psi(2S)K^*(892)^+$	(6.7 ± 1.4)	$\times 10^{-4}$	S=1.3
Γ_{200}	$\psi(2S)K^+\pi^+\pi^-$	(1.9 ± 1.2)	$\times 10^{-3}$	
Γ_{201}	$\psi(3770)K^+$	(4.9 ± 1.3)	$\times 10^{-4}$	
Γ_{202}	$\psi(3770)K^+ \times B(\psi \rightarrow D^0\bar{D}^0)$	(1.6 ± 0.4)	$\times 10^{-4}$	S=1.1
Γ_{203}	$\psi(3770)K^+ \times B(\psi \rightarrow D^+D^-)$	(9.4 ± 3.5)	$\times 10^{-5}$	
Γ_{204}	$\chi_{c0}\pi^+ \times B(\chi_{c0} \rightarrow \pi^+\pi^-)$	< 3	$\times 10^{-7}$	CL=90%
Γ_{205}	$\chi_{c0}(1P)K^+$	(1.43 ± 0.21)	$\times 10^{-4}$	
Γ_{206}	$\chi_{c0}K^*(892)^+$	< 2.1	$\times 10^{-4}$	CL=90%
Γ_{207}	$\chi_{c2}K^+$	< 2.9	$\times 10^{-5}$	CL=90%
Γ_{208}	$\chi_{c2}K^*(892)^+$	< 1.2	$\times 10^{-5}$	CL=90%
Γ_{209}	$\chi_{c1}(1P)\pi^+$	(2.2 ± 0.5)	$\times 10^{-5}$	
Γ_{210}	$\chi_{c1}(1P)K^+$	(5.1 ± 0.5)	$\times 10^{-4}$	S=1.6
Γ_{211}	$\chi_{c1}(1P)K^*(892)^+$	(3.6 ± 0.9)	$\times 10^{-4}$	
Γ_{212}	$h_c(1P)K^+$	< 3.8	$\times 10^{-5}$	

K or K* modes

Γ_{213}	$K^0\pi^+$	(2.31 ± 0.10)	$\times 10^{-5}$	
Γ_{214}	$K^+\pi^0$	(1.29 ± 0.06)	$\times 10^{-5}$	
Γ_{215}	$\eta'K^+$	(7.00 ± 0.24)	$\times 10^{-5}$	
Γ_{216}	$\eta'K^*(892)^+$	(4.9 ± 2.0)	$\times 10^{-6}$	
Γ_{217}	ηK^+	(2.6 ± 0.5)	$\times 10^{-6}$	S=2.4
Γ_{218}	$\eta K^*(892)^+$	(1.93 ± 0.16)	$\times 10^{-5}$	
Γ_{219}	$\eta K_0^*(1430)^+$	(1.8 ± 0.4)	$\times 10^{-5}$	
Γ_{220}	$\eta K_2^*(1430)^+$	(9.1 ± 3.0)	$\times 10^{-6}$	
Γ_{221}	$\eta(1295)K^+ \times B(\eta(1295) \rightarrow \eta\pi\pi)$	(2.9 ± 0.8)	$\times 10^{-6}$	
Γ_{222}	$\eta(1405)K^+ \times B(\eta(1405) \rightarrow \eta\pi\pi)$	< 1.3	$\times 10^{-6}$	CL=90%
Γ_{223}	$\eta(1405)K^+ \times B(\eta(1405) \rightarrow K^*K)$	< 1.2	$\times 10^{-6}$	CL=90%
Γ_{224}	$\eta(1475)K^+ \times B(\eta(1475) \rightarrow K^*K)$	(1.38 ± 0.21)	$\times 10^{-5}$	
Γ_{225}	$f_1(1285)K^+$	< 2.0	$\times 10^{-6}$	CL=90%
Γ_{226}	$f_1(1420)K^+ \times B(f_1(1420) \rightarrow \eta\pi\pi)$	< 2.9	$\times 10^{-6}$	CL=90%
Γ_{227}	$f_1(1420)K^+ \times B(f_1(1420) \rightarrow K^*K)$	< 4.1	$\times 10^{-6}$	CL=90%

Γ_{228}	$\phi(1680) K^+ \times B(\phi(1680) \rightarrow K^* K)$	<	3.4	$\times 10^{-6}$	CL=90%
Γ_{229}	ωK^+	(6.7 ± 0.8) $\times 10^{-6}$	S=1.8
Γ_{230}	$\omega K^*(892)^+$	<	7.4	$\times 10^{-6}$	CL=90%
Γ_{231}	$\omega (K\pi)_0^{*+}$	(2.7 ± 0.4) $\times 10^{-5}$	
Γ_{232}	$\omega K_0^*(1430)^+$	(2.4 ± 0.5) $\times 10^{-5}$	
Γ_{233}	$\omega K_2^*(1430)^+$	(2.1 ± 0.4) $\times 10^{-5}$	
Γ_{234}	$a_0(980)^+ K^0 \times B(a_0(980)^+ \rightarrow \eta \pi^+)$	<	3.9	$\times 10^{-6}$	CL=90%
Γ_{235}	$a_0(980)^0 K^+ \times B(a_0(980)^0 \rightarrow \eta \pi^0)$	<	2.5	$\times 10^{-6}$	CL=90%
Γ_{236}	$K^*(892)^0 \pi^+$	(1.01 ± 0.09) $\times 10^{-5}$	
Γ_{237}	$K^*(892)^+ \pi^0$	(6.9 ± 2.4) $\times 10^{-6}$	
Γ_{238}	$K^+ \pi^- \pi^+$	(5.10 ± 0.29) $\times 10^{-5}$	
Γ_{239}	$K^+ \pi^- \pi^+$ nonresonant	(1.63 $^{+0.21}_{-0.15}$) $\times 10^{-5}$	
Γ_{240}	$\omega(782) K^+$	(6 ± 9) $\times 10^{-6}$	
Γ_{241}	$K^+ f_0(980) \times B(f_0(980) \rightarrow \pi^+ \pi^-)$	(9.4 $^{+1.0}_{-1.2}$) $\times 10^{-6}$	
Γ_{242}	$f_2(1270)^0 K^+$	(1.07 ± 0.27) $\times 10^{-6}$	
Γ_{243}	$f_0(1370)^0 K^+ \times B(f_0(1370)^0 \rightarrow \pi^+ \pi^-)$	<	1.07	$\times 10^{-5}$	CL=90%
Γ_{244}	$\rho^0(1450) K^+ \times B(\rho^0(1450) \rightarrow \pi^+ \pi^-)$	<	1.17	$\times 10^{-5}$	CL=90%
Γ_{245}	$K^+ f_X(1300) \times B(f_X \rightarrow \pi^+ \pi^-)$	(7 ± 5) $\times 10^{-7}$	
Γ_{246}	$f_0(1500) K^+ \times B(f_0(1500) \rightarrow \pi^+ \pi^-)$	<	4.4	$\times 10^{-6}$	CL=90%
Γ_{247}	$f_2'(1525) K^+ \times B(f_2'(1525) \rightarrow \pi^+ \pi^-)$	<	3.4	$\times 10^{-6}$	CL=90%
Γ_{248}	$K^+ \rho^0$	(3.7 ± 0.5) $\times 10^{-6}$	
Γ_{249}	$K_0^*(1430)^0 \pi^+$	(4.5 $^{+0.9}_{-0.7}$) $\times 10^{-5}$	S=1.5
Γ_{250}	$K_2^*(1430)^0 \pi^+$	(5.6 $^{+2.2}_{-1.5}$) $\times 10^{-6}$	
Γ_{251}	$K^*(1410)^0 \pi^+$	<	4.5	$\times 10^{-5}$	CL=90%
Γ_{252}	$K^*(1680)^0 \pi^+$	<	1.2	$\times 10^{-5}$	CL=90%
Γ_{253}	$K^- \pi^+ \pi^+$	<	9.5	$\times 10^{-7}$	CL=90%
Γ_{254}	$K^- \pi^+ \pi^+$ nonresonant	<	5.6	$\times 10^{-5}$	CL=90%
Γ_{255}	$K_1(1400)^0 \pi^+$	<	2.6	$\times 10^{-3}$	CL=90%
Γ_{256}	$K^0 \pi^+ \pi^0$	<	6.6	$\times 10^{-5}$	CL=90%
Γ_{257}	$K^0 \rho^+$	(8.0 ± 1.5) $\times 10^{-6}$	
Γ_{258}	$K^*(892)^+ \pi^+ \pi^-$	(7.5 ± 1.0) $\times 10^{-5}$	
Γ_{259}	$K^*(892)^+ \rho^0$	<	6.1	$\times 10^{-6}$	CL=90%
Γ_{260}	$K^*(892)^+ f_0(980)$	(5.2 ± 1.3) $\times 10^{-6}$	

Γ_{261}	$a_1^+ K^0$	$(3.5 \pm 0.7) \times 10^{-5}$	
Γ_{262}	$b_1^+ K^0 \times B(b_1^+ \rightarrow \omega \pi^+)$	$(9.6 \pm 1.9) \times 10^{-6}$	
Γ_{263}	$K^*(892)^0 \rho^+$	$(9.2 \pm 1.5) \times 10^{-6}$	
Γ_{264}	$K_1(1400)^+ \rho^0$	$< 7.8 \times 10^{-4}$	CL=90%
Γ_{265}	$K_2^*(1430)^+ \rho^0$	$< 1.5 \times 10^{-3}$	CL=90%
Γ_{266}	$b_1^0 K^+ \times B(b_1^0 \rightarrow \omega \pi^0)$	$(9.1 \pm 2.0) \times 10^{-6}$	
Γ_{267}	$K^+ \bar{K}^0$	$(1.36 \pm 0.27) \times 10^{-6}$	
Γ_{268}	$\bar{K}^0 K^+ \pi^0$	$< 2.4 \times 10^{-5}$	CL=90%
Γ_{269}	$K^+ K_S^0 K_S^0$	$(1.15 \pm 0.13) \times 10^{-5}$	
Γ_{270}	$K_S^0 K_S^0 \pi^+$	$< 5.1 \times 10^{-7}$	CL=90%
Γ_{271}	$K^+ K^- \pi^+$	$(5.0 \pm 0.7) \times 10^{-6}$	
Γ_{272}	$K^+ K^- \pi^+$ nonresonant	$< 7.5 \times 10^{-5}$	CL=90%
Γ_{273}	$K^+ \bar{K}^*(892)^0$	$< 1.1 \times 10^{-6}$	CL=90%
Γ_{274}	$K^+ \bar{K}_0^*(1430)^0$	$< 2.2 \times 10^{-6}$	CL=90%
Γ_{275}	$K^+ K^+ \pi^-$	$< 1.6 \times 10^{-7}$	CL=90%
Γ_{276}	$K^+ K^+ \pi^-$ nonresonant	$< 8.79 \times 10^{-5}$	CL=90%
Γ_{277}	$K^+ f_J(2220)$		
Γ_{278}	$K^{*+} \pi^+ K^-$	$< 1.18 \times 10^{-5}$	CL=90%
Γ_{279}	$K^*(892)^+ K^*(892)^0$	$(1.2 \pm 0.5) \times 10^{-6}$	
Γ_{280}	$K^{*+} K^+ \pi^-$	$< 6.1 \times 10^{-6}$	CL=90%
Γ_{281}	$K^+ K^- K^+$	$(3.37 \pm 0.22) \times 10^{-5}$	S=1.4
Γ_{282}	$K^+ \phi$	$(8.3 \pm 0.7) \times 10^{-6}$	
Γ_{283}	$f_0(980) K^+ \times B(f_0(980) \rightarrow K^+ K^-)$	$< 2.9 \times 10^{-6}$	CL=90%
Γ_{284}	$a_2(1320) K^+ \times B(a_2(1320) \rightarrow K^+ K^-)$	$< 1.1 \times 10^{-6}$	CL=90%
Γ_{285}	$f_2'(1525) K^+ \times B(f_2'(1525) \rightarrow K^+ K^-)$	$< 4.9 \times 10^{-6}$	CL=90%
Γ_{286}	$X_0(1550) K^+ \times B(X_0(1550) \rightarrow K^+ K^-)$	$(4.3 \pm 0.7) \times 10^{-6}$	
Γ_{287}	$\phi(1680) K^+ \times B(\phi(1680) \rightarrow K^+ K^-)$	$< 8 \times 10^{-7}$	CL=90%
Γ_{288}	$f_0(1710) K^+ \times B(f_0(1710) \rightarrow K^+ K^-)$	$(1.7 \pm 1.0) \times 10^{-6}$	
Γ_{289}	$K^+ K^- K^+$ nonresonant	$(2.8 \pm_{-1.6}^{+0.9}) \times 10^{-5}$	S=3.3
Γ_{290}	$K^*(892)^+ K^+ K^-$	$(3.6 \pm 0.5) \times 10^{-5}$	
Γ_{291}	$K^*(892)^+ \phi$	$(1.05 \pm 0.15) \times 10^{-5}$	S=1.4
Γ_{292}	$\phi(K\pi)_0^{*+}$	$(8.3 \pm 1.6) \times 10^{-6}$	
Γ_{293}	$\phi K_1(1270)^+$	$(6.1 \pm 1.9) \times 10^{-6}$	
Γ_{294}	$\phi K_1(1400)^+$	$< 3.2 \times 10^{-6}$	CL=90%
Γ_{295}	$\phi K^*(1410)^+$	$< 4.3 \times 10^{-6}$	CL=90%
Γ_{296}	$\phi K_0^*(1430)^+$	$(7.0 \pm 1.6) \times 10^{-6}$	
Γ_{297}	$\phi K_2^*(1430)^+$	$(8.4 \pm 2.1) \times 10^{-6}$	

Γ_{298}	$\phi K_2^*(1770)^+$	< 1.50	$\times 10^{-5}$	CL=90%
Γ_{299}	$\phi K_2^*(1820)^+$	< 1.63	$\times 10^{-5}$	CL=90%
Γ_{300}	$K^+ \phi \phi$	$(4.9 \begin{smallmatrix} +2.4 \\ -2.2 \end{smallmatrix})$	$\times 10^{-6}$	S=2.9
Γ_{301}	$\eta' \eta' K^+$	< 2.5	$\times 10^{-5}$	CL=90%
Γ_{302}	$K^*(892)^+ \gamma$	(4.03 ± 0.26)	$\times 10^{-5}$	
Γ_{303}	$K_1(1270)^+ \gamma$	(4.3 ± 1.3)	$\times 10^{-5}$	
Γ_{304}	$\eta K^+ \gamma$	(7.9 ± 0.9)	$\times 10^{-6}$	
Γ_{305}	$\eta' K^+ \gamma$	< 4.2	$\times 10^{-6}$	CL=90%
Γ_{306}	$\phi K^+ \gamma$	(3.5 ± 0.6)	$\times 10^{-6}$	
Γ_{307}	$K^+ \pi^- \pi^+ \gamma$	(2.76 ± 0.22)	$\times 10^{-5}$	S=1.2
Γ_{308}	$K^*(892)^0 \pi^+ \gamma$	$(2.0 \begin{smallmatrix} +0.7 \\ -0.6 \end{smallmatrix})$	$\times 10^{-5}$	
Γ_{309}	$K^+ \rho^0 \gamma$	< 2.0	$\times 10^{-5}$	CL=90%
Γ_{310}	$K^+ \pi^- \pi^+ \gamma$ nonresonant	< 9.2	$\times 10^{-6}$	CL=90%
Γ_{311}	$K^0 \pi^+ \pi^0 \gamma$	(4.6 ± 0.5)	$\times 10^{-5}$	
Γ_{312}	$K_1(1400)^+ \gamma$	< 1.5	$\times 10^{-5}$	CL=90%
Γ_{313}	$K_2^*(1430)^+ \gamma$	(1.4 ± 0.4)	$\times 10^{-5}$	
Γ_{314}	$K^*(1680)^+ \gamma$	< 1.9	$\times 10^{-3}$	CL=90%
Γ_{315}	$K_3^*(1780)^+ \gamma$	< 3.9	$\times 10^{-5}$	CL=90%
Γ_{316}	$K_4^*(2045)^+ \gamma$	< 9.9	$\times 10^{-3}$	CL=90%

Light unflavored meson modes

Γ_{317}	$\rho^+ \gamma$	(9.8 ± 2.5)	$\times 10^{-7}$	
Γ_{318}	$\pi^+ \pi^0$	(5.7 ± 0.5)	$\times 10^{-6}$	S=1.4
Γ_{319}	$\pi^+ \pi^+ \pi^-$	(1.62 ± 0.15)	$\times 10^{-5}$	
Γ_{320}	$\rho^0 \pi^+$	(8.7 ± 1.1)	$\times 10^{-6}$	
Γ_{321}	$\pi^+ f_0(980) \times B(f_0(980) \rightarrow \pi^+ \pi^-)$	< 3.0	$\times 10^{-6}$	CL=90%
Γ_{322}	$\pi^+ f_2(1270)$	(8.2 ± 2.5)	$\times 10^{-6}$	
Γ_{323}	$\rho(1450)^0 \pi^+$	< 2.3	$\times 10^{-6}$	CL=90%
Γ_{324}	$f_0(1370) \pi^+ \times B(f_0(1370) \rightarrow \pi^+ \pi^-)$	< 3.0	$\times 10^{-6}$	CL=90%
Γ_{325}	$f_0(600) \pi^+ \times B(f_0(600) \rightarrow \pi^+ \pi^-)$	< 4.1	$\times 10^{-6}$	CL=90%
Γ_{326}	$\pi^+ \pi^- \pi^+$ nonresonant	< 4.6	$\times 10^{-6}$	CL=90%
Γ_{327}	$\pi^+ \pi^0 \pi^0$	< 8.9	$\times 10^{-4}$	CL=90%
Γ_{328}	$\rho^+ \pi^0$	(1.09 ± 0.14)	$\times 10^{-5}$	
Γ_{329}	$\pi^+ \pi^- \pi^+ \pi^0$	< 4.0	$\times 10^{-3}$	CL=90%
Γ_{330}	$\rho^+ \rho^0$	(2.40 ± 0.19)	$\times 10^{-5}$	
Γ_{331}	$\rho^+ f_0(980) \times B(f_0(980) \rightarrow \pi^+ \pi^-)$	< 2.0	$\times 10^{-6}$	CL=90%
Γ_{332}	$a_1(1260)^+ \pi^0$	(2.6 ± 0.7)	$\times 10^{-5}$	
Γ_{333}	$a_1(1260)^0 \pi^+$	(2.0 ± 0.6)	$\times 10^{-5}$	

Γ_{334}	$\omega\pi^+$	$(6.9 \pm 0.5) \times 10^{-6}$	
Γ_{335}	$\omega\rho^+$	$(1.59 \pm 0.21) \times 10^{-5}$	
Γ_{336}	$\eta\pi^+$	$(4.4 \pm 0.4) \times 10^{-6}$	S=1.1
Γ_{337}	$\eta\rho^+$	$(7.0 \pm 2.9) \times 10^{-6}$	S=2.8
Γ_{338}	$\eta'\pi^+$	$(2.7 \pm 1.0) \times 10^{-6}$	S=2.1
Γ_{339}	$\eta'\rho^+$	$(8.7 \begin{smallmatrix} +4.0 \\ -3.1 \end{smallmatrix}) \times 10^{-6}$	
Γ_{340}	$\phi\pi^+$	< 2.4	$\times 10^{-7}$ CL=90%
Γ_{341}	$\phi\rho^+$	< 3.0	$\times 10^{-6}$ CL=90%
Γ_{342}	$a_0(980)^0\pi^+ \times B(a_0(980)^0 \rightarrow \eta\pi^0)$	< 5.8	$\times 10^{-6}$ CL=90%
Γ_{343}	$a_0(980)^+\pi^0 \times B(a_0^+ \rightarrow \eta\pi^+)$	< 1.4	$\times 10^{-6}$ CL=90%
Γ_{344}	$\pi^+\pi^+\pi^+\pi^-\pi^-$	< 8.6	$\times 10^{-4}$ CL=90%
Γ_{345}	$\rho^0 a_1(1260)^+$	< 6.2	$\times 10^{-4}$ CL=90%
Γ_{346}	$\rho^0 a_2(1320)^+$	< 7.2	$\times 10^{-4}$ CL=90%
Γ_{347}	$b_1^0\pi^+ \times B(b_1^0 \rightarrow \omega\pi^0)$	$(6.7 \pm 2.0) \times 10^{-6}$	
Γ_{348}	$b_1^+\pi^0 \times B(b_1^+ \rightarrow \omega\pi^+)$	< 3.3	$\times 10^{-6}$ CL=90%
Γ_{349}	$\pi^+\pi^+\pi^+\pi^-\pi^-\pi^0$	< 6.3	$\times 10^{-3}$ CL=90%
Γ_{350}	$a_1(1260)^+ a_1(1260)^0$	< 1.3	% CL=90%

Charged particle (h^\pm) modes

$$h^\pm = K^\pm \text{ or } \pi^\pm$$

Γ_{351}	$h^+\pi^0$	$(1.6 \begin{smallmatrix} +0.7 \\ -0.6 \end{smallmatrix}) \times 10^{-5}$	
Γ_{352}	ωh^+	$(1.38 \begin{smallmatrix} +0.27 \\ -0.24 \end{smallmatrix}) \times 10^{-5}$	
Γ_{353}	$h^+ X^0$ (Familon)	< 4.9	$\times 10^{-5}$ CL=90%

Baryon modes

Γ_{354}	$\rho\bar{p}\pi^+$	$(1.62 \pm 0.20) \times 10^{-6}$	
Γ_{355}	$\rho\bar{p}\pi^+$ nonresonant	< 5.3	$\times 10^{-5}$ CL=90%
Γ_{356}	$\rho\bar{p}\pi^+\pi^+\pi^-$		
Γ_{357}	$\rho\bar{p}K^+$	$(5.9 \pm 0.5) \times 10^{-6}$	S=1.5
Γ_{358}	$\Theta(1710)^{++}\bar{p} \times B(\Theta(1710)^{++} \rightarrow \rho K^+)$	[g] < 9.1	$\times 10^{-8}$ CL=90%
Γ_{359}	$f_J(2220)K^+ \times B(f_J(2220) \rightarrow \rho\bar{p})$	[g] < 4.1	$\times 10^{-7}$ CL=90%
Γ_{360}	$\rho\bar{\Lambda}(1520)$	< 1.5	$\times 10^{-6}$ CL=90%
Γ_{361}	$\rho\bar{p}K^+$ nonresonant	< 8.9	$\times 10^{-5}$ CL=90%
Γ_{362}	$\rho\bar{p}K^*(892)^+$	$(3.6 \begin{smallmatrix} +0.8 \\ -0.7 \end{smallmatrix}) \times 10^{-6}$	
Γ_{363}	$f_J(2220)K^{*+} \times B(f_J(2220) \rightarrow \rho\bar{p})$	< 7.7	$\times 10^{-7}$ CL=90%
Γ_{364}	$\rho\bar{\Lambda}$	< 3.2	$\times 10^{-7}$ CL=90%
Γ_{365}	$\rho\bar{\Lambda}\gamma$	$(2.5 \begin{smallmatrix} +0.5 \\ -0.4 \end{smallmatrix}) \times 10^{-6}$	

Γ_{366}	$\rho \bar{\Lambda} \pi^0$		$(3.0 \begin{smallmatrix} +0.7 \\ -0.6 \end{smallmatrix}) \times 10^{-6}$	
Γ_{367}	$\rho \bar{\Sigma}(1385)^0$	$<$	4.7×10^{-7}	CL=90%
Γ_{368}	$\Delta^+ \bar{\Lambda}$	$<$	8.2×10^{-7}	CL=90%
Γ_{369}	$\rho \bar{\Sigma} \gamma$	$<$	4.6×10^{-6}	CL=90%
Γ_{370}	$\rho \bar{\Lambda} \pi^+ \pi^-$	$<$	2.0×10^{-4}	CL=90%
Γ_{371}	$\Lambda \bar{\Lambda} \pi^+$	$<$	2.8×10^{-6}	CL=90%
Γ_{372}	$\Lambda \bar{\Lambda} K^+$		$(2.9 \begin{smallmatrix} +1.0 \\ -0.8 \end{smallmatrix}) \times 10^{-6}$	
Γ_{373}	$\bar{\Delta}^0 \rho$	$<$	1.38×10^{-6}	CL=90%
Γ_{374}	$\Delta^{++} \bar{p}$	$<$	1.4×10^{-7}	CL=90%
Γ_{375}	$D^+ p \bar{p}$	$<$	1.5×10^{-5}	CL=90%
Γ_{376}	$D^*(2010)^+ p \bar{p}$	$<$	1.5×10^{-5}	CL=90%
Γ_{377}	$\bar{\Lambda}_c^- p \pi^+$		$(2.8 \pm 0.8) \times 10^{-4}$	
Γ_{378}	$\bar{\Lambda}_c^- \Delta(1232)^{++}$	$<$	1.9×10^{-5}	CL=90%
Γ_{379}	$\bar{\Lambda}_c^- \Delta_X(1600)^{++}$		$(5.9 \pm 1.9) \times 10^{-5}$	
Γ_{380}	$\bar{\Lambda}_c^- \Delta_X(2420)^{++}$		$(4.7 \pm 1.6) \times 10^{-5}$	
Γ_{381}	$(\bar{\Lambda}_c^- p)_s \pi^+$	[h]	$(3.9 \pm 1.3) \times 10^{-5}$	
Γ_{382}	$\bar{\Sigma}_c(2520)^0 p$	$<$	3×10^{-6}	CL=90%
Γ_{383}	$\bar{\Sigma}_c(2800)^0 p$		$(3.3 \pm 1.3) \times 10^{-5}$	
Γ_{384}	$\bar{\Lambda}_c^- p \pi^+ \pi^0$		$(1.8 \pm 0.6) \times 10^{-3}$	
Γ_{385}	$\bar{\Lambda}_c^- p \pi^+ \pi^+ \pi^-$		$(2.3 \pm 0.7) \times 10^{-3}$	
Γ_{386}	$\bar{\Lambda}_c^- p \pi^+ \pi^+ \pi^- \pi^0$	$<$	1.34%	CL=90%
Γ_{387}	$\Lambda_c^+ \Lambda_c^- K^+$		$(8.7 \pm 3.5) \times 10^{-4}$	
Γ_{388}	$\bar{\Sigma}_c(2455)^0 p$		$(3.7 \pm 1.3) \times 10^{-5}$	
Γ_{389}	$\bar{\Sigma}_c(2455)^0 p \pi^0$		$(4.4 \pm 1.8) \times 10^{-4}$	
Γ_{390}	$\bar{\Sigma}_c(2455)^0 p \pi^- \pi^+$		$(4.4 \pm 1.7) \times 10^{-4}$	
Γ_{391}	$\bar{\Sigma}_c(2455)^{--} p \pi^+ \pi^+$		$(2.8 \pm 1.2) \times 10^{-4}$	
Γ_{392}	$\bar{\Lambda}_c(2593)^- / \bar{\Lambda}_c(2625)^- p \pi^+$	$<$	1.9×10^{-4}	CL=90%
Γ_{393}	$\Xi_c^0 \Lambda_c^+ \times B(\Xi_c^0 \rightarrow \Xi^+ \pi^-)$		$(3.0 \pm 1.1) \times 10^{-5}$	
Γ_{394}	$\Xi_c^0 \Lambda_c^+ \times B(\Xi_c^0 \rightarrow \Lambda K^+ \pi^-)$		$(2.6 \pm 1.1) \times 10^{-5}$	S=1.1

Lepton Family number (LF) or Lepton number (L) violating modes, or $\Delta B = 1$ weak neutral current (B1) modes

Γ_{395}	$\pi^+ \ell^+ \ell^-$	B1	$<$	4.9×10^{-8}	CL=90%
Γ_{396}	$\pi^+ e^+ e^-$	B1	$<$	8.0×10^{-8}	CL=90%
Γ_{397}	$\pi^+ \mu^+ \mu^-$	B1	$<$	6.9×10^{-8}	CL=90%
Γ_{398}	$\pi^+ \nu \bar{\nu}$	B1	$<$	1.0×10^{-4}	CL=90%
Γ_{399}	$K^+ \ell^+ \ell^-$	B1	[a]	$(4.4 \begin{smallmatrix} +0.8 \\ -0.7 \end{smallmatrix}) \times 10^{-7}$	S=1.1
Γ_{400}	$K^+ e^+ e^-$	B1		$(4.9 \pm 1.0) \times 10^{-7}$	
Γ_{401}	$K^+ \mu^+ \mu^-$	B1		$(4.5 \begin{smallmatrix} +0.9 \\ -0.8 \end{smallmatrix}) \times 10^{-7}$	
Γ_{402}	$K^+ \bar{\nu} \nu$	B1	$<$	1.4×10^{-5}	CL=90%
Γ_{403}	$\rho^+ \nu \bar{\nu}$	B1	$<$	1.5×10^{-4}	CL=90%

Γ_{404}	$K^*(892)^+ \ell^+ \ell^-$	$B1$	[a] (7 ± 5)	$\times 10^{-7}$	
Γ_{405}	$K^*(892)^+ \nu \bar{\nu}$	$B1$	< 8	$\times 10^{-5}$	CL=90%
Γ_{406}	$K^*(892)^+ e^+ e^-$	$B1$	(8 ± 8)	$\times 10^{-7}$	
Γ_{407}	$K^*(892)^+ \mu^+ \mu^-$	$B1$	(8 $\begin{smallmatrix} +6 \\ -4 \end{smallmatrix}$)	$\times 10^{-7}$	
Γ_{408}	$\pi^+ e^+ \mu^-$	LF	< 6.4	$\times 10^{-3}$	CL=90%
Γ_{409}	$\pi^+ e^- \mu^+$	LF	< 6.4	$\times 10^{-3}$	CL=90%
Γ_{410}	$\pi^+ e^\pm \mu^\mp$	LF	< 1.7	$\times 10^{-7}$	CL=90%
Γ_{411}	$K^+ e^+ \mu^-$	LF	< 9.1	$\times 10^{-8}$	CL=90%
Γ_{412}	$K^+ e^- \mu^+$	LF	< 1.3	$\times 10^{-7}$	CL=90%
Γ_{413}	$K^+ e^\pm \mu^\mp$	LF	< 9.1	$\times 10^{-8}$	CL=90%
Γ_{414}	$K^+ \mu^\pm \tau^\mp$	LF	< 7.7	$\times 10^{-5}$	CL=90%
Γ_{415}	$K^*(892)^+ e^+ \mu^-$	LF	< 1.3	$\times 10^{-6}$	CL=90%
Γ_{416}	$K^*(892)^+ e^- \mu^+$	LF	< 9.9	$\times 10^{-7}$	CL=90%
Γ_{417}	$K^*(892)^+ e^\pm \mu^\mp$	LF	< 1.4	$\times 10^{-7}$	CL=90%
Γ_{418}	$\pi^- e^+ e^+$	L	< 1.6	$\times 10^{-6}$	CL=90%
Γ_{419}	$\pi^- \mu^+ \mu^+$	L	< 1.4	$\times 10^{-6}$	CL=90%
Γ_{420}	$\pi^- e^+ \mu^+$	L	< 1.3	$\times 10^{-6}$	CL=90%
Γ_{421}	$\rho^- e^+ e^+$	L	< 2.6	$\times 10^{-6}$	CL=90%
Γ_{422}	$\rho^- \mu^+ \mu^+$	L	< 5.0	$\times 10^{-6}$	CL=90%
Γ_{423}	$\rho^- e^+ \mu^+$	L	< 3.3	$\times 10^{-6}$	CL=90%
Γ_{424}	$K^- e^+ e^+$	L	< 1.0	$\times 10^{-6}$	CL=90%
Γ_{425}	$K^- \mu^+ \mu^+$	L	< 1.8	$\times 10^{-6}$	CL=90%
Γ_{426}	$K^- e^+ \mu^+$	L	< 2.0	$\times 10^{-6}$	CL=90%
Γ_{427}	$K^*(892)^- e^+ e^+$	L	< 2.8	$\times 10^{-6}$	CL=90%
Γ_{428}	$K^*(892)^- \mu^+ \mu^+$	L	< 8.3	$\times 10^{-6}$	CL=90%
Γ_{429}	$K^*(892)^- e^+ \mu^+$	L	< 4.4	$\times 10^{-6}$	CL=90%

[a] An ℓ indicates an e or a μ mode, not a sum over these modes.

[b] An $CP(\pm 1)$ indicates the $CP=+1$ and $CP=-1$ eigenstates of the D^0 - \bar{D}^0 system.

[c] D denotes D^0 or \bar{D}^0 .

[d] D_{CP+}^{*0} decays into $D^0 \pi^0$ with the D^0 reconstructed in CP -even eigenstates $K^+ K^-$ and $\pi^+ \pi^-$.

[e] \bar{D}^{**} represents an excited state with mass $2.2 < M < 2.8$ GeV/ c^2 .

[f] $X(3872)^+$ is a hypothetical charged partner of the $X(3872)$.

[g] $\Theta(1710)^{++}$ is a possible narrow pentaquark state and $G(2220)$ is a possible glueball resonance.

[h] $(\bar{\Lambda}_c^- \rho)_s$ denotes a low-mass enhancement near 3.35 GeV/ c^2 .

CONSTRAINED FIT INFORMATION

An overall fit to 3 branching ratios uses 11 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 7.2$ for 9 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_{189}	24
x_{166}	

B^+ BRANCHING RATIOS

$\Gamma(\ell^+ \nu_\ell \text{ anything}) / \Gamma_{\text{total}}$ Γ_1 / Γ

"OUR EVALUATION" is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements.

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
10.99 ± 0.28 OUR EVALUATION			
10.76 ± 0.32 OUR AVERAGE	Error includes scale factor of 1.1.		
11.17 ± 0.25 ± 0.28	¹ URQUIJO	07	BELL $e^+ e^- \rightarrow \gamma(4S)$
10.28 ± 0.26 ± 0.39	² AUBERT,B	06Y	BABR $e^+ e^- \rightarrow \gamma(4S)$
10.25 ± 0.57 ± 0.65	³ ARTUSO	97	CLE2 $e^+ e^- \rightarrow \gamma(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
11.15 ± 0.26 ± 0.41	⁴ OKABE	05	BELL Repl. by URQUIJO 07
10.1 ± 1.8 ± 1.5	ATHANAS	94	CLE2 Sup. by ARTUSO 97

¹ URQUIJO 07 report a measurement of $(10.34 \pm 0.23 \pm 0.25)\%$ for the partial branching fraction of $B^+ \rightarrow e^+ \nu_e X_c$ decay with electron energy above 0.6 GeV. We converted the result to $B^+ \rightarrow e^+ \nu_e X$ branching fraction.

² The measurements are obtained for charged and neutral B mesons partial rates of semileptonic decay to electrons with momentum above 0.6 GeV/c in the B rest frame. The best precision on the ratio is achieved for a momentum threshold of 1.0 GeV: $B(B^+ \rightarrow e^+ \nu_e X) / B(B^0 \rightarrow e^+ \nu_e X) = 1.074 \pm 0.041 \pm 0.026$.

³ ARTUSO 97 uses partial reconstruction of $B \rightarrow D^* \ell \nu_\ell$ and inclusive semileptonic branching ratio from BARISH 96B ($0.1049 \pm 0.0017 \pm 0.0043$).

⁴ The measurements are obtained for charged and neutral B mesons partial rates of semileptonic decay to electrons with momentum above 0.6 GeV/c in the B rest frame, and their ratio of $B(B^+ \rightarrow e^+ \nu_e X) / B(B^0 \rightarrow e^+ \nu_e X) = 1.08 \pm 0.05 \pm 0.02$.

$\Gamma(e^+ \nu_e X_c) / \Gamma_{\text{total}}$ Γ_2 / Γ

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
10.79 ± 0.25 ± 0.27	¹ URQUIJO	07	BELL $e^+ e^- \rightarrow \gamma(4S)$

¹ Measure the independent B^+ and B^0 partial branching fractions with electron threshold energies of 0.4 GeV.

$\Gamma(\bar{D}^0 \ell^+ \nu_\ell) / \Gamma_{\text{total}}$ Γ_4 / Γ

“OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements. $\ell = e$ or μ , not sum over e and μ modes.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

0.0224 ± 0.0011 OUR EVALUATION

0.0229 ± 0.0011 OUR AVERAGE

0.0234 ± 0.0003 ± 0.0013	AUBERT	09A	BABR $e^+ e^- \rightarrow \gamma(4S)$
--------------------------	--------	-----	---------------------------------------

0.0221 ± 0.0013 ± 0.0019	¹ BARTELT	99	CLE2 $e^+ e^- \rightarrow \gamma(4S)$
--------------------------	----------------------	----	---------------------------------------

0.016 ± 0.006 ± 0.003	² FULTON	91	CLEO $e^+ e^- \rightarrow \gamma(4S)$
-----------------------	---------------------	----	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0233 ± 0.0009 ± 0.0009	³ AUBERT	08Q	BABR Repl. by AUBERT 09A
--------------------------	---------------------	-----	--------------------------

0.0194 ± 0.0015 ± 0.0034	⁴ ATHANAS	97	CLE2 Repl. by BARTELT 99
--------------------------	----------------------	----	--------------------------

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² FULTON 91 assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at the $\gamma(4S)$.

³ Uses a fully reconstructed B meson as a tag on the recoil side.

⁴ ATHANAS 97 uses missing energy and missing momentum to reconstruct neutrino.

$\Gamma(\bar{D}^0 \tau^+ \nu_\tau) / \Gamma_{\text{total}}$ Γ_5 / Γ

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

0.67 ± 0.37 ± 0.13	¹ AUBERT	08N	BABR $e^+ e^- \rightarrow \gamma(4S)$
---------------------------	---------------------	-----	---------------------------------------

¹ Uses a fully reconstructed B meson as a tag on the recoil side.

$\Gamma(\bar{D}^0 \ell^+ \nu_\ell) / \Gamma(D \ell^+ \nu_\ell \text{ anything})$ Γ_4 / Γ_3

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

0.227 ± 0.014 ± 0.016	¹ AUBERT	07AN	BABR $e^+ e^- \rightarrow \gamma(4S)$
------------------------------	---------------------	------	---------------------------------------

¹ Uses a fully reconstructed B meson on the recoil side.

$\Gamma(\bar{D}^{*0(2007)} \ell^+ \nu_\ell) / \Gamma_{\text{total}}$ Γ_6 / Γ

“OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements. $\ell = e$ or μ , not sum over e and μ modes.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

0.0568 ± 0.0019 OUR EVALUATION

0.0558 ± 0.0026 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.

0.0540 ± 0.0002 ± 0.0021	AUBERT	09A	BABR $e^+ e^- \rightarrow \gamma(4S)$
--------------------------	--------	-----	---------------------------------------

0.0556 ± 0.0008 ± 0.0041	¹ AUBERT	08AT	BABR $e^+ e^- \rightarrow \gamma(4S)$
--------------------------	---------------------	------	---------------------------------------

0.0650 ± 0.0020 ± 0.0043	² ADAM	03	CLE2 $e^+ e^- \rightarrow \gamma(4S)$
--------------------------	-------------------	----	---------------------------------------

0.066 ± 0.016 ± 0.015	³ ALBRECHT	92C	ARG $e^+ e^- \rightarrow \gamma(4S)$
-----------------------	-----------------------	-----	--------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0583 ± 0.0015 ± 0.0030	⁴ AUBERT	08Q	BABR Repl. by AUBERT 09A
--------------------------	---------------------	-----	--------------------------

0.0650 ± 0.0020 ± 0.0043	⁵ BRIERE	02	CLE2 $e^+ e^- \rightarrow \gamma(4S)$
--------------------------	---------------------	----	---------------------------------------


0.0513 ± 0.0054 ± 0.0064	⁶ BARISH	95	CLE2 Repl. by ADAM 03
--------------------------	---------------------	----	-----------------------

seen	⁷ SANGHERA	93	CLE2 $e^+ e^- \rightarrow \gamma(4S)$
------	-----------------------	----	---------------------------------------

0.041 ± 0.008 ^{+0.008} / _{-0.009}	⁸ FULTON	91	CLEO $e^+ e^- \rightarrow \gamma(4S)$
---	---------------------	----	---------------------------------------

0.070 ± 0.018 ± 0.014	⁹ ANTREASYAN	90B	CBAL $e^+ e^- \rightarrow \gamma(4S)$
-----------------------	-------------------------	-----	---------------------------------------

- ¹ Measured using the dependence of $B^- \rightarrow D^{*0} e^- \bar{\nu}_e$ decay differential rate and the form factor description by CAPRINI 98.
- ² Simultaneous measurements of both $B^0 \rightarrow D^*(2010)^- \ell \nu$ and $B^+ \rightarrow \bar{D}^*(2007)^0 \ell \nu$.
- ³ ALBRECHT 92C reports $0.058 \pm 0.014 \pm 0.013$. We rescale using the method described in STONE 94 but with the updated PDG 94 $B(D^0 \rightarrow K^- \pi^+)$. Assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at the $\Upsilon(4S)$.
- ⁴ Uses a fully reconstructed B meson as a tag on the recoil side.
- ⁵ The results are based on the same analysis and data sample reported in ADAM 03.
- ⁶ BARISH 95 use $B(D^0 \rightarrow K^- \pi^+) = (3.91 \pm 0.08 \pm 0.17)\%$ and $B(D^{*0} \rightarrow D^0 \pi^0) = (63.6 \pm 2.3 \pm 3.3)\%$.
- ⁷ Combining $\bar{D}^{*0} \ell^+ \nu_\ell$ and $\bar{D}^{*-} \ell^+ \nu_\ell$ SANGHERA 93 test $V-A$ structure and fit the decay angular distributions to obtain $A_{FB} = 3/4 * (\Gamma^- - \Gamma^+) / \Gamma = 0.14 \pm 0.06 \pm 0.03$. Assuming a value of V_{cb} , they measure V , A_1 , and A_2 , the three form factors for the $D^* \ell \nu_\ell$ decay, where results are slightly dependent on model assumptions.
- ⁸ Assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at the $\Upsilon(4S)$. Uncorrected for D and D^* branching ratio assumptions.
- ⁹ ANTREASYAN 90B is average over B and $\bar{D}^*(2010)$ charge states.



$\Gamma(\bar{D}^*(2007)^0 \tau^+ \nu_\tau) / \Gamma_{\text{total}}$ Γ_7 / Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$2.25 \pm 0.48 \pm 0.28$	¹ AUBERT	08N BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

- ¹ Uses a fully reconstructed B meson as a tag on the recoil side.

$\Gamma(\bar{D}^*(2007)^0 \ell^+ \nu_\ell) / \Gamma(D \ell^+ \nu_\ell \text{ anything})$ Γ_6 / Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
0.582 ± 0.018 ± 0.030	¹ AUBERT	07AN BABR	$e^+ e^- \rightarrow \gamma(4S)$

¹ Uses a fully reconstructed B meson on the recoil side.

$\Gamma(D^{(*)} n \pi \ell^+ \nu_\ell (n \geq 1)) / \Gamma(D \ell^+ \nu_\ell \text{ anything})$ Γ_{11} / Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
0.191 ± 0.013 ± 0.019	¹ AUBERT	07AN BABR	$e^+ e^- \rightarrow \gamma(4S)$

¹ Uses a fully reconstructed B meson on the recoil side.

$\Gamma(D^- \pi^+ \ell^+ \nu_\ell) / \Gamma_{\text{total}}$ Γ_8 / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
4.2 ± 0.5 OUR AVERAGE			

4.2 ± 0.6 ± 0.3 ¹ AUBERT 08Q BABR $e^+ e^- \rightarrow \gamma(4S)$

4.2 ± 0.6 ± 0.2 ^{1,2} LIVENTSEV 08 BELL $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.4 ± 0.9 ± 0.3 ³ LIVENTSEV 05 BELL Repl. by LIVENTSEV 08

¹ Uses a fully reconstructed B meson as a tag on the recoil side.

² LIVENTSEV 08 reports $(4.0 \pm 0.4 \pm 0.6) \times 10^{-3}$ from a measurement of $[\Gamma(B^+ \rightarrow D^- \pi^+ \ell^+ \nu_\ell) / \Gamma_{\text{total}}] / [B(B^+ \rightarrow \bar{D}^0 \ell^+ \nu_\ell)]$ assuming $B(B^+ \rightarrow \bar{D}^0 \ell^+ \nu_\ell) = (2.15 \pm 0.22) \times 10^{-2}$. We rescale to our best value $B(B^+ \rightarrow \bar{D}^0 \ell^+ \nu_\ell) = (2.24 \pm 0.11) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ LIVENTSEV 05 reports $[\Gamma(B^+ \rightarrow D^- \pi^+ \ell^+ \nu_\ell) / \Gamma_{\text{total}}] / [B(B^0 \rightarrow D^- \ell^+ \nu_\ell)] = 0.25 \pm 0.03 \pm 0.03$. We multiply by our best value $B(B^0 \rightarrow D^- \ell^+ \nu_\ell) = (2.17 \pm 0.12) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\bar{D}_0^*(2420)^0 \ell^+ \nu_\ell \times B(\bar{D}_0^* \rightarrow D^+ \pi^-)) / \Gamma_{\text{total}}$ Γ_9 / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.5 ± 0.5 OUR AVERAGE			

2.6 ± 0.5 ± 0.4 ¹ AUBERT 08BL BABR $e^+ e^- \rightarrow \gamma(4S)$

2.4 ± 0.4 ± 0.6 ¹ LIVENTSEV 08 BELL $e^+ e^- \rightarrow \gamma(4S)$

¹ Uses a fully reconstructed B meson as a tag on the recoil side.

$\Gamma(\bar{D}_2^*(2460)^0 \ell^+ \nu_\ell \times B(\bar{D}_2^* \rightarrow D^+ \pi^-)) / \Gamma_{\text{total}}$ Γ_{10} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.67 ± 0.30 OUR AVERAGE			Error includes scale factor of 1.2.

1.5 ± 0.2 ± 0.2 ¹ AUBERT 08BL BABR $e^+ e^- \rightarrow \gamma(4S)$

2.2 ± 0.3 ± 0.4 ¹ LIVENTSEV 08 BELL $e^+ e^- \rightarrow \gamma(4S)$

¹ Uses a fully reconstructed B meson as a tag on the recoil side.

$\Gamma(D^{*-} \pi^+ \ell^+ \nu_\ell) / \Gamma_{\text{total}}$ Γ_{12} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
6.1 ± 0.6 OUR AVERAGE			

5.9 ± 0.5 ± 0.4 ¹ AUBERT 08Q BABR $e^+ e^- \rightarrow \gamma(4S)$

6.7 ± 1.1 ± 0.3 ^{1,2} LIVENTSEV 08 BELL $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.2 ± 1.5 ± 0.1 ^{3,4} LIVENTSEV 05 BELL Repl. by LIVENTSEV 08

¹ Uses a fully reconstructed B meson as a tag on the recoil side.

² LIVENTSEV 08 reports $(6.4 \pm 0.8 \pm 0.9) \times 10^{-3}$ from a measurement of $[\Gamma(B^+ \rightarrow D^{*-} \pi^+ \ell^+ \nu_\ell) / \Gamma_{\text{total}}] / [B(B^+ \rightarrow \bar{D}^0 \ell^+ \nu_\ell)]$ assuming $B(B^+ \rightarrow \bar{D}^0 \ell^+ \nu_\ell) = (2.15 \pm 0.22) \times 10^{-2}$. We rescale to our best value $B(B^+ \rightarrow \bar{D}^0 \ell^+ \nu_\ell) = (2.24 \pm 0.11) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Excludes D^{*+} contribution to $D\pi$ modes.

⁴ LIVENTSEV 05 reports $[\Gamma(B^+ \rightarrow D^{*-} \pi^+ \ell^+ \nu_\ell) / \Gamma_{\text{total}}] / [B(B^0 \rightarrow D^*(2010)^- \ell^+ \nu_\ell)] = 0.12 \pm 0.02 \pm 0.02$. We multiply by our best value $B(B^0 \rightarrow D^*(2010)^- \ell^+ \nu_\ell) = (5.16 \pm 0.11) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\bar{D}_1(2420)^0 \ell^+ \nu_\ell \times B(\bar{D}_1^0 \rightarrow D^{*+} \pi^-)) / \Gamma_{\text{total}} \quad \Gamma_{13}/\Gamma$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
3.2 ± 0.4 OUR AVERAGE			
2.9 ± 0.3 ± 0.3	¹ AUBERT 08BL	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
4.2 ± 0.7 ± 0.7	¹ LIVENTSEV 08	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
3.73 ± 0.85 ± 0.57	² ANASTASSOV 98	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Uses a fully reconstructed B meson as a tag on the recoil side.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}_1'(2430)^0 \ell^+ \nu_\ell \times B(\bar{D}_1'^0 \rightarrow D^{*+} \pi^-)) / \Gamma_{\text{total}} \quad \Gamma_{14}/\Gamma$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
2.7 ± 0.4 ± 0.5		¹ AUBERT 08BL	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 0.7	90	¹ LIVENTSEV 08	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Uses a fully reconstructed B meson as a tag on the recoil side.

$\Gamma(\bar{D}_2^*(2460)^0 \ell^+ \nu_\ell \times B(\bar{D}_2^{*0} \rightarrow D^{*+} \pi^-)) / \Gamma_{\text{total}} \quad \Gamma_{15}/\Gamma$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
1.8 ± 0.6 ± 0.3		¹ LIVENTSEV 08	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 1.6	90	² ANASTASSOV 98	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Uses a fully reconstructed B meson as a tag on the recoil side.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\pi^0 \ell^+ \nu_\ell) / \Gamma_{\text{total}} \quad \Gamma_{16}/\Gamma$

"OUR EVALUATION" is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements.

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
0.77 ± 0.10 ± 0.07 OUR EVALUATION			
0.81 ± 0.09 OUR AVERAGE			
0.82 ± 0.09 ± 0.05	^{1,2} AUBERT 08AV	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
0.77 ± 0.14 ± 0.08	³ HOKUUE 07	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.74 ± 0.05 ± 0.10	⁴ AUBERT,B 05O	BABR	Repl. by AUBERT 08AV

- ¹ Using isospin relation, B^+ and B^0 branching fractions are combined.
- ² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
- ³ The signal events are tagged by a second B meson reconstructed in the semileptonic mode $B \rightarrow D^{(*)} \ell \nu_\ell$.
- ⁴ B^+ and B^0 decays combined assuming isospin symmetry. Systematic errors include both experimental and form-factor uncertainties.

$\Gamma(\pi^0 e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{17}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

- • • We do not use the following data for averages, fits, limits, etc. • • •
- $0.9 \pm 0.2 \pm 0.2$ ¹ ALEXANDER 96T CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$
- < 22 90 ANTREASYAN 90B CBAL $e^+ e^- \rightarrow \Upsilon(4S)$
- ¹ Derived based in the reported B^0 result by assuming isospin symmetry: $\Gamma(B^0 \rightarrow \pi^- \ell^+ \nu) = 2\Gamma(B^+ \rightarrow \pi^0 \ell^+ \nu)$.

$\Gamma(\eta \ell^+ \nu_\ell)/\Gamma_{\text{total}}$ Γ_{18}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

- • • We do not use the following data for averages, fits, limits, etc. • • •
- $0.64 \pm 0.20 \pm 0.30$** ¹ AUBERT 08AV BABR $e^+ e^- \rightarrow \Upsilon(4S)$
- < 1.01 90 ² ADAM 07 CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$
- $0.84 \pm 0.31 \pm 0.18$ ³ ATHAR 03 CLE2 Repl. by ADAM 07
- ¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
- ² The B^0 and B^+ results are combined assuming the isospin, B lifetimes, and relative charged/neutral B production at the $\Upsilon(4S)$.
- ³ ATHAR 03 reports systematic errors 0.16 ± 0.09 , which are experimental systematic and systematic due to model dependence. We combine these in quadrature.

$\Gamma(\eta' \ell^+ \nu_\ell)/\Gamma_{\text{total}}$ Γ_{19}/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

- 0.17 ± 0.22 OUR AVERAGE**
- $0.04 \pm 0.22^{+0.05}_{-0.02}$ ¹ AUBERT 08AV BABR $e^+ e^- \rightarrow \Upsilon(4S)$
- $2.66 \pm 0.80 \pm 0.56$ ² ADAM 07 CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$
- ¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
- ² The B^0 and B^+ results are combined assuming the isospin, B lifetimes, and relative charged/neutral B production at the $\Upsilon(4S)$. Corresponds to 90% CL interval $(1.20-4.46) \times 10^{-4}$.

$\Gamma(\omega \ell^+ \nu_\ell)/\Gamma_{\text{total}}$ Γ_{20}/Γ

$\ell = e$ or μ , not sum over e and μ modes.

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

- • • We do not use the following data for averages, fits, limits, etc. • • •
- $1.3 \pm 0.4 \pm 0.4$** ¹ SCHWANDA 04 BELL $e^+ e^- \rightarrow \Upsilon(4S)$
- < 2.1 90 ² BEAN 93B CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$
- ¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
- ² BEAN 93B limit set using ISGW Model. Using isospin and the quark model to combine $\Gamma(\rho^0 \ell^+ \nu_\ell)$ and $\Gamma(\rho^- \ell^+ \nu_\ell)$ with this result, they obtain a limit $< (1.6-2.7) \times 10^{-4}$ at 90% CL for $B^+ \rightarrow \omega \ell^+ \nu_\ell$. The range corresponds to the ISGW, WSB, and KS models. An upper limit on $|V_{ub}/V_{cb}| < 0.8-0.13$ at 90% CL is derived as well.

$\Gamma(\omega\mu^+\nu_\mu)/\Gamma_{\text{total}}$

Γ_{21}/Γ

VALUE DOCUMENT ID TECN

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen ¹ ALBRECHT 91C ARG

¹In ALBRECHT 91C, one event is fully reconstructed providing evidence for the $b \rightarrow u$ transition.

$\Gamma(\rho^0\ell^+\nu_\ell)/\Gamma_{\text{total}}$

Γ_{22}/Γ

$\ell = e$ or μ , not sum over e and μ modes.

VALUE (units 10^{-4}) CL% DOCUMENT ID TECN COMMENT

1.28 ± 0.18 OUR AVERAGE

1.33 ± 0.23 ± 0.18 ¹ HOKUUE 07 BELL $e^+e^- \rightarrow \gamma(4S)$

1.16 ± 0.11 ± 0.30 ² AUBERT,B 050 BABR $e^+e^- \rightarrow \gamma(4S)$

1.34 ± 0.15 ^{+0.28}_{-0.32} ³ BEHRENS 00 CLE2 $e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.40 ± 0.21 ^{+0.32}_{-0.33} ³ BEHRENS 00 CLE2 $e^+e^- \rightarrow \gamma(4S)$

1.2 ± 0.2 ^{+0.3}_{-0.4} ³ ALEXANDER 96T CLE2 $e^+e^- \rightarrow \gamma(4S)$

<2.1 90 ⁴ BEAN 93B CLE2 $e^+e^- \rightarrow \gamma(4S)$

¹The signal events are tagged by a second B meson reconstructed in the semileptonic mode $B \rightarrow D^{(*)}\ell\nu_\ell$.

² B^+ and B^0 decays combined assuming isospin symmetry. Systematic errors include both experimental and form-factor uncertainties.

³Derived based in the reported B^0 result by assuming isospin symmetry: $\Gamma(B^0 \rightarrow \rho^-\ell^+\nu) \approx 2\Gamma(B^+ \rightarrow \omega\ell^+\nu)$.

⁴BEAN 93B limit set using ISGW Model. Using isospin and the quark model to combine $\Gamma(\omega^0\ell^+\nu_\ell)$ and $\Gamma(\rho^-\ell^+\nu_\ell)$ with this result, they obtain a limit $<(1.6-2.7) \times 10^{-4}$ at 90% CL for $B^+ \rightarrow \rho^0\ell^+\nu_\ell$. The range corresponds to the ISGW, WSB, and KS models. An upper limit on $|V_{ub}/V_{cb}| < 0.8-0.13$ at 90% CL is derived as well.

$\Gamma(\rho\bar{p}e^+\nu_e)/\Gamma_{\text{total}}$

Γ_{23}/Γ

VALUE CL% DOCUMENT ID TECN COMMENT

<5.2 × 10⁻³ 90 ¹ ADAM 03B CLE2 $e^+e^- \rightarrow \gamma(4S)$

¹Based on phase-space model; if $V-A$ model is used, the 90% CL upper limit becomes $< 1.2 \times 10^{-3}$.

$\Gamma(e^+\nu_e)/\Gamma_{\text{total}}$

Γ_{24}/Γ

VALUE (units 10^{-6}) CL% DOCUMENT ID TECN COMMENT

< 5.2 90 ¹ AUBERT 08AD BABR $e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 9.8 90 ¹ SATOYAMA 07 BELL $e^+e^- \rightarrow \gamma(4S)$

<15 90 ARTUSO 95 CLE2 $e^+e^- \rightarrow \gamma(4S)$

¹Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\mu^+ \nu_\mu)/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.7	90	¹ SATOYAMA 07	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
< 5.6	90	¹ AUBERT 08AD	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
< 6.6	90	AUBERT 04O	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
<21	90	ARTUSO 95	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\tau^+ \nu_\tau)/\Gamma_{\text{total}}$ Γ_{26}/Γ

See the note on "Decay Constants of Charged Pseudoscalar Mesons" in the D_s^+ Listings.

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
1.4 ± 0.4 OUR AVERAGE				
1.8 $^{+0.9}_{-0.8}$ ± 0.45		1,2 AUBERT 08D	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
0.9 ± 0.6 ± 0.1		2,3 AUBERT 07AL	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
1.79 $^{+0.56}_{-0.49}$ $^{+0.46}_{-0.51}$		1,2 IKADO 06	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

< 2.6	90	² AUBERT 06K	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
< 4.2	90	² AUBERT,B 05B	BABR	Repl. by AUBERT 06K
< 8.3	90	⁴ BARATE 01E	ALEP	$e^+ e^- \rightarrow Z$
< 8.4	90	² BROWDER 01	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
< 5.7	90	⁵ ACCIARRI 97F	L3	$e^+ e^- \rightarrow Z$
<104	90	⁶ ALBRECHT 95D	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
< 22	90	ARTUSO 95	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
< 18	90	⁷ BUSKULIC 95	ALEP	$e^+ e^- \rightarrow Z$

¹ The analysis is based on a sample of events with one fully reconstructed tag B in a hadronic decay mode $B^- \rightarrow D^{(*)0} X^-$.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ Requires one reconstructed semileptonic B decay $B^- \rightarrow D^0 \ell^- \bar{\nu}_\ell X$ in the recoil.

⁴ The energy-flow and b -tagging algorithms were used.

⁵ ACCIARRI 97F uses missing-energy technique and $f(b \rightarrow B^-) = (38.2 \pm 2.5)\%$.

⁶ ALBRECHT 95D uses full reconstruction of one B decay as tag.

⁷ BUSKULIC 95 uses same missing-energy technique as in $\bar{b} \rightarrow \tau^+ \nu_\tau X$, but analysis is restricted to endpoint region of missing-energy distribution.

$\Gamma(e^+ \nu_e \gamma)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 2.0 × 10⁻⁴	90	¹ BROWDER 97	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ BROWDER 97 uses the hermiticity of the CLEO II detector to reconstruct the neutrino energy and momentum.

$\Gamma(\mu^+ \nu_\mu \gamma)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 5.2 × 10⁻⁵	90	¹ BROWDER 97	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ BROWDER 97 uses the hermiticity of the CLEO II detector to reconstruct the neutrino energy and momentum.

$\Gamma(D^0 X)/\Gamma_{\text{total}}$ Γ_{29}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.086 ± 0.006 ± 0.004	¹ AUBERT	07N	BABR $e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.098 ± 0.009 ± 0.006	¹ AUBERT,BE	04B	BABR Repl. by AUBERT 07N

¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(\overline{D}^0 X)/\Gamma_{\text{total}}$ Γ_{30}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.786 ± 0.016^{+0.034}_{-0.033}	¹ AUBERT	07N	BABR $e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.793 ± 0.025 ^{+0.045} _{-0.044}	¹ AUBERT,BE	04B	BABR Repl. by AUBERT 07N

¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(D^0 X)/[\Gamma(D^0 X) + \Gamma(\overline{D}^0 X)]$ $\Gamma_{29}/(\Gamma_{29} + \Gamma_{30})$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.098 ± 0.007 ± 0.001	AUBERT	07N	BABR $e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.110 ± 0.010 ± 0.003	AUBERT,BE	04B	BABR Repl. by AUBERT 07N

$\Gamma(D^+ X)/\Gamma_{\text{total}}$ Γ_{31}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.025 ± 0.005 ± 0.002	¹ AUBERT	07N	BABR $e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.038 ± 0.009 ± 0.005	¹ AUBERT,BE	04B	BABR Repl. by AUBERT 07N

¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(D^- X)/\Gamma_{\text{total}}$ Γ_{32}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.099 ± 0.008 ± 0.009	¹ AUBERT	07N	BABR $e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.098 ± 0.012 ± 0.014	¹ AUBERT,BE	04B	BABR Repl. by AUBERT 07N

¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(D^+ X)/[\Gamma(D^+ X) + \Gamma(D^- X)]$ $\Gamma_{31}/(\Gamma_{31} + \Gamma_{32})$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.204 ± 0.035 ± 0.001	AUBERT	07N	BABR $e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.278 ± 0.052 ± 0.009	AUBERT,BE	04B	BABR Repl. by AUBERT 07N

$\Gamma(D_s^+ X)/\Gamma_{\text{total}}$ Γ_{33}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.079 \pm 0.006^{+0.013}_{-0.011}$	¹ AUBERT	07N	BABR $e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.143 \pm 0.016^{+0.051}_{-0.034}$	¹ AUBERT,BE	04B	BABR Repl. by AUBERT 07N
-------------------------------------	------------------------	-----	--------------------------

¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(D_s^- X)/\Gamma_{\text{total}}$ Γ_{34}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.011^{+0.004}_{-0.003}^{+0.002}_{-0.001}$		¹ AUBERT	07N	BABR $e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.022	90	¹ AUBERT,BE	04B	BABR Repl. by AUBERT 07N
--------	----	------------------------	-----	--------------------------

¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(D_s^+ X)/[\Gamma(D_s^+ X) + \Gamma(D_s^- X)]$ $\Gamma_{33}/(\Gamma_{33} + \Gamma_{34})$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.884 \pm 0.038 \pm 0.002$	AUBERT	07N	BABR $e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.966 \pm 0.039 \pm 0.012$	AUBERT,BE	04B	BABR Repl. by AUBERT 07N
-----------------------------	-----------	-----	--------------------------

$\Gamma(D_s^- X)/[\Gamma(D_s^+ X) + \Gamma(D_s^- X)]$ $\Gamma_{34}/(\Gamma_{33} + \Gamma_{34})$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.126	90	AUBERT,BE	04B	BABR $e^+e^- \rightarrow \gamma(4S)$

$\Gamma(\Lambda_c^+ X)/\Gamma_{\text{total}}$ Γ_{35}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.021 \pm 0.005^{+0.008}_{-0.004}$	¹ AUBERT	07N	BABR $e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.029 \pm 0.008^{+0.011}_{-0.007}$	¹ AUBERT,BE	04B	BABR Repl. by AUBERT 07N
-------------------------------------	------------------------	-----	--------------------------

¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(\bar{\Lambda}_c^- X)/\Gamma_{\text{total}}$ Γ_{36}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.028 \pm 0.005^{+0.010}_{-0.007}$	¹ AUBERT	07N	BABR $e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.035 \pm 0.008^{+0.013}_{-0.009}$	¹ AUBERT,BE	04B	BABR Repl. by AUBERT 07N
-------------------------------------	------------------------	-----	--------------------------

¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(\Lambda_c^+ X) / [\Gamma(\Lambda_c^+ X) + \Gamma(\bar{\Lambda}_c^- X)]$ $\Gamma_{35}/(\Gamma_{35}+\Gamma_{36})$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.427±0.071±0.001	AUBERT	07N BABR	$e^+e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.452±0.090±0.003	AUBERT,BE	04B BABR	Repl. by AUBERT 07N

$\Gamma(\tau X) / \Gamma_{total}$ Γ_{37}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.968±0.019^{+0.041}_{-0.039}	¹ AUBERT	07N BABR	$e^+e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.983±0.030 ^{+0.054} _{-0.051}	¹ AUBERT,BE	04B BABR	Repl. by AUBERT 07N

¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(c X) / \Gamma_{total}$ Γ_{38}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.234±0.012^{+0.018}_{-0.014}	¹ AUBERT	07N BABR	$e^+e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.330±0.022 ^{+0.055} _{-0.037}	¹ AUBERT,BE	04B BABR	Repl. by AUBERT 07N

¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(\tau c X) / \Gamma_{total}$ Γ_{39}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.202±0.023^{+0.053}_{-0.049}	¹ AUBERT	07N BABR	$e^+e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1.313±0.037 ^{+0.088} _{-0.075}	¹ AUBERT,BE	04B BABR	Repl. by AUBERT 07N

¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(\bar{D}^0 \pi^+) / \Gamma_{total}$ Γ_{40}/Γ

<u>VALUE (units 10⁻³)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.84±0.15 OUR AVERAGE				
4.90±0.07±0.22		¹ AUBERT	07H BABR	$e^+e^- \rightarrow \Upsilon(4S)$
5.3 ±0.6 ±0.3		² ABULENCIA	06J CDF	$p\bar{p}$ at 1.96 TeV
4.49±0.21±0.23		³ AUBERT,BE	06J BABR	$e^+e^- \rightarrow \Upsilon(4S)$
4.97±0.12±0.29		^{1,4} AHMED	02B CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
5.0 ±0.7 ±0.6	54	⁵ BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
5.4 ^{+1.8} ^{+1.2} _{-1.5 -0.9}	14	⁶ BEBEK	87 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.72 \pm 0.26^{+0.05}_{-0.06}$		⁷ AUBERT,B	04P	BABR	Repl. by AUBERT 07H
$5.5 \pm 0.4 \pm 0.5$	304	⁸ ALAM	94	CLE2	Repl. by AHMED 02B
$2.0 \pm 0.8 \pm 0.6$	12	⁵ ALBRECHT	90J	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
$1.9 \pm 1.0 \pm 0.6$	7	⁹ ALBRECHT	88K	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ABULENCIA 06J reports $[\Gamma(B^+ \rightarrow \bar{D}^0 \pi^+)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow D^- \pi^+)] = 1.97 \pm 0.10 \pm 0.21$. We multiply by our best value $B(B^0 \rightarrow D^- \pi^+) = (2.68 \pm 0.13) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Uses a missing-mass method. Does not depend on D branching fractions or B^+/B^0 production rates.

⁴ AHMED 02B reports an additional uncertainty on the branching ratios to account for 4.5% uncertainty on relative production of B^0 and B^+ , which is not included here.

⁵ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses the Mark III branching fractions for the D .

⁶ BEBEK 87 value has been updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92.

⁷ AUBERT,B 04P reports $[\Gamma(B^+ \rightarrow \bar{D}^0 \pi^+)/\Gamma_{\text{total}}] \times [B(D^0 \rightarrow K^- \pi^+)] = (1.846 \pm 0.032 \pm 0.097) \times 10^{-4}$. We divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.91 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁸ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

⁹ ALBRECHT 88K assumes $B^0 \bar{B}^0 : B^+ B^-$ ratio is 45:55. Superseded by ALBRECHT 90J.

$\Gamma(\bar{D}^0 \rho^+)/\Gamma_{\text{total}}$ Γ_{43}/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

0.0134 ± 0.0018 OUR AVERAGE

$0.0135 \pm 0.0012 \pm 0.0015$	212	¹ ALAM	94	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
$0.013 \pm 0.004 \pm 0.004$	19	² ALBRECHT	90J	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.021 \pm 0.008 \pm 0.009$	10	³ ALBRECHT	88K	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
-----------------------------	----	-----------------------	-----	-----	-----------------------------------

¹ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses the Mark III branching fractions for the D .

³ ALBRECHT 88K assumes $B^0 \bar{B}^0 : B^+ B^-$ ratio is 45:55.

$\Gamma(\bar{D}^0 K^+)/\Gamma(\bar{D}^0 \pi^+)$ Γ_{44}/Γ_{40}

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------


7.6 ± 0.6 OUR AVERAGE Error includes scale factor of 2.3. See the ideogram below.

$6.77 \pm 0.23 \pm 0.30$	HORII	08	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$8.31 \pm 0.35 \pm 0.20$	AUBERT	04N	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$9.9^{+1.4}_{-1.2}^{+0.7}_{-0.6}$	BORNHEIM	03	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

9.4 ±0.9 ±0.7	ABE	03D	BELL	Repl. by SWAIN 03
7.7 ±0.5 ±0.6	SWAIN	03	BELL	Repl. by HORII 08
7.9 ±0.9 ±0.6	ABE	01I	BELL	Repl. by ABE 03D
5.5 ±1.4 ±0.5	ATHANAS	98	CLE2	Repl. by BORNHEIM 03

WEIGHTED AVERAGE
7.6±0.6 (Error scaled by 2.3)



$\Gamma(D_{CP(+1)} K^+) / \Gamma(D_{CP(+1)} \pi^+)$

$\Gamma_{45} / \Gamma_{41}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.086±0.009 OUR AVERAGE			
0.086±0.008±0.007	^{1,2} ABE	06	BELL $e^+ e^- \rightarrow \Upsilon(4S)$
0.088±0.016±0.005	³ AUBERT	04N	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.125±0.036±0.010	³ ABE	03D	BELL	Repl. by SWAIN 03
0.093±0.018±0.008	³ SWAIN	03	BELL	Repl. by ABE 06

¹ Reports a double ratio of $B(B^+ \rightarrow D_{CP(+1)} K^+) / B(B^+ \rightarrow D_{CP(+1)} \pi^+)$ and $B(B^+ \rightarrow \bar{D}^0 K^+) / B(B^+ \rightarrow \bar{D}^0 \pi^+)$, $1.13 \pm 0.16 \pm 0.08$. We multiply by our best value of $B(B^+ \rightarrow \bar{D}^0 K^+) / B(B^+ \rightarrow \bar{D}^0 \pi^+) = 0.083 \pm 0.006$. Our first error is their experiment's error and the second error is systematic error from using our best value.

² ABE 06 reports $[\Gamma(B^+ \rightarrow D_{CP(+1)} K^+) / \Gamma(B^+ \rightarrow D_{CP(+1)} \pi^+)] / [\Gamma(B^+ \rightarrow \bar{D}^0 K^+) / \Gamma(B^+ \rightarrow \bar{D}^0 \pi^+)] = 1.13 \pm 0.06 \pm 0.08$. We multiply by our best value $\Gamma(B^+ \rightarrow \bar{D}^0 K^+) / \Gamma(B^+ \rightarrow \bar{D}^0 \pi^+) = (7.6 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ $CP=+1$ eigenstate of $D^0 \bar{D}^0$ system is reconstructed via $K^+ K^-$ and $\pi^+ \pi^-$.

$\Gamma(D_{CP(+1)}K^+)/\Gamma(\bar{D}^0K^+)$ Γ_{45}/Γ_{44}

VALUE	DOCUMENT ID	TECN	COMMENT
0.53±0.05±0.025	AUBERT	08AA BABR	$e^+e^- \rightarrow \Upsilon(4S)$
••• We do not use the following data for averages, fits, limits, etc. •••			
0.45±0.06±0.02	AUBERT	06J BABR	Repl. by AUBERT 08AA

$\Gamma(D_{CP(-1)}K^+)/\Gamma(D_{CP(-1)}\pi^+)$ Γ_{46}/Γ_{42}

VALUE	DOCUMENT ID	TECN	COMMENT
0.097±0.016±0.007	¹ ABE	06 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
••• We do not use the following data for averages, fits, limits, etc. •••			
0.119±0.028±0.006	² ABE	03D BELL	Repl. by SWAIN 03
0.108±0.019±0.007	² SWAIN	03 BELL	Repl. by ABE 06

¹ Reports a double ratio of $B(B^+ \rightarrow D_{CP(-1)}K^+)/B(B^+ \rightarrow D_{CP(-1)}\pi^+)$ and $B(B^+ \rightarrow \bar{D}^0K^+)/B(B^+ \rightarrow \bar{D}^0\pi^+)$, $1.17 \pm 0.14 \pm 0.14$. We multiply by our best value of $B(B^+ \rightarrow \bar{D}^0K^+)/B(B^+ \rightarrow \bar{D}^0\pi^+) = 0.083 \pm 0.006$. Our first error is their experiment's error and the second error is systematic error from using our best value.

² $CP=-1$ eigenstate of $D^0\bar{D}^0$ system is reconstructed via $K_S^0\pi^0$, $K_S^0\omega$, $K_S^0\phi$, $K_S^0\eta$, and $K_S^0\eta'$.

$\Gamma(D_{CP(-1)}K^+)/\Gamma(\bar{D}^0K^+)$ Γ_{46}/Γ_{44}

VALUE	DOCUMENT ID	TECN	COMMENT
0.515±0.05±0.025	AUBERT	08AA BABR	$e^+e^- \rightarrow \Upsilon(4S)$
••• We do not use the following data for averages, fits, limits, etc. •••			
0.43 ±0.05±0.02	AUBERT	06J BABR	Repl. by AUBERT 08AA

$\Gamma([K^-\pi^+]_D K^+)/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.8 × 10⁻⁷	90	HORII	08 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
••• We do not use the following data for averages, fits, limits, etc. •••				
<6.3 × 10 ⁻⁷	90	SAIGO	05 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma([K^-\pi^+]_D K^+)/\Gamma([K^+\pi^-]_D K^+)$ Γ_{47}/Γ_{48}

VALUE (units 10 ⁻³)	CL%	DOCUMENT ID	TECN	COMMENT
7.8^{+6.2+2.0}_{-5.7-2.8}		HORII	08 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
••• We do not use the following data for averages, fits, limits, etc. •••				
<29	90	¹ AUBERT	05G BABR	$e^+e^- \rightarrow \Upsilon(4S)$
<44	90	² SAIGO	05 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
<26	90	³ AUBERT,B	04L BABR	Repl. by AUBERT 05G

¹ AUBERT 05G extract a constraint on the magnitude of the ratio of amplitudes $|A(B^+ \rightarrow D^0K^+)/A(B^+ \rightarrow \bar{D}^0K^+)| < 0.23$ at 90% CL (Bayesian). Similar measurements from $B^+ \rightarrow D^{*0}K^+$ are also reported.

² SAIGO 05 extract a constraint on the magnitude of the ratio of amplitudes $|A(B^+ \rightarrow D^0K^+)/A(B^+ \rightarrow \bar{D}^0K^+)| < 0.27$ at 90% CL.

³ AUBERT,B 04L extract a constraint on the magnitude of the ratio of amplitudes $|A(B^+ \rightarrow D^0K^+)/A(B^+ \rightarrow \bar{D}^0K^+)| < 0.22$ at 90% CL.

$\Gamma([K^- \pi^+ \pi^0]_D K^+)/\Gamma([K^+ \pi^- \pi^0]_D K^+)$ Γ_{49}/Γ_{50}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.039	95	¹ AUBERT	07BN BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Extracts a constraint on the magnitude of the ratio of amplitudes $|A(B^+ \rightarrow D^0 K^+)/A(B^+ \rightarrow \bar{D}^0 K^+)| < 0.19$ at 95% CL.

$\Gamma([K^- \pi^+]_D K^*(892)^+)/\Gamma([K^+ \pi^-]_D K^*(892)^+)$ Γ_{51}/Γ_{52}

VALUE	DOCUMENT ID	TECN	COMMENT
0.046 ± 0.031 ± 0.008	AUBERT,B	05V BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma([K^- \pi^+]_D \pi^+)/\Gamma_{\text{total}}$ Γ_{53}/Γ

VALUE (units 10 ⁻⁷)	DOCUMENT ID	TECN	COMMENT
6.29^{+1.02+0.37}_{-0.98-0.48}	HORII	08 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.6 ^{+1.9} _{-1.7} ± 0.5	SAIGO	05 BELL	Repl. by HORII 08
---	-------	---------	-------------------

$\Gamma([K^- \pi^+]_D \pi^+)/\Gamma([K^+ \pi^-]_D \pi^+)$ Γ_{53}/Γ_{54}

VALUE (units 10 ⁻³)	DOCUMENT ID	TECN	COMMENT
3.40^{+0.55+0.15}_{-0.53-0.22}	HORII	08 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.5 ^{+1.0} _{-0.9} ± 0.2	SAIGO	05 BELL	Repl. by HORII 08
---	-------	---------	-------------------

$\Gamma([\pi^+ \pi^- \pi^0]_D K^-)/\Gamma_{\text{total}}$ Γ_{55}/Γ

VALUE (units 10 ⁻⁶)	DOCUMENT ID	TECN	COMMENT
4.6 ± 0.8 ± 0.4	¹ AUBERT	07BJ BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.5 ± 1.0 ± 0.7	¹ AUBERT,B	05T BABR	Repl. by AUBERT 07BJ
-----------------	-----------------------	----------	----------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^0 K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{56}/Γ

VALUE (units 10 ⁻⁴)	DOCUMENT ID	TECN	COMMENT
5.3 ± 0.4 OUR AVERAGE			

5.29 ± 0.30 ± 0.34	¹ AUBERT	06Z BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
--------------------	---------------------	----------	------------------------------------

6.1 ± 1.6 ± 1.7	¹ MAHAPATRA	02 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
-----------------	------------------------	---------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.3 ± 0.7 ± 0.5	¹ AUBERT	04Q BABR	Repl. by AUBERT 06Z
-----------------	---------------------	----------	---------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D_{CP(-1)} K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{57}/Γ

VALUE (units 10 ⁻⁴)	DOCUMENT ID	TECN	COMMENT
1.7 ± 0.7 ± 0.1	¹ AUBERT,B	05U BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ AUBERT,B 05U reports $[\Gamma(B^+ \rightarrow D_{CP(-1)} K^*(892)^+)/\Gamma_{\text{total}}] / [B(B^+ \rightarrow \bar{D}^0 K^*(892)^+)] = 0.325 \pm 0.13 \pm 0.04$. We multiply by our best value $B(B^+ \rightarrow \bar{D}^0 K^*(892)^+) = (5.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_{CP(+)} K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{58}/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$5.2 \pm 1.1 \pm 0.4$	¹ AUBERT,B	05U	BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ AUBERT,B 05U reports $[\Gamma(B^+ \rightarrow D_{CP(+)} K^*(892)^+)/\Gamma_{\text{total}}] / [B(B^+ \rightarrow \bar{D}^0 K^*(892)^+)] = 0.98 \pm 0.20 \pm 0.055$. We multiply by our best value $B(B^+ \rightarrow \bar{D}^0 K^*(892)^+) = (5.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\bar{D}^0 K^+ \bar{K}^0)/\Gamma_{\text{total}}$ Γ_{59}/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$5.5 \pm 1.4 \pm 0.8$	¹ DRUTSKOY	02	BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^0 K^+ \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{60}/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$7.5 \pm 1.3 \pm 1.1$	¹ DRUTSKOY	02	BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^0 \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{61}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.0115 \pm 0.0029 \pm 0.0021$	¹ BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

$\Gamma(\bar{D}^0 \pi^+ \pi^+ \pi^- \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{62}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.0051 \pm 0.0034 \pm 0.0023$	¹ BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

$\Gamma(\bar{D}^0 \pi^+ \rho^0)/\Gamma_{\text{total}}$ Γ_{63}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.0042 \pm 0.0023 \pm 0.0020$	¹ BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

$\Gamma(\bar{D}^0 a_1(1260)^+)/\Gamma_{\text{total}}$ Γ_{64}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.0045 \pm 0.0019 \pm 0.0031$	¹ BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

$\Gamma(\bar{D}^0 \omega \pi^+)/\Gamma_{\text{total}}$ Γ_{65}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.0041 \pm 0.0007 \pm 0.0006$	¹ ALEXANDER	01B	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. The signal is consistent with all observed $\omega \pi^+$ having proceeded through the ρ'^+ resonance at mass $1349 \pm 25^{+10}_{-5}$ MeV and width $547 \pm 86^{+46}_{-45}$ MeV.

$\Gamma(D^*(2010)^- \pi^+ \pi^+)/\Gamma_{\text{total}}$ Γ_{66}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.35 ± 0.22 OUR AVERAGE					
1.25 ± 0.08 ± 0.22			1 ABE	04D BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
1.9 ± 0.7 ± 0.3		14	2 ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
2.6 ± 1.4 ± 0.7		11	3 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
2.4 $\begin{smallmatrix} +1.7 & +1.0 \\ -1.6 & -0.6 \end{smallmatrix}$		3	4 BEBEK	87 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4.	90		5 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
5. ± 2. ± 3.		7	6 ALBRECHT	87C ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

³ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses the Mark III branching fractions for the D .

⁴ BEBEK 87 value has been updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92.

⁵ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D and $D^*(2010)$. The authors also find the product branching fraction into $D^{**} \pi$ followed by $D^{**} \rightarrow D^*(2010) \pi$ to be $0.0014^{+0.0008}_{-0.0006} \pm 0.0003$ where D^{**} represents all orbitally excited D mesons.

⁶ ALBRECHT 87C use PDG 86 branching ratios for D and $D^*(2010)$ and assume $B(\Upsilon(4S) \rightarrow B^+ B^-) = 55\%$ and $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 45\%$. Superseded by ALBRECHT 90J.

$\Gamma(D^- \pi^+ \pi^+)/\Gamma_{\text{total}}$ Γ_{67}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.02 ± 0.04 ± 0.15					
			1 ABE	04D BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
<1.4	90		2 ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
<7	90		3 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
2.5 $\begin{smallmatrix} +4.1 & +2.4 \\ -2.3 & -0.8 \end{smallmatrix}$		1	4 BEBEK	87 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the Mark III $B(D^+ \rightarrow K^- 2\pi^+)$.

³ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D . The product branching fraction into $D_0^*(2340) \pi$ followed by $D_0^*(2340) \rightarrow D \pi$ is < 0.005 at 90%CL and into $D_2^*(2460)$ followed by $D_2^*(2460) \rightarrow D \pi$ is < 0.004 at 90%CL.

⁴ BEBEK 87 assume the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. $B(D^- \rightarrow K^+ \pi^- \pi^-) = (9.1 \pm 1.3 \pm 0.4)\%$ is assumed.

$\Gamma(D^+ K^0)/\Gamma_{\text{total}}$ Γ_{68}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<5.0				
	90	1 AUBERT,B	05E BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{D}^*(2007)^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{69}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.19±0.26 OUR AVERAGE				
5.52±0.17±0.42		¹ AUBERT	07H BABR	$e^+e^- \rightarrow \Upsilon(4S)$
5.5 ±0.4 ±0.2		^{2,3} AUBERT,BE	06J BABR	$e^+e^- \rightarrow \Upsilon(4S)$
4.34±0.47±0.18		⁴ BRANDENB...	98 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
5.2 ±0.7 ±0.7	71	⁵ ALAM	94 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
7.2 ±1.8 ±1.6		⁶ BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
4.0 ±1.4 ±1.2	9	⁶ ALBRECHT	90J ARG	$e^+e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2.7 ±4.4		⁷ BEBEK	87 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² AUBERT,BE 06J reports $[\Gamma(B^+ \rightarrow \overline{D}^*(2007)^0 \pi^+)/\Gamma_{\text{total}}] / [B(B^+ \rightarrow \overline{D}^0 \pi^+)] = 1.14 \pm 0.07 \pm 0.04$. We multiply by our best value $B(B^+ \rightarrow \overline{D}^0 \pi^+) = (4.84 \pm 0.15) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Uses a missing-mass method. Does not depend on D branching fractions or B^+/B^0 production rates.

⁴ BRANDENBURG 98 assume equal production of B^+ and B^0 at $\Upsilon(4S)$ and use the D^* reconstruction technique. The first error is their experiment's error and the second error is the systematic error from the PDG 96 value of $B(D^* \rightarrow D\pi)$.

⁵ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

⁶ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D and $D^*(2010)$.

⁷ This is a derived branching ratio, using the inclusive pion spectrum and other two-body B decays. BEBEK 87 assume the $\Upsilon(4S)$ decays 43% to $B^0 \overline{B}^0$.

$\Gamma(\overline{D}^*(2007)^0 \omega \pi^+)/\Gamma_{\text{total}}$ Γ_{72}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0045±0.0010±0.0007			
	¹ ALEXANDER	01B CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. The signal is consistent with all observed $\omega \pi^+$ having proceeded through the ρ'^+ resonance at mass $1349 \pm 25^{+10}_{-5}$ MeV and width $547 \pm 86^{+46}_{-45}$ MeV.

$\Gamma(\overline{D}^*(2007)^0 \rho^+)/\Gamma_{\text{total}}$ Γ_{73}/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0098±0.0017 OUR AVERAGE				
0.0098±0.0006±0.0017		¹ CSORNA	03 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
0.010 ±0.006 ±0.004	7	² ALBRECHT	90J ARG	$e^+e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.0168±0.0021±0.0028	86	³ ALAM	94 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$ resonance. The second error combines the systematic and theoretical uncertainties in quadrature. CSORNA 03 includes data used in ALAM 94. A full angular fit to three complex helicity amplitudes is performed.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses MarkIII branching fractions for the D and $D^*(2010)$.

³ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$. The nonresonant $\pi^+ \pi^0$ contribution under the ρ^+ is negligible.

$\Gamma(\bar{D}^*(2007)^0 K^+)/\Gamma_{\text{total}}$ Γ_{74}/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

4.16 ± 0.33 OUR AVERAGE

4.22^{+0.30}_{-0.26} ± 0.21 ¹ AUBERT 05N BABR $e^+ e^- \rightarrow \Upsilon(4S)$

3.59 ± 0.97 ± 0.31 ² ABE 01I BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ AUBERT 05N reports $[\Gamma(B^+ \rightarrow \bar{D}^*(2007)^0 K^+)/\Gamma_{\text{total}}] / [B(B^+ \rightarrow \bar{D}^*(2007)^0 \pi^+)] = 0.0813 \pm 0.0040^{+0.0042}_{-0.0031}$. We multiply by our best value $B(B^+ \rightarrow \bar{D}^*(2007)^0 \pi^+) = (5.19 \pm 0.26) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABE 01I reports $B(B^+ \rightarrow \bar{D}^*(2007)^0 K^+)/B(B^+ \rightarrow \bar{D}^*(2007)^0 \pi^+) = 0.078 \pm 0.019 \pm 0.009$. We multiply by our best value $B(B^+ \rightarrow \bar{D}^*(2007)^0 \pi^+) = (4.6 \pm 0.4) \times 10^{-3}$. Our first error is their experiment's error and the second error is systematic error from using our best value.

$\Gamma(\bar{D}_{CP(+1)}^{*0} K^+)/\Gamma_{\text{total}}$ Γ_{75}/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

2.72 ± 0.28^{+0.23}_{-0.21} ¹ AUBERT 08BF BABR $e^+ e^- \rightarrow \Upsilon(4S)$

¹ AUBERT 08BF reports $[\Gamma(B^+ \rightarrow \bar{D}_{CP(+1)}^{*0} K^+)/\Gamma_{\text{total}}] / [B(B^+ \rightarrow \bar{D}^*(2007)^0 K^+)] = 0.655 \pm 0.065 \pm 0.020$. We multiply by our best value $B(B^+ \rightarrow \bar{D}^*(2007)^0 K^+) = (4.16 \pm 0.33) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\bar{D}_{CP(+1)}^{*0} K^+)/\Gamma(\bar{D}_{CP(+1)}^{*0} \pi^+)$ Γ_{75}/Γ_{70}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

0.095 ± 0.017 OUR AVERAGE

0.11 ± 0.02 ± 0.02 ¹ ABE 06 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

0.086 ± 0.021 ± 0.007 ² AUBERT 05N BABR $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Reports a double ratio of $B(B^+ \rightarrow (D_{CP(+1)}^*)^0 K^+)/B(B^+ \rightarrow (D_{CP(+1)}^*)^0 \pi^+)$ and $B(B^+ \rightarrow \bar{D}^{*0} K^+)/B(B^+ \rightarrow \bar{D}^{*0} \pi^+)$, $1.41 \pm 0.25 \pm 0.06$. We multiply by our best value of $B(B^+ \rightarrow \bar{D}^{*0} K^+)/B(B^+ \rightarrow \bar{D}^{*0} \pi^+) = 0.080 \pm 0.011$. Our first error is their experiment's error and the second error is systematic error from using our best value.

² Uses $D^{*0} \rightarrow D^0 \pi^0$ with D^0 reconstructed in the CP -even eigenstates $K^+ K^-$ and $\pi^+ \pi^-$.

$\Gamma(\overline{D}_{CP(-1)}^{*0} K^+)/\Gamma_{\text{total}}$ Γ_{76}/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.29 \pm 0.26 \pm_{-0.18}^{+0.19}$	¹ AUBERT	08BF BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ AUBERT 08BF reports $[\Gamma(B^+ \rightarrow \overline{D}_{CP(-1)}^{*0} K^+)/\Gamma_{\text{total}}] / [B(B^+ \rightarrow \overline{D}^*(2007)^0 K^+)] = 0.55 \pm 0.06 \pm 0.02$. We multiply by our best value $B(B^+ \rightarrow \overline{D}^*(2007)^0 K^+) = (4.16 \pm 0.33) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\overline{D}_{CP(-1)}^{*0} K^+)/\Gamma(D_{CP(-1)}^{*0} \pi^+)$ Γ_{76}/Γ_{71}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.09 \pm 0.03 \pm 0.01$	¹ ABE	06 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Reports a double ratio of $B(B^+ \rightarrow (D_{CP(-1)}^{*0})^0 K^+)/B(B^+ \rightarrow (D_{CP(-1)}^{*0})^0 \pi^+)$ and $B(B^+ \rightarrow \overline{D}^{*0} K^+)/B(B^+ \rightarrow \overline{D}^{*0} \pi^+)$, $1.15 \pm 0.31 \pm 0.12$. We multiply by our best value of $B(B^+ \rightarrow \overline{D}^{*0} K^+)/B(B^+ \rightarrow \overline{D}^{*0} \pi^+) = 0.080 \pm 0.011$. Our first error is their experiment's error and the second error is systematic error from using our best value.

$\Gamma(\overline{D}^*(2007)^0 K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{77}/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
8.1 ± 1.4 OUR AVERAGE			
$8.3 \pm 1.1 \pm 1.0$	¹ AUBERT	04K BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$7.2 \pm 2.2 \pm 2.6$	² MAHAPATRA	02 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and an unpolarized final state.

$\Gamma(\overline{D}^*(2007)^0 K^+ \overline{K}^0)/\Gamma_{\text{total}}$ Γ_{78}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<10.6	90	¹ DRUTSKOY	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{D}^*(2007)^0 K^+ K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{79}/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$15.3 \pm 3.1 \pm 2.9$	¹ DRUTSKOY	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{D}^*(2007)^0 \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{80}/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.03 ± 0.12 OUR AVERAGE				
$1.055 \pm 0.047 \pm 0.129$		¹ MAJUMDER	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$0.94 \pm 0.20 \pm 0.17$	48	^{2,3} ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
² ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.
³ The three pion mass is required to be between 1.0 and 1.6 GeV consistent with an a_1 meson. (If this channel is dominated by a_1^+ , the branching ratio for $\overline{D}^{*0} a_1^+$ is twice that for $\overline{D}^{*0} \pi^+ \pi^+ \pi^-$.)

$\Gamma(\bar{D}^*(2007)^0 a_1(1260)^+)/\Gamma_{\text{total}}$ Γ_{81}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0188 ± 0.0040 ± 0.0034	1,2 ALAM 94	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ ALAM 94 value is twice their $\Gamma(\bar{D}^*(2007)^0 \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ value based on their observation that the three pions are dominantly in the $a_1(1260)$ mass range 1.0 to 1.6 GeV.

² ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

$\Gamma(\bar{D}^*(2007)^0 \pi^- \pi^+ \pi^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{82}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0180 ± 0.0024 ± 0.0027	¹ ALEXANDER 01B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. The signal is consistent with all observed $\omega \pi^+$ having proceeded through the ρ'^+ resonance at mass $1349 \pm 25^{+10}_{-5}$ MeV and width $547 \pm 86^{+46}_{-45}$ MeV.

$\Gamma(\bar{D}^{*0} 3\pi^+ 2\pi^-)/\Gamma_{\text{total}}$ Γ_{83}/Γ

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.67 ± 0.91 ± 0.85	¹ MAJUMDER 04	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^*(2010)^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{84}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 3.6 × 10⁻⁶		¹ IWABUCHI 08	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.7 × 10 ⁻⁴	90	² BRANDENB... 98	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
--------------------------	----	-----------------------------	------	-----------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² BRANDENBURG 98 assume equal production of B^+ and B^0 at $\Upsilon(4S)$ and use the D^* partial reconstruction technique. The first error is their experiment's error and the second error is the systematic error from the PDG 96 value of $B(D^* \rightarrow D\pi)$.

$\Gamma(D^*(2010)^+ K^0)/\Gamma_{\text{total}}$ Γ_{85}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 9.0 × 10⁻⁶	90	¹ AUBERT,B 05E	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 9.5 × 10 ⁻⁵	90	¹ GRITSAN 01	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
--------------------------	----	-------------------------	------	-----------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^*(2010)^- \pi^+ \pi^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{86}/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0152 ± 0.0071 ± 0.0001	26	¹ ALBRECHT 90J	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.043 ± 0.013 ± 0.026	24	² ALBRECHT 87C	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
-----------------------	----	---------------------------	-----	-----------------------------------

¹ ALBRECHT 90J reports $0.018 \pm 0.007 \pm 0.005$ from a measurement of $[\Gamma(B^+ \rightarrow D^*(2010)^- \pi^+ \pi^+ \pi^0) / \Gamma_{\text{total}}] \times [B(D^*(2010)^+ \rightarrow D^0 \pi^+)]$ assuming $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

² ALBRECHT 87C use PDG 86 branching ratios for D and $D^*(2010)$ and assume $B(\Upsilon(4S) \rightarrow B^+ B^-) = 55\%$ and $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 45\%$. Superseded by ALBRECHT 90J.

$\Gamma(D^*(2010)^- \pi^+ \pi^+ \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{87} / Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

2.56 ± 0.26 ± 0.33		¹ MAJUMDER 04	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
---------------------------	--	--------------------------	------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<10	90	² ALBRECHT 90J	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
-----	----	---------------------------	-----	------------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D and $D^*(2010)$.

$\Gamma(\bar{D}^{*0} \pi^+) / \Gamma_{\text{total}}$ Γ_{88} / Γ

\bar{D}^{*0} represents an excited state with mass $2.2 < M < 2.8 \text{ GeV}/c^2$.

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

5.9 ± 1.3 ± 0.2	^{1,2} AUBERT, BE 06J	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
------------------------	-------------------------------	------	------------------------------------

¹ AUBERT, BE 06J reports $[\Gamma(B^+ \rightarrow \bar{D}^{*0} \pi^+) / \Gamma_{\text{total}}] / [B(B^+ \rightarrow \bar{D}^0 \pi^+)] = 1.22 \pm 0.13 \pm 0.23$. We multiply by our best value $B(B^+ \rightarrow \bar{D}^0 \pi^+) = (4.84 \pm 0.15) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Uses a missing-mass method. Does not depend on D branching fractions or B^+ / B^0 production rates.

$\Gamma(\bar{D}_1^*(2420)^0 \pi^+) / \Gamma_{\text{total}}$ Γ_{89} / Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.0015 ± 0.0006 OUR AVERAGE		Error includes scale factor of 1.3.		
------------------------------------	--	-------------------------------------	--	--

0.0011 ± 0.0005 ± 0.0002	8	¹ ALAM 94	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
--------------------------	---	----------------------	------	------------------------------------

0.0025 ± 0.0007 ± 0.0006		² ALBRECHT 94D	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
--------------------------	--	---------------------------	-----	------------------------------------

¹ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0) / B(D^0 \rightarrow K^- \pi^+)$ and assuming $B(D_1(2420)^0 \rightarrow D^*(2010)^+ \pi^-) = 67\%$.

² ALBRECHT 94D assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ assuming $B(D_1(2420)^0 \rightarrow D^*(2010)^+ \pi^-) = 67\%$.

$\Gamma(\bar{D}_1(2420)^0 \pi^+ \times B(\bar{D}_1^0 \rightarrow \bar{D}^0 \pi^+ \pi^-)) / \Gamma_{\text{total}}$ Γ_{90} / Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

1.85 ± 0.29^{+0.35}_{-0.55}	¹ ABE 05A	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
--	----------------------	------	------------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}_2^*(2462)^0 \pi^+ \times B(\bar{D}_2^*(2462)^0 \rightarrow D^- \pi^+))/\Gamma_{\text{total}}$ Γ_{91}/Γ

<u>VALUE (units 10⁻⁴)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.4±0.3±0.72	¹ ABE	04D BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}_0^*(2400)^0 \pi^+ \times B(\bar{D}_0^*(2400)^0 \rightarrow D^- \pi^+))/\Gamma_{\text{total}}$ Γ_{92}/Γ

<u>VALUE (units 10⁻⁴)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.1±0.6±1.8	¹ ABE	04D BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}_1(2421)^0 \pi^+ \times B(\bar{D}_1(2421)^0 \rightarrow D^{*-} \pi^+))/\Gamma_{\text{total}}$ Γ_{93}/Γ

<u>VALUE (units 10⁻⁴)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.8±0.7±1.3	¹ ABE	04D BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}_2^*(2462)^0 \pi^+ \times B(\bar{D}_2^*(2462)^0 \rightarrow D^{*-} \pi^+))/\Gamma_{\text{total}}$ Γ_{94}/Γ

<u>VALUE (units 10⁻⁴)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.8±0.3±0.4	¹ ABE	04D BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}'_1(2427)^0 \pi^+ \times B(\bar{D}'_1(2427)^0 \rightarrow D^{*-} \pi^+))/\Gamma_{\text{total}}$ Γ_{95}/Γ

<u>VALUE (units 10⁻⁴)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.0±0.4±1.1	¹ ABE	04D BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}_1(2420)^0 \pi^+ \times B(\bar{D}_1^0 \rightarrow \bar{D}^{*0} \pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{96}/Γ

<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.06	90	¹ ABE	05A BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}_1^*(2420)^0 \rho^+)/\Gamma_{\text{total}}$ Γ_{97}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0014	90	¹ ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ assuming $B(D_1(2420)^0 \rightarrow D^*(2010)^+ \pi^-) = 67\%$.

$\Gamma(\bar{D}_2^*(2460)^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{98}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0013	90	¹ ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0028	90	² ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
---------	----	-------------------	---------	------------------------------------

<0.0023	90	³ ALBRECHT	94D ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
---------	----	-----------------------	---------	------------------------------------

¹ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the Mark III $B(D^+ \rightarrow K^- 2\pi^+)$ and $B(D_2^*(2460)^0 \rightarrow D^+ \pi^-) = 30\%$.

² ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the Mark III $B(D^+ \rightarrow K^- 2\pi^+)$, the CLEO II $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ and $B(D_2^*(2460)^0 \rightarrow D^*(2010)^+ \pi^-) = 20\%$.

³ ALBRECHT 94D assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ and $B(D_2^*(2460)^0 \rightarrow D^*(2010)^+ \pi^-) = 30\%$.

$\Gamma(\overline{D}_2^*(2460)^0 \pi^+ \times B(\overline{D}_2^0 \rightarrow \overline{D}^{*0} \pi^+ \pi^-)) / \Gamma_{\text{total}}$ Γ_{99} / Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.22	90	¹ ABE	05A	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{D}_2^*(2460)^0 \rho^+) / \Gamma_{\text{total}}$ Γ_{100} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0047	90	¹ ALAM	94	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$
<0.005	90	² ALAM	94	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

¹ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the Mark III $B(D^+ \rightarrow K^- 2\pi^+)$ and $B(D_2^*(2460)^0 \rightarrow D^+ \pi^-) = 30\%$.

² ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the Mark III $B(D^+ \rightarrow K^- 2\pi^+)$, the CLEO II $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ and $B(D_2^*(2460)^0 \rightarrow D^*(2010)^+ \pi^-) = 20\%$.

$\Gamma(\overline{D}^0 D_s^+) / \Gamma_{\text{total}}$ Γ_{101} / Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0102 ± 0.0017 OUR AVERAGE				
0.0097 ± 0.0020 ± 0.0008		¹ AUBERT	06N	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
0.0100 ± 0.0027 ± 0.0008		² GIBAUT	96	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$
0.015 ± 0.008 ± 0.001		³ ALBRECHT	92G	ARG $e^+ e^- \rightarrow \Upsilon(4S)$
0.013 ± 0.006 ± 0.001	5	⁴ BORTOLETTO	90	CLEO $e^+ e^- \rightarrow \Upsilon(4S)$

¹ AUBERT 06N reports $(0.92 \pm 0.14 \pm 0.18) \times 10^{-2}$ from a measurement of $[\Gamma(B^+ \rightarrow \overline{D}^0 D_s^+) / \Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.0462 \pm 0.0062$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.39 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² GIBAUT 96 reports $0.0126 \pm 0.0022 \pm 0.0025$ from a measurement of $[\Gamma(B^+ \rightarrow \overline{D}^0 D_s^+) / \Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.39 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ALBRECHT 92G reports $0.024 \pm 0.012 \pm 0.004$ from a measurement of $[\Gamma(B^+ \rightarrow \overline{D}^0 D_s^+) / \Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.39 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^0 branching ratios, e.g., $B(D^0 \rightarrow K^- \pi^+) = 3.71 \pm 0.25\%$.

⁴ BORTOLETTO 90 reports 0.029 ± 0.013 from a measurement of $[\Gamma(B^+ \rightarrow \overline{D}^0 D_s^+) / \Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.02$. We rescale to our best

value $B(D_s^+ \rightarrow \phi\pi^+) = (4.39 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_{s0}(2317)^+ \bar{D}^0 \times B(D_{s0}(2317)^+ \rightarrow D_s^+ \pi^0))/\Gamma_{\text{total}}$ **Γ_{102}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

$0.74^{+0.22}_{-0.17}$ OUR AVERAGE

$0.82^{+0.35}_{-0.21} \pm 0.06$ ^{1,2} AUBERT,B 04s BABR $e^+e^- \rightarrow \Upsilon(4S)$

$0.66^{+0.27}_{-0.25} \pm 0.05$ ^{1,3} KROKOVNY 03B BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² AUBERT,B 04s reports $(1.0 \pm 0.3^{+0.4}_{-0.2}) \times 10^{-3}$ from a measurement of $[\Gamma(B^+ \rightarrow D_{s0}(2317)^+ \bar{D}^0 \times B(D_{s0}(2317)^+ \rightarrow D_s^+ \pi^0))/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.036 \pm 0.009$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.39 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ KROKOVNY 03B reports $(0.81^{+0.30}_{-0.27} \pm 0.24) \times 10^{-3}$ from a measurement of $[\Gamma(B^+ \rightarrow D_{s0}(2317)^+ \bar{D}^0 \times B(D_{s0}(2317)^+ \rightarrow D_s^+ \pi^0))/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.036 \pm 0.009$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.39 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_{s0}(2317)^+ \bar{D}^0 \times B(D_{s0}(2317)^+ \rightarrow D_s^{*+} \gamma))/\Gamma_{\text{total}}$ **Γ_{103}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<0.76 90 ¹ KROKOVNY 03B BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D_{s0}(2317)^+ \bar{D}^*(2007)^0 \times B(D_{s0}(2317)^+ \rightarrow D_s^+ \pi^0))/\Gamma_{\text{total}}$ **Γ_{104}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

$0.9 \pm 0.6^{+0.4}_{-0.3}$ ¹ AUBERT,B 04s BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D_{sJ}(2457)^+ \bar{D}^0)/\Gamma_{\text{total}}$ **Γ_{105}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

$3.1^{+1.0}_{-0.9}$ OUR AVERAGE

$4.3 \pm 1.6 \pm 1.3$ ¹ AUBERT 06N BABR $e^+e^- \rightarrow \Upsilon(4S)$

$4.6^{+1.8}_{-1.6} \pm 1.0$ ^{2,3} AUBERT,B 04s BABR $e^+e^- \rightarrow \Upsilon(4S)$

$2.1^{+1.1}_{-0.9} \pm 0.5$ ^{2,4} KROKOVNY 03B BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Uses a missing-mass method in the events that one of the B mesons is fully reconstructed.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ AUBERT,B 04s reports $[\Gamma(B^+ \rightarrow D_{sJ}(2457)^+\bar{D}^0)/\Gamma_{\text{total}}] \times [B(D_{s1}(2460)^+ \rightarrow D_s^{*+}\pi^0)] = (2.2^{+0.8}_{-0.7} \pm 0.3) \times 10^{-3}$. We divide by our best value $B(D_{s1}(2460)^+ \rightarrow D_s^{*+}\pi^0) = (48 \pm 11) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ KROKOVNY 03B reports $[\Gamma(B^+ \rightarrow D_{sJ}(2457)^+\bar{D}^0)/\Gamma_{\text{total}}] \times [B(D_{s1}(2460)^+ \rightarrow D_s^{*+}\pi^0)] = (1.0^{+0.5}_{-0.4} \pm 0.1) \times 10^{-3}$. We divide by our best value $B(D_{s1}(2460)^+ \rightarrow D_s^{*+}\pi^0) = (48 \pm 11) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_{sJ}(2457)^+\bar{D}^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^+\gamma))/\Gamma_{\text{total}}$ Γ_{106}/Γ

VALUE (units 10^{-3}) DOCUMENT ID TECN COMMENT

$0.47^{+0.13}_{-0.11}$ OUR AVERAGE

$0.49^{+0.20}_{-0.14} \pm 0.04$ ^{1,2} AUBERT,B 04s BABR $e^+e^- \rightarrow \Upsilon(4S)$

$0.46 \pm 0.15 \pm 0.04$ ^{1,3} KROKOVNY 03B BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² AUBERT,B 04s reports $(0.6 \pm 0.2^{+0.2}_{-0.1}) \times 10^{-3}$ from a measurement of $[\Gamma(B^+ \rightarrow D_{sJ}(2457)^+\bar{D}^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^+\gamma))/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.036 \pm 0.009$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.39 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ KROKOVNY 03B reports $(0.56^{+0.16}_{-0.15} \pm 0.17) \times 10^{-3}$ from a measurement of $[\Gamma(B^+ \rightarrow D_{sJ}(2457)^+\bar{D}^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^+\gamma))/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.036 \pm 0.009$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.39 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_{sJ}(2457)^+\bar{D}^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^+\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{107}/Γ

VALUE (units 10^{-3}) CL% DOCUMENT ID TECN COMMENT

<0.22 90 ¹ KROKOVNY 03B BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D_{sJ}(2457)^+\bar{D}^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^+\pi^0))/\Gamma_{\text{total}}$ Γ_{108}/Γ

VALUE (units 10^{-3}) CL% DOCUMENT ID TECN COMMENT

<0.27 90 ¹ KROKOVNY 03B BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D_{sJ}(2457)^+\bar{D}^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^{*+}\gamma))/\Gamma_{\text{total}}$ Γ_{109}/Γ

VALUE (units 10^{-3}) CL% DOCUMENT ID TECN COMMENT

<0.98 90 ¹ KROKOVNY 03B BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D_{sJ}(2457)^+ \bar{D}^*(2007)^0) / \Gamma_{\text{total}}$ Γ_{110} / Γ

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
12.0 ± 3.0 OUR AVERAGE			
11.2 ± 2.6 ± 2.0	¹ AUBERT	06N BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
16 $^{+8}_{-6}$ ± 4	^{2,3} AUBERT,B	04S BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Uses a missing-mass method in the events that one of the B mesons is fully reconstructed.
² AUBERT,B 04S reports $[\Gamma(B^+ \rightarrow D_{sJ}(2457)^+ \bar{D}^*(2007)^0) / \Gamma_{\text{total}}] \times [B(D_{s1}(2460)^+ \rightarrow D_s^{*+} \pi^0)] = (7.6 \pm 1.7^{+3.2}_{-2.4}) \times 10^{-3}$. We divide by our best value $B(D_{s1}(2460)^+ \rightarrow D_s^{*+} \pi^0) = (48 \pm 11) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
³ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D_{sJ}(2457)^+ \bar{D}^*(2007)^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^+ \gamma)) / \Gamma_{\text{total}}$ Γ_{111} / Γ

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.4 ± 0.4 $^{+0.6}_{-0.4}$	¹ AUBERT,B	04S BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^0 D_{s1}(2536)^+ \times B(D_{s1}(2536)^+ \rightarrow D^*(2007)^0 K^+)) / \Gamma_{\text{total}}$ Γ_{112} / Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.16 ± 0.52 ± 0.45		¹ AUBERT	08B BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2 90 AUBERT 03X BABR Repl. by AUBERT 08B

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^*(2007)^0 D_{s1}(2536)^+ \times B(D_{s1}(2536)^+ \rightarrow D^*(2007)^0 K^+)) / \Gamma_{\text{total}}$ Γ_{113} / Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.46 ± 1.17 ± 1.04		¹ AUBERT	08B BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<7 90 AUBERT 03X BABR Repl. by AUBERT 08B

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^0 D_{s1}(2536)^+ \times B(D_{s1}(2536)^+ \rightarrow D^{*+} K^0)) / \Gamma_{\text{total}}$ Γ_{114} / Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.30 ± 0.98 ± 0.43	¹ AUBERT	08B BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^0 D_{sJ}(2700)^+ \times B(D_{sJ}(2700)^+ \rightarrow D^0 K^+)) / \Gamma_{\text{total}}$ Γ_{115} / Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
11.3 ± 2.2 $^{+1.4}_{-2.8}$	¹ BRODZICKA	08 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^{*0} D_{s1}(2536)^+ \times B(D_{s1}(2536)^+ \rightarrow D^{*+} K^0))/\Gamma_{\text{total}}$ Γ_{116}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$3.92 \pm 2.46 \pm 0.83$	¹ AUBERT	08B BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^{*0} D_{sJ}(2573)^+ \times B(D_{sJ}(2573)^+ \rightarrow D^0 K^+))/\Gamma_{\text{total}}$ Γ_{117}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	AUBERT	03X BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(\bar{D}^*(2007)^0 D_{sJ}(2573)^+ \times B(D_{sJ}(2573)^+ \rightarrow D^0 K^+))/\Gamma_{\text{total}}$ Γ_{118}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	AUBERT	03X BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(\bar{D}^0 D_s^{*+})/\Gamma_{\text{total}}$ Γ_{119}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0078 ± 0.0016 OUR AVERAGE			
$0.0081 \pm 0.0018 \pm 0.0006$	¹ AUBERT	06N BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$0.0069 \pm 0.0025 \pm 0.0005$	² GIBAUT	96 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
$0.010 \pm 0.008 \pm 0.001$	³ ALBRECHT	92G ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ AUBERT 06N reports $(0.77 \pm 0.15 \pm 0.13) \times 10^{-2}$ from a measurement of $[\Gamma(B^+ \rightarrow \bar{D}^0 D_s^{*+})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.0462 \pm 0.0062$.

We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.39 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² GIBAUT 96 reports $0.0087 \pm 0.0027 \pm 0.0017$ from a measurement of $[\Gamma(B^+ \rightarrow \bar{D}^0 D_s^{*+})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.035$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.39 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ALBRECHT 92G reports $0.016 \pm 0.012 \pm 0.003$ from a measurement of $[\Gamma(B^+ \rightarrow \bar{D}^0 D_s^{*+})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.39 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^0 branching ratios, e.g., $B(D^0 \rightarrow K^- \pi^+) = 3.71 \pm 0.25\%$.

$\Gamma(\bar{D}^*(2007)^0 D_s^+)/\Gamma_{\text{total}}$ Γ_{120}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0084 ± 0.0017 OUR AVERAGE			
$0.0080 \pm 0.0018 \pm 0.0006$	¹ AUBERT	06N BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$0.011 \pm 0.004 \pm 0.001$	² GIBAUT	96 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
$0.008 \pm 0.006 \pm 0.001$	³ ALBRECHT	92G ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ AUBERT 06N reports $(0.76 \pm 0.15 \pm 0.13) \times 10^{-2}$ from a measurement of $[\Gamma(B^+ \rightarrow \bar{D}^*(2007)^0 D_s^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.0462 \pm 0.0062$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.39 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² GIBAUT 96 reports $0.0140 \pm 0.0043 \pm 0.0035$ from a measurement of $[\Gamma(B^+ \rightarrow \bar{D}^*(2007)^0 D_s^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.035$.

We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.39 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ALBRECHT 92G reports $0.013 \pm 0.009 \pm 0.002$ from a measurement of $[\Gamma(B^+ \rightarrow \bar{D}^*(2007)^0 D_s^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We

rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.39 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^0 and $D^*(2007)^0$ branching ratios, e.g., $B(D^0 \rightarrow K^-\pi^+) = 3.71 \pm 0.25\%$ and $B(D^*(2007)^0 \rightarrow D^0\pi^0) = 55 \pm 6\%$.

$\Gamma(\bar{D}^*(2007)^0 D_s^{*+})/\Gamma_{\text{total}}$ Γ_{121}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0174 ± 0.0023 OUR AVERAGE			
0.0170 ± 0.0019 ± 0.0013	¹ AUBERT	06N BABR	$e^+e^- \rightarrow \Upsilon(4S)$
0.025 ± 0.009 ± 0.002	² GIBAUT	96 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
0.019 ± 0.010 ± 0.001	³ ALBRECHT	92G ARG	$e^+e^- \rightarrow \Upsilon(4S)$

¹ AUBERT 06N reports $(1.62 \pm 0.22 \pm 0.18) \times 10^{-2}$ from a measurement of $[\Gamma(B^+ \rightarrow \bar{D}^*(2007)^0 D_s^{*+})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.0462 \pm 0.0062$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.39 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² GIBAUT 96 reports $0.0310 \pm 0.0088 \pm 0.0065$ from a measurement of $[\Gamma(B^+ \rightarrow \bar{D}^*(2007)^0 D_s^{*+})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.035$.

We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.39 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ALBRECHT 92G reports $0.031 \pm 0.016 \pm 0.005$ from a measurement of $[\Gamma(B^+ \rightarrow \bar{D}^*(2007)^0 D_s^{*+})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$.

We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.39 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^0 and $D^*(2007)^0$ branching ratios, e.g., $B(D^0 \rightarrow K^-\pi^+) = 3.71 \pm 0.25\%$ and $B(D^*(2007)^0 \rightarrow D^0\pi^0) = 55 \pm 6\%$.

$\Gamma(D_s^{(*)+} \bar{D}^{*0})/\Gamma_{\text{total}}$ Γ_{122}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$(2.73 \pm 0.93 \pm 0.68) \times 10^{-2}$	¹ AHMED	00B CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ AHMED 00B reports their experiment's uncertainties $(\pm 0.78 \pm 0.48 \pm 0.68)\%$, where the first error is statistical, the second is systematic, and the third is the uncertainty in the $D_s \rightarrow \phi\pi$ branching fraction. We combine the first two in quadrature.

$\Gamma(\bar{D}^*(2007)^0 D^*(2010)^+)/\Gamma_{\text{total}}$ Γ_{123}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
8.1 ± 1.2 ± 1.2		¹ AUBERT,B	06A BABR	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<110	90	BARATE	98Q ALEP	$e^+e^- \rightarrow Z$
------	----	--------	----------	------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$[\Gamma(\bar{D}^0 D^*(2010)^+) + \Gamma(\bar{D}^*(2007)^0 D^+)]/\Gamma_{\text{total}}$ Γ_{124}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<130	90	BARATE	98Q	ALEP $e^+ e^- \rightarrow Z$

$\Gamma(\bar{D}^0 D^*(2010)^+)/\Gamma_{\text{total}}$ Γ_{125}/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.9 ± 0.5 OUR AVERAGE			
3.6 ± 0.5 ± 0.4	¹ AUBERT,B	06A	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
4.57 ± 0.71 ± 0.56	¹ MAJUMDER	05	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^0 D^+)/\Gamma_{\text{total}}$ Γ_{126}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.8 ± 0.4 OUR AVERAGE				
3.85 ± 0.31 ± 0.38		¹ ADACHI	08	BELL $e^+ e^- \rightarrow \Upsilon(4S)$
3.8 ± 0.6 ± 0.5		¹ AUBERT,B	06A	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.83 ± 0.78 ± 0.58		¹ MAJUMDER	05	BELL Repl. by ADACHI 08
<67	90	BARATE	98Q	ALEP $e^+ e^- \rightarrow Z$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^0 D^+ K^0)/\Gamma_{\text{total}}$ Γ_{127}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.8	90	¹ AUBERT	03X	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^+ \bar{D}^*(2007)^0)/\Gamma_{\text{total}}$ Γ_{128}/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.3 ± 1.4 ± 1.0			
	¹ AUBERT,B	06A	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^*(2007)^0 D^+ K^0)/\Gamma_{\text{total}}$ Γ_{129}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<6.1	90	¹ AUBERT	03X	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^0 \bar{D}^*(2010)^+ K^0)/\Gamma_{\text{total}}$ Γ_{130}/Γ

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.2^{+1.0}_{-0.9} ± 0.7			
	¹ AUBERT	03X	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^*(2007)^0 D^*(2010)^+ K^0)/\Gamma_{\text{total}}$ Γ_{131}/Γ

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
7.8^{+2.3}_{-2.1} ± 1.4			
	¹ AUBERT	03X	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^0 D^0 K^+)/\Gamma_{\text{total}}$ Γ_{132}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

2.10±0.26 OUR AVERAGE

2.22±0.22^{+0.26}_{-0.24} ¹ BRODZICKA 08 BELL $e^+e^- \rightarrow \Upsilon(4S)$

1.9 ±0.3 ±0.3 ¹ AUBERT 03X BABR $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.17±0.21±0.15 ¹ CHISTOV 04 BELL Repl. by BRODZICKA 08

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^*(2007)^0 D^0 K^+)/\Gamma_{\text{total}}$ Γ_{133}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<3.8 90 ¹ AUBERT 03X BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^0 D^*(2007)^0 K^+)/\Gamma_{\text{total}}$ Γ_{134}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

4.7±0.7±0.7 ¹ AUBERT 03X BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^*(2007)^0 D^*(2007)^0 K^+)/\Gamma_{\text{total}}$ Γ_{135}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

5.3^{+1.1}_{-1.0}±1.2 ¹ AUBERT 03X BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^- D^+ K^+)/\Gamma_{\text{total}}$ Γ_{136}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<0.4 90 ¹ AUBERT 03X BABR $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.90 90 ¹ CHISTOV 04 BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^- D^*(2010)^+ K^+)/\Gamma_{\text{total}}$ Γ_{137}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<0.7 90 ¹ AUBERT 03X BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^*(2010)^- D^+ K^+)/\Gamma_{\text{total}}$ Γ_{138}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

1.5±0.3±0.2 ¹ AUBERT 03X BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^*(2010)^- D^*(2010)^+ K^+)/\Gamma_{\text{total}}$ Γ_{139}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<1.8 90 ¹ AUBERT 03X BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma((\bar{D}+\bar{D}^*)(D+D^*)K)/\Gamma_{\text{total}}$ Γ_{140}/Γ

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
$3.5 \pm 0.3 \pm 0.5$		¹ AUBERT	03X BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D_s^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{141}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
$1.6^{+0.6}_{-0.5} \pm 0.1$		¹ AUBERT	07M BABR	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<17 90 ² ALEXANDER 93B CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

¹ AUBERT 07M reports $[\Gamma(B^+ \rightarrow D_s^+ \pi^0)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)] = (7.0^{+2.4+0.6}_{-2.1-0.8}) \times 10^{-7}$. We divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.39 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ALEXANDER 93B reports $< 2.0 \times 10^{-4}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^+ \pi^0)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.39 \times 10^{-2}$.

$[\Gamma(D_s^+ \pi^0) + \Gamma(D_s^{*+} \pi^0)]/\Gamma_{\text{total}}$ $(\Gamma_{141} + \Gamma_{142})/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0006	90	¹ ALBRECHT 93E	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

¹ ALBRECHT 93E reports $< 0.9 \times 10^{-3}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^+ \pi^0) + \Gamma(B^+ \rightarrow D_s^{*+} \pi^0)]/\Gamma_{\text{total}} \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.39 \times 10^{-2}$.

$\Gamma(D_s^{*+} \pi^0)/\Gamma_{\text{total}}$ Γ_{142}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.00027	90	¹ ALEXANDER 93B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ ALEXANDER 93B reports $< 3.2 \times 10^{-4}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^{*+} \pi^0)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.39 \times 10^{-2}$.

$\Gamma(D_s^+ \eta)/\Gamma_{\text{total}}$ Γ_{143}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0004	90	¹ ALEXANDER 93B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ ALEXANDER 93B reports $< 4.6 \times 10^{-4}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^+ \eta)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.39 \times 10^{-2}$.

$\Gamma(D_s^{*+} \eta) / \Gamma_{\text{total}}$ Γ_{144} / Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0006	90	¹ ALEXANDER 93B	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

¹ ALEXANDER 93B reports $< 7.5 \times 10^{-4}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^{*+} \eta) / \Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 4.39 \times 10^{-2}$.

$\Gamma(D_s^+ \rho^0) / \Gamma_{\text{total}}$ Γ_{145} / Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.00031	90	¹ ALEXANDER 93B	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

¹ ALEXANDER 93B reports $< 3.7 \times 10^{-4}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^+ \rho^0) / \Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 4.39 \times 10^{-2}$.

$[\Gamma(D_s^+ \rho^0) + \Gamma(D_s^+ \bar{K}^*(892)^0)] / \Gamma_{\text{total}}$ $(\Gamma_{145} + \Gamma_{155}) / \Gamma$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0021	90	¹ ALBRECHT 93E	ARG	$e^+ e^- \rightarrow \gamma(4S)$

¹ ALBRECHT 93E reports $< 3.4 \times 10^{-3}$ from a measurement of $[[\Gamma(B^+ \rightarrow D_s^+ \rho^0) + \Gamma(B^+ \rightarrow D_s^+ \bar{K}^*(892)^0)] / \Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 4.39 \times 10^{-2}$.

$\Gamma(D_s^{*+} \rho^0) / \Gamma_{\text{total}}$ Γ_{146} / Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0004	90	¹ ALEXANDER 93B	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

¹ ALEXANDER 93B reports $< 4.8 \times 10^{-4}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^{*+} \rho^0) / \Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 4.39 \times 10^{-2}$.

$[\Gamma(D_s^{*+} \rho^0) + \Gamma(D_s^{*+} \bar{K}^*(892)^0)] / \Gamma_{\text{total}}$ $(\Gamma_{146} + \Gamma_{156}) / \Gamma$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0012	90	¹ ALBRECHT 93E	ARG	$e^+ e^- \rightarrow \gamma(4S)$

¹ ALBRECHT 93E reports $< 2.0 \times 10^{-3}$ from a measurement of $[[\Gamma(B^+ \rightarrow D_s^{*+} \rho^0) + \Gamma(B^+ \rightarrow D_s^{*+} \bar{K}^*(892)^0)] / \Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 4.39 \times 10^{-2}$.

$\Gamma(D_s^+ \omega) / \Gamma_{\text{total}}$ Γ_{147} / Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0004	90	¹ ALEXANDER 93B	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0021	90	² ALBRECHT 93E	ARG	$e^+ e^- \rightarrow \gamma(4S)$
---------	----	---------------------------	-----	----------------------------------

¹ ALEXANDER 93B reports $< 4.8 \times 10^{-4}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^+ \omega)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.39 \times 10^{-2}$.

² ALBRECHT 93E reports $< 3.4 \times 10^{-3}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^+ \omega)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.39 \times 10^{-2}$.

$\Gamma(D_s^{*+} \omega)/\Gamma_{\text{total}}$ Γ_{148}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0006	90	¹ ALEXANDER 93B	CLE2	$e^+e^- \rightarrow \gamma(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.0012	90	² ALBRECHT 93E	ARG	$e^+e^- \rightarrow \gamma(4S)$
¹ ALEXANDER 93B reports $< 6.8 \times 10^{-4}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^{*+} \omega)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.39 \times 10^{-2}$.				
² ALBRECHT 93E reports $< 1.9 \times 10^{-3}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^{*+} \omega)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.39 \times 10^{-2}$.				

$\Gamma(D_s^+ a_1(1260)^0)/\Gamma_{\text{total}}$ Γ_{149}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0018	90	¹ ALBRECHT 93E	ARG	$e^+e^- \rightarrow \gamma(4S)$
¹ ALBRECHT 93E reports $< 3.0 \times 10^{-3}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^+ a_1(1260)^0)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.39 \times 10^{-2}$.				

$\Gamma(D_s^{*+} a_1(1260)^0)/\Gamma_{\text{total}}$ Γ_{150}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0014	90	¹ ALBRECHT 93E	ARG	$e^+e^- \rightarrow \gamma(4S)$
¹ ALBRECHT 93E reports $< 2.2 \times 10^{-3}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^{*+} a_1(1260)^0)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.39 \times 10^{-2}$.				

$\Gamma(D_s^+ \phi)/\Gamma_{\text{total}}$ Γ_{151}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.9 $\times 10^{-6}$	90	¹ AUBERT 06F	BABR	$e^+e^- \rightarrow \gamma(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.0010	90	² ALBRECHT 93E	ARG	$e^+e^- \rightarrow \gamma(4S)$
<0.00026	90	³ ALEXANDER 93B	CLE2	$e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ALBRECHT 93E reports $< 1.7 \times 10^{-3}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^+ \phi)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.39 \times 10^{-2}$.

³ ALEXANDER 93B reports $< 3.1 \times 10^{-4}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^+ \phi)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.39 \times 10^{-2}$.

$\Gamma(D_s^{*+} \phi)/\Gamma_{\text{total}}$ Γ_{152}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.2 × 10⁻⁵	90	¹ AUBERT 06F BABR	$e^+e^- \rightarrow \Upsilon(4S)$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.0013	90	² ALBRECHT 93E ARG	$e^+e^- \rightarrow \Upsilon(4S)$	
<0.00035	90	³ ALEXANDER 93B CLE2	$e^+e^- \rightarrow \Upsilon(4S)$	

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ALBRECHT 93E reports $< 2.1 \times 10^{-3}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^{*+} \phi)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.39 \times 10^{-2}$.

³ ALEXANDER 93B reports $< 4.2 \times 10^{-4}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^{*+} \phi)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.39 \times 10^{-2}$.

$\Gamma(D_s^+ \bar{K}^0)/\Gamma_{\text{total}}$ Γ_{153}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0009	90	¹ ALEXANDER 93B CLE2	$e^+e^- \rightarrow \Upsilon(4S)$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.0015	90	² ALBRECHT 93E ARG	$e^+e^- \rightarrow \Upsilon(4S)$	

¹ ALEXANDER 93B reports $< 10.3 \times 10^{-4}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^+ \bar{K}^0)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.39 \times 10^{-2}$.

² ALBRECHT 93E reports $< 2.5 \times 10^{-3}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^+ \bar{K}^0)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.39 \times 10^{-2}$.

$\Gamma(D_s^{*+} \bar{K}^0)/\Gamma_{\text{total}}$ Γ_{154}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0009	90	¹ ALEXANDER 93B CLE2	$e^+e^- \rightarrow \Upsilon(4S)$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.0019	90	² ALBRECHT 93E ARG	$e^+e^- \rightarrow \Upsilon(4S)$	

¹ ALEXANDER 93B reports $< 10.9 \times 10^{-4}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^{*+} \bar{K}^0)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.39 \times 10^{-2}$.

² ALBRECHT 93E reports $< 3.1 \times 10^{-3}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^{*+} \bar{K}^0)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.39 \times 10^{-2}$.

$\Gamma(D_s^+ \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{155}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0004	90	¹ ALEXANDER 93B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ ALEXANDER 93B reports $< 4.4 \times 10^{-4}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^+ \bar{K}^*(892)^0)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.39 \times 10^{-2}$.

$\Gamma(D_s^{*+} \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{156}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0004	90	¹ ALEXANDER 93B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ ALEXANDER 93B reports $< 4.3 \times 10^{-4}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^{*+} \bar{K}^*(892)^0)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.39 \times 10^{-2}$.

$\Gamma(D_s^- \pi^+ K^+)/\Gamma_{\text{total}}$ Γ_{157}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
$2.02 \pm 0.13 \pm 0.38$		¹ AUBERT 08G	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<7	90	² ALBRECHT 93E	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
----	----	---------------------------	-----	-----------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ALBRECHT 93E reports $< 1.1 \times 10^{-3}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^- \pi^+ K^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.39 \times 10^{-2}$.

$\Gamma(D_s^{*-} \pi^+ K^+)/\Gamma_{\text{total}}$ Γ_{158}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
$1.67 \pm 0.16 \pm 0.35$		¹ AUBERT 08G	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<10	90	² ALBRECHT 93E	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
-----	----	---------------------------	-----	-----------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ALBRECHT 93E reports $< 1.6 \times 10^{-3}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^{*-} \pi^+ K^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.39 \times 10^{-2}$.

$\Gamma(D_s^- \pi^+ K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{159}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.005	90	¹ ALBRECHT	93E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ ALBRECHT 93E reports $< 8.6 \times 10^{-3}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^- \pi^+ K^*(892)^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 4.39 \times 10^{-2}$.

$\Gamma(D_s^{*-} \pi^+ K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{160}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.007	90	¹ ALBRECHT	93E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ ALBRECHT 93E reports $< 1.1 \times 10^{-2}$ from a measurement of $[\Gamma(B^+ \rightarrow D_s^{*-} \pi^+ K^*(892)^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 4.39 \times 10^{-2}$.

$\Gamma(D_s^- K^+ K^+)/\Gamma_{\text{total}}$ Γ_{161}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$0.11 \pm 0.04 \pm 0.02$	¹ AUBERT	08G BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D_s^{*-} K^+ K^+)/\Gamma_{\text{total}}$ Γ_{162}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.15	90	¹ AUBERT	08G BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta_c K^+)/\Gamma_{\text{total}}$ Γ_{163}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.91 ± 0.13 OUR AVERAGE			

0.87 ± 0.15 ^{1,2} AUBERT 06E BABR $e^+ e^- \rightarrow \Upsilon(4S)$

1.4 $^{+0.3}_{-0.2} \pm 0.4$ ³ AUBERT,B 05L BABR $e^+ e^- \rightarrow \Upsilon(4S)$

1.25 ± 0.14 $^{+0.39}_{-0.40}$ ⁴ FANG 03 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

0.69 $^{+0.26}_{-0.21} \pm 0.22$ ⁵ EDWARDS 01 CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.06 ± 0.12 ± 0.18 ^{2,6} AUBERT,B 04B BABR $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Perform measurements of absolute branching fractions using a missing mass technique.

² The ratio of $B(B^\pm \rightarrow K^\pm \eta_c) B(\eta_c \rightarrow K \bar{K} \pi) = (7.4 \pm 0.5 \pm 0.7) \times 10^{-5}$ reported in AUBERT,B 04B and $B(B^\pm \rightarrow K^\pm \eta_c) = (8.7 \pm 1.5) \times 10^{-3}$ reported in AUBERT 06E contribute to the determination of $B(\eta_c \rightarrow K \bar{K} \pi)$, which is used by others for normalization.

³ AUBERT,B 05L reports $[\Gamma(B^+ \rightarrow \eta_c K^+)/\Gamma_{\text{total}}] \times [B(\eta_c(1S) \rightarrow p \bar{p})] = (1.8^{+0.3}_{-0.2} \pm 0.2) \times 10^{-6}$. We divide by our best value $B(\eta_c(1S) \rightarrow p \bar{p}) = (1.3 \pm 0.4) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

⁵ EDWARDS 01 assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$. The correlated uncertainties (28.3)% from $B(J/\psi(1S) \rightarrow \gamma\eta_c)$ in those modes have been accounted for.

⁶ AUBERT, B 04B reports $[\Gamma(B^+ \rightarrow \eta_c K^+)/\Gamma_{\text{total}}] \times [B(\eta_c(1S) \rightarrow K\bar{K}\pi)] = (0.074 \pm 0.005 \pm 0.007) \times 10^{-3}$. We divide by our best value $B(\eta_c(1S) \rightarrow K\bar{K}\pi) = (7.0 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(B^+ \rightarrow \eta_c K^+)/\Gamma_{\text{total}} \times \Gamma(\eta_c(1S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}} \qquad \Gamma_{163}/\Gamma \times \Gamma_{28}^{\eta_c(1S)}/\Gamma_{\eta_c(1S)}$$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$0.22^{+0.09+0.04}_{-0.07-0.02}$	¹ WICHT	08	BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(\eta_c K^*(892)^+)/\Gamma_{\text{total}} \qquad \Gamma_{164}/\Gamma$$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$1.2^{+0.6}_{-0.4} \pm 0.4$	^{1,2} AUBERT	07AV	BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ AUBERT 07AV reports $[\Gamma(B^+ \rightarrow \eta_c K^*(892)^+)/\Gamma_{\text{total}}] \times [B(\eta_c(1S) \rightarrow p\bar{p})] = (1.57^{+0.56+0.45}_{-0.46-0.36}) \times 10^{-6}$. We divide by our best value $B(\eta_c(1S) \rightarrow p\bar{p}) = (1.3 \pm 0.4) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(\eta_c(2S) K^+)/\Gamma_{\text{total}} \qquad \Gamma_{165}/\Gamma$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$3.4 \pm 1.8 \pm 0.3$	¹ AUBERT	06E	BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ Perform measurements of absolute branching fractions using a missing mass technique.

$$\Gamma(B^+ \rightarrow h_c(1P) K^+)/\Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \eta_c \gamma)/\Gamma_{\text{total}} \qquad \Gamma_{212}/\Gamma \times \Gamma_4^{h_c(1P)}/\Gamma_{h_c(1P)}$$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.48	90	¹ AUBERT	08AB	BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ Uses the production ratio of $(B^+ B^-)/(B^0 \bar{B}^0) = 1.026 \pm 0.032$ at $\Upsilon(4S)$.

$$\Gamma(B^+ \rightarrow \eta_c(2S) K^+)/\Gamma_{\text{total}} \times \Gamma(\eta_c(2S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}} \qquad \Gamma_{165}/\Gamma \times \Gamma_7^{\eta_c(2S)}/\Gamma_{\eta_c(2S)}$$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<0.18	90	¹ WICHT	08	BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)K^+)/\Gamma_{\text{total}}$ Γ_{166}/Γ

VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT

10.07 ± 0.35 OUR FIT
10.22 ± 0.35 OUR AVERAGE

8.1 ± 1.3 ± 0.7		¹ AUBERT	06E	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
10.61 ± 0.15 ± 0.48		² AUBERT	05J	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
10.1 ± 1.0 ± 0.3		³ AUBERT,B	05L	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
10.1 ± 0.2 ± 0.7		² ABE	03B	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
10.2 ± 0.8 ± 0.7		² JESSOP	97	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
9.3 ± 3.1 ± 0.1		⁴ BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	
8.1 ± 3.5 ± 0.1	6	⁵ ALBRECHT	90J	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
10.1 ± 0.3 ± 0.5		² AUBERT	02	BABR	Repl. by AUBERT 05J
11.0 ± 1.5 ± 0.9	59	² ALAM	94	CLE2	Repl. by JESSOP 97
22 ± 10 ± 2		BUSKULIC	92G	ALEP	$e^+e^- \rightarrow Z$
7 ± 4	3	⁶ ALBRECHT	87D	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
10 ± 7 ± 2	3	⁷ BEBEK	87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
9 ± 5	3	⁸ ALAM	86	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Perform measurements of absolute branching fractions using a missing mass technique.
² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
³ AUBERT,B 05L reports $[\Gamma(B^+ \rightarrow J/\psi(1S)K^+)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow p\bar{p})] = (2.2 \pm 0.2 \pm 0.1) \times 10^{-6}$. We divide by our best value $B(J/\psi(1S) \rightarrow p\bar{p}) = (2.17 \pm 0.07) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
⁴ BORTOLETTO 92 reports $(8 \pm 2 \pm 2) \times 10^{-4}$ from a measurement of $[\Gamma(B^+ \rightarrow J/\psi(1S)K^+)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow e^+e^-)]$ assuming $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.94 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
⁵ ALBRECHT 90J reports $(7 \pm 3 \pm 1) \times 10^{-4}$ from a measurement of $[\Gamma(B^+ \rightarrow J/\psi(1S)K^+)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow e^+e^-)]$ assuming $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.94 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
⁶ ALBRECHT 87D assume $B^+B^-/B^0\bar{B}^0$ ratio is 55/45. Superseded by ALBRECHT 90J.
⁷ BEBEK 87 value has been updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92.
⁸ ALAM 86 assumes B^\pm/B^0 ratio is 60/40.

$\Gamma(\eta_c K^+)/\Gamma(J/\psi(1S)K^+)$ $\Gamma_{163}/\Gamma_{166}$

VALUE DOCUMENT ID TECN COMMENT

1.33 ± 0.10 ± 0.43 ¹ AUBERT,B 04B BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ Uses BABAR measurement of $B(B^+ \rightarrow J/\psi K^+) = (10.1 \pm 0.3 \pm 0.5) \times 10^{-4}$.

$$\Gamma(B^+ \rightarrow J/\psi(1S)K^+)/\Gamma_{\text{total}} \times \Gamma(J/\psi(1S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{166}/\Gamma \times \Gamma_{182}^{J/\psi(1S)}/\Gamma_{J/\psi(1S)}$$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<0.16	90	¹ WICHT 08	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(J/\psi(1S)K^+\pi^+\pi^-)/\Gamma_{\text{total}} \quad \Gamma_{167}/\Gamma$$

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

1.07 ± 0.19 OUR AVERAGE Error includes scale factor of 1.9.

1.16 ± 0.07 ± 0.09 ¹ AUBERT 05R BABR $e^+e^- \rightarrow \Upsilon(4S)$

0.69 ± 0.18 ± 0.12 ² ACOSTA 02F CDF $p\bar{p}$ 1.8 TeV

1.39 ± 0.82 ± 0.01 ³ BORTOLETTO92 CLEO $e^+e^- \rightarrow \Upsilon(4S)$

1.39 ± 0.91 ± 0.01 ⁶ ⁴ ALBRECHT 87D ARG $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.9 90 ⁵ ALBRECHT 90J ARG $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ACOSTA 02F uses as reference of $B(B \rightarrow J/\psi(1S)K^+) = (10.1 \pm 0.6) \times 10^{-4}$. The second error includes the systematic error and the uncertainties of the branching ratio.

³ BORTOLETTO 92 reports $(1.2 \pm 0.6 \pm 0.4) \times 10^{-3}$ from a measurement of $[\Gamma(B^+ \rightarrow J/\psi(1S)K^+\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow e^+e^-)]$ assuming $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.94 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

⁴ ALBRECHT 87D reports $(1.2 \pm 0.8) \times 10^{-3}$ from a measurement of $[\Gamma(B^+ \rightarrow J/\psi(1S)K^+\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow e^+e^-)]$ assuming $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.94 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. They actually report 0.0011 ± 0.0007 assuming $B^+B^-/B^0\bar{B}^0$ ratio is 55/45. We rescale to 50/50. Analysis explicitly removes $B^+ \rightarrow \psi(2S)K^+$.

⁵ ALBRECHT 90J reports $< 1.6 \times 10^{-3}$ from a measurement of $[\Gamma(B^+ \rightarrow J/\psi(1S)K^+\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow e^+e^-)]$ assuming $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = 5.94 \times 10^{-2}$. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(h_c(1P)K^+ \times B(h_c(1P) \rightarrow J/\psi\pi^+\pi^-))/\Gamma_{\text{total}} \quad \Gamma_{168}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.4 × 10⁻⁶	90	¹ AUBERT 05R	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(X(3872)K^+)/\Gamma_{\text{total}} \quad \Gamma_{169}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.2 × 10⁻⁴	90	¹ AUBERT 06E	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Perform measurements of absolute branching fractions using a missing mass technique.

$$\Gamma(B^+ \rightarrow X(3872)K^+)/\Gamma_{\text{total}} \times \Gamma(X(3872) \rightarrow \gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{169}/\Gamma \times \Gamma_4^{X(3872)}/\Gamma_{X(3872)}$$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<0.24	90	¹ WICHT 08	BELL	$e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$$\Gamma(X(3872)K^+ \times B(X \rightarrow J/\psi\pi^+\pi^-))/\Gamma_{\text{total}} \quad \Gamma_{170}/\Gamma$$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
9.5±1.9 OUR AVERAGE	Error includes scale factor of 1.3.		
8.4±1.5±0.7	¹ AUBERT 08Y	BABR	$e^+e^- \rightarrow \gamma(4S)$
13.0±2.9±0.7	² CHOI 03	BELL	$e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

10.1±2.5±1.0	¹ AUBERT 06	BABR	Repl. by AUBERT 08Y
12.8±4.1	¹ AUBERT 05R	BABR	Repl. by AUBERT 06

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² CHOI 03 reports $[\Gamma(B^+ \rightarrow X(3872)K^+ \times B(X \rightarrow J/\psi\pi^+\pi^-))/\Gamma_{\text{total}}] / [B(B^+ \rightarrow \psi(2S)K^+)] = 0.0200 \pm 0.0038 \pm 0.0023$. We multiply by our best value $B(B^+ \rightarrow \psi(2S)K^+) = (6.48 \pm 0.35) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(X(3872)K^+ \times B(X \rightarrow J/\psi\gamma))/\Gamma_{\text{total}} \quad \Gamma_{171}/\Gamma$$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
3.3±1.0±0.3	¹ AUBERT,BE 06M	BABR	$e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$$\Gamma(X(3872)K^+ \times B(X \rightarrow D^0\bar{D}^0))/\Gamma_{\text{total}} \quad \Gamma_{172}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<6.0 × 10⁻⁵	90	¹ CHISTOV 04	BELL	$e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$$\Gamma(X(3872)K^+ \times B(X \rightarrow D^+D^-))/\Gamma_{\text{total}} \quad \Gamma_{173}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<4.0 × 10⁻⁵	90	¹ CHISTOV 04	BELL	$e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$$\Gamma(X(3872)K^+ \times B(X \rightarrow D^0\bar{D}^0\pi^0))/\Gamma_{\text{total}} \quad \Gamma_{174}/\Gamma$$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
1.02±0.31^{+0.21}_{-0.29}		¹ GOKHROO 06	BELL	$e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.6	90	² CHISTOV 04	BELL	Repl. by GOKHROO 06
------	----	-------------------------	------	---------------------

¹ Measure the near-threshold enhancements in the $(D^0\bar{D}^0\pi^0)$ system at a mass $3875.2 \pm 0.7^{+0.3}_{-1.6} \pm 0.8$ MeV/ c^2 .

² Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(X(3872)K^+ \times B(X \rightarrow \bar{D}^{*0}D^0))/\Gamma_{\text{total}}$ Γ_{175}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
$1.67 \pm 0.36 \pm 0.47$		¹ AUBERT	08B BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(X(3872)K^+ \times B(X(3872) \rightarrow J/\psi(1S)\eta))/\Gamma_{\text{total}}$ Γ_{176}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 7.7 \times 10^{-6}$	90	¹ AUBERT	04Y BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(X(3872)^+K^0 \times B(X(3872)^+ \rightarrow J/\psi(1S)\pi^+\pi^0))/\Gamma_{\text{total}}$ Γ_{177}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 22	90	¹ AUBERT	05B BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. The isovector- X hypothesis is excluded with a likelihood test at 1×10^{-4} level.

$\Gamma(X(4260)^0K^+ \times B(X^0 \rightarrow J/\psi\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{178}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 29	95	¹ AUBERT	06 BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(X(3945)^0K^+ \times B(X^0 \rightarrow J/\psi\gamma))/\Gamma_{\text{total}}$ Γ_{179}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 14	90	¹ AUBERT, BE	06M BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(B^+ \rightarrow X(3945)K^+)/\Gamma_{\text{total}} \times \Gamma(X(3945) \rightarrow \omega J/\psi)/\Gamma_{\text{total}}$
 $\Gamma_{181}/\Gamma \times \Gamma_1^{X(3945)}/\Gamma_{X(3945)}$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$4.9^{+1.0}_{-0.9} \pm 0.5$	¹ AUBERT	08W BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(Z(3930)^0K^+ \times B(Z^0 \rightarrow J/\psi\gamma))/\Gamma_{\text{total}}$ Γ_{180}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.5	90	¹ AUBERT, BE	06M BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{182}/Γ

For polarization information see the Listings at the end of the “ B^0 Branching Ratios” section.

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.43 ± 0.08 OUR AVERAGE				
1.74 $^{+0.36}_{-0.31} \pm 0.06$		^{1,2} AUBERT	07AV BABR	$e^+e^- \rightarrow \Upsilon(4S)$
1.454 $\pm 0.047 \pm 0.097$		² AUBERT	05J BABR	$e^+e^- \rightarrow \Upsilon(4S)$
1.28 $\pm 0.07 \pm 0.14$		² ABE	02N BELL	$e^+e^- \rightarrow \Upsilon(4S)$

1.41 ± 0.23 ± 0.24		² JESSOP	97	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
1.58 ± 0.47 ± 0.27		³ ABE	96H	CDF	$p\bar{p}$ at 1.8 TeV
1.51 ± 1.08 ± 0.02		⁴ BORTOLETTO	92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
1.86 ± 1.30 ± 0.02	2	⁵ ALBRECHT	90J	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.37 ± 0.09 ± 0.11		² AUBERT	02	BABR	Repl. by AUBERT 05J
1.78 ± 0.51 ± 0.23	13	² ALAM	94	CLE2	Sup. by JESSOP 97

¹ AUBERT 07AV reports $[\Gamma(B^+ \rightarrow J/\psi(1S)K^*(892)^+)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow p\bar{p})] = (3.78^{+0.72+0.28}_{-0.64-0.23}) \times 10^{-6}$. We divide by our best value $B(J/\psi(1S) \rightarrow p\bar{p}) = (2.17 \pm 0.07) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ ABE 96H assumes that $B(B^+ \rightarrow J/\psi K^+) = (1.02 \pm 0.14) \times 10^{-3}$.

⁴ BORTOLETTO 92 reports $(1.3 \pm 0.9 \pm 0.3) \times 10^{-3}$ from a measurement of $[\Gamma(B^+ \rightarrow J/\psi(1S)K^*(892)^+)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow e^+e^-)]$ assuming $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.94 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

⁵ ALBRECHT 90J reports $(1.6 \pm 1.1 \pm 0.3) \times 10^{-3}$ from a measurement of $[\Gamma(B^+ \rightarrow J/\psi(1S)K^*(892)^+)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow e^+e^-)]$ assuming $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.94 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)K^*(892)^+)/\Gamma(J/\psi(1S)K^+)$ $\Gamma_{182}/\Gamma_{166}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

1.39 ± 0.09 OUR AVERAGE

1.37 ± 0.05 ± 0.08	AUBERT	05J	BABR $e^+e^- \rightarrow \Upsilon(4S)$
1.45 ± 0.20 ± 0.17	¹ JESSOP	97	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$
1.92 ± 0.60 ± 0.17	ABE	96Q	CDF $p\bar{p}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.37 ± 0.10 ± 0.08	² AUBERT	02	BABR Repl. by AUBERT 05J
--------------------	---------------------	----	--------------------------

¹ JESSOP 97 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. The measurement is actually measured as an average over kaon charged and neutral states.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)K(1270)^+)/\Gamma_{\text{total}}$ Γ_{183}/Γ

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

1.80 ± 0.34 ± 0.39	¹ ABE	01L	BELL $e^+e^- \rightarrow \Upsilon(4S)$
---------------------------	------------------	-----	--

¹ Uses the PDG value of $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.00 \pm 0.10) \times 10^{-3}$.

$\Gamma(J/\psi(1S)K(1400)^+)/\Gamma(J/\psi(1S)K(1270)^+)$ $\Gamma_{184}/\Gamma_{183}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

< 0.30	90	ABE	01L	BELL $e^+e^- \rightarrow \Upsilon(4S)$
------------------	----	-----	-----	--

$\Gamma(J/\psi(1S)\eta K^+)/\Gamma_{\text{total}}$ Γ_{185}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$10.8 \pm 2.3 \pm 2.4$	¹ AUBERT	04Y BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)\eta' K^+)/\Gamma_{\text{total}}$ Γ_{186}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<8.8	90	¹ XIE	07 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)\phi K^+)/\Gamma_{\text{total}}$ Γ_{187}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$(5.2 \pm 1.7) \times 10^{-5}$ OUR AVERAGE Error includes scale factor of 1.2.			
$(4.4 \pm 1.4 \pm 0.5) \times 10^{-5}$	¹ AUBERT	03O BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$(8.8^{+3.5}_{-3.0} \pm 1.3) \times 10^{-5}$	² ANASTASSOV 00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ANASTASSOV 00 finds 10 events on a background of 0.5 ± 0.2 . Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$, a uniform Dalitz plot distribution, isotropic $J/\psi(1S)$ and ϕ decays, and $B(B^+ \rightarrow J/\psi(1S)\phi K^+) = B(B^0 \rightarrow J/\psi(1S)\phi K^0)$.

$\Gamma(J/\psi(1S)\omega K^+ \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{188}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$3.5 \pm 0.2 \pm 0.4$	¹ AUBERT	08W BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)\pi^+)/\Gamma_{\text{total}}$ Γ_{189}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$(4.9 \pm 0.6) \times 10^{-5}$ OUR FIT Error includes scale factor of 1.5.			
$(3.8 \pm 0.6 \pm 0.3) \times 10^{-5}$	¹ ABE	03B BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)\pi^+)/\Gamma(J/\psi(1S)K^+)$ $\Gamma_{189}/\Gamma_{166}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.049 ± 0.006 OUR FIT Error includes scale factor of 1.5.				
0.053 ± 0.004 OUR AVERAGE				

0.0537 \pm 0.0045 \pm 0.0011 AUBERT 04P BABR $e^+e^- \rightarrow \Upsilon(4S)$

0.05 $^{+0.019}_{-0.017} \pm 0.001$ ABE 96R CDF $p\bar{p}$ 1.8 TeV

0.052 \pm 0.024 BISHAI 96 CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0391 \pm 0.0078 \pm 0.0019 AUBERT 02F BABR Repl. by AUBERT 04P

0.043 \pm 0.023 5 ¹ ALEXANDER 95 CLE2 Sup. by BISHAI 96

¹ Assumes equal production of $B^+ B^-$ and $B^0 \bar{B}^0$ on $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)\rho^+)/\Gamma_{\text{total}}$ Γ_{190}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

$5.0 \pm 0.7 \pm 0.3$ ¹ AUBERT 07AC BABR $e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<77 90 BISHAI 96 CLE2 $e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(J/\psi(1S)\pi^+\pi^0\text{nonresonant})/\Gamma_{\text{total}}$ Γ_{191}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<0.73 ¹ AUBERT 07AC BABR $e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(J/\psi(1S)a_1(1260)^+)/\Gamma_{\text{total}}$ Γ_{192}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 1.2×10^{-3} 90 BISHAI 96 CLE2 $e^+e^- \rightarrow \gamma(4S)$

$\Gamma(J/\psi(1S)\rho\bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{193}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

11.8 ± 3.1 OUR AVERAGE

$11.7 \pm 2.8^{+1.8}_{-2.3}$ ¹ XIE 05 BELL $e^+e^- \rightarrow \gamma(4S)$

12 $^{+9}_{-6}$ ¹ AUBERT 03K BABR $e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<41 90 ZANG 04 BELL $e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(J/\psi(1S)\bar{\Sigma}^0\rho)/\Gamma_{\text{total}}$ Γ_{194}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 1.1×10^{-5} 90 ¹ XIE 05 BELL $e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(J/\psi(1S)D^+)/\Gamma_{\text{total}}$ Γ_{195}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<12 90 ¹ AUBERT 05U BABR $e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(J/\psi(1S)\bar{D}^0\pi^+)/\Gamma_{\text{total}}$ Γ_{196}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<2.5 90 ¹ ZHANG 05B BELL $e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<5.2 90 ¹ AUBERT 05R BABR $e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\psi(2S)\pi^+)/\Gamma_{\text{total}}$ Γ_{197}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
2.44 ± 0.22 ± 0.20	¹ BHARDWAJ 08	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\psi(2S)\pi^+)/\Gamma(\psi(2S)K^+)$ $\Gamma_{197}/\Gamma_{198}$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
3.99 ± 0.36 ± 0.17	BHARDWAJ 08	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\psi(2S)K^+)/\Gamma_{\text{total}}$ Γ_{198}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
6.48 ± 0.35 OUR AVERAGE					
4.9 ± 1.6 ± 0.4			¹ AUBERT 06E	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
6.17 ± 0.32 ± 0.44			² AUBERT 05J	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
6.9 ± 0.6			² ABE 03B	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
7.8 ± 0.7 ± 0.9			² RICHICHI 01	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
5.5 ± 1.0 ± 0.6			³ ABE 98O	CDF	$p\bar{p}$ 1.8 TeV
18 ± 8 ± 4		5	² ALBRECHT 90J	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.4 ± 0.5 ± 0.8			² AUBERT 02	BABR	Repl. by AUBERT 05J
6.1 ± 2.3 ± 0.9		7	² ALAM 94	CLE2	Repl. by RICHICHI 01
<5		90	² BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
22 ± 17		3	⁴ ALBRECHT 87D	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Perform measurements of absolute branching fractions using a missing mass technique.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ ABE 98O reports $[B(B^+ \rightarrow \psi(2S)K^+)/B(B^+ \rightarrow J/\psi(1S)K^+)] = 0.558 \pm 0.082 \pm 0.056$. We multiply by our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = (9.9 \pm 1.0) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ ALBRECHT 87D assume $B^+B^-/B^0\bar{B}^0$ ratio is 55/45. Superseded by ALBRECHT 90J.

$\Gamma(\psi(2S)K^+)/\Gamma(J/\psi(1S)K^+)$ $\Gamma_{198}/\Gamma_{166}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.64 ± 0.06 ± 0.07	¹ AUBERT 02	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\psi(2S)K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{199}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
6.7 ± 1.4 OUR AVERAGE				Error includes scale factor of 1.3.
5.92 ± 0.85 ± 0.89		¹ AUBERT 05J	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
9.2 ± 1.9 ± 1.2		¹ RICHICHI 01	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<30		90	¹ ALAM 94	CLE2	Repl. by RICHICHI 01
<35		90	¹ BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
<49		90	¹ ALBRECHT 90J	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\psi(2S)K^*(892)^+)/\Gamma(\psi(2S)K^+)$ $\Gamma_{199}/\Gamma_{198}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.96±0.15±0.09	AUBERT	05J	BABR $e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\psi(2S)K^+\pi^+\pi^-)/\Gamma_{total}$ Γ_{200}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0019±0.0011±0.0004	3	¹ ALBRECHT	90J	ARG $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\psi(3770)K^+)/\Gamma_{total}$ Γ_{201}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.49±0.13 OUR AVERAGE			

3.5 ±2.5 ±0.3 ¹ AUBERT 06E BABR $e^+e^- \rightarrow \Upsilon(4S)$

0.48±0.11±0.07 ² CHISTOV 04 BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Perform measurements of absolute branching fractions using a missing mass technique.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\psi(3770)K^+ \times B(\psi \rightarrow D^0\bar{D}^0))/\Gamma_{total}$ Γ_{202}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
1.6 ±0.4 OUR AVERAGE	Error includes scale factor of 1.1.		

1.41±0.30±0.22 ¹ AUBERT 08B BABR $e^+e^- \rightarrow \Upsilon(4S)$

2.2 ±0.5 ±0.3 ¹ BRODZICKA 08 BELL $e^+e^- \rightarrow \Upsilon(4S)$

••• We do not use the following data for averages, fits, limits, etc. •••

3.4 ±0.8 ±0.5 ¹ CHISTOV 04 BELL Repl. by BRODZICKA 08

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\psi(3770)K^+ \times B(\psi \rightarrow D^+D^-))/\Gamma_{total}$ Γ_{203}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
0.94±0.35 OUR AVERAGE			

0.84±0.32±0.21 ¹ AUBERT 08B BABR $e^+e^- \rightarrow \Upsilon(4S)$

1.4 ±0.8 ±0.2 ¹ CHISTOV 04 BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\chi_{c0}\pi^+ \times B(\chi_{c0} \rightarrow \pi^+\pi^-))/\Gamma_{total}$ Γ_{204}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<0.3	90	¹ AUBERT,B	05G	BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\chi_{c0}(1P)K^+)/\Gamma_{total}$ Γ_{205}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
1.43^{+0.21}_{-0.17} OUR AVERAGE				

1.46^{+0.32}_{-0.29}±0.11 ^{1,2} AUBERT 08AI BABR $e^+e^- \rightarrow \Upsilon(4S)$

1.84±0.32±0.31 ^{1,3} AUBERT 06O BABR $e^+e^- \rightarrow \Upsilon(4S)$

5.3 ±2.5 ±0.4 ⁴ AUBERT,BE 06M BABR $e^+e^- \rightarrow \Upsilon(4S)$

1.12±0.12^{+0.30}_{-0.20} ¹ GARMASH 06 BELL $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<5	90	^{1,5} WICHT	08	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
<1.8	90	⁶ AUBERT	06E	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
<8.9	90	¹ AUBERT	05K	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$1.39 \pm 0.49 \pm 0.11$		⁷ AUBERT,B	05N	BABR	Repl. by AUBERT 08AI
$1.96 \pm 0.35^{+2.00}_{-0.42}$		¹ GARMASH	05	BELL	Repl. by GARMASH 06
2.7 ± 0.7		⁸ AUBERT	04T	BABR	Repl. by AUBERT,B 04P
$3.0 \pm 0.8 \pm 0.3$		⁹ AUBERT,B	04P	BABR	Repl. by AUBERT,B 05N
$6.0^{+2.1}_{-1.8} \pm 1.1$		¹⁰ ABE	02B	BELL	Repl. by GARMASH 05
<4.8	90	¹¹ EDWARDS	01	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² AUBERT 08AI reports $(0.70 \pm 0.10^{+0.12}_{-0.10}) \times 10^{-6}$ for $B(B^+ \rightarrow \chi_{c0} K^+) \times B(\chi_{c0} \rightarrow \pi^+ \pi^-)$. We compute $B(B^+ \rightarrow \chi_{c0} K^+)$ using the PDG value $B(\chi_{c0} \rightarrow \pi\pi) = (7.2 \pm 0.6) \times 10^{-3}$ and 2/3 for the $\pi^+ \pi^-$ fraction. Our first error is their experiment's error and the second error is systematic error from using our best value.

³ Measured in the $B^+ \rightarrow K^+ K^- K^+$ decay.

⁴ AUBERT, BE 06M reports $[\Gamma(B^+ \rightarrow \chi_{c0}(1P) K^+)/\Gamma_{\text{total}}] \times [B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S))] = (6.1 \pm 2.6 \pm 1.1) \times 10^{-6}$. We divide by our best value $B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (1.14 \pm 0.08) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. The significance of the observed signal is 2.4 σ .

⁵ WICHT 08 reports $[\Gamma(B^+ \rightarrow \chi_{c0}(1P) K^+)/\Gamma_{\text{total}}] \times [B(\chi_{c0}(1P) \rightarrow \gamma\gamma)] < 0.11 \times 10^{-6}$. We divide by our best value $B(\chi_{c0}(1P) \rightarrow \gamma\gamma) = 2.27 \times 10^{-4}$.

⁶ Perform measurements of absolute branching fractions using a missing mass technique.

⁷ AUBERT,B 05N reports $(0.66 \pm 0.22 \pm 0.08) \times 10^{-6}$ for $B(B^+ \rightarrow \chi_c^0 K^+) \times B(\chi_c^0 \rightarrow \pi^+ \pi^-)$. We compute $B(B^+ \rightarrow \chi_c^0 K^+)$ using the PDG value $B(\chi_c^0 \rightarrow \pi^+ \pi^-) = (7.1 \pm 0.6) \times 10^{-3}$ and 2/3 for the $\pi^+ \pi^-$ fraction.

⁸ The measurement performed using decay channels $\chi_c^0 \rightarrow \pi^+ \pi^-$ and $\chi_c^0 \rightarrow K^+ K^-$. The ratio of the branching ratios for these channels is found to be consistent with world average.

⁹ AUBERT 04P reports $B(B^+ \rightarrow \chi_c^0 K^+) \times B(\chi_c^0 \rightarrow \pi^+ \pi^-) = (1.5 \pm 0.4 \pm 0.1) \times 10^{-6}$ and used PDG value of $B(\chi_c^0 \rightarrow \pi\pi) = (7.4 \pm 0.8) \times 10^{-3}$ and Clebsh-Gordan coefficient to compute $B(B^{\pm} \rightarrow \chi_c^0 K^+)$.

¹⁰ ABE 02B measures the ratio of $B(B^+ \rightarrow \chi_c^0 K^+)/B(B^+ \rightarrow J/\psi(1S) K^+) = 0.60 + 0.21 - 0.18 \pm 0.05 \pm 0.08$, where the third error is due to the uncertainty in the $B(\chi_c^0 \rightarrow \pi^+ \pi^-)$, and uses $B(B^+ \rightarrow J/\psi(1S) K^+) = (10.0 \pm 1.0) \times 10^{-4}$ to obtain the result.

¹¹ EDWARDS 01 assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$. The correlated uncertainties (28.3)% from $B(J/\psi(1S) \rightarrow \gamma \eta_c)$ in those modes have been accounted for.

$\Gamma(\chi_{c0} K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{206}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.1	90	¹ AUBERT	08BD	BABR $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<28.6	90	¹ AUBERT	05K	BABR	Repl. by AUBERT 08BD
-------	----	---------------------	-----	------	----------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\chi_{c2} K^+)/\Gamma_{\text{total}}$ Γ_{207}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<2.9 \times 10^{-5}$	90	¹ SONI	06 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
-----------------------	----	-------------------	---------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<2.0 \times 10^{-4}$	90	² AUBERT	06E BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
-----------------------	----	---------------------	----------	------------------------------------

$<3.0 \times 10^{-5}$	90	¹ AUBERT	05K BABR	Repl. by AUBERT 06E
-----------------------	----	---------------------	----------	---------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Perform measurements of absolute branching fractions using a missing mass technique.

$\Gamma(B^+ \rightarrow \chi_{c2} K^+)/\Gamma_{\text{total}} \times \Gamma(\chi_{c2}(1P) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{207}/\Gamma \times \Gamma_{54}^{\chi_{c2}(1P)}/\Gamma_{\chi_{c2}(1P)}$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<0.09	90	¹ WICHT	08 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
---------	----	--------------------	---------	------------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\chi_{c2} K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{208}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<1.2 \times 10^{-5}$	90	¹ AUBERT	05K BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
-----------------------	----	---------------------	----------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.27 \times 10^{-4}$	90	¹ SONI	06 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
------------------------	----	-------------------	---------	------------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\chi_{c1}(1P)\pi^+)/\Gamma_{\text{total}}$ Γ_{209}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$2.2 \pm 0.4 \pm 0.3$	¹ KUMAR	06 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
-----------------------	--------------------	---------	------------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\chi_{c1}(1P)K^+)/\Gamma_{\text{total}}$ Γ_{210}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

5.1 ± 0.5 OUR AVERAGE		Error includes scale factor of 1.6. See the ideogram below.		
------------------------------	--	---	--	--

$8.1 \pm 1.4 \pm 0.7$		¹ AUBERT	06E BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
-----------------------	--	---------------------	----------	------------------------------------

$5.2 \pm 0.4 \pm 0.2$		² AUBERT, BE	06M BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
-----------------------	--	-------------------------	----------	------------------------------------

$4.49 \pm 0.19 \pm 0.53$		³ SONI	06 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
--------------------------	--	-------------------	---------	------------------------------------

$15.5 \pm 5.4 \pm 2.0$		⁴ ACOSTA	02F CDF	$p\bar{p}$ 1.8 TeV
------------------------	--	---------------------	---------	--------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.79 \pm 0.26 \pm 0.65$		³ AUBERT	05J BABR	Repl. by AUBERT, BE 06M
--------------------------	--	---------------------	----------	-------------------------

$6.0 \pm 0.9 \pm 0.3$		⁵ AUBERT	02 BABR	Repl. by AUBERT 05J
-----------------------	--	---------------------	---------	---------------------

$9.7 \pm 4.0 \pm 0.9$	6	³ ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
-----------------------	---	-------------------	---------	------------------------------------


$19 \pm 13 \pm 6$		⁶ ALBRECHT	92E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
-------------------	--	-----------------------	---------	------------------------------------

¹ Perform measurements of absolute branching fractions using a missing mass technique.

² AUBERT, BE 06M reports $[\Gamma(B^+ \rightarrow \chi_{c1}(1P)K^+)/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))] = (1.76 \pm 0.07 \pm 0.12) \times 10^{-4}$. We divide by our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (34.1 \pm 1.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

- ⁴ ACOSTA 02F uses as reference of $B(B \rightarrow J/\psi(1S)K^+) = (10.1 \pm 0.6) \times 10^{-4}$. The second error includes the systematic error and the uncertainties of the branching ratio.
- ⁵ AUBERT 02 reports $(7.5 \pm 0.9 \pm 0.8) \times 10^{-4}$ from a measurement of $[\Gamma(B^+ \rightarrow \chi_{c1}(1P)K^+)/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$. We rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (34.1 \pm 1.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
- ⁶ ALBRECHT 92E assumes no $\chi_{c2}(1P)$ production and $B(\Upsilon(4S) \rightarrow B^+ B^-) = 50\%$.



$\Gamma(\chi_{c1}(1P)K^+)/\Gamma(J/\psi(1S)K^+)$ $\Gamma_{210}/\Gamma_{166}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.60±0.07±0.03	¹ AUBERT	02	BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ AUBERT 02 reports $0.75 \pm 0.08 \pm 0.05$ from a measurement of $[\Gamma(B^+ \rightarrow \chi_{c1}(1P)K^+)/\Gamma(B^+ \rightarrow J/\psi(1S)K^+)] \times [B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$. We rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (34.1 \pm 1.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\chi_{c1}(1P)\pi^+)/\Gamma(\chi_{c1}(1P)K^+)$ $\Gamma_{209}/\Gamma_{210}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.043±0.008±0.003	¹ KUMAR	06	BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\chi_{c1}(1P)K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{211}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

3.6 ± 0.9 OUR AVERAGE

4.05 ± 0.59 ± 0.95		¹ SONI	06	BELL $e^+e^- \rightarrow \Upsilon(4S)$
--------------------	--	-------------------	----	--

2.94 ± 0.95 ± 0.98		¹ AUBERT	05J	BABR $e^+e^- \rightarrow \Upsilon(4S)$
--------------------	--	---------------------	-----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

<21	90	¹ ALAM	94	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$
-------	----	-------------------	----	--

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\chi_{c1}(1P)K^*(892)^+)/\Gamma(\chi_{c1}(1P)K^+)$ $\Gamma_{211}/\Gamma_{210}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.51 ± 0.17 ± 0.16	AUBERT	05J	BABR $e^+e^- \rightarrow \Upsilon(4S)$
---------------------------	--------	-----	--

$\Gamma(h_c(1P)K^+)/\Gamma_{\text{total}}$ Γ_{212}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

<3.8	90	¹ FANG	06	BELL $e^+e^- \rightarrow \Upsilon(4S)$
-----------------------------	----	-------------------	----	--

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and $B(h_c \rightarrow \eta_c \gamma) = 50\%$.

$\Gamma(K^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{213}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

23.1 ± 1.0 OUR AVERAGE

22.8 ⁺ ₋ 0.8 ⁺ ₋ ± 1.3		¹ LIN	07	BELL $e^+e^- \rightarrow \Upsilon(4S)$
--	--	------------------	----	--

23.9 ± 1.1 ± 1.0		¹ AUBERT, BE	06C	BABR $e^+e^- \rightarrow \Upsilon(4S)$
------------------	--	-------------------------	-----	--

18.8 ⁺ ₋ 3.7 ⁺ ₋ ± 2.1 ₋ 1.8		¹ BORNHEIM	03	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$
---	--	-----------------------	----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

26.0 ± 1.3 ± 1.0		¹ AUBERT, BE	05E	BABR Repl. by AUBERT, BE 06C
------------------	--	-------------------------	-----	------------------------------

22.3 ± 1.7 ± 1.1		¹ AUBERT	04M	BABR Repl. by AUBERT, BE 05E
------------------	--	---------------------	-----	------------------------------

22.0 ± 1.9 ± 1.1		¹ CHAO	04	BELL Repl. by LIN 07
------------------	--	-------------------	----	----------------------

19.4 ⁺ ₋ 3.1 ⁺ ₋ ± 1.6		¹ CASEY	02	BELL Repl. by CHAO 04
--	--	--------------------	----	-----------------------

13.7 ⁺ ₋ 5.7 ⁺ ₋ ± 1.9 ₋ 1.8		¹ ABE	01H	BELL Repl. by CASEY 02
---	--	------------------	-----	------------------------

18.2 ⁺ ₋ 3.3 ⁺ ₋ ± 2.0		¹ AUBERT	01E	BABR Repl. by AUBERT 04M
--	--	---------------------	-----	--------------------------

18.2 ⁺ ₋ 4.6 ⁺ ₋ ± 1.6		¹ CRONIN-HEN..00	CLE2	Repl. by BORNHEIM 03
--	--	-----------------------------	------	----------------------

23 ⁺ ₋ 11 ⁺ ₋ ± 3.6		GODANG	98	CLE2 Repl. by CRONIN-HENNESSY 00
---	--	--------	----	----------------------------------

<48	90	ASNER	96	CLE2 Repl. by GODANG 98
-------	----	-------	----	-------------------------

<190	90	ALBRECHT	91B	ARG $e^+e^- \rightarrow \Upsilon(4S)$
--------	----	----------	-----	---------------------------------------

<100	90	² AVERY	89B	CLEO $e^+e^- \rightarrow \Upsilon(4S)$
--------	----	--------------------	-----	--

<680	90	AVERY	87	CLEO $e^+e^- \rightarrow \Upsilon(4S)$
--------	----	-------	----	--

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² AVERY 89B reports $<9 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(K^+\pi^0)/\Gamma_{\text{total}}$ Γ_{214}/Γ

VALUE (units 10^{-6}) CL% DOCUMENT ID TECN COMMENT

12.9 ± 0.6 OUR AVERAGE

13.6 ± 0.6 ± 0.7		¹ AUBERT	07BC	BABR	$e^+e^- \rightarrow \gamma(4S)$
12.4 ± 0.5 ± 0.6		¹ LIN	07A	BELL	$e^+e^- \rightarrow \gamma(4S)$
12.9 ^{+2.4+1.2} _{-2.2-1.1}		¹ BORNHEIM	03	CLE2	$e^+e^- \rightarrow \gamma(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
12.0 ± 0.7 ± 0.6		¹ AUBERT	05L	BABR	Repl. by AUBERT 07BC
12.0 ± 1.3 ^{+1.3} _{-0.9}		¹ CHAO	04	BELL	Repl. by LIN 07A
12.8 ^{+1.2} _{-1.1} ± 1.0		¹ AUBERT	03L	BABR	Repl. by AUBERT 05L
13.0 ^{+2.5} _{-2.4} ± 1.3		¹ CASEY	02	BELL	Repl. by CHAO 04
16.3 ^{+3.5+1.6} _{-3.3-1.8}		¹ ABE	01H	BELL	Repl. by CASEY 02
10.8 ^{+2.1} _{-1.9} ± 1.0		¹ AUBERT	01E	BABR	Repl. by AUBERT 03L
11.6 ^{+3.0+1.4} _{-2.7-1.3}		¹ CRONIN-HEN..00		CLE2	Repl. by BORNHEIM 03
<16	90	GODANG	98	CLE2	Repl. by CRONIN-HENNESSY 00
<14	90	ASNER	96	CLE2	Repl. by GODANG 98

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K^+\pi^0)/\Gamma(K^0\pi^+)$ $\Gamma_{214}/\Gamma_{213}$

VALUE DOCUMENT ID TECN COMMENT

0.54 ± 0.03 ± 0.04 LIN 07A BELL $e^+e^- \rightarrow \gamma(4S)$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

2.38 ^{+0.98+0.39} _{-1.10-0.26}		ABE	01H	BELL	Repl. by LIN 07A
--	--	-----	-----	------	------------------

$\Gamma(\eta' K^+)/\Gamma_{\text{total}}$ Γ_{215}/Γ

VALUE (units 10^{-6}) DOCUMENT ID TECN COMMENT

70.0 ± 2.4 OUR AVERAGE

67 ⁺¹⁰ ₋₉ ± 4		^{1,2} WICHT	08	BELL	$e^+e^- \rightarrow \gamma(4S)$
70.0 ± 1.5 ± 2.8		¹ AUBERT	07AE	BABR	$e^+e^- \rightarrow \gamma(4S)$
69.2 ± 2.2 ± 3.7		¹ SCHUEMANN	06	BELL	$e^+e^- \rightarrow \gamma(4S)$
80 ⁺¹⁰ ₋₉ ± 7		¹ RICHICHI	00	CLE2	$e^+e^- \rightarrow \gamma(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
68.9 ± 2.0 ± 3.2		¹ AUBERT	05M	BABR	Repl. by AUBERT 07AE
76.9 ± 3.5 ± 4.4		¹ AUBERT	03W	BABR	Repl. by AUBERT 05M
79 ⁺¹² ₋₁₁ ± 9		¹ ABE	01M	BELL	Repl. by SCHUEMANN 06
70 ± 8 ± 5		¹ AUBERT	01G	BABR	Repl. by AUBERT 03W
65 ⁺¹⁵ ₋₁₄ ± 9		BEHRENS	98	CLE2	Repl. by RICHICHI 00

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² WICHT 08 reports $[\Gamma(B^+ \rightarrow \eta' K^+)/\Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \gamma\gamma)] = (1.40^{+0.16+0.15}_{-0.15-0.12}) \times 10^{-6}$. We divide by our best value $B(\eta'(958) \rightarrow \gamma\gamma) = (2.10 \pm 0.12) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\eta' K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{216}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$4.9^{+1.9}_{-1.7} \pm 0.8$		¹ AUBERT	07E BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 2.9	90	¹ SCHUEMANN	07 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
<14	90	¹ AUBERT,B	04D BABR	Repl. by AUBERT 07E
<35	90	¹ RICHICHI	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
<13	90	BEHRENS	98 CLE2	Repl. by RICHICHI 00

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta K^+)/\Gamma_{\text{total}}$ Γ_{217}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.6 ± 0.5 OUR AVERAGE		Error includes scale factor of 2.4. See the ideogram below.		


$2.21^{+0.48}_{-0.42} \pm 0.01$		^{1,2} WICHT	08 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$3.7 \pm 0.4 \pm 0.1$		¹ AUBERT	07AE BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$1.9 \pm 0.3^{+0.2}_{-0.1}$		¹ CHANG	07B BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$2.2^{+2.8}_{-2.2}$		¹ RICHICHI	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.3 \pm 0.6 \pm 0.3$		¹ AUBERT,B	05K BABR	Repl. by AUBERT 07AE
$2.1 \pm 0.6 \pm 0.2$		¹ CHANG	05A BELL	Repl. by CHANG 07B
$3.4 \pm 0.8 \pm 0.2$		¹ AUBERT	04H BABR	Repl. by AUBERT,B 05K
<14	90	BEHRENS	98 CLE2	Repl. by RICHICHI 00

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² WICHT 08 reports $[\Gamma(B^+ \rightarrow \eta K^+)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = (0.87^{+0.16+0.10}_{-0.15-0.07}) \times 10^{-6}$. We divide by our best value $B(\eta \rightarrow 2\gamma) = (39.30 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.



$\Gamma(\eta K^*(892)^+)/\Gamma_{\text{total}}$ **Γ_{218}/Γ**

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
19.3±1.6 OUR AVERAGE				
19.3 ^{+2.0} _{-1.9} ±1.5		¹ WANG	07B BELL	$e^+e^- \rightarrow \Upsilon(4S)$
18.9±1.8±1.3		¹ AUBERT,B	06H BABR	$e^+e^- \rightarrow \Upsilon(4S)$
26.4 ^{+9.6} _{-8.2} ±3.3		¹ RICHICHI	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
25.6±4.0±2.4		¹ AUBERT,B	04D BABR	Repl. by AUBERT,B 06H
<30	90	BEHRENS	98 CLE2	Repl. by RICHICHI 00

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta K_0^*(1430)^+)/\Gamma_{\text{total}}$ **Γ_{219}/Γ**

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
18.2±2.6±2.6	¹ AUBERT,B	06H BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta K_2^*(1430)^+)/\Gamma_{\text{total}}$ **Γ_{220}/Γ**

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
9.1±2.7±1.4	¹ AUBERT,B	06H BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta(1295)K^+ \times B(\eta(1295) \rightarrow \eta\pi\pi))/\Gamma_{\text{total}}$ Γ_{221}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

$2.9^{+0.8}_{-0.7} \pm 0.2$	90	¹ AUBERT	08X BABR	$e^+e^- \rightarrow \Upsilon(4S)$
-----------------------------	----	---------------------	----------	-----------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta(1405)K^+ \times B(\eta(1405) \rightarrow \eta\pi\pi))/\Gamma_{\text{total}}$ Γ_{222}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<1.3	90	¹ AUBERT	08X BABR	$e^+e^- \rightarrow \Upsilon(4S)$
--------	----	---------------------	----------	-----------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta(1405)K^+ \times B(\eta(1405) \rightarrow K^*K))/\Gamma_{\text{total}}$ Γ_{223}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<1.2	90	¹ AUBERT	08X BABR	$e^+e^- \rightarrow \Upsilon(4S)$
--------	----	---------------------	----------	-----------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta(1475)K^+ \times B(\eta(1475) \rightarrow K^*K))/\Gamma_{\text{total}}$ Γ_{224}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

$13.8^{+1.8+1.0}_{-1.7-0.6}$	90	¹ AUBERT	08X BABR	$e^+e^- \rightarrow \Upsilon(4S)$
------------------------------	----	---------------------	----------	-----------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(f_1(1285)K^+)/\Gamma_{\text{total}}$ Γ_{225}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<2.0	90	¹ AUBERT	08X BABR	$e^+e^- \rightarrow \Upsilon(4S)$
--------	----	---------------------	----------	-----------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(f_1(1420)K^+ \times B(f_1(1420) \rightarrow \eta\pi\pi))/\Gamma_{\text{total}}$ Γ_{226}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<2.9	90	¹ AUBERT	08X BABR	$e^+e^- \rightarrow \Upsilon(4S)$
--------	----	---------------------	----------	-----------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(f_1(1420)K^+ \times B(f_1(1420) \rightarrow K^*K))/\Gamma_{\text{total}}$ Γ_{227}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<4.1	90	¹ AUBERT	08X BABR	$e^+e^- \rightarrow \Upsilon(4S)$
--------	----	---------------------	----------	-----------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\phi(1680)K^+ \times B(\phi(1680) \rightarrow K^*K))/\Gamma_{\text{total}}$ Γ_{228}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<3.4	90	¹ AUBERT	08X BABR	$e^+e^- \rightarrow \Upsilon(4S)$
--------	----	---------------------	----------	-----------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\omega K^+)/\Gamma_{\text{total}}$ Γ_{229}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
6.7 ± 0.8	OUR AVERAGE	Error includes scale factor of 1.8.		
$6.3 \pm 0.5 \pm 0.3$		¹ AUBERT	07AE BABR	$e^+ e^- \rightarrow \gamma(4S)$
$8.1 \pm 0.6 \pm 0.6$		¹ JEN	06 BELL	$e^+ e^- \rightarrow \gamma(4S)$
$3.2^{+2.4}_{-1.9} \pm 0.8$		¹ JESSOP	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$6.1 \pm 0.6 \pm 0.4$		¹ AUBERT,B	06E BABR	AUBERT 07AE
$4.8 \pm 0.8 \pm 0.4$		¹ AUBERT	04H BABR	Repl. by AUBERT,B 06E
$6.5^{+1.3}_{-1.2} \pm 0.6$		¹ WANG	04A BELL	Repl. by JEN 06
$9.2^{+2.6}_{-2.3} \pm 1.0$		¹ LU	02 BELL	Repl. by WANG 04A
<4	90	¹ AUBERT	01G BABR	$e^+ e^- \rightarrow \gamma(4S)$
$1.5^{+7}_{-6} \pm 2$		¹ BERGFELD	98 CLE2	Repl. by JESSOP 00

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\omega K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{230}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 7.4	90	¹ AUBERT	09H BABR	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.4	90	¹ AUBERT,B	06T BABR	Repl. by AUBERT 09H
< 7.4	90	¹ AUBERT	05O BABR	Repl. by AUBERT,B 06T
<87	90	¹ BERGFELD	98 CLE2	

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\omega(K\pi)_0^{*+})/\Gamma_{\text{total}}$ Γ_{231}/Γ

$(K\pi)_0^{*+}$ is the total S-wave composed of $K_0^*(1430)$ and nonresonant that are described using LASS shape.

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$27.5 \pm 3.0 \pm 2.6$	¹ AUBERT	09H BABR	$e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\omega K_0^*(1430)^+)/\Gamma_{\text{total}}$ Γ_{232}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$24.0 \pm 2.6 \pm 4.4$	¹ AUBERT	09H BABR	$e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\omega K_2^*(1430)^+)/\Gamma_{\text{total}}$ Γ_{233}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$21.5 \pm 3.6 \pm 2.4$	¹ AUBERT	09H BABR	$e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(a_0(980)^0 K^+ \times B(a_0(980)^0 \rightarrow \eta\pi^0))/\Gamma_{\text{total}}$ Γ_{235}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<2.5	90	¹ AUBERT,BE 04	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of charged and neutral B mesons from $\Upsilon(4S)$ decays.

$\Gamma(a_0(980)^+ K^0 \times B(a_0(980)^+ \rightarrow \eta\pi^+))/\Gamma_{\text{total}}$ Γ_{234}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<3.9	90	¹ AUBERT,BE 04	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of charged and neutral B mesons from $\Upsilon(4S)$ decays.

$\Gamma(K^*(892)^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{236}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
10.1 ± 0.9 OUR AVERAGE				
10.8 ± 0.6 ^{+1.2} / _{-1.4}		¹ AUBERT 08AI	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
9.67 ± 0.64 ^{+0.81} / _{-0.89}		¹ GARMASH 06	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

13.5 ± 1.2 ^{+0.8} / _{-0.9}		¹ AUBERT,B 05N	BABR	Repl. by AUBERT 08AI
9.8 ± 0.9 ^{+1.1} / _{-1.2}		¹ GARMASH 05	BELL	Repl. by GARMASH 06
15.5 ± 1.8 ^{+1.5} / _{-4.0}		^{1,2} AUBERT,B 04P	BABR	Repl. by AUBERT,B 05N
19.4 ^{+4.2} / _{-3.9} ^{+4.1} / _{-7.1}		³ GARMASH 02	BELL	Repl. by GARMASH 05
<119	90	⁴ ABE 00C	SLD	$e^+e^- \rightarrow Z$
< 16	90	¹ JESSOP 00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
<390	90	⁵ ADAM 96D	DLPH	$e^+e^- \rightarrow Z$
< 41	90	ASNER 96	CLE2	Repl. by JESSOP 00
<480	90	⁵ ABREU 95N	DLPH	Sup. by ADAM 96D
<170	90	ALBRECHT 91B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
<150	90	⁶ AVERY 89B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
<260	90	AVERY 87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² AUBERT 04P also report a branching ratio for $B^+ \rightarrow$ "higher K^* resonances" π^+ , $K^* \rightarrow K^+\pi^-$, $(25.1 \pm 2.0^{+11.0}_{-5.7}) \times 10^{-6}$.

³ Uses a reference decay mode $B^+ \rightarrow \bar{D}^0 \pi^+$ and $\bar{D}^0 \rightarrow K^+\pi^-$ with $B(B^+ \rightarrow \bar{D}^0 \pi^+) \cdot B(\bar{D}^0 \rightarrow K^+\pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.

⁴ ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

⁵ Assumes a B^0 , B^- production fraction of 0.39 and a B_s production fraction of 0.12.

⁶ AVERY 89B reports $< 1.3 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(K^*(892)^+\pi^0)/\Gamma_{\text{total}}$ Γ_{237}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$6.9 \pm 2.0 \pm 1.3$		¹ AUBERT	05X BABR	$e^+e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<31	90	¹ JESSOP	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
<99	90	ASNER	96 CLE2	Repl. by JESSOP 00
¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.				

$\Gamma(K^+\pi^-\pi^+)/\Gamma_{\text{total}}$ Γ_{238}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
51.0 ± 2.9 OUR AVERAGE			
$54.4 \pm 1.1 \pm 4.6$	¹ AUBERT	08AI BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$48.8 \pm 1.1 \pm 3.6$	¹ GARMASH	06 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$64.1 \pm 2.4 \pm 4.0$	¹ AUBERT,B	05N BABR	Repl. by AUBERT 08AI
$46.6 \pm 2.1 \pm 4.3$	¹ GARMASH	05 BELL	Repl. by GARMASH 06
$53.6 \pm 3.1 \pm 5.1$	¹ GARMASH	04 BELL	Repl. by GARMASH 05
$59.1 \pm 3.8 \pm 3.2$	² AUBERT	03M BABR	Repl. by AUBERT,B 05N
$55.6 \pm 5.8 \pm 7.7$	³ GARMASH	02 BELL	Repl. by GARMASH 04
¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.			
² Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$; charm and charmonium contributions are subtracted, otherwise no assumptions about intermediate resonances.			
³ Uses a reference decay mode $B^+ \rightarrow \bar{D}^0\pi^+$ and $\bar{D}^0 \rightarrow K^+\pi^-$ with $B(B^+ \rightarrow \bar{D}^0\pi^+) \cdot B(\bar{D}^0 \rightarrow K^+\pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.			

$\Gamma(K^+\pi^-\pi^+ \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{239}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$16.3^{+2.1}_{-1.5}$ OUR AVERAGE				
$9.3 \pm 1.0^{+6.9}_{-1.7}$		^{1,2} AUBERT	08AI BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$16.9 \pm 1.3^{+1.7}_{-1.6}$		¹ GARMASH	06 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$2.9 \pm 0.6^{+0.8}_{-0.5}$		¹ AUBERT,B	05N BABR	Repl. by AUBERT 08AI
$17.3 \pm 1.7^{+17.2}_{-8.0}$		¹ GARMASH	05 BELL	Repl. by GARMASH 06
< 17	90	¹ AUBERT,B	04P BABR	Repl. by AUBERT,B 05N
<330	90	³ ADAM	96D DLPH	$e^+e^- \rightarrow Z$
< 28	90	BERGFELD	96B CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
<400	90	³ ABREU	95N DLPH	Sup. by ADAM 96D
<330	90	ALBRECHT	91E ARG	$e^+e^- \rightarrow \Upsilon(4S)$
<190	90	⁴ AVERY	89B CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.				
² Calculate the total nonresonant contribution by combining the S-wave composed of $K_0^*(1430)$ and nonresonant that are described using LASS shape.				
³ Assumes a B^0, B^- production fraction of 0.39 and a B_s^- production fraction of 0.12.				
⁴ AVERY 89B reports $< 1.7 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.				

$\Gamma(\omega(782)K^+)/\Gamma_{\text{total}}$ **Γ_{240}/Γ**

<u>VALUE (units 10^{-6})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.9^{+8.8+0.5}_{-9.0-0.4}	1,2 AUBERT	08AI BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² AUBERT 08AI reports $[\Gamma(B^+ \rightarrow \omega(782)K^+)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-)] = (0.09 \pm 0.13^{+0.036}_{-0.045}) \times 10^{-6}$. We divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-) = (1.53^{+0.11}_{-0.13}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ f_0(980) \times B(f_0(980) \rightarrow \pi^+\pi^-))/\Gamma_{\text{total}}$ **Γ_{241}/Γ**

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
9.4^{+1.0}_{-1.2}		OUR AVERAGE		

10.3 ± 0.5 ^{+2.0} _{-1.4}	1	AUBERT	08AI BABR	$e^+e^- \rightarrow \Upsilon(4S)$
--	---	--------	-----------	-----------------------------------

8.78 ± 0.82 ^{+0.85} _{-1.76}	1	GARMASH	06 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
---	---	---------	---------	-----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

9.47 ± 0.97 ^{+0.62} _{-0.88}	1	AUBERT,B	05N BABR	Repl. by AUBERT 08AI
---	---	----------	----------	----------------------

7.55 ± 1.24 ^{+1.63} _{-1.18}	1	GARMASH	05 BELL	Repl. by GARMASH 06
---	---	---------	---------	---------------------

9.2 ± 1.2 ^{+2.1} _{-2.6}	2	AUBERT,B	04P BABR	Repl. by AUBERT,B 05N
---	---	----------	----------	-----------------------

9.6 ^{+2.5} _{-2.3} ^{+3.7} _{-1.7}	3	GARMASH	02 BELL	Repl. by GARMASH 05
---	---	---------	---------	---------------------

<80	90	4 AVERY	89B CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
-----	----	---------	----------	-----------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² AUBERT,B 04P also reports $B(B^+ \rightarrow \text{"higher } f^0 \text{ resonances"} \pi^+, f(980)^0 \rightarrow \pi^+\pi^-) = (3.2 \pm 1.2^{+6.0}_{-2.9}) \times 10^{-6}$.

³ Uses a reference decay mode $B^+ \rightarrow \bar{D}^0 \pi^+$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ with $B(B^+ \rightarrow \bar{D}^0 \pi^+) \times B(\bar{D}^0 \rightarrow K^+ \pi^-) = (20.3 \pm 2.0) \times 10^{-5}$. Only charged pions from the $f_0(980)$ are used.

⁴ AVERY 89B reports $< 7 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(f_2(1270)^0 K^+)/\Gamma_{\text{total}}$ **Γ_{242}/Γ**

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.07 ± 0.27		OUR AVERAGE		

0.88 ^{+0.38} _{-0.33} ^{+0.01} _{-0.03}	1,2	AUBERT	08AI BABR	$e^+e^- \rightarrow \Upsilon(4S)$
--	-----	--------	-----------	-----------------------------------

1.33 ± 0.30 ^{+0.23} _{-0.34}	1	GARMASH	06 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
---	---	---------	---------	-----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<16	90	3 AUBERT,B	05N BABR	Repl. by AUBERT 08AI
-----	----	------------	----------	----------------------

< 2.3	90	4 GARMASH	05 BELL	Repl. by GARMASH 06
-------	----	-----------	---------	---------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² AUBERT 08AI reports $(0.50 \pm 0.15^{+0.15}_{-0.11}) \times 10^{-6}$ for $B(B^+ \rightarrow f_2(1270) K^+) \times B(f_2 \rightarrow \pi^+ \pi^-)$. We compute $B(B^+ \rightarrow f_2(1270) K^+)$ using the PDG value $B(f_2(1270) \rightarrow \pi\pi) = (84.8^{+2.4}_{-1.2}) \times 10^{-2}$ and 2/3 for the $\pi^+ \pi^-$ fraction. Our first error is their experiment's error and the second error is systematic error from using our best value.

³ AUBERT,B 05N reports 8.9×10^{-6} at 90% CL for $B(B^+ \rightarrow f_2(1270) K^+) \times B(f_2(1270) \rightarrow \pi^+ \pi^-)$. We rescaled it using the PDG value $B(f_2(1270) \rightarrow \pi\pi) = 84.7\%$ and 2/3 for the $\pi^+ \pi^-$ fraction.

⁴ GARMASH 05 reports 1.3×10^{-6} at 90% CL for $B(B^+ \rightarrow f_2(1270) K^+) \times B(f_2(1270) \rightarrow \pi^+ \pi^-)$. We rescaled it using the PDG value $B(f_2(1270) \rightarrow \pi\pi) = 84.7\%$ and 2/3 for the $\pi^+ \pi^-$ fraction.

$\Gamma(f_0(1370)^0 K^+ \times B(f_0(1370)^0 \rightarrow \pi^+ \pi^-)) / \Gamma_{\text{total}} \quad \Gamma_{243} / \Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 10.7 \times 10^{-6}$	90	¹ AUBERT,B 05N	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\rho^0(1450) K^+ \times B(\rho^0(1450) \rightarrow \pi^+ \pi^-)) / \Gamma_{\text{total}} \quad \Gamma_{244} / \Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 11.7 \times 10^{-6}$	90	¹ AUBERT,B 05N	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+ f_X(1300) \times B(f_X \rightarrow \pi^+ \pi^-)) / \Gamma_{\text{total}} \quad \Gamma_{245} / \Gamma$

$f_X(1300)$ is a possible spin zero state with $\pi^+ \pi^-$ invariant mass near $1.3 \text{ GeV}/c^2$.

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$0.73 \pm 0.21^{+0.47}_{-0.48}$	¹ AUBERT 08AI	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(f_0(1500) K^+ \times B(f_0(1500) \rightarrow \pi^+ \pi^-)) / \Gamma_{\text{total}} \quad \Gamma_{246} / \Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 4.4 \times 10^{-6}$	90	¹ AUBERT,B 05N	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(f'_2(1525) K^+ \times B(f'_2(1525) \rightarrow \pi^+ \pi^-)) / \Gamma_{\text{total}} \quad \Gamma_{247} / \Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.4 \times 10^{-6}$	90	¹ AUBERT,B 05N	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+ \rho^0) / \Gamma_{\text{total}} \quad \Gamma_{248} / \Gamma$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
3.7 ± 0.5 OUR AVERAGE				
$3.56 \pm 0.45^{+0.57}_{-0.46}$		¹ AUBERT 08AI	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$3.89 \pm 0.47^{+0.43}_{-0.41}$		¹ GARMASH 06	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.07 \pm 0.75^{+0.55}_{-0.88}$		1	AUBERT,B	05N	BABR	Repl. by AUBERT 08AI
$4.78 \pm 0.75^{+1.01}_{-0.97}$		1	GARMASH	05	BELL	Repl. by GARMASH 06
< 6.2	90	2	AUBERT,B	04P	BABR	Repl. by AUBERT,B 05N
< 12	90	3	GARMASH	02	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
< 86	90	4	ABE	00C	SLD	$e^+e^- \rightarrow Z$
< 17	90	1	JESSOP	00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
<120	90	5	ADAM	96D	DLPH	$e^+e^- \rightarrow Z$
< 19	90		ASNER	96	CLE2	Repl. by JESSOP 00
<190	90	5	ABREU	95N	DLPH	Sup. by ADAM 96D
<180	90		ALBRECHT	91B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
< 80	90	6	AVERY	89B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
<260	90		AVERY	87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² AUBERT 04P reports a central value of $(3.9 \pm 1.2^{+1.3}_{-3.5}) \times 10^{-6}$ for this branching ratio.

³ Uses a reference decay mode $B^+ \rightarrow \bar{D}^0 \pi^+$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ with $B(B^+ \rightarrow \bar{D}^0 \pi^+) \cdot B(\bar{D}^0 \rightarrow K^+ \pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.

⁴ ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

⁵ Assumes production fractions $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

⁶ AVERY 89B reports $< 7 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(K_0^*(1430)^0 \pi^+) / \Gamma_{\text{total}}$

Γ_{249} / Γ

<u>VALUE (units 10^{-6})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

45 $^{+9}_{-7}$ OUR AVERAGE Error includes scale factor of 1.5.

$32.0 \pm 1.2^{+10.8}_{-6.0}$	1	AUBERT	08AI	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
-------------------------------	---	--------	------	------	-----------------------------------

$51.6 \pm 1.7^{+7.0}_{-7.5}$	1	GARMASH	06	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
------------------------------	---	---------	----	------	-----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$44.4 \pm 2.2 \pm 5.3$	1,2	AUBERT,B	05N	BABR	Repl. by AUBERT 08AI
------------------------	-----	----------	-----	------	----------------------

$45.0 \pm 2.9^{+15.0}_{-10.7}$	1	GARMASH	05	BELL	Repl. by GARMASH 06
--------------------------------	---	---------	----	------	---------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² See erratum: AUBERT, BE 06A.

$\Gamma(K_2^*(1430)^0 \pi^+) / \Gamma_{\text{total}}$

Γ_{250} / Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

$5.6^{+2.2}_{-1.5} \pm 0.1$	1,2	AUBERT	08AI	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
---	-----	--------	------	------	-----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 23	90	3	AUBERT,B	05N	BABR	Repl. by AUBERT 08AI
------	----	---	----------	-----	------	----------------------

< 6.9	90	4	GARMASH	05	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
-------	----	---	---------	----	------	-----------------------------------

<680	90		ALBRECHT	91B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
------	----	--	----------	-----	-----	-----------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² AUBERT 08AI reports $(1.85 \pm 0.41_{-0.29}^{+0.61}) \times 10^{-6}$ for $B(B^+ \rightarrow K_2^*(1430)^0 \pi^+) \times B(K_2^*(1430)^0 \rightarrow K^+ \pi^-)$. We compute $B(B^+ \rightarrow K_2^*(1430)^0 \pi^+)$ using the PDG value $B(K_2^*(1430)^0 \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$ and 2/3 for the $K^+ \pi^-$ fraction. Our first error is their experiment's error and the second error is systematic error from using our best value.

³ AUBERT,B 05N reports 7.7×10^{-6} at 90% CL for $B(B^+ \rightarrow K_2^*(1430)^0 \pi^+) \times B(K_2^*(1430)^0 \rightarrow K^+ \pi^-)$. We rescaled it using the PDG value $B(K_2^*(1430)^0 \rightarrow K\pi) = 49.9\%$ and 2/3 for the $K^+ \pi^-$ fraction.

⁴ GARMASH 05 reports 2.3×10^{-6} at 90% CL for $B(B^+ \rightarrow K_2^*(1430)^0 \pi^+) \times B(K_2^*(1430)^0 \rightarrow K^+ \pi^-)$. We rescaled it using the PDG value $B(K_2^*(1430)^0 \rightarrow K\pi) = 49.9\%$ and 2/3 for the $K^+ \pi^-$ mode.

$\Gamma(K^*(1410)^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{251}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<45	90	¹ GARMASH 05	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ GARMASH 05 reports 2.0×10^{-6} at 90% CL for $B(B^+ \rightarrow K^*(1410)^0 \pi^+) \times B(K^*(1410)^0 \rightarrow K^+ \pi^-)$. We rescaled it using the PDG value $B(K^*(1410)^0 \rightarrow K\pi) = 6.6\%$ and 2/3 for the $K^+ \pi^-$ mode.

$\Gamma(K^*(1680)^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{252}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<12	90	¹ GARMASH 05	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<15	90	² AUBERT,B 05N	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
-----	----	---------------------------	------	------------------------------------

¹ GARMASH 05 reports 3.1×10^{-6} at 90% CL for $B(B^+ \rightarrow K^*(1680)^0 \pi^+) \times B(K^*(1680)^0 \rightarrow K^+ \pi^-)$. We rescaled it using the PDG value $B(K^*(1680)^0 \rightarrow K\pi) = 38.7\%$ and 2/3 for the $K^+ \pi^-$ mode.

² AUBERT,B 05N reports 3.8×10^{-6} at 90% CL for $B(B^+ \rightarrow K^*(1680)^0 \pi^+) \times B(K^*(1680)^0 \rightarrow K^+ \pi^-)$. We rescaled it using the PDG value $B(K^*(1680)^0 \rightarrow K\pi) = 38.7\%$ and 2/3 for the $K^+ \pi^-$ fraction.

$\Gamma(K^- \pi^+ \pi^+)/\Gamma_{\text{total}}$ Γ_{253}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<0.95	90	¹ AUBERT 08BE	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4.5	90	¹ GARMASH 04	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
<1.8	90	² AUBERT 03M	BABR	Repl. by AUBERT 08BE
<7.0	90	³ GARMASH 02	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$; charm and charmonium contributions are subtracted, otherwise no assumptions about intermediate resonances.

³ Uses a reference decay mode $B^+ \rightarrow \bar{D}^0 \pi^+$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ with $B(B^+ \rightarrow \bar{D}^0 \pi^+) \cdot B(\bar{D}^0 \rightarrow K^+ \pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.

$\Gamma(K^- \pi^+ \pi^+ \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{254}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<56	90	BERGFELD	96B	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(K_1(1400)^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{255}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.6 $\times 10^{-3}$	90	ALBRECHT	91B	ARG $e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^0 \pi^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{256}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<66 $\times 10^{-6}$	90	¹ ECKHART	02	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^0 \rho^+)/\Gamma_{\text{total}}$ Γ_{257}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$8.0^{+1.4}_{-1.3} \pm 0.6$		AUBERT	07Z	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<48	90	ASNER	96	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$
-----	----	-------	----	---

$\Gamma(K^*(892)^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{258}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$75.3 \pm 6.0 \pm 8.1$		¹ AUBERT,B	06U	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1100	90	ALBRECHT	91E	ARG $e^+ e^- \rightarrow \Upsilon(4S)$
-------	----	----------	-----	--

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)^+ \rho^0)/\Gamma_{\text{total}}$ Γ_{259}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 6.1	90	¹ AUBERT,B	06G	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$10.6^{+3.0}_{-2.6} \pm 2.4$		¹ AUBERT	03V	BABR Repl. by AUBERT,B 06G
< 74	90	² GODANG	02	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$
<900	90	ALBRECHT	91B	ARG $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Assumes a helicity 00 configuration. For a helicity 11 configuration, the limit decreases to 4.9×10^{-5} .

$\Gamma(K^*(892)^+ f_0(980))/\Gamma_{\text{total}}$ Γ_{260}/Γ

<u>VALUE (units 10^{-6})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$5.2 \pm 1.2 \pm 0.5$	¹ AUBERT,B	06G	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(a_1^+ K^0)/\Gamma_{\text{total}}$ Γ_{261}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
34.9±5.0±4.4	1,2 AUBERT	08F BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Assumes a_1^\pm decays only to 3π and $B(a_1^\pm \rightarrow \pi^\pm \pi^\mp \pi^\pm) = 0.5$.

$\Gamma(b_1^+ K^0 \times B(b_1^+ \rightarrow \omega \pi^+))/\Gamma_{\text{total}}$ Γ_{262}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
9.6±1.7±0.9	¹ AUBERT	08AG BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)^0 \rho^+)/\Gamma_{\text{total}}$ Γ_{263}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
9.2±1.5 OUR AVERAGE			
9.6±1.7±1.5	¹ AUBERT,B	06G BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
8.9±1.7±1.2	¹ ZHANG	05D BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K_1(1400)^+ \rho^0)/\Gamma_{\text{total}}$ Γ_{264}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<7.8 × 10⁻⁴	90	ALBRECHT	91B ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(K_2^*(1430)^+ \rho^0)/\Gamma_{\text{total}}$ Γ_{265}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.5 × 10⁻³	90	ALBRECHT	91B ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(b_1^0 K^+ \times B(b_1^0 \rightarrow \omega \pi^0))/\Gamma_{\text{total}}$ Γ_{266}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
9.1±1.7±1.0	¹ AUBERT	07BI BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+ \bar{K}^0)/\Gamma_{\text{total}}$ Γ_{267}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
1.36±0.27 OUR AVERAGE				
1.22 ^{+0.32+0.13} _{-0.28-0.16}		¹ LIN	07 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
1.61±0.44±0.09		¹ AUBERT,BE	06C BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1.0 ±0.4 ±0.1		¹ ABE	05G BELL	Repl. by LIN 07
1.5 ±0.5 ±0.1		¹ AUBERT,BE	05E BABR	Repl. by AUBERT,BE 06C
< 2.5	90	¹ AUBERT	04M BABR	Repl. by AUBERT,BE 05E
< 3.3	90	¹ CHAO	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
< 3.3	90	¹ BORNHEIM	03 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
< 2.0	90	¹ CASEY	02 BELL	Repl. by CHAO 04
< 5.0	90	¹ ABE	01H BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
< 2.4	90	¹ AUBERT	01E BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
< 5.1	90	¹ CRONIN-HEN..00	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
<21	90	GODANG	98 CLE2	Repl. by CRONIN-HENNESSY 00

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{K}^0 K^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{268}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<24 \times 10^{-6}$	90	¹ ECKHART	02	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+ K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{269}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
11.5 ± 1.3 OUR AVERAGE			
10.7 ± 1.2 ± 1.0	¹ AUBERT,B	04V	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
13.4 ± 1.9 ± 1.5	¹ GARMASH	04	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K_S^0 K_S^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{270}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<0.51	90	¹ AUBERT	09J	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<3.2	90	¹ GARMASH	04	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+ K^- \pi^+)/\Gamma_{\text{total}}$ Γ_{271}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
5.0 ± 0.5 ± 0.5		¹ AUBERT	07BB	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<13	90	¹ GARMASH	04	BELL $e^+ e^- \rightarrow \Upsilon(4S)$
< 6.3	90	^{1,2} AUBERT	03M	BABR Repl. by AUBERT 07BB
<12	90	³ GARMASH	02	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Charm and charmonium contributions are subtracted, otherwise no assumptions about intermediate resonances.

³ Uses a reference decay mode $B^+ \rightarrow \overline{D}^0 \pi^+$ and $\overline{D}^0 \rightarrow K^+ \pi^-$ with $B(B^+ \rightarrow \overline{D}^0 \pi^+) \cdot B(\overline{D}^0 \rightarrow K^+ \pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.

$\Gamma(K^+ K^- \pi^+ \text{nonresonant})/\Gamma_{\text{total}}$ Γ_{272}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<75	90	BERGFELD	96B	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^+ \overline{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{273}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.1	90	¹ AUBERT	07AR	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<129	90	ABBIENDI	00B	OPAL $e^+ e^- \rightarrow Z$
<138	90	² ABE	00C	SLD $e^+ e^- \rightarrow Z$
< 5.3	90	¹ JESSOP	00	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ABE 00C assumes $B(Z \rightarrow b\overline{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

$\Gamma(K^+ \bar{K}_0^*(1430)^0)/\Gamma_{\text{total}}$ Γ_{274}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<2.2	90	¹ AUBERT	07AR BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+ K^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{275}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.6 $\times 10^{-7}$	90	¹ AUBERT	08BE BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.4 $\times 10^{-6}$	90	¹ GARMASH	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
<1.3 $\times 10^{-6}$	90	² AUBERT	03M BABR	Repl. by AUBERT 08BE
<3.2 $\times 10^{-6}$	90	³ GARMASH	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$; charm and charmonium contributions are subtracted, otherwise no assumptions about intermediate resonances.

³ Uses a reference decay mode $B^+ \rightarrow \bar{D}^0 \pi^+$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ with $B(B^+ \rightarrow \bar{D}^0 \pi^+) \cdot B(\bar{D}^0 \rightarrow K^+ \pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.

$\Gamma(K^+ K^+ \pi^- \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{276}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<87.9	90	ABBIENDI	00B OPAL	$e^+ e^- \rightarrow Z$

$\Gamma(K^+ f_J(2220))/\Gamma_{\text{total}}$ Γ_{277}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
not seen		¹ HUANG	03 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ No evidence is found for such decay and set a limit on $B(B^+ \rightarrow f_J(2220)) \times B(f_J(2220) \rightarrow \phi\phi) < 1.2 \times 10^{-6}$ at 90%CL where the $f_J(2220)$ is a possible glueball state.

$\Gamma(K^{*+} \pi^+ K^-)/\Gamma_{\text{total}}$ Γ_{278}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<11.8	90	¹ AUBERT,B	06U BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)^+ K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{279}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
1.2 \pm 0.5 \pm 0.1		AUBERT	09F BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<71	90	¹ GODANG	02 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
-----	----	---------------------	---------	------------------------------------

¹ Assumes a helicity 00 configuration. For a helicity 11 configuration, the limit decreases to 4.8×10^{-5} .

$\Gamma(K^{*+} K^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{280}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<6.1	90	¹ AUBERT,B	06U BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+ K^- K^+)/\Gamma_{\text{total}}$ Γ_{281}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
33.7 ± 2.2 OUR AVERAGE Error includes scale factor of 1.4.				
35.2 ± 0.9 ± 1.6		¹ AUBERT	06O BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
30.6 ± 1.2 ± 2.3		¹ GARMASH	05 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
32.8 ± 1.8 ± 2.8		¹ GARMASH	04 BELL	Repl. by GARMASH 05
29.6 ± 2.1 ± 1.6		² AUBERT	03M BABR	Repl. by AUBERT 06O
35.3 ± 3.7 ± 4.5		³ GARMASH	02 BELL	Repl. by GARMASH 04
<200	90	⁴ ADAM	96D DLPH	$e^+ e^- \rightarrow Z$
<320	90	⁴ ABREU	95N DLPH	Sup. by ADAM 96D
<350	90	ALBRECHT	91E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

- ¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
- ² Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$; charm and charmonium contributions are subtracted, otherwise no assumptions about intermediate resonances.
- ³ Uses a reference decay mode $B^+ \rightarrow \bar{D}^0 \pi^+$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ with $B(B^+ \rightarrow \bar{D}^0 \pi^+) \cdot B(\bar{D}^0 \rightarrow K^+ \pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.
- ⁴ Assumes B^0 and B^- production fractions of 0.39, and B_s production fraction of 0.12.

$\Gamma(K^+ \phi)/\Gamma_{\text{total}}$ Γ_{282}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
8.3 ± 0.7 OUR AVERAGE				
8.4 ± 0.7 ± 0.7		¹ AUBERT	06O BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
7.6 ± 1.3 ± 0.6		² ACOSTA	05J CDF	$p\bar{p}$ at 1.96 TeV
9.60 ± 0.92 ^{+1.05} _{-0.85}		¹ GARMASH	05 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
5.5 ^{+2.1} _{-1.8} ± 0.6		¹ BRIERE	01 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
10.0 ^{+0.9} _{-0.8} ± 0.5		¹ AUBERT	04A BABR	Repl. by AUBERT 06O
9.4 ± 1.1 ± 0.7		¹ CHEN	03B BELL	Repl. by GARMASH 05
14.6 ^{+3.0} _{-2.8} ± 2.0		³ GARMASH	02 BELL	Repl. by CHEN 03B
7.7 ^{+1.6} _{-1.4} ± 0.8		¹ AUBERT	01D BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
<144	90	⁴ ABE	00C SLD	$e^+ e^- \rightarrow Z$
< 5	90	¹ BERGFELD	98 CLE2	
<280	90	⁵ ADAM	96D DLPH	$e^+ e^- \rightarrow Z$
< 12	90	ASNER	96 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
<440	90	⁶ ABREU	95N DLPH	Sup. by ADAM 96D
<180	90	ALBRECHT	91B ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
< 90	90	⁷ AVERY	89B CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
<210	90	AVERY	87 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

- ¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
- ² Uses $B(B^+ \rightarrow J/\psi K^+) = (1.00 \pm 0.04) \times 10^{-3}$ and $B(J/\psi \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$.
- ³ Uses a reference decay mode $B^+ \rightarrow \bar{D}^0 \pi^+$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ with $B(B^+ \rightarrow \bar{D}^0 \pi^+) \cdot B(\bar{D}^0 \rightarrow K^+ \pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.

⁴ ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

⁵ ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

⁶ Assumes a B^0 , B^- production fraction of 0.39 and a B_s production fraction of 0.12.

⁷ AVERY 89B reports $< 8 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(f_0(980)K^+ \times B(f_0(980) \rightarrow K^+K^-))/\Gamma_{\text{total}}$ Γ_{283}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<2.9	90	¹ GARMASH 05	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$6.5 \pm 2.5 \pm 1.6$ ¹ AUBERT 060 BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(a_2(1320)K^+ \times B(a_2(1320) \rightarrow K^+K^-))/\Gamma_{\text{total}}$ Γ_{284}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.1 $\times 10^{-6}$	90	¹ GARMASH 05	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(f'_2(1525)K^+ \times B(f'_2(1525) \rightarrow K^+K^-))/\Gamma_{\text{total}}$ Γ_{285}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<4.9 $\times 10^{-6}$	90	¹ GARMASH 05	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(X_0(1550)K^+ \times B(X_0(1550) \rightarrow K^+K^-))/\Gamma_{\text{total}}$ Γ_{286}/Γ

$X_0(1550)$ is a possible spin zero state near 1.55 GeV/ c^2 invariant mass of K^+K^- .

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
4.3 $\pm 0.6 \pm 0.3$	¹ AUBERT 060 BABR	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\phi(1680)K^+ \times B(\phi(1680) \rightarrow K^+K^-))/\Gamma_{\text{total}}$ Γ_{287}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.8 $\times 10^{-6}$	90	¹ GARMASH 05	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(f_0(1710)K^+ \times B(f_0(1710) \rightarrow K^+K^-))/\Gamma_{\text{total}}$ Γ_{288}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
1.7 $\pm 1.0 \pm 0.3$	¹ AUBERT 060 BABR	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+ K^- K^+ \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{289}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

28 \pm 9 \pm 16 OUR AVERAGE Error includes scale factor of 3.3.

50.0 \pm 6.0 \pm 4.0 ¹ AUBERT 06O BABR $e^+ e^- \rightarrow \gamma(4S)$

24.0 \pm 1.5 \pm 2.6 \pm 6.0 ¹ GARMASH 05 BELL $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<38 90 BERGFELD 96B CLE2 $e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K^*(892)^+ K^+ K^-)/\Gamma_{\text{total}}$ Γ_{290}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

36.2 \pm 3.3 \pm 3.6 ¹ AUBERT,B 06U BABR $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1600 90 ALBRECHT 91E ARG $e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K^*(892)^+ \phi)/\Gamma_{\text{total}}$ Γ_{291}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

10.5 \pm 1.5 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.

11.2 \pm 1.0 \pm 0.9 ¹ AUBERT 07BA BABR $e^+ e^- \rightarrow \gamma(4S)$

12.7 \pm 2.2 \pm 1.1 \pm 2.0 ¹ AUBERT 03V BABR $e^+ e^- \rightarrow \gamma(4S)$

6.7 \pm 2.1 \pm 0.7 \pm 1.9 \pm 1.0 ¹ CHEN 03B BELL $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

9.7 \pm 4.2 \pm 1.7 \pm 3.4 ¹ AUBERT 01D BABR Repl. by AUBERT 03v

< 22.5 90 ¹ BRIERE 01 CLE2 $e^+ e^- \rightarrow \gamma(4S)$


< 41 90 ¹ BERGFELD 98 CLE2

< 70 90 ASNER 96 CLE2 $e^+ e^- \rightarrow \gamma(4S)$

<1300 90 ALBRECHT 91B ARG $e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

WEIGHTED AVERAGE
 10.5 ± 1.5 (Error scaled by 1.4)



$\Gamma(\phi(K\pi)_0^{*+})/\Gamma_{\text{total}}$

Γ_{292}/Γ

$(K\pi)_0^{*+}$ is the total S-wave composed of $K_0^*(1430)$ and nonresonant that are described using LASS shape.

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$8.3 \pm 1.4 \pm 0.8$	¹ AUBERT	08BI BABR	$e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\phi K_1(1270)^+)/\Gamma_{\text{total}}$

Γ_{293}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$6.1 \pm 1.6 \pm 1.1$	¹ AUBERT	08BI BABR	$e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\phi K_1(1400)^+)/\Gamma_{\text{total}}$

Γ_{294}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.2	90	¹ AUBERT	08BI BABR	$e^+e^- \rightarrow \gamma(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<1100	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\phi K^*(1410)^+)/\Gamma_{\text{total}}$

Γ_{295}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<4.3	90	¹ AUBERT	08BI BABR	$e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\phi K_0^*(1430)^+)/\Gamma_{\text{total}}$ Γ_{296}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$7.0 \pm 1.3 \pm 0.9$		¹ AUBERT	08BI BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\phi K_2^*(1430)^+)/\Gamma_{\text{total}}$ Γ_{297}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$8.4 \pm 1.8 \pm 1.0$		¹ AUBERT	08BI BABR	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3400	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow \Upsilon(4S)$
-------	----	----------	---------	-----------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\phi K_2^*(1770)^+)/\Gamma_{\text{total}}$ Γ_{298}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<15.0	90	¹ AUBERT	08BI BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\phi K_2^*(1820)^+)/\Gamma_{\text{total}}$ Γ_{299}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<16.3	90	¹ AUBERT	08BI BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+\phi\phi)/\Gamma_{\text{total}}$ Γ_{300}/Γ

<u>VALUE (units 10^{-6})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$4.9^{+2.4}_{-2.2}$ OUR AVERAGE			Error includes scale factor of 2.9.

$7.5 \pm 1.0 \pm 0.7$	¹ AUBERT, BE	06H BABR	$e^+e^- \rightarrow \Upsilon(4S)$
-----------------------	-------------------------	----------	-----------------------------------

$2.6^{+1.1}_{-0.9} \pm 0.3$	¹ HUANG	03 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
-----------------------------	--------------------	---------	-----------------------------------

¹ Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$ and for a $\phi\phi$ invariant mass below $2.85 \text{ GeV}/c^2$.

$\Gamma(\eta'\eta'K^+)/\Gamma_{\text{total}}$ Γ_{301}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<25	90	¹ AUBERT, B	06P BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)^+\gamma)/\Gamma_{\text{total}}$ Γ_{302}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.03 ± 0.26 OUR AVERAGE				
$3.87 \pm 0.28 \pm 0.26$		¹ AUBERT, BE	04A BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$4.25 \pm 0.31 \pm 0.24$		² NAKAO	04 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$3.76^{+0.89}_{-0.83} \pm 0.28$		² COAN	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.83 \pm 0.62 \pm 0.22$		² AUBERT	02C	BABR	Repl. by AUBERT, BE 04A
$5.7 \pm 3.1 \pm 1.1$		³ AMMAR	93	CLE2	Repl. by COAN 00
< 55	90	⁴ ALBRECHT	89G	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
< 55	90	⁴ AVERY	89B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
< 180	90	AVERY	87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Uses the production ratio of charged and neutral B from $\Upsilon(4S)$ decays $R^{+/0} = 1.006 \pm 0.048$.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ AMMAR 93 observed 4.1 ± 2.3 events above background.

⁴ Assumes the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$.

$\Gamma(K_1(1270)^+\gamma)/\Gamma_{\text{total}}$ Γ_{303}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
$4.3 \pm 0.9 \pm 0.9$		¹ YANG	05	BELL $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 9.9	90	¹ NISHIDA	02	BELL	Repl. by YANG 05
< 730	90	² ALBRECHT	89G	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ALBRECHT 89G reports < 0.0066 assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\eta K^+\gamma)/\Gamma_{\text{total}}$ Γ_{304}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
7.9 ± 0.9 OUR AVERAGE			
$7.7 \pm 1.0 \pm 0.4$	^{1,2} AUBERT	09	BABR $e^+e^- \rightarrow \Upsilon(4S)$
$8.4 \pm 1.5^{+1.2}_{-0.9}$	^{2,3} NISHIDA	05	BELL $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$10.0 \pm 1.3 \pm 0.5$	^{1,2} AUBERT, B	06M	BABR	Repl. by AUBERT 09
------------------------	--------------------------	-----	------	--------------------

¹ $m_{\eta K} < 3.25 \text{ GeV}/c^2$.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ $m_{\eta K} < 2.4 \text{ GeV}/c^2$

$\Gamma(\eta' K^+\gamma)/\Gamma_{\text{total}}$ Γ_{305}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.2	90	^{1,2} AUBERT, B	06M	BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ $m_{\eta' K} < 3.25 \text{ GeV}/c^2$.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\phi K^+\gamma)/\Gamma_{\text{total}}$ Γ_{306}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
3.5 ± 0.6 OUR AVERAGE			
$3.5 \pm 0.6 \pm 0.4$	¹ AUBERT	07Q	BABR $e^+e^- \rightarrow \Upsilon(4S)$
$3.4 \pm 0.9 \pm 0.4$	¹ DRUTSKOY	04	BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at $\Upsilon(4S)$.

$\Gamma(K^+ \pi^- \pi^+ \gamma)/\Gamma_{\text{total}}$ Γ_{307}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.76 ± 0.22 OUR AVERAGE	Error includes scale factor of 1.2.		
$2.95 \pm 0.13 \pm 0.20$	1,2 AUBERT	07R BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$2.50 \pm 0.18 \pm 0.22$	2,3 YANG	05 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$2.4 \pm 0.5 \begin{smallmatrix} +0.4 \\ -0.2 \end{smallmatrix}$	2,4 NISHIDA	02 BELL	Repl. by YANG 05

¹ $M_{K \pi \pi} < 1.8 \text{ GeV}/c^2$.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ $M_{K \pi \pi} < 2.0 \text{ GeV}/c^2$.

⁴ $M_{K \pi \pi} < 2.4 \text{ GeV}/c^2$.

$\Gamma(K^*(892)^0 \pi^+ \gamma)/\Gamma_{\text{total}}$ Γ_{308}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$(2.0^{+0.7}_{-0.6} \pm 0.2) \times 10^{-5}$	1,2 NISHIDA	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² $M_{K \pi \pi} < 2.4 \text{ GeV}/c^2$.

$\Gamma(K^+ \rho^0 \gamma)/\Gamma_{\text{total}}$ Γ_{309}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 2.0 \times 10^{-5}$	90	1,2 NISHIDA	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² $M_{K \pi \pi} < 2.4 \text{ GeV}/c^2$.

$\Gamma(K^+ \pi^- \pi^+ \gamma \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{310}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 9.2 \times 10^{-6}$	90	1,2 NISHIDA	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² $M_{K \pi \pi} < 2.4 \text{ GeV}/c^2$.

$\Gamma(K^0 \pi^+ \pi^0 \gamma)/\Gamma_{\text{total}}$ Γ_{311}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$4.56 \pm 0.42 \pm 0.31$	1,2 AUBERT	07R BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ $M_{K \pi \pi} < 1.8 \text{ GeV}/c^2$.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K_1(1400)^+ \gamma)/\Gamma_{\text{total}}$ Γ_{312}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 1.5	90	1 YANG	05 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 5.0	90	1 NISHIDA	02 BELL	Repl. by YANG 05
< 220	90	2 ALBRECHT	89G ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ALBRECHT 89G reports < 0.0020 assuming the $\Upsilon(4S)$ decays 45% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(K_2^*(1430)^+\gamma)/\Gamma_{\text{total}}$ Γ_{313}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

$1.45 \pm 0.40 \pm 0.15$ ¹ AUBERT,B 04U BABR $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<140 90 ² ALBRECHT 89G ARG $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ALBRECHT 89G reports < 0.0013 assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K^*(1680)^+\gamma)/\Gamma_{\text{total}}$ Γ_{314}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.0019 90 ¹ ALBRECHT 89G ARG $e^+e^- \rightarrow \Upsilon(4S)$

¹ ALBRECHT 89G reports < 0.0017 assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K_3^*(1780)^+\gamma)/\Gamma_{\text{total}}$ Γ_{315}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 39 90 ^{1,2} NISHIDA 05 BELL $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<5500 90 ³ ALBRECHT 89G ARG $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Uses $B(K_3^*(1780) \rightarrow \eta K) = 0.11^{+0.05}_{-0.04}$.

³ ALBRECHT 89G reports < 0.005 assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K_4^*(2045)^+\gamma)/\Gamma_{\text{total}}$ Γ_{316}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.0099 90 ¹ ALBRECHT 89G ARG $e^+e^- \rightarrow \Upsilon(4S)$

¹ ALBRECHT 89G reports < 0.0090 assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\rho^+\gamma)/\Gamma_{\text{total}}$ Γ_{317}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

0.98 ± 0.25 OUR AVERAGE

$1.20^{+0.42}_{-0.37} \pm 0.20$ ¹ AUBERT 08BH BABR $e^+e^- \rightarrow \Upsilon(4S)$

$0.87^{+0.29+0.09}_{-0.27-0.11}$ ¹ TANIGUCHI 08 BELL $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.10^{+0.37}_{-0.33} \pm 0.09$ ¹ AUBERT 07L BABR Repl. by AUBERT 08BH

$0.55^{+0.42+0.09}_{-0.36-0.08}$ ¹ MOHAPATRA 06 BELL Repl. by TANIGUCHI 08

$0.9^{+0.6}_{-0.5} \pm 0.1$ 90 ¹ AUBERT 05 BABR Repl. by AUBERT 07L

< 2.2 90 ¹ MOHAPATRA 05 BELL $e^+e^- \rightarrow \Upsilon(4S)$

< 2.1 90 ¹ AUBERT 04C BABR $e^+e^- \rightarrow \Upsilon(4S)$

<13 90 ^{1,2} COAN 00 CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at $\Upsilon(4S)$.

² No evidence for a nonresonant $K\pi\gamma$ contamination was seen; the central value assumes no contamination.

$\Gamma(\pi^+\pi^0)/\Gamma_{\text{total}}$

Γ_{318}/Γ

VALUE (units 10^{-6}) CL% DOCUMENT ID TECN COMMENT


5.7 ± 0.5 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.

5.02 ± 0.46 ± 0.29		¹ AUBERT	07BC	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
6.5 ± 0.4 ± 0.4		¹ LIN	07A	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
4.6 ^{+1.8} _{-1.6} ^{+0.6} _{-0.7}		¹ BORNHEIM	03	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
5.8 ± 0.6 ± 0.4		¹ AUBERT	05L	BABR	Repl. by AUBERT 07BC
5.0 ± 1.2 ± 0.5		¹ CHAO	04	BELL	Repl. by LIN 07A
5.5 ^{+1.0} _{-1.9} ± 0.6		¹ AUBERT	03L	BABR	Repl. by AUBERT 05L
7.4 ^{+2.3} _{-2.2} ± 0.9		¹ CASEY	02	BELL	Repl. by CHAO 04
< 13.4	90	¹ ABE	01H	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
< 9.6	90	¹ AUBERT	01E	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
< 12.7	90	¹ CRONIN-HEN..00		CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
< 20	90	GODANG	98	CLE2	Repl. by CRONIN-HENNESSY 00
< 17	90	ASNER	96	CLE2	Repl. by GODANG 98
< 240	90	¹ ALBRECHT	90B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
< 2300	90	² BEBEK	87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² BEBEK 87 assume the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$.

WEIGHTED AVERAGE
5.7±0.5 (Error scaled by 1.4)



$\Gamma(\pi^+\pi^0)/\Gamma(K^0\pi^+)$		$\Gamma_{318}/\Gamma_{213}$		
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.285±0.02 ±0.02		LIN	07A BELL	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\pi^+\pi^+\pi^-)/\Gamma_{total}$		Γ_{319}/Γ		
<u>VALUE (units 10⁻⁶)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
16.2±1.2±0.9		¹ AUBERT,B	05G BABR	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

10.9±3.3±1.6		¹ AUBERT	03M BABR	Repl. by AUBERT 05G
<130	90	² ADAM	96D DLPH	$e^+e^- \rightarrow Z$
<220	90	³ ABREU	95N DLPH	Sup. by ADAM 96D
<450	90	⁴ ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$
<190	90	⁵ BORTOLETTO89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$; charm and charmonium contributions are subtracted, otherwise no assumptions about intermediate resonances.

² ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

³ Assumes a B^0 , B^- production fraction of 0.39 and a B_s production fraction of 0.12.

⁴ ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.

⁵ BORTOLETTO 89 reports $< 1.7 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\rho^0\pi^+)/\Gamma_{total}$		Γ_{320}/Γ		
<u>VALUE (units 10⁻⁶)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
8.7±1.1 OUR AVERAGE				

8.8±1.0 ^{+0.6} _{-0.9}		¹ AUBERT,B	05G BABR	$e^+e^- \rightarrow \Upsilon(4S)$
8.0 ^{+2.3} _{-2.0} ±0.7		¹ GORDON	02 BELL	$e^+e^- \rightarrow \Upsilon(rS)$
10.4 ^{+3.3} _{-3.4} ±2.1		¹ JESSOP	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

9.5±1.1±0.9		¹ AUBERT	04Z BABR	Repl. by AUBERT 05G
< 83	90	² ABE	00C SLD	$e^+e^- \rightarrow Z$
<160	90	³ ADAM	96D DLPH	$e^+e^- \rightarrow Z$
< 43	90	ASNER	96 CLE2	Repl. by JESSOP 00
<260	90	⁴ ABREU	95N DLPH	Sup. by ADAM 96D
<150	90	¹ ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$
<170	90	⁵ BORTOLETTO89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
<230	90	⁵ BEBEK	87 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
<600	90	GILES	84 CLEO	Repl. by BEBEK 87

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

³ ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

⁴ Assumes a B^0 , B^- production fraction of 0.39 and a B_s production fraction of 0.12.

⁵ Papers assume the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

$[\Gamma(K^*(892)^0 \pi^+) + \Gamma(\rho^0 \pi^+)]/\Gamma_{\text{total}}$ $(\Gamma_{236} + \Gamma_{320})/\Gamma$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$170_{-80}^{+120} \pm 20$		¹ ADAM	96D DLPH	$e^+ e^- \rightarrow Z$

¹ ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

$\Gamma(\pi^+ f_0(980) \times B(f_0(980) \rightarrow \pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{321}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.0	90	¹ AUBERT,B	05G BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
< 140	90	² BORTOLETTO89	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² BORTOLETTO 89 reports $< 1.2 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(\pi^+ f_2(1270))/\Gamma_{\text{total}}$ Γ_{322}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$8.2 \pm 2.1 \pm 1.4$		^{1,2} AUBERT,B	05G BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
< 240	90	³ BORTOLETTO89	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Reported $B(B^+ \rightarrow f_2(1270) \pi^+) \times B(f_2(1270) \rightarrow \pi^+ \pi^-) = (2.3 \pm 0.6 \pm 0.4) \times 10^{-6}$ and rescaled using $B(f_2(1270) \rightarrow \pi^+ \pi^-) = 0.28$.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ BORTOLETTO 89 reports $< 2.1 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(\rho(1450)^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{323}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.3	90	¹ AUBERT,B	05G BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(f_0(1370) \pi^+ \times B(f_0(1370) \rightarrow \pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{324}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.0	90	¹ AUBERT,B	05G BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(f_0(600) \pi^+ \times B(f_0(600) \rightarrow \pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{325}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.1	90	¹ AUBERT,B	05G BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\pi^+ \pi^- \pi^+ \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{326}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.6	90	¹ AUBERT,B	05G BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
< 41	90	BERGFELD	96B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\pi^+\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{327}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<8.9 × 10⁻⁴	90	¹ ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$

¹ ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(\rho^+\pi^0)/\Gamma_{\text{total}}$ Γ_{328}/Γ

<u>VALUE (units 10⁻⁶)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
10.9 ± 1.4 OUR AVERAGE				
10.2 ± 1.4 ± 0.9		¹ AUBERT	07X BABR	$e^+e^- \rightarrow \Upsilon(4S)$
13.2 ± 2.3 ^{+1.4} _{-1.9}		¹ ZHANG	05A BELL	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

10.9 ± 1.9 ± 1.9		¹ AUBERT	04Z BABR	Repl. by AUBERT 07X
< 43	90	^{1,2} JESSOP	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
< 77	90	ASNER	96 CLE2	Repl. by JESSOP 00
< 550	90	¹ ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Assumes no nonresonant contributions of $B^+ \rightarrow \pi^+\pi^0\pi^0$.

$\Gamma(\pi^+\pi^-\pi^+\pi^0)/\Gamma_{\text{total}}$ Γ_{329}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4.0 × 10⁻³	90	¹ ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$

¹ ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(\rho^+\rho^0)/\Gamma_{\text{total}}$ Γ_{330}/Γ

<u>VALUE (units 10⁻⁶)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
24.0 ± 1.9 OUR AVERAGE				
23.7 ± 1.4 ± 1.4		¹ AUBERT	09G BABR	$e^+e^- \rightarrow \Upsilon(4S)$
31.7 ± 7.1 ^{+3.8} _{-6.7}		^{1,2} ZHANG	03B BELL	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

16.8 ± 2.2 ± 2.3		¹ AUBERT, BE	06G BABR	Repl. by AUBERT 09G
22.5 ^{+5.7} _{-5.4} ± 5.8		¹ AUBERT	03V BABR	Repl. by AUBERT, BE 06G
< 1000	90	¹ ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² The systematic error includes the error associated with the helicity-mix uncertainty.

$\Gamma(\rho^+f_0(980) \times B(f_0(980) \rightarrow \pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{331}/Γ

<u>VALUE (units 10⁻⁶)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.0	90	¹ AUBERT	09G BABR	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.9	90	¹ AUBERT, BE	06G BABR	Repl. by AUBERT 09G
------	----	-------------------------	----------	---------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(a_1(1260)^+\pi^0)/\Gamma_{\text{total}}$ Γ_{332}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

26.4±5.4±4.1 1,2 AUBERT 07BL BABR $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1700 90 1 ALBRECHT 90B ARG $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Assumes a_1^+ decays only to 3π and $B(a_1^+ \rightarrow \pi^\pm \pi^\mp \pi^+) = 0.5$.

$\Gamma(a_1(1260)^0\pi^+)/\Gamma_{\text{total}}$ Γ_{333}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

20.4±4.7±3.4 1,2 AUBERT 07BL BABR $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<900 90 1 ALBRECHT 90B ARG $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Assumes a_1^0 decays only to 3π and $B(a_1^+ \rightarrow \pi^\pm \pi^\mp \pi^0) = 1.0$.

$\Gamma(\omega\pi^+)/\Gamma_{\text{total}}$ Γ_{334}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

6.9±0.5 OUR AVERAGE

6.7±0.5±0.4 1 AUBERT 07AE BABR $e^+e^- \rightarrow \Upsilon(4S)$

6.9±0.6±0.5 1 JEN 06 BELL $e^+e^- \rightarrow \Upsilon(4S)$

11.3^{+3.3}_{-2.9}±1.4 1 JESSOP 00 CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.1±0.7±0.4 1 AUBERT,B 06E BABR Repl. by AUBERT 07AE

5.5±0.9±0.5 1 AUBERT 04H BABR Repl. by AUBERT,B 06E

5.7^{+1.4}_{-1.3}±0.6 1 WANG 04A BELL Repl. by JEN 06

4.2^{+2.0}_{-1.8}±0.5 1 LU 02 BELL Repl. by WANG 04A

6.6^{+2.1}_{-1.8}±0.7 1 AUBERT 01G BABR Repl. by AUBERT 04H

< 23 90 1 BERGFELD 98 CLE2 Repl. by JESSOP 00

<400 90 1 ALBRECHT 90B ARG $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\omega\rho^+)/\Gamma_{\text{total}}$ Γ_{335}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

15.9±1.6±1.4 1 AUBERT 09H BABR $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

10.6±2.1^{+1.6}_{-1.0} 1 AUBERT,B 06T BABR Repl. by AUBERT 09H

12.6^{+3.7}_{-3.3}±1.6 1 AUBERT 05O BABR Repl. by AUBERT,B 06T

<61 90 1 BERGFELD 98 CLE2

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta\pi^+)/\Gamma_{\text{total}}$ Γ_{336}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.4 ± 0.4 OUR AVERAGE		Error includes scale factor of 1.1.		
5.0 ± 0.5 ± 0.3		¹ AUBERT	07AE BABR	$e^+e^- \rightarrow \Upsilon(4S)$
4.2 ± 0.4 ± 0.2		¹ CHANG	07B BELL	$e^+e^- \rightarrow \Upsilon(4S)$
1.2 ^{+2.8} _{-1.2}		¹ RICHICHI	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.1 ± 0.6 ± 0.3		¹ AUBERT,B	05K BABR	Repl. by AUBERT 07AE
4.8 ± 0.7 ± 0.3		¹ CHANG	05A BELL	Repl. by CHANG 07B
5.3 ± 1.0 ± 0.3		¹ AUBERT	04H BABR	Repl. by AUBERT,B 05K
< 15	90	BEHRENS	98 CLE2	Repl. by RICHICHI 00
< 700	90	¹ ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta\rho^+)/\Gamma_{\text{total}}$ Γ_{337}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
7.0 ± 2.9 OUR AVERAGE		Error includes scale factor of 2.8.		
9.9 ± 1.2 ± 0.8		¹ AUBERT	08AH BABR	$e^+e^- \rightarrow \Upsilon(4S)$
4.1 ^{+1.4} _{-1.3} ± 0.4		¹ WANG	07B BELL	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.4 ± 1.9 ± 1.1		¹ AUBERT,B	05K BABR	Repl. by AUBERT 08AH
< 14	90	¹ AUBERT,B	04D BABR	Repl. by AUBERT,B 05K
< 15	90	¹ RICHICHI	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
< 32	90	BEHRENS	98 CLE2	Repl. by RICHICHI 00

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta'\pi^+)/\Gamma_{\text{total}}$ Γ_{338}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.7 ± 1.0 OUR AVERAGE		Error includes scale factor of 2.1.		
3.9 ± 0.7 ± 0.3		¹ AUBERT	07AE BABR	$e^+e^- \rightarrow \Upsilon(4S)$
1.76 ^{+0.67+0.15} _{-0.62-0.14}		¹ SCHUEMANN	06 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.0 ± 0.8 ± 0.4		¹ AUBERT,B	05K BABR	Repl. by AUBERT 07AE
< 4.5	90	¹ AUBERT	04H BABR	Repl. by AUBERT,B 05K
< 7.0	90	¹ ABE	01M BELL	$e^+e^- \rightarrow \Upsilon(4S)$
< 12	90	¹ AUBERT	01G BABR	$e^+e^- \rightarrow \Upsilon(4S)$
< 12	90	¹ RICHICHI	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
< 31	90	BEHRENS	98 CLE2	Repl. by RICHICHI 00

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta' \rho^+)/\Gamma_{\text{total}}$ Γ_{339}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$8.7^{+3.1+2.3}_{-2.8-1.3}$		¹ AUBERT	07E BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 5.8	90	¹ SCHUEMANN	07 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
< 22	90	¹ AUBERT,B	04D BABR	Repl. by AUBERT 07E
< 33	90	¹ RICHICHI	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
< 47	90	BEHRENS	98 CLE2	Repl. by RICHICHI 00

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\phi \pi^+)/\Gamma_{\text{total}}$ Γ_{340}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.24	90	¹ AUBERT,B	06C BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.41	90	¹ AUBERT	04A BABR	Repl. by AUBERT,B 06C
< 1.4	90	¹ AUBERT	01D BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
< 153	90	² ABE	00C SLD	$e^+ e^- \rightarrow Z$
< 5	90	¹ BERGFELD	98 CLE2	

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

$\Gamma(\phi \rho^+)/\Gamma_{\text{total}}$ Γ_{341}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.0	90	¹ AUBERT	08BK BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 16		¹ BERGFELD	98 CLE2	
------	--	-----------------------	---------	--

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(a_0(980)^0 \pi^+ \times B(a_0(980)^0 \rightarrow \eta \pi^0))/\Gamma_{\text{total}}$ Γ_{342}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 5.8	90	¹ AUBERT,BE	04 BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of charged and neutral B mesons from $\Upsilon(4S)$ decays.

$\Gamma(a_0(980)^+ \pi^0 \times B(a_0^+ \rightarrow \eta \pi^+))/\Gamma_{\text{total}}$ Γ_{343}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.4	90	¹ AUBERT	08A BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\pi^+ \pi^+ \pi^+ \pi^- \pi^-)/\Gamma_{\text{total}}$ Γ_{344}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 8.6×10^{-4}	90	¹ ALBRECHT	90B ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\Upsilon(4S)$.

$\Gamma(\rho^0 a_1(1260)^+)/\Gamma_{\text{total}}$ Γ_{345}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.2 \times 10^{-4}$	90	¹ BORTOLETTO89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
$<6.0 \times 10^{-4}$	90	² ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$
$<3.2 \times 10^{-3}$	90	¹ BEBEK	87 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

- • • We do not use the following data for averages, fits, limits, etc. • • •
- ¹ BORTOLETTO 89 reports $< 5.4 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.
- ² ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(\rho^0 a_2(1320)^+)/\Gamma_{\text{total}}$ Γ_{346}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.2 \times 10^{-4}$	90	¹ BORTOLETTO89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
$<2.6 \times 10^{-3}$	90	² BEBEK	87 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

- • • We do not use the following data for averages, fits, limits, etc. • • •
- ¹ BORTOLETTO 89 reports $< 6.3 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.
- ² BEBEK 87 reports $< 2.3 \times 10^{-3}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(b_1^0 \pi^+ \times B(b_1^0 \rightarrow \omega \pi^0))/\Gamma_{\text{total}}$ Γ_{347}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$6.7 \pm 1.7 \pm 1.0$	¹ AUBERT	07BI BABR	$e^+e^- \rightarrow \Upsilon(4S)$

- ¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(b_1^+ \pi^0 \times B(b_1^+ \rightarrow \omega \pi^+))/\Gamma_{\text{total}}$ Γ_{348}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
<3.3	¹ AUBERT	08AG BABR	$e^+e^- \rightarrow \Upsilon(4S)$

- ¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{349}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.3 \times 10^{-3}$	90	¹ ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$

- ¹ ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(a_1(1260)^+ a_1(1260)^0)/\Gamma_{\text{total}}$ Γ_{350}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.3 \times 10^{-2}$	90	¹ ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$

- ¹ ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(h^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{351}/Γ

$h^+ = K^+ \text{ or } \pi^+$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$16 \pm 6 \pm 3.6$	GODANG	98 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\omega h^+)/\Gamma_{\text{total}}$ Γ_{352}/Γ
 $h^+ = K^+ \text{ or } \pi^+$

<u>VALUE (units 10^{-6})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

13.8^{+2.7}_{-2.4} OUR AVERAGE

13.4 ^{+3.3} _{-2.9} ± 1.1	¹ LU	02	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
14.3 ^{+3.6} _{-3.2} ± 2.0	¹ JESSOP	00	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

25 ⁺⁸ ₋₇ ± 3	¹ BERGFELD	98	CLE2	Repl. by JESSOP 00
------------------------------------	-----------------------	----	------	--------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(h^+ X^0(\text{Familon}))/\Gamma_{\text{total}}$ Γ_{353}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<49	90	¹ AMMAR	01B	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$
---------------	----	--------------------	-----	---

¹ AMMAR 01B searched for the two-body decay of the B meson to a massless neutral feebly-interacting particle X^0 such as the familon, the Nambu-Goldstone boson associated with a spontaneously broken global family symmetry.

$\Gamma(\rho\bar{\rho}\pi^+)/\Gamma_{\text{total}}$ Γ_{354}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

1.62 ± 0.20 OUR AVERAGE

1.60 ⁺ ₋	0.22 [±] 0.19	0.12	1,2,3 WEI	08 BELL $e^+ e^- \rightarrow \Upsilon(4S)$
1.69 ±	0.29 ±	0.26	¹ AUBERT	07AV BABR $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.06 ⁺ ₋	0.73 [±] 0.62	0.37	1,3 WANG	04 BELL Repl. by WEI 08
< 3.7	90	1,2 ABE	02K BELL	Repl. by WANG 04
<500	90	⁴ ABREU	95N DLPH	Repl. by ADAM 96D
<160	90	⁵ BEBEK	89 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
570 ± 150 ± 210		⁶ ALBRECHT	88F ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Explicitly vetoes resonant production of $\rho\bar{\rho}$ from Charmonium states.

³ Also provides results with $m_{\rho\bar{\rho}} < 2.85 \text{ GeV}/c^2$ and angular asymmetry of $\rho\bar{\rho}$ system.

⁴ Assumes a B^0 , B^- production fraction of 0.39 and a B_s production fraction of 0.12.

⁵ BEBEK 89 reports $< 1.4 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

⁶ ALBRECHT 88F reports $(5.2 \pm 1.4 \pm 1.9) \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\rho\bar{\rho}\pi^+ \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{355}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<53	90	BERGFELD	96B	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$
---------------	----	----------	-----	---

$\Gamma(\rho\bar{\rho}\pi^+\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{356}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<5.2 \times 10^{-4}$	90	¹ ALBRECHT 88F	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
-----------------------	----	---------------------------	-----	-----------------------------------

¹ ALBRECHT 88F reports $< 4.7 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\rho\bar{\rho}K^+)/\Gamma_{\text{total}}$ Γ_{357}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

5.9 ± 0.5 OUR AVERAGE Error includes scale factor of 1.5.

$5.54^{+0.27}_{-0.25} \pm 0.36$	1,2,3 WEI	08 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
---------------------------------	-----------	---------	-----------------------------------

$6.7 \pm 0.5 \pm 0.4$	^{1,3} AUBERT,B	05L BABR	$e^+e^- \rightarrow \Upsilon(4S)$
-----------------------	-------------------------	----------	-----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.59^{+0.38}_{-0.34} \pm 0.50$	1,2,3 WANG	05A BELL	Repl. by WEI 08
---------------------------------	------------	----------	-----------------

$5.66^{+0.67}_{-0.57} \pm 0.62$	1,2,3 WANG	04 BELL	Repl. by WANG 05A
---------------------------------	------------	---------	-------------------

$4.3^{+1.1}_{-0.9} \pm 0.5$	^{1,2} ABE	02K BELL	Repl. by WANG 04
-----------------------------	--------------------	----------	------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Explicitly vetoes resonant production of $\rho\bar{\rho}$ from Charmonium states.

³ Provides also results with $m_{\rho\bar{\rho}} < 2.85 \text{ GeV}/c^2$ and angular asymmetry of $\rho\bar{\rho}$ system.

$\Gamma(\Theta(1710)^{++}\bar{\rho} \times B(\Theta(1710)^{++} \rightarrow \rho K^+))/\Gamma_{\text{total}}$ Γ_{358}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<0.091	90	¹ WANG 05A	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
------------------	----	-----------------------	------	-----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.1	90	^{1,2} AUBERT,B	05L BABR	$e^+e^- \rightarrow \Upsilon(4S)$
--------	----	-------------------------	----------	-----------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Provides upper limits depending on the pentaquark masses between 1.43 to 2.0 GeV/c^2 .

$\Gamma(f_J(2220)K^+ \times B(f_J(2220) \rightarrow \rho\bar{\rho}))/\Gamma_{\text{total}}$ Γ_{359}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<0.41	90	¹ WANG 05A	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
-----------------	----	-----------------------	------	-----------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\rho\bar{\Lambda}(1520))/\Gamma_{\text{total}}$ Γ_{360}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<1.5	90	¹ AUBERT,B	05L BABR	$e^+e^- \rightarrow \Upsilon(4S)$
----------------	----	-----------------------	----------	-----------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\rho\bar{\rho}K^+ \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{361}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<89	90	BERGFELD 96B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
---------------	----	--------------	------	-----------------------------------

$\Gamma(\rho\bar{p}K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{362}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

3.6 $^{+0.8}_{-0.7}$ OUR AVERAGE

3.38 $^{+0.73}_{-0.60} \pm 0.39$ 1,2 CHEN 08C BELL $e^+e^- \rightarrow \gamma(4S)$

5.3 $\pm 1.5 \pm 1.3$ 2 AUBERT 07AV BABR $e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

10.3 $^{+3.6}_{-2.8} \ ^{+1.3}_{-1.7}$ 2,3 WANG 04 BELL Repl. by CHEN 08C

¹ Explicitly vetoes resonant production of $\rho\bar{p}$ from charmonium states.

² Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

³ Explicitly vetoes resonant production of $\rho\bar{p}$ from charmonium states. The branching fraction for $M_{\rho\bar{p}} < 2.85 \text{ GeV}/c^2$ is also reported.

$\Gamma(f_J(2220)K^{*+} \times B(f_J(2220) \rightarrow p\bar{p}))/\Gamma_{\text{total}}$ Γ_{363}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<0.77 90 1 AUBERT 07AV BABR $e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\rho\bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{364}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 0.32 90 1 TSAI 07 BELL $e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.49 90 1 CHANG 05 BELL Repl. by TSAI 2007

< 1.5 90 1 BORNHEIM 03 CLE2 $e^+e^- \rightarrow \gamma(4S)$

< 2.2 90 1 ABE 02O BELL $e^+e^- \rightarrow \gamma(4S)$

< 2.6 90 1 COAN 99 CLE2 $e^+e^- \rightarrow \gamma(4S)$

<60 90 2 AVERY 89B CLEO $e^+e^- \rightarrow \gamma(4S)$

<93 90 3 ALBRECHT 88F ARG $e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² AVERY 89B reports $< 5 \times 10^{-5}$ assuming the $\gamma(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

³ ALBRECHT 88F reports $< 8.5 \times 10^{-5}$ assuming the $\gamma(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\rho\bar{\Lambda}\gamma)/\Gamma_{\text{total}}$ Γ_{365}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

2.45 $^{+0.44}_{-0.38} \pm 0.22$ 1 WANG 07C BELL $e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.16 $^{+0.58}_{-0.53} \pm 0.20$ 1 LEE 05 BELL Repl. by WANG 07C

<3.9 90 2 EDWARDS 03 CLE2 $e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² Corresponds to $E_\gamma > 1.5 \text{ GeV}$. The limit changes to 3.3×10^{-6} for $E_\gamma > 2.0 \text{ GeV}$.

$\Gamma(\rho\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$ Γ_{366}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.00^{+0.61}_{-0.53} \pm 0.33$		¹ WANG	07C BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\rho\bar{\Sigma}(1385)^0)/\Gamma_{\text{total}}$ Γ_{367}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.47	90	¹ WANG	07C BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\Delta^+\bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{368}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.82	90	¹ WANG	07C BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\rho\bar{\Sigma}\gamma)/\Gamma_{\text{total}}$ Γ_{369}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4.6	90	¹ LEE	05 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<7.9	90	² EDWARDS	03 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
--------	----	----------------------	---------	-----------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Corresponds to $E_\gamma > 1.5$ GeV. The limit changes to 6.4×10^{-6} for $E_\gamma > 2.0$ GeV.

$\Gamma(\rho\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{370}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.0 \times 10^{-4}$	90	¹ ALBRECHT	88F ARG	$e^+e^- \rightarrow \Upsilon(4S)$

¹ ALBRECHT 88F reports $< 1.8 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\Lambda\bar{\Lambda}\pi^+)/\Gamma_{\text{total}}$ Γ_{371}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.8	90	¹ LEE	04 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\Lambda\bar{\Lambda}K^+)/\Gamma_{\text{total}}$ Γ_{372}/Γ

<u>VALUE (units 10^{-6})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.91^{+0.9}_{-0.70} \pm 0.38$	¹ LEE	04 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{\Delta^0} p)/\Gamma_{\text{total}}$ Γ_{373}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.38	90	¹ WEI	08	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<380	90	² BORTOLETTO89	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
------	----	---------------------------	------	------------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² BORTOLETTO 89 reports $< 3.3 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \overline{B}^0$. We rescale to 50%.

$\Gamma(\Delta^{++} \overline{p})/\Gamma_{\text{total}}$ Γ_{374}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.14	90	¹ WEI	08	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<150	90	² BORTOLETTO89	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
------	----	---------------------------	------	------------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² BORTOLETTO 89 reports $< 1.3 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \overline{B}^0$. We rescale to 50%.

$\Gamma(D^+ \rho \overline{p})/\Gamma_{\text{total}}$ Γ_{375}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 1.5×10^{-5}	90	¹ ABE	02W	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^*(2010)^+ \rho \overline{p})/\Gamma_{\text{total}}$ Γ_{376}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 1.5×10^{-5}	90	¹ ABE	02W	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{\Lambda}_c^- \rho \pi^+)/\Gamma_{\text{total}}$ Γ_{377}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
2.8 ± 0.8 OUR AVERAGE			

$3.4 \pm 0.2 \pm 0.9$	^{1,2} AUBERT	08BN	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
-----------------------	-----------------------	------	---

$2.0 \pm 0.2 \pm 0.5$	^{1,3} GABYSHEV	06A	BELL $e^+ e^- \rightarrow \Upsilon(4S)$
-----------------------	-------------------------	-----	---

$2.4 \pm 0.6 \pm 0.6$	^{1,4} DYTMAN	02	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$
-----------------------	-----------------------	----	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.9 \pm 0.5 \pm 0.5$	^{1,5} GABYSHEV	02	BELL Repl. by GABYSHEV 06A
-----------------------	-------------------------	----	----------------------------

$6.2^{+2.3}_{-2.0} \pm 1.6$	^{1,6} FU	97	CLE2 Repl. by DYTMAN 02
-----------------------------	-------------------	----	-------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² AUBERT 08BN reports $(3.4 \pm 0.1 \pm 0.9) \times 10^{-4}$ from a measurement of $[\Gamma(B^+ \rightarrow \overline{\Lambda}_c^- \rho \pi^+)/\Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow \rho K^- \pi^+)]$ assuming $B(\Lambda_c^+ \rightarrow \rho K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. We rescale to our best value $B(\Lambda_c^+ \rightarrow \rho K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ GABYSHEV 06A reports $(2.01 \pm 0.15 \pm 0.20) \times 10^{-4}$ from a measurement of $[\Gamma(B^+ \rightarrow \overline{\Lambda}_c^- \rho \pi^+)/\Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow \rho K^- \pi^+)]$ assuming $B(\Lambda_c^+ \rightarrow \rho K^- \pi^+) = 0.05$. We

rescale to our best value $B(\Lambda_c^+ \rightarrow pK^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ DYTMAN 02 reports $(2.4^{+0.63}_{-0.62}) \times 10^{-4}$ from a measurement of $[\Gamma(B^+ \rightarrow \bar{\Lambda}_c^- p \pi^+)/\Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow pK^- \pi^+)]$ assuming $B(\Lambda_c^+ \rightarrow pK^- \pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow pK^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ GABYSHEV 02 reports $(1.87^{+0.51}_{-0.49}) \times 10^{-4}$ from a measurement of $[\Gamma(B^+ \rightarrow \bar{\Lambda}_c^- p \pi^+)/\Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow pK^- \pi^+)]$ assuming $B(\Lambda_c^+ \rightarrow pK^- \pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow pK^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁶ FU 97 uses PDG 96 values of Λ_c branching fraction.

$\Gamma(\bar{\Lambda}_c^- \Delta(1232)^{++})/\Gamma_{\text{total}}$ **Γ_{378}/Γ**

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.9	90	GABYSHEV 06A	BELL	$e^+ e^- \rightarrow \gamma(4S)$

$\Gamma(\bar{\Lambda}_c^- \Delta_X(1600)^{++})/\Gamma_{\text{total}}$ **Γ_{379}/Γ**

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$5.9 \pm 1.2 \pm 1.5$	¹ GABYSHEV 06A	BELL	$e^+ e^- \rightarrow \gamma(4S)$

¹ GABYSHEV 06A reports $(5.9 \pm 1.0 \pm 0.6) \times 10^{-5}$ from a measurement of $[\Gamma(B^+ \rightarrow \bar{\Lambda}_c^- \Delta_X(1600)^{++})/\Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow pK^- \pi^+)]$ assuming $B(\Lambda_c^+ \rightarrow pK^- \pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow pK^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\bar{\Lambda}_c^- \Delta_X(2420)^{++})/\Gamma_{\text{total}}$ **Γ_{380}/Γ**

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$4.7^{+1.1}_{-1.0} \pm 1.2$	¹ GABYSHEV 06A	BELL	$e^+ e^- \rightarrow \gamma(4S)$

¹ GABYSHEV 06A reports $(4.7^{+1.0}_{-0.9} \pm 0.4) \times 10^{-5}$ from a measurement of $[\Gamma(B^+ \rightarrow \bar{\Lambda}_c^- \Delta_X(2420)^{++})/\Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow pK^- \pi^+)]$ assuming $B(\Lambda_c^+ \rightarrow pK^- \pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow pK^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma((\bar{\Lambda}_c^- p)_s \pi^+)/\Gamma_{\text{total}}$ **Γ_{381}/Γ**

$(\bar{\Lambda}_c^- p)_s$ denotes a low-mass enhancement near $3.35 \text{ GeV}/c^2$.

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.9^{+0.9}_{-0.8} \pm 1.0$	¹ GABYSHEV 06A	BELL	$e^+ e^- \rightarrow \gamma(4S)$

¹ GABYSHEV 06A reports $(3.9^{+0.8}_{-0.7} \pm 0.4) \times 10^{-5}$ from a measurement of $[\Gamma(B^+ \rightarrow (\bar{\Lambda}_c^- p)_s \pi^+)/\Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow pK^- \pi^+)]$ assuming $B(\Lambda_c^+ \rightarrow pK^- \pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow pK^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\overline{\Sigma}_c(2520)^0 p)/\Gamma_{\text{total}}$ Γ_{382}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<0.3	90	1,2 AUBERT	08BN BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
----------------	----	------------	-----------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.7	90	1,2 GABYSHEV	06A BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
------	----	--------------	----------	------------------------------------

<4.6	90	1,2 GABYSHEV	02 BELL	Repl. by GABYSHEV 06A
------	----	--------------	---------	-----------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Uses the value for $\Lambda_c \rightarrow p K^- \pi^+$ branching ratio (5.0 ± 1.3)%.

$\Gamma(\overline{\Sigma}_c(2520)^0 p)/\Gamma(\overline{\Lambda}_c^- p \pi^+)$ $\Gamma_{382}/\Gamma_{377}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<9	90	AUBERT	08BN BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
--------------	----	--------	-----------	------------------------------------

$\Gamma(\overline{\Sigma}_c(2800)^0 p)/\Gamma_{\text{total}}$ Γ_{383}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$3.3 \pm 0.9 \pm 0.9$	1 AUBERT	08BN BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
---	----------	-----------	------------------------------------

¹ AUBERT 08BN reports $[\Gamma(B^+ \rightarrow \overline{\Sigma}_c(2800)^0 p)/\Gamma_{\text{total}}] / [B(B^+ \rightarrow \overline{\Lambda}_c^- p \pi^+)] = 0.117 \pm 0.023 \pm 0.024$. We multiply by our best value $B(B^+ \rightarrow \overline{\Lambda}_c^- p \pi^+) = (2.8 \pm 0.8) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\overline{\Lambda}_c^- p \pi^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{384}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

$1.81 \pm 0.29^{+0.52}_{-0.50}$	1,2 DYTMAN	02 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
---	------------	---------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.12	90	³ FU	97 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
-------	----	-----------------	---------	------------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² DYTMAN 02 measurement uses $B(\Lambda_c^- \rightarrow \overline{p} K^+ \pi^-) = 5.0 \pm 1.3\%$. The second error includes the systematic and the uncertainty of the branching ratio.

³ FU 97 uses PDG 96 values of Λ_c branching ratio.

$\Gamma(\overline{\Lambda}_c^- p \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{385}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

$2.25 \pm 0.25^{+0.63}_{-0.61}$	1,2 DYTMAN	02 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
---	------------	---------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.46	90	³ FU	97 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
-------	----	-----------------	---------	------------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² DYTMAN 02 measurement uses $B(\Lambda_c^- \rightarrow \overline{p} K^+ \pi^-) = 5.0 \pm 1.3\%$. The second error includes the systematic and the uncertainty of the branching ratio.

³ FU 97 uses PDG 96 values of Λ_c branching ratio.

$\Gamma(\bar{\Lambda}_c^- p \pi^+ \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{386}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.34 \times 10^{-2}$	90	¹ FU	97	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

¹ FU 97 uses PDG 96 values of Λ_c branching ratio.

$\Gamma(\Lambda_c^+ \Lambda_c^- K^+)/\Gamma_{\text{total}}$ Γ_{387}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

8.7 ± 3.5 OUR AVERAGE

11 ± 6 ± 3 ^{1,2} AUBERT 08H BABR $e^+e^- \rightarrow \Upsilon(4S)$

7.9 ± 3.1 ± 2.1 ^{2,3} GABYSHEV 06 BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ AUBERT 08H reports $(1.14 \pm 0.15 \pm 0.62) \times 10^{-3}$ from a measurement of $[\Gamma(B^+ \rightarrow \Lambda_c^+ \Lambda_c^- K^+)/\Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow p K^- \pi^+)]$ assuming $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ GABYSHEV 06 reports $(7.9_{-0.9}^{+1.0} \pm 3.6) \times 10^{-4}$ from a measurement of $[\Gamma(B^+ \rightarrow \Lambda_c^+ \Lambda_c^- K^+)/\Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow p K^- \pi^+)]$ assuming $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\bar{\Sigma}_c(2455)^0 p)/\Gamma_{\text{total}}$ Γ_{388}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

3.7 ± 0.8 ± 1.0 ^{1,2} GABYSHEV 06A BELL $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<8 90 ^{1,3} DYTMAN 02 CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

<9.3 90 ^{1,4} GABYSHEV 02 BELL Repl. by GABYSHEV 06A

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² GABYSHEV 06A reports $(3.7 \pm 0.7 \pm 0.4) \times 10^{-5}$ from a measurement of $[\Gamma(B^+ \rightarrow \bar{\Sigma}_c(2455)^0 p)/\Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow p K^- \pi^+)]$ assuming $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ DYTMAN 02 measurement uses $B(\Lambda_c^- \rightarrow \bar{p} K^+ \pi^-) = 5.0 \pm 1.3\%$. The second error includes the systematic and the uncertainty of the branching ratio.

⁴ Uses the value for $\Lambda_c \rightarrow p K^- \pi^+$ branching ratio $(5.0 \pm 1.3)\%$.

$\Gamma(\bar{\Sigma}_c(2455)^0 p)/\Gamma(\bar{\Lambda}_c^- p \pi^+)$ $\Gamma_{388}/\Gamma_{377}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.123 ± 0.012 ± 0.008 ¹ AUBERT 08BN BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{\Sigma}_c(2455)^0 p \pi^0)/\Gamma_{\text{total}}$ Γ_{389}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
4.4±1.4±1.1	1,2 DYTMAN	02	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

¹ DYTMAN 02 reports $(4.4 \pm 1.4) \times 10^{-4}$ from a measurement of $[\Gamma(B^+ \rightarrow \overline{\Sigma}_c(2455)^0 p \pi^0)/\Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow p K^- \pi^+)]$ assuming $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{\Sigma}_c(2455)^0 p \pi^- \pi^+)/\Gamma_{\text{total}}$ Γ_{390}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
4.4±1.3±1.1	1,2 DYTMAN	02	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

¹ DYTMAN 02 reports $(4.4 \pm 1.3) \times 10^{-4}$ from a measurement of $[\Gamma(B^+ \rightarrow \overline{\Sigma}_c(2455)^0 p \pi^- \pi^+)/\Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow p K^- \pi^+)]$ assuming $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{\Sigma}_c(2455)^{-} p \pi^+ \pi^+)/\Gamma_{\text{total}}$ Γ_{391}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
2.8±1.0±0.7	1,2 DYTMAN	02	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

¹ DYTMAN 02 reports $(2.8 \pm 1.0) \times 10^{-4}$ from a measurement of $[\Gamma(B^+ \rightarrow \overline{\Sigma}_c(2455)^{-} p \pi^+ \pi^+)/\Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow p K^- \pi^+)]$ assuming $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{\Lambda}_c(2593)^- / \overline{\Lambda}_c(2625)^- p \pi^+)/\Gamma_{\text{total}}$ Γ_{392}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.9 × 10⁻⁴	90	1,2 DYTMAN	02	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² DYTMAN 02 measurement uses $B(\Lambda_c^- \rightarrow \overline{p} K^+ \pi^-) = 5.0 \pm 1.3\%$. The second error includes the systematic and the uncertainty of the branching ratio.

$\Gamma(\Xi_c^0 \Lambda_c^+ \times B(\Xi_c^0 \rightarrow \Xi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{393}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
3.0±1.1 OUR AVERAGE			
2.5±0.9±0.7	1,2 AUBERT	08H	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
5.6 ^{+2.3} _{-1.9} ±1.5	2,3 CHISTOV	06A	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ AUBERT 08H reports $(2.51 \pm 0.89 \pm 0.61) \times 10^{-5}$ from a measurement of $[\Gamma(B^+ \rightarrow \Xi_c^0 \Lambda_c^+ \times B(\Xi_c^0 \rightarrow \Xi^+ \pi^-)) / \Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow p K^- \pi^+)]$ assuming $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ CHISTOV 06A reports $(5.6^{+1.9}_{-1.5} \pm 1.9) \times 10^{-5}$ from a measurement of $[\Gamma(B^+ \rightarrow \Xi_c^0 \Lambda_c^+ \times B(\Xi_c^0 \rightarrow \Xi^+ \pi^-)) / \Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow p K^- \pi^+)]$ assuming $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Xi_c^0 \Lambda_c^+ \times B(\Xi_c^0 \rightarrow \Lambda K^+ \pi^-)) / \Gamma_{\text{total}}$ Γ_{394} / Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.6 ± 1.1 OUR AVERAGE	Error includes scale factor of 1.1.		
$1.7 \pm 1.0 \pm 0.4$	1,2 AUBERT 08H	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$4.0^{+1.3}_{-1.2} \pm 1.0$	2,3 CHISTOV 06A	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ AUBERT 08H reports $(1.70 \pm 0.93 \pm 0.53) \times 10^{-5}$ from a measurement of $[\Gamma(B^+ \rightarrow \Xi_c^0 \Lambda_c^+ \times B(\Xi_c^0 \rightarrow \Lambda K^+ \pi^-)) / \Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow p K^- \pi^+)]$ assuming $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ CHISTOV 06A reports $(4.0^{+1.1}_{-0.9} \pm 1.3) \times 10^{-5}$ from a measurement of $[\Gamma(B^+ \rightarrow \Xi_c^0 \Lambda_c^+ \times B(\Xi_c^0 \rightarrow \Lambda K^+ \pi^-)) / \Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow p K^- \pi^+)]$ assuming $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^+ \ell^+ \ell^-) / \Gamma_{\text{total}}$ Γ_{395} / Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 4.9 \times 10^{-8}$	90	1 WEI 08A	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$< 1.2 \times 10^{-7}$	90	1 AUBERT 07AG	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\pi^+ e^+ e^-) / \Gamma_{\text{total}}$ Γ_{396} / Γ

Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 8.0 \times 10^{-8}$	90	1 WEI 08A	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$< 1.8 \times 10^{-7}$	90	1 AUBERT 07AG	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$< 3.9 \times 10^{-3}$	90	2 WEIR 90B	MRK2	$e^+ e^-$ 29 GeV

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(\pi^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{397}/Γ**

Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.9 \times 10^{-8}$	90	¹ WEI	08A BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$<2.8 \times 10^{-7}$	90	¹ AUBERT	07AG BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$<9.1 \times 10^{-3}$	90	² WEIR	90B MRK2	$e^+ e^-$ 29 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(\pi^+ \nu \bar{\nu})/\Gamma_{\text{total}}$ **Γ_{398}/Γ**

Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.0 \times 10^{-4}$	90	¹ AUBERT	05H BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$<1.7 \times 10^{-4}$	90	¹ CHEN	07D BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+ \ell^+ \ell^-)/\Gamma_{\text{total}}$ **Γ_{399}/Γ**

Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$4.4^{+0.8}_{-0.7}$ OUR AVERAGE Error includes scale factor of 1.1.

$3.8^{+0.9}_{-0.8} \pm 0.2$ ¹ AUBERT,B 06J BABR $e^+ e^- \rightarrow \Upsilon(4S)$

$5.3^{+1.1}_{-1.0} \pm 0.3$ ¹ ISHIKAWA 03 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+ e^+ e^-)/\Gamma_{\text{total}}$ **Γ_{400}/Γ**

Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

4.9 ± 1.0 OUR AVERAGE

$4.2^{+1.2}_{-1.1} \pm 0.2$ ¹ AUBERT,B 06J BABR $e^+ e^- \rightarrow \Upsilon(4S)$

$6.3^{+1.9}_{-1.7} \pm 0.3$ ² ISHIKAWA 03 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$10.5^{+2.5}_{-2.2} \pm 0.7$ ¹ AUBERT 03U BABR Repl. by
AUBERT,B 06J

< 14 90 ¹ ABE 02 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

< 9 90 ¹ AUBERT 02L BABR $e^+ e^- \rightarrow \Upsilon(4S)$

< 24 90 ³ ANDERSON 01B CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

< 990 90 ⁴ ALBRECHT 91E ARG $e^+ e^- \rightarrow \Upsilon(4S)$

< 68000 90 ⁵ WEIR 90B MRK2 $e^+ e^-$ 29 GeV

< 600 90 ⁶ AVERY 89B CLEO $e^+ e^- \rightarrow \Upsilon(4S)$

< 2500 90 ⁷ AVERY 87 CLEO $e^+ e^- \rightarrow \Upsilon(4S)$

- ¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
- ² Assumes equal production of B^0 and B^+ at $\Upsilon(4S)$. The second error is a total of systematic uncertainties including model dependence.
- ³ The result is for di-lepton masses above 0.5 GeV.
- ⁴ ALBRECHT 91E reports $< 9.0 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.
- ⁵ WEIR 90B assumes B^+ production cross section from LUND.
- ⁶ AVERY 89B reports $< 5 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.
- ⁷ AVERY 87 reports $< 2.1 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 40% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K^+\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{401}/Γ
 Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-6}) CL% DOCUMENT ID TECN COMMENT

$0.45^{+0.09}_{-0.08}$ OUR AVERAGE

$0.59 \pm 0.15 \pm 0.02$		¹ AALTONEN	09B	CDF	$p\bar{p}$ at 1.96 TeV
$0.31^{+0.15}_{-0.12} \pm 0.03$		² AUBERT,B	06J	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$0.45^{+0.14}_{-0.12} \pm 0.03$		³ ISHIKAWA	03	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$0.07^{+0.19}_{-0.11} \pm 0.02$		² AUBERT	03U	BABR	Repl. by AUBERT,B 06J
$0.98^{+0.46}_{-0.36} \pm 0.16$		² ABE	02	BELL	Repl. by ISHIKAWA 03
< 1.2	90	² AUBERT	02L	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
< 3.68	90	⁴ ANDERSON	01B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
< 5.2	90	⁵ AFFOLDER	99B	CDF	$p\bar{p}$ at 1.8 TeV
< 10	90	⁶ ABE	96L	CDF	Repl. by AF-FOLDER 99B
< 240	90	⁷ ALBRECHT	91E	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
< 6400	90	⁸ WEIR	90B	MRK2	e^+e^- 29 GeV
< 170	90	⁹ AVERY	89B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
< 380	90	¹⁰ AVERY	87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹ AALTONEN 09B reports $[\Gamma(B^+ \rightarrow K^+\mu^+\mu^-)/\Gamma_{\text{total}}] / [B(B^+ \rightarrow J/\psi(1S)K^+)] = (0.59 \pm 0.15 \pm 0.03) \times 10^{-3}$. We multiply by our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.007 \pm 0.035) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

- ² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
- ³ Assumes equal production of B^0 and B^+ at $\Upsilon(4S)$. The second error is a total of systematic uncertainties including model dependence.
- ⁴ The result is for di-lepton masses above 0.5 GeV.
- ⁵ AFFOLDER 99B measured relative to $B^+ \rightarrow J/\psi(1S)K^+$.
- ⁶ ABE 96L measured relative to $B^+ \rightarrow J/\psi(1S)K^+$ using PDG 94 branching ratios.
- ⁷ ALBRECHT 91E reports $< 2.2 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.
- ⁸ WEIR 90B assumes B^+ production cross section from LUND.
- ⁹ AVERY 89B reports $< 1.5 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.
- ¹⁰ AVERY 87 reports $< 3.2 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 40% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K^+\bar{\nu})/\Gamma_{\text{total}}$ Γ_{402}/Γ

Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.4 \times 10^{-5}$	90	¹ CHEN	07D BELL	$e^+e^- \rightarrow \gamma(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<5.2 \times 10^{-5}$	90	¹ AUBERT	05H BABR	$e^+e^- \rightarrow \gamma(4S)$
$<2.4 \times 10^{-4}$	90	¹ BROWDER	01 CLE2	$e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\rho^+\nu\bar{\nu})/\Gamma_{\text{total}}$ Γ_{403}/Γ

Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interaction.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.5 \times 10^{-4}$	90	¹ CHEN	07D BELL	$e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K^*(892)^+\ell^+\ell^-)/\Gamma_{\text{total}}$ Γ_{404}/Γ

Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
$7.3^{+5.0}_{-4.2} \pm 2.1$		¹ AUBERT,B	06J BABR	$e^+e^- \rightarrow \gamma(4S)$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

<22	90	¹ ISHIKAWA	03 BELL	$e^+e^- \rightarrow \gamma(4S)$
-------	----	-----------------------	---------	---------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K^*(892)^+\nu\bar{\nu})/\Gamma_{\text{total}}$ Γ_{405}/Γ

Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interaction.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8 \times 10^{-5}$	90	AUBERT	08BC BABR	$e^+e^- \rightarrow \gamma(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<1.4 \times 10^{-4}$	90	¹ CHEN	07D BELL	$e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K^*(892)^+e^+e^-)/\Gamma_{\text{total}}$ Γ_{406}/Γ

Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$0.75^{+0.76}_{-0.65} \pm 0.38$		¹ AUBERT,B	06J BABR	$e^+e^- \rightarrow \gamma(4S)$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

$0.20^{+1.34}_{-0.87} \pm 0.28$		¹ AUBERT	03U BABR	$e^+e^- \rightarrow \gamma(4S)$
< 4.6	90	² ISHIKAWA	03 BELL	$e^+e^- \rightarrow \gamma(4S)$
< 8.9	90	¹ ABE	02 BELL	Repl. by ISHIKAWA 03
< 9.5	90	¹ AUBERT	02L BABR	$e^+e^- \rightarrow \gamma(4S)$
<690	90	³ ALBRECHT	91E ARG	$e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² Assumes equal production of B^0 and B^+ at $\gamma(4S)$. The second error is a total of systematic uncertainties including model dependence.

³ ALBRECHT 91E reports $< 6.3 \times 10^{-4}$ assuming the $\gamma(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K^*(892)^+ \mu^+ \mu^-) / \Gamma_{\text{total}}$ **Γ_{407} / Γ**

Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

0.8 $^{+0.6}_{-0.4}$ OUR AVERAGE

0.97 $^{+0.94}_{-0.69} \pm 0.14$	1	AUBERT,B	06J	BABR $e^+ e^- \rightarrow \gamma(4S)$
----------------------------------	---	----------	-----	---------------------------------------

0.65 $^{+0.69+0.15}_{-0.53-0.16}$	2	ISHIKAWA	03	BELL $e^+ e^- \rightarrow \gamma(4S)$
-----------------------------------	---	----------	----	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.07 $^{+2.58}_{-1.78} \pm 0.42$	1	AUBERT	03U	BABR $e^+ e^- \rightarrow \gamma(4S)$
----------------------------------	---	--------	-----	---------------------------------------

< 3.9	90	1 ABE	02	BELL Repl. by ISHIKAWA 03
---------	----	-------	----	---------------------------

< 17.0	90	1 AUBERT	02L	BABR $e^+ e^- \rightarrow \gamma(4S)$
----------	----	----------	-----	---------------------------------------

< 1200	90	3 ALBRECHT	91E	ARG $e^+ e^- \rightarrow \gamma(4S)$
----------	----	------------	-----	--------------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² Assumes equal production of B^0 and B^+ at $\gamma(4S)$. The second error is a total of systematic uncertainties including model dependence. The 90% C.L. upper limit is 2.2×10^{-6} .

³ ALBRECHT 91E reports $< 1.1 \times 10^{-3}$ assuming the $\gamma(4S)$ decays 45% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(\pi^+ e^+ \mu^-) / \Gamma_{\text{total}}$ **Γ_{408} / Γ**

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 0.0064	90	1 WEIR	90B	MRK2 $e^+ e^-$ 29 GeV
---------------------------------	----	--------	-----	-----------------------

¹ WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(\pi^+ e^- \mu^+) / \Gamma_{\text{total}}$ **Γ_{409} / Γ**

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 0.0064	90	1 WEIR	90B	MRK2 $e^+ e^-$ 29 GeV
---------------------------------	----	--------	-----	-----------------------

¹ WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(\pi^+ e^\pm \mu^\mp) / \Gamma_{\text{total}}$ **Γ_{410} / Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$< 1.7 \times 10^{-7}$	90	1 AUBERT	07AG	BABR $e^+ e^- \rightarrow \gamma(4S)$
---	----	----------	------	---------------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K^+ e^+ \mu^-) / \Gamma_{\text{total}}$ **Γ_{411} / Γ**

Test of lepton family number conservation.

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 0.91	90	1 AUBERT,B	06J	BABR $e^+ e^- \rightarrow \gamma(4S)$
-------------------------------	----	------------	-----	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 8	90	1 AUBERT	02L	BABR Repl. by AUBERT,B 06J
-------	----	----------	-----	-------------------------------

$< 6.4 \times 10^4$	90	2 WEIR	90B	MRK2 $e^+ e^-$ 29 GeV
---------------------	----	--------	-----	-----------------------

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(K^+ e^- \mu^+)/\Gamma_{\text{total}}$ Γ_{412}/Γ

Test of lepton family number conservation.

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
<1.3	90	¹ AUBERT,B 06J	BABR	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<6.4 \times 10^4$	90	² WEIR 90B	MRK2	$e^+ e^-$ 29 GeV
--------------------	----	-----------------------	------	------------------

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(K^+ e^\pm \mu^\mp)/\Gamma_{\text{total}}$ Γ_{413}/Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
<0.91	90	¹ AUBERT,B 06J	BABR	$e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K^+ \mu^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{414}/Γ

Test of lepton family number conservation.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<77	90	¹ AUBERT 07AZ	BABR	$e^+ e^- \rightarrow \gamma(4S)$

¹ Uses a fully reconstructed hadronic B decay as a tag on the recoil side.

$\Gamma(K^*(892)^+ e^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{415}/Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
<13	90	¹ AUBERT,B 06J	BABR	$e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K^*(892)^+ e^- \mu^+)/\Gamma_{\text{total}}$ Γ_{416}/Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
<9.9	90	¹ AUBERT,B 06J	BABR	$e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K^*(892)^+ e^\pm \mu^\mp)/\Gamma_{\text{total}}$ Γ_{417}/Γ

Test of lepton family number conservation.

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.4	90	¹ AUBERT,B 06J	BABR	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<79	90	¹ AUBERT 02L	BABR	Repl. by AUBERT,B 06J
-------	----	-------------------------	------	--------------------------

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\pi^- e^+ e^+)/\Gamma_{\text{total}}$ Γ_{418}/Γ

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.6 $\times 10^{-6}$	90	¹ EDWARDS 02B	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0039	90	² WEIR 90B	MRK2	$e^+ e^-$ 29 GeV
-----------	----	-----------------------	------	------------------

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(\pi^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ **Γ_{419}/Γ**

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.4 $\times 10^{-6}$	90	¹ EDWARDS	02B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0091	90	² WEIR	90B MRK2	$e^+ e^-$ 29 GeV
---------	----	-------------------	----------	------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(\pi^- e^+ \mu^+)/\Gamma_{\text{total}}$ **Γ_{420}/Γ**

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.3 $\times 10^{-6}$	90	¹ EDWARDS	02B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0064	90	² WEIR	90B MRK2	$e^+ e^-$ 29 GeV
---------	----	-------------------	----------	------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(\rho^- e^+ e^+)/\Gamma_{\text{total}}$ **Γ_{421}/Γ**

Test of total lepton number conservation.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<2.6	90	¹ EDWARDS	02B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\rho^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ **Γ_{422}/Γ**

Test of total lepton number conservation.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<5.0	90	¹ EDWARDS	02B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\rho^- e^+ \mu^+)/\Gamma_{\text{total}}$ **Γ_{423}/Γ**

Test of total lepton number conservation.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<3.3	90	¹ EDWARDS	02B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^- e^+ e^+)/\Gamma_{\text{total}}$ **Γ_{424}/Γ**

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.0 $\times 10^{-6}$	90	¹ EDWARDS	02B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0039	90	² WEIR	90B MRK2	$e^+ e^-$ 29 GeV
---------	----	-------------------	----------	------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(K^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{425}/Γ

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.8 \times 10^{-6}$	90	¹ EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$
<0.0091	90	² WEIR	90B MRK2	$e^+ e^-$ 29 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(K^- e^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{426}/Γ

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.0 \times 10^{-6}$	90	¹ EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$
<0.0064	90	² WEIR	90B MRK2	$e^+ e^-$ 29 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(K^*(892)^- e^+ e^+)/\Gamma_{\text{total}}$ Γ_{427}/Γ

Test of total lepton number conservation.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<2.8	90	¹ EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K^*(892)^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{428}/Γ

Test of total lepton number conservation.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<8.3	90	¹ EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K^*(892)^- e^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{429}/Γ

Test of total lepton number conservation.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<4.4	90	¹ EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

POLARIZATION IN B^+ DECAY

In decays involving two vector mesons, one can distinguish among the states in which meson polarizations are both longitudinal (L) or both are transverse and parallel (\parallel) or perpendicular (\perp) to each other with the parameters Γ_L/Γ , Γ_{\perp}/Γ , and the relative phases ϕ_{\parallel} and ϕ_{\perp} . See the definitions in the note on "Polarization in B Decays" review in the B^0 Particle Listings.

Γ_L/Γ in $B^+ \rightarrow \bar{D}^{*0} \rho^+$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.892 \pm 0.018 \pm 0.016$	CSORNA	03 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

Γ_L/Γ in $B^+ \rightarrow \bar{D}^{*0} K^{*+}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.86±0.06±0.03	AUBERT	04K	BABR $e^+e^- \rightarrow \Upsilon(4S)$

Γ_L/Γ in $B^+ \rightarrow J/\psi K^{*+}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.604±0.015±0.018	ITOH	05	BELL $e^+e^- \rightarrow \Upsilon(4S)$

Γ_{\perp}/Γ in $B^+ \rightarrow J/\psi K^{*+}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.180±0.014±0.010	ITOH	05	BELL $e^+e^- \rightarrow \Upsilon(4S)$

Γ_L/Γ in $B^+ \rightarrow \omega K^{*+}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.41±0.18±0.05	AUBERT	09H	BABR $e^+e^- \rightarrow \Upsilon(4S)$

Γ_L/Γ in $B^+ \rightarrow \omega K_2^*(1430)^+$

VALUE	DOCUMENT ID	TECN	COMMENT
0.56±0.10±0.04	AUBERT	09H	BABR $e^+e^- \rightarrow \Upsilon(4S)$

Γ_L/Γ in $B^+ \rightarrow K^{*+} \bar{K}^{*0}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.75^{+0.16}_{-0.26}±0.03	¹ AUBERT	09F	BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

Γ_L/Γ in $B^+ \rightarrow \phi K^{*+}(892)^+$

VALUE	DOCUMENT ID	TECN	COMMENT
0.50±0.05 OUR AVERAGE			
0.49±0.05±0.03	AUBERT	07BA	BABR $e^+e^- \rightarrow \Upsilon(4S)$
0.52±0.08±0.03	CHEN	05A	BELL $e^+e^- \rightarrow \Upsilon(4S)$
••• We do not use the following data for averages, fits, limits, etc. •••			
0.46±0.12±0.03	AUBERT	03V	BABR Repl. by AUBERT 07BA

Γ_{\perp}/Γ in $B^+ \rightarrow \phi K^{*+}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.20±0.05 OUR AVERAGE			
0.21±0.05±0.02	AUBERT	07BA	BABR $e^+e^- \rightarrow \Upsilon(4S)$
0.19±0.08±0.02	CHEN	05A	BELL $e^+e^- \rightarrow \Upsilon(4S)$

ϕ_{\parallel} in $B^+ \rightarrow \phi K^{*+}$

VALUE (°)	DOCUMENT ID	TECN	COMMENT
2.34±0.18 OUR AVERAGE			
2.47±0.20±0.07	AUBERT	07BA	BABR $e^+e^- \rightarrow \Upsilon(4S)$
2.10±0.28±0.04	CHEN	05A	BELL $e^+e^- \rightarrow \Upsilon(4S)$

ϕ_{\perp} in $B^+ \rightarrow \phi K^{*+}$

VALUE (°)	DOCUMENT ID	TECN	COMMENT
2.58±0.17 OUR AVERAGE			
2.69±0.20±0.03	AUBERT	07BA	BABR $e^+e^- \rightarrow \Upsilon(4S)$
2.31±0.30±0.07	CHEN	05A	BELL $e^+e^- \rightarrow \Upsilon(4S)$

$\delta_0(B^+ \rightarrow \phi K^{*+})$

<u>VALUE (rad)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.07 \pm 0.18 \pm 0.06$	AUBERT	07BA BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

$A_{CP}^0(B^+ \rightarrow \phi K^{*+})$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.17 \pm 0.11 \pm 0.02$	AUBERT	07BA BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

$A_{CP}^\perp(B^+ \rightarrow \phi K^{*+})$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.22 \pm 0.24 \pm 0.08$	AUBERT	07BA BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Delta\phi_{\parallel}(B^+ \rightarrow \phi K^{*+})$

<u>VALUE (rad)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.07 \pm 0.20 \pm 0.05$	AUBERT	07BA BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Delta\phi_\perp(B^+ \rightarrow \phi K^{*+})$

<u>VALUE (rad)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.19 \pm 0.20 \pm 0.07$	AUBERT	07BA BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Delta\delta_0(B^+ \rightarrow \phi K^{*+})$

<u>VALUE (rad)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.20 \pm 0.18 \pm 0.03$	AUBERT	07BA BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

Γ_L/Γ in $B^+ \rightarrow \phi K_1(1270)^+$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.46^{+0.12+0.06}_{-0.13-0.07}$	AUBERT	08BI BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

Γ_L/Γ in $B^+ \rightarrow \phi K_2^*(1430)^+$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.80^{+0.09}_{-0.10} \pm 0.03$	AUBERT	08BI BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

$\delta_0(B^+ \rightarrow \phi K_2^*(1430)^+)$

<u>VALUE (rad)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.59 \pm 0.19 \pm 0.12$	AUBERT	08BI BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Delta\delta_0(B^+ \rightarrow \phi K_2^*(1430)^+)$

<u>VALUE (rad)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.05 \pm 0.19 \pm 0.06$	AUBERT	08BI BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

Γ_L/Γ in $B^+ \rightarrow \rho^0 K^*(892)^+$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.96^{+0.04}_{-0.15} \pm 0.04$	AUBERT	03V BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma_L/\Gamma(B^+ \rightarrow K^*(892)^0 \rho^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.48±0.08 OUR AVERAGE			
0.52±0.10±0.04	AUBERT,B	06G	BABR $e^+e^- \rightarrow \Upsilon(4S)$
0.43±0.11 ^{+0.05} _{-0.02}	ZHANG	05D	BELL $e^+e^- \rightarrow \Upsilon(4S)$

Γ_L/Γ in $B^+ \rightarrow \rho^+ \rho^0$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.950±0.016 OUR AVERAGE			
0.950±0.015±0.006	AUBERT	09G	BABR $e^+e^- \rightarrow \Upsilon(4S)$
0.948±0.106±0.021	ZHANG	03B	BELL $e^+e^- \rightarrow \Upsilon(4S)$
••• We do not use the following data for averages, fits, limits, etc. •••			
0.905±0.042 ^{+0.023} _{-0.027}	AUBERT,BE	06G	BABR Repl. by AUBERT 09G
0.97 ^{+0.03} _{-0.07} ±0.04	AUBERT	03V	BABR Repl. by AUBERT,BE 06G

Γ_L/Γ in $B^+ \rightarrow \omega \rho^+$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.90±0.05±0.03	AUBERT	09H	BABR $e^+e^- \rightarrow \Upsilon(4S)$
••• We do not use the following data for averages, fits, limits, etc. •••			
0.82±0.11±0.02	AUBERT,B	06T	BABR Repl. by AUBERT 09H
0.88 ^{+0.12} _{-0.15} ±0.03	AUBERT	05O	BABR Repl. by AUBERT,B 06T

Γ_L/Γ in $B^+ \rightarrow \rho^+ \bar{p} K^*(892)^+$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.32±0.17±0.09	CHEN	08C	BELL $e^+e^- \rightarrow \Upsilon(4S)$

CP VIOLATION

A_{CP} is defined as


$$\frac{B(B^- \rightarrow \bar{f}) - B(B^+ \rightarrow f)}{B(B^- \rightarrow \bar{f}) + B(B^+ \rightarrow f)},$$

the CP-violation charge asymmetry of exclusive B^- and B^+ decay.

$A_{CP}(B^+ \rightarrow J/\psi(1S) K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.009 ±0.008 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.		
0.0075±0.0061±0.0030	1 ABAZOV	08O	D0 $\rho^+ \bar{p}$ at 1.96 TeV
0.09 ±0.07 ±0.02	2 WEI	08	BELL $e^+e^- \rightarrow \Upsilon(4S)$
0.030 ±0.014 ±0.010	3 AUBERT	05J	BABR $e^+e^- \rightarrow \Upsilon(4S)$
-0.026 ±0.022 ±0.017	ABE	03B	BELL $e^+e^- \rightarrow \Upsilon(4S)$
0.018 ±0.043 ±0.004	4 BONVICINI	00	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$
••• We do not use the following data for averages, fits, limits, etc. •••			
0.03 ±0.015 ±0.006	AUBERT	04P	BABR Repl. by AUBERT 05J
0.003 ±0.030 ±0.004	AUBERT	02F	BABR Repl. by AUBERT 04P


- ¹ Uses $J/\psi \rightarrow \mu^+ \mu^-$ decay.
- ² Uses $B^+ \rightarrow J/\psi K^+$, where $J/\psi \rightarrow \rho \bar{p}$.
- ³ The result reported corresponds to $-A_{CP}$.
- ⁴ A +0.3% correction is applied due to a slightly higher reconstruction efficiency for the positive kaons.



$A_{CP}(B^+ \rightarrow J/\psi(1S) \pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.01 ± 0.07 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.		
$-0.09 \pm 0.08 \pm 0.03$	¹ ABAZOV	08O D0	$\rho \bar{p}$ at 1.96 TeV
$+0.123 \pm 0.085 \pm 0.004$	AUBERT	04P BABR	$e^+ e^- \rightarrow \gamma(4S)$
$-0.023 \pm 0.164 \pm 0.015$	ABE	03B BELL	$e^+ e^- \rightarrow \gamma(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$+0.01 \pm 0.22 \pm 0.01$	AUBERT	02F BABR	Repl. by AUBERT 04P

- ¹ Uses $J/\psi \rightarrow \mu^+ \mu^-$ decay.



$A_{CP}(B^+ \rightarrow J/\psi\rho^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.11 \pm 0.12 \pm 0.08$	AUBERT	07AC	BABR $e^+e^- \rightarrow \Upsilon(4S)$

$A_{CP}(B^+ \rightarrow J/\psi K^*(892)^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.048 \pm 0.029 \pm 0.016$	¹ AUBERT	05J	BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹The result reported corresponds to $-A_{CP}$.

$A_{CP}(B^+ \rightarrow \eta_c K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.16 \pm 0.08 \pm 0.02$	¹ WEI	08	BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹Uses $B^+ \rightarrow \eta_c K^+$, where $\eta_c \rightarrow p\bar{p}$.

$A_{CP}(B^+ \rightarrow \psi(2S)\pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.022 \pm 0.085 \pm 0.016$	BHARDWAJ	08	BELL $e^+e^- \rightarrow \Upsilon(4S)$

$A_{CP}(B^+ \rightarrow \psi(2S)K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.025 ± 0.024 OUR AVERAGE			
$0.052 \pm 0.059 \pm 0.020$	AUBERT	05J	BABR $e^+e^- \rightarrow \Upsilon(4S)$
$-0.042 \pm 0.020 \pm 0.017$	ABE	03B	BELL $e^+e^- \rightarrow \Upsilon(4S)$
$0.02 \pm 0.091 \pm 0.01$	¹ BONVICINI	00	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

¹A +0.3% correction is applied due to a slightly higher reconstruction efficiency for the positive kaons.

$A_{CP}(B^+ \rightarrow \psi(2S)K^*(892)^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.077 \pm 0.207 \pm 0.051$	¹ AUBERT	05J	BABR $e^+e^- \rightarrow \gamma(4S)$

¹ The result reported corresponds to $-A_{CP}$.

$A_{CP}(B^+ \rightarrow \chi_{c1}(1P)\pi^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.07 \pm 0.18 \pm 0.02$	KUMAR	06	BELL $e^+e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow \chi_{c0}K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.11 ± 0.12 OUR AVERAGE			
$-0.14 \pm 0.15^{+0.03}_{-0.06}$	AUBERT	08AI	BABR $e^+e^- \rightarrow \gamma(4S)$
$-0.065 \pm 0.20^{+0.035}_{-0.024}$	GARMASH	06	BELL $e^+e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow \chi_{c1}K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.009 ± 0.033 OUR AVERAGE			
$-0.01 \pm 0.03 \pm 0.02$	KUMAR	06	BELL $e^+e^- \rightarrow \gamma(4S)$
$-0.003 \pm 0.076 \pm 0.017$	¹ AUBERT	05J	BABR $e^+e^- \rightarrow \gamma(4S)$

¹ The result reported corresponds to $-A_{CP}$.

$A_{CP}(B^+ \rightarrow \chi_{c1}K^*(892)^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.471 \pm 0.378 \pm 0.268$	¹ AUBERT	05J	BABR $e^+e^- \rightarrow \gamma(4S)$

¹ The result reported corresponds to $-A_{CP}$.

$A_{CP}(B^+ \rightarrow \bar{D}^0\pi^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.008 ± 0.008	ABE	06	BELL $e^+e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow D_{CP(+1)}\pi^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.035 ± 0.024	ABE	06	BELL $e^+e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow D_{CP(-1)}\pi^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.017 ± 0.026	ABE	06	BELL $e^+e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow \bar{D}^0K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.066 ± 0.036	ABE	06	BELL $e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.003 \pm 0.080 \pm 0.037$	¹ ABE	03D	BELL	Repl. by SWAIN 03
$0.04 \pm 0.06 \pm 0.03$	² SWAIN	03	BELL	Repl. by ABE 06

¹ Corresponds to 90% confidence range $-0.15 < A_{CP} < 0.16$.

² Corresponds to 90% confidence range $-0.07 < A_{CP} < 0.15$.

$r_B(B^+ \rightarrow D^0 K^+)$

$r_B^{(*)}$ and $\delta_B^{(*)}$ are the amplitude ratios and relative strong phases between the amplitudes of $A(B^+ \rightarrow D^{(*)0} K^+)$ and $A(B^+ \rightarrow \bar{D}^{(*)0} K^+)$,

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

0.101 ± 0.032 OUR AVERAGE

0.086 ± 0.032 ± 0.015		1 AUBERT	08AL BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
-----------------------	--	----------	-----------	------------------------------------

0.159 ^{+0.054} _{-0.050} ± 0.050		2 POLUEKTOV	06 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
---	--	-------------	---------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.19	90	HORII	08 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
-------	----	-------	---------	------------------------------------

0.12 ± 0.08 ± 0.05		3 AUBERT,B	05Y BABR	Repl. by AUBERT 08AL
--------------------	--	------------	----------	----------------------

¹ Uses Dalitz plot analysis of $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ and $\bar{D}^0 \rightarrow K_S^0 K^+ K^-$ decays coming from $B^\pm \rightarrow D^{(*)} K^{(*)\pm}$ modes.

² Uses a Dalitz plot analysis of the $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ decays; Combines the DK^+ , $D^* K^+$ and DK^{*+} modes.

³ Uses a Dalitz analysis of neutral D decays to $K_S^0 \pi^+ \pi^-$ in the processes $B^\pm \rightarrow D^{(*)} K^\pm$, $D^* \rightarrow D\pi^0$, $D\gamma$.

$\delta_B(B^+ \rightarrow D^0 K^+)$

VALUE (degrees)	DOCUMENT ID	TECN	COMMENT
-----------------	-------------	------	---------

126 ± 21 OUR AVERAGE

109 ⁺²⁷ ₋₃₀ ± 8	1 AUBERT	08AL BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
---------------------------------------	----------	-----------	------------------------------------

145.7 ^{+19.0} _{-19.7} ± 23.1	2 POLUEKTOV	06 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
--	-------------	---------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

104 ± 45 ⁺²³ ₋₃₂	3 AUBERT,B	05Y BABR	Repl. by AUBERT 08AL
--	------------	----------	----------------------

¹ Uses Dalitz plot analysis of $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ and $\bar{D}^0 \rightarrow K_S^0 K^+ K^-$ decays coming from $B^\pm \rightarrow D^{(*)} K^{(*)\pm}$ modes.

² Uses a Dalitz plot analysis of the $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ decays; Combines the DK^+ , $D^* K^+$ and DK^{*+} modes.

³ Uses a Dalitz analysis of neutral D decays to $K_S^0 \pi^+ \pi^-$ in the processes $B^\pm \rightarrow D^{(*)} K^\pm$, $D^* \rightarrow D\pi^0$, $D\gamma$.

$r_B(B^+ \rightarrow DK^{*+})$

r_B and δ_B are the amplitude ratios and relative strong phases between the amplitudes of $A_{CP}(B^+ \rightarrow DK^{*+})$ and $A_{CP}(B^+ \rightarrow \bar{D}K^{*+})$,

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.27 ± 0.17 OUR AVERAGE Error includes scale factor of 1.9.

0.181 ^{+0.088} _{-0.108} ± 0.042	1 AUBERT	08AL BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
---	----------	-----------	------------------------------------

0.564 ^{+0.216} _{-0.155} ± 0.093	2 POLUEKTOV	06 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
---	-------------	---------	------------------------------------

¹ Uses Dalitz plot analysis of $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ and $\bar{D}^0 \rightarrow K_S^0 K^+ K^-$ decays coming from $B^\pm \rightarrow D^{(*)} K^{(*)\pm}$ modes.

² Uses a Dalitz plot analysis of the $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ decays; Combines the DK^+ , $D^* K^+$ and DK^{*+} modes.

$\delta_B(B^+ \rightarrow DK^{*+})$

VALUE (degrees)	DOCUMENT ID	TECN	COMMENT
157 ±70 OUR AVERAGE Error	includes scale factor of 2.0.		
104 ⁺³⁹ / ₋₃₇ ±18	¹ AUBERT	08AL BABR	$e^+e^- \rightarrow \gamma(4S)$
242.6 ^{+20.2} / _{-23.2} ±49.4	² POLUEKTOV	06 BELL	$e^+e^- \rightarrow \gamma(4S)$

¹ Uses Dalitz plot analysis of $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ and $\bar{D}^0 \rightarrow K_S^0 K^+ K^-$ decays coming from $B^\pm \rightarrow D^{(*)} K^{(*)\pm}$ modes.

² Uses a Dalitz plot analysis of the $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ decays; Combines the DK^+ , $D^* K^+$ and DK^{*+} modes.

$A_{CP}(B^+ \rightarrow [K^- \pi^+]_D K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.1 ^{+0.8}/_{-1.0} ±0.4	HORII	08 BELL	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
+0.88 ^{+0.77} / _{-0.62} ±0.06	SAIGO	05 BELL	Repl. by HORII 08

$A_{CP}(B^+ \rightarrow [K^- \pi^+]_{\bar{D}} K^*(892)^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.22 ±0.61 ±0.17	AUBERT,B	05V BABR	$e^+e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow [K^- \pi^+]_D \pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.02 ^{+0.15}/_{-0.16} ±0.04	HORII	08 BELL	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
+0.30 ^{+0.29} / _{-0.25} ±0.06	SAIGO	05 BELL	Repl. by HORII 08

$A_{CP}(B^+ \rightarrow [\pi^+ \pi^- \pi^0]_D K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.02 ±0.15 ±0.03	¹ AUBERT	07BJ BABR	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.02 ±0.16 ±0.03	AUBERT,B	05T BABR	Repl. by AUBERT 07BJ

¹ Uses a Dalitz plot analysis of $D^0 \rightarrow \pi^+ \pi^- \pi^0$. Also reports the one-sigma regions: $0.06 < r_B < 0.78$, $-30^\circ < \gamma < 76^\circ$, and $-27^\circ < \delta < 78^\circ$.

$A_{CP}(B^+ \rightarrow D_{CP(+1)} K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.21 ±0.10 OUR AVERAGE Error	includes scale factor of 1.2.		
0.27 ±0.09 ±0.04	AUBERT	08AA BABR	$e^+e^- \rightarrow \gamma(4S)$
0.06 ±0.14 ±0.05	ABE	06 BELL	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.35 ±0.13 ±0.04	AUBERT	06J BABR	Repl. by AUBERT 08AA
0.07 ±0.17 ±0.06	AUBERT	04N BABR	Repl. by AUBERT 06J
0.29 ±0.26 ±0.05	¹ ABE	03D BELL	Repl. by SWAIN 03
0.06 ±0.19 ±0.04	² SWAIN	03 BELL	Repl. by ABE 06

¹ Corresponds to 90% confidence range $-0.14 < A_{CP} < 0.73$.

² Corresponds to 90% confidence range $-0.26 < A_{CP} < 0.38$.

$A_{CP}(B^+ \rightarrow D_{CP(-1)} K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.10 ± 0.08 OUR AVERAGE			
$-0.09 \pm 0.09 \pm 0.02$	AUBERT	08AA BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$-0.12 \pm 0.14 \pm 0.05$	ABE	06 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$-0.06 \pm 0.13 \pm 0.04$	AUBERT	06J BABR	Repl. by AUBERT 08AA
$-0.22 \pm 0.24 \pm 0.04$	¹ ABE	03D BELL	Repl. by SWAIN 03
$-0.19 \pm 0.17 \pm 0.05$	² SWAIN	03 BELL	Repl. by ABE 06
¹ Corresponds to 90% confidence range $-0.62 < A_{CP} < 0.18$.			
² Corresponds to 90% confidence range $-0.47 < A_{CP} < 0.11$.			

$A_{CP}(B^+ \rightarrow \bar{D}^{*0} \pi^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.014 ± 0.015	ABE	06 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

$A_{CP}(B^+ \rightarrow (D_{CP(+1)}^*)^0 \pi^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.021 ± 0.045	ABE	06 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

$A_{CP}(B^+ \rightarrow (D_{CP(-1)}^*)^0 \pi^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.090 ± 0.051	ABE	06 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

$A_{CP}(B^+ \rightarrow D^{*0} K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.07 ± 0.04 OUR AVERAGE			
$-0.06 \pm 0.04 \pm 0.01$	AUBERT	08BF BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
-0.089 ± 0.086	ABE	06 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

$r_B^*(B^+ \rightarrow D^{*0} K^+)$

$r_B^{(*)}$ and $\delta_B^{(*)}$ are the amplitude ratios and relative strong phases between the amplitudes of $A(B^+ \rightarrow D^{(*)0} K^+)$ and $A(B^+ \rightarrow \bar{D}^{(*)0} K^+)$,

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.14 ± 0.05 OUR AVERAGE			
$0.135 \pm 0.050 \pm 0.012$	¹ AUBERT	08AL BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$0.175^{+0.108}_{-0.099} \pm 0.050$	² POLUEKTOV	06 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

$0.17 \pm 0.10 \pm 0.04$	³ AUBERT,B	05Y BABR	Repl. by AUBERT 08AL
¹ Uses Dalitz plot analysis of $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ and $\bar{D}^0 \rightarrow K_S^0 K^+ K^-$ decays coming from $B^\pm \rightarrow D^{(*)} K^{(*)\pm}$ modes.			
² Uses a Dalitz plot analysis of the $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ decays; Combines the DK^+ , $D^* K^+$ and DK^{*+} modes.			
³ Uses a Dalitz analysis of neutral D decays to $K_S^0 \pi^+ \pi^-$ in the processes $B^\pm \rightarrow D^{(*)} K^\pm$, $D^* \rightarrow D \pi^0$, $D \gamma$.			

$\delta_B^*(B^+ \rightarrow D^{*0} K^+)$

<u>VALUE (degrees)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
299 ± 24 OUR AVERAGE			
297 $^{+27}_{-29} \pm 6.4$	¹ AUBERT	08AL BABR	$e^+ e^- \rightarrow \gamma(4S)$
302.0 $^{+33.8}_{-35.1} \pm 23.7$	² POLUEKTOV	06 BELL	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

296 ± 41 $^{+20}_{-19}$	³ AUBERT,B	05Y BABR	Repl. by AUBERT 08AL
-----------------------------	-----------------------	----------	----------------------

¹ Uses Dalitz plot analysis of $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ and $\bar{D}^0 \rightarrow K_S^0 K^+ K^-$ decays coming from $B^\pm \rightarrow D^{(*)} K^{(*)\pm}$ modes.

² Uses a Dalitz plot analysis of the $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ decays; Combines the DK^+ , $D^* K^+$ and DK^{*+} modes.

³ Uses a Dalitz analysis of neutral D decays to $K_S^0 \pi^+ \pi^-$ in the processes $B^\pm \rightarrow D^{(*)} K^\pm$, $D^* \rightarrow D\pi^0$, $D\gamma$.

$A_{CP}(B^+ \rightarrow D_{CP(+1)}^{*0} K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.12 ± 0.08 OUR AVERAGE			
-0.11 ± 0.09 ± 0.01	AUBERT	08BF BABR	$e^+ e^- \rightarrow \gamma(4S)$
-0.20 ± 0.22 ± 0.04	ABE	06 BELL	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.10 ± 0.23 $^{+0.03}_{-0.04}$	AUBERT	05N BABR	Repl. by AUBERT 08BF
---------------------------------	--------	----------	----------------------

$A_{CP}(B^+ \rightarrow D_{CP(-1)}^{*0} K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.07 ± 0.10 OUR AVERAGE			
+0.06 ± 0.10 ± 0.02	AUBERT	08BF BABR	$e^+ e^- \rightarrow \gamma(4S)$
+0.13 ± 0.30 ± 0.08	ABE	06 BELL	$e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow D_{CP(+1)} K^*(892)^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.08 ± 0.19 ± 0.08	AUBERT,B	05U BABR	$e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow D_{CP(-1)} K^*(892)^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.26 ± 0.40 ± 0.12	AUBERT,B	05U BABR	$e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow D^{*+} \bar{D}^{*0})$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.15 ± 0.11 ± 0.02	AUBERT,B	06A BABR	$e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow D^{*+} \bar{D}^0)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.06 ± 0.13 ± 0.02	AUBERT,B	06A BABR	$e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow D^+ \bar{D}^{*0})$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.13±0.18±0.04	AUBERT,B	06A	BABR $e^+e^- \rightarrow \Upsilon(4S)$

$A_{CP}(B^+ \rightarrow D^+ \bar{D}^0)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.03±0.07 OUR AVERAGE			
0.00±0.08±0.02	ADACHI	08	BELL $e^+e^- \rightarrow \Upsilon(4S)$
-0.13±0.14±0.02	AUBERT,B	06A	BABR $e^+e^- \rightarrow \Upsilon(4S)$

$A_{CP}(B^+ \rightarrow K_S^0 \pi^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.009±0.029 OUR AVERAGE	Error includes scale factor of 1.2.		
0.03 ±0.03 ±0.01	LIN	07	BELL $e^+e^- \rightarrow \Upsilon(4S)$
-0.029±0.039±0.010	¹ AUBERT,BE	06c	BABR $e^+e^- \rightarrow \Upsilon(4S)$
0.18 ±0.24	² CHEN	00	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
-0.09 ±0.05 ±0.01	³ AUBERT,BE	05E	BABR Repl. by AUBERT,BE 06C
0.05 ±0.05 ±0.01	⁴ CHAO	05A	BELL Repl. by LIN 07
-0.05 ±0.08 ±0.01	⁵ AUBERT	04M	BABR Repl. by AUBERT,BE 05E
0.07 +0.09 +0.01 -0.08 -0.03	⁶ UNNO	03	BELL Repl. by CHAO 05A
0.46 ±0.15 ±0.02	⁷ CASEY	02	BELL Repl. by UNNO 03
0.098 +0.430 +0.020 -0.343 -0.063	⁸ ABE	01K	BELL Repl. by CASEY 02
-0.21 ±0.18 ±0.03	⁹ AUBERT	01E	BABR Repl. by AUBERT 04M

¹ Corresponds to 90% confidence range $-0.092 < A_{CP} < 0.036$.

² Corresponds to 90% confidence range $-0.22 < A_{CP} < 0.56$.

³ Corresponds to 90% confidence range $-0.16 < A_{CP} < -0.02$.

⁴ Corresponds to a 90% CL interval of $-0.04 < A_{CP} < 0.13$.

⁵ 90% CL interval $-0.18 < A_{CP} < 0.08$

⁶ Corresponds to 90% confidence range $-0.10 < A_{CP} < +0.22$.

⁷ Corresponds to 90% confidence range $+0.19 < A_{CP} < +0.72$.

⁸ Corresponds to 90% confidence range $-0.53 < A_{CP} < 0.82$.

⁹ Corresponds to 90% confidence range $-0.51 < A_{CP} < 0.09$.

$A_{CP}(B^+ \rightarrow K^+ \pi^0)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.051±0.025 OUR AVERAGE			
0.07 ±0.03 ±0.01	LIN	08	BELL $e^+e^- \rightarrow \Upsilon(4S)$
0.030±0.039±0.010	AUBERT	07BC	BABR $e^+e^- \rightarrow \Upsilon(4S)$
-0.29 ±0.23	¹ CHEN	00	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.06 ±0.06 ±0.01	² AUBERT	05L	BABR Repl. by AUBERT 07BC
0.06 ±0.06 ±0.02	² CHAO	05A	BELL Repl. by CHAO 04B
0.04 ±0.05 ±0.02	³ CHAO	04B	BELL Repl. by LIN 08
-0.09 ±0.09 ±0.01	⁴ AUBERT	03L	BABR Repl. by AUBERT 05L
-0.02 ±0.19 ±0.02	⁵ CASEY	02	BELL Repl. by CHAO 04B
-0.059 +0.222 +0.055 -0.196 -0.017	⁶ ABE	01K	BELL Repl. by CASEY 02
0.00 ±0.18 ±0.04	⁷ AUBERT	01E	BABR Repl. by AUBERT 03L

- ¹ Corresponds to 90% confidence range $-0.67 < A_{CP} < 0.09$.
² Corresponds to a 90% CL interval of $-0.06 < A_{CP} < 0.18$.
³ Corresponds to 90% CL interval of $-0.05 < A_{CP} < 0.13$.
⁴ Corresponds to 90% confidence range $-0.24 < A_{CP} < 0.06$.
⁵ Corresponds to 90% confidence range $-0.35 < A_{CP} < +0.30$.
⁶ Corresponds to 90% confidence range $-0.40 < A_{CP} < 0.36$.
⁷ Corresponds to 90% confidence range $-0.30 < A_{CP} < +0.30$.

$A_{CP}(B^+ \rightarrow \eta' K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.016±0.019 OUR AVERAGE			
0.010±0.022±0.006	AUBERT	07AE BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
0.028±0.028±0.021	SCHUEMANN	06 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
0.03 ±0.12	¹ CHEN	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.033±0.028±0.005	² AUBERT	05M BABR	Repl. by AUBERT 07AE
0.037±0.045±0.011	³ AUBERT	03W BABR	Repl. by AUBERT 05M
-0.11 ±0.11 ±0.02	⁴ AUBERT	02E BABR	Repl. by AUBERT 05M
-0.015±0.070±0.009	⁵ CHEN	02B BELL	Repl. by SCHUE- MANN 06
0.06 ±0.15 ±0.01	⁶ ABE	01M BELL	Repl. by CHEN 02B

- ¹ Corresponds to 90% confidence range $-0.17 < A_{CP} < 0.23$.
² Corresponds to 90% confidence range $-0.012 < A_{CP} < 0.078$.
³ Corresponds to 90% confidence range $-0.04 < A_{CP} < 0.11$.
⁴ Corresponds to 90% confidence range $-0.28 < A_{CP} < 0.07$.
⁵ Corresponds to 90% confidence range $-0.13 < A_{CP} < 0.10$.
⁶ Corresponds to 90% confidence range $-0.20 < A_{CP} < 0.32$.

$A_{CP}(B^+ \rightarrow \eta' K^*(892)^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.30^{+0.33}_{-0.37} \pm 0.02$	¹ AUBERT	07E BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Reports A_{CP} with the opposite sign convention.

$A_{CP}(B^+ \rightarrow \eta K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.27±0.09 OUR AVERAGE			
-0.22±0.11±0.01	AUBERT	07AE BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
-0.39±0.16±0.03	CHANG	07B BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
-0.20±0.15±0.01	AUBERT,B	05K BABR	Repl. by AUBERT 07AE
-0.49±0.31±0.07	CHANG	05A BELL	Repl. by CHANG 07B
-0.52±0.24±0.01	AUBERT	04H BABR	Repl. by AUBERT,B 05K

$A_{CP}(B^+ \rightarrow \eta K^*(892)^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.02±0.06 OUR AVERAGE			
0.03±0.10±0.01	WANG	07B BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
0.01±0.08±0.02	AUBERT,B	06H BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.13±0.14±0.02	AUBERT,B	04D BABR	Repl. by AUBERT,B 06H

$A_{CP}(B^+ \rightarrow \eta K_0^*(1430)^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.05 \pm 0.13 \pm 0.02$	AUBERT,B	06H	BABR $e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow \eta K_2^*(1430)^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.45 \pm 0.30 \pm 0.02$	AUBERT,B	06H	BABR $e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow \omega K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.02 ± 0.05 OUR AVERAGE			

$-0.01 \pm 0.07 \pm 0.01$ AUBERT 07AE BABR $e^+ e^- \rightarrow \gamma(4S)$

$0.05^{+0.08}_{-0.07} \pm 0.01$ JEN 06 BELL $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.05 \pm 0.09 \pm 0.01$ AUBERT,B 06E BABR Repl. by AUBERT 07AE

$-0.09 \pm 0.17 \pm 0.01$ AUBERT 04H BABR Repl. by AUBERT,B 06E

$0.06^{+0.21}_{-0.18} \pm 0.01$ ¹ WANG 04A BELL Repl. by JEN 06

$-0.21 \pm 0.28 \pm 0.03$ ² LU 02 BELL Repl. by WANG 04A

¹ Corresponds to 90% CL interval $0.15 < A_{CP} < 0.90$

² Corresponds to 90% confidence range $-0.70 < A_{CP} < +0.38$.

$A_{CP}(B^+ \rightarrow \omega K^{*+})$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$+0.29 \pm 0.35 \pm 0.02$	AUBERT	09H	BABR $e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow \omega (K\pi)_0^{*+})$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.10 \pm 0.09 \pm 0.02$	AUBERT	09H	BABR $e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow \omega K_2^*(1430)^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$+0.14 \pm 0.15 \pm 0.02$	AUBERT	09H	BABR $e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow K^*(892)^+ \pi^0)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.04 \pm 0.29 \pm 0.05$	AUBERT	05X	BABR $e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow K^{*0} \pi^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.04 ± 0.09 OUR AVERAGE			Error includes scale factor of 2.1.

$0.032 \pm 0.052^{+0.016}_{-0.013}$ AUBERT 08AI BABR $e^+ e^- \rightarrow \gamma(4S)$

$-0.149 \pm 0.064 \pm 0.022$ GARMASH 06 BELL $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.068 \pm 0.078^{+0.070}_{-0.067}$ AUBERT,B 05N BABR Repl. by AUBERT 08AI

$A_{CP}(B^+ \rightarrow K^+ \pi^- \pi^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.038 ± 0.022 OUR AVERAGE			
$0.028 \pm 0.020 \pm 0.023$	AUBERT	08AI BABR	$e^+ e^- \rightarrow \gamma(4S)$
$0.049 \pm 0.026 \pm 0.020$	GARMASH	06 BELL	$e^+ e^- \rightarrow \gamma(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$-0.013 \pm 0.037 \pm 0.011$	AUBERT,B	05N BABR	Repl. by AUBERT 08AI
$0.01 \pm 0.07 \pm 0.03$	AUBERT	03M BABR	Repl. by AUBERT,B 05N

$A_{CP}(B^+ \rightarrow f_0(980) K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.10^{+0.05}_{-0.04}$ OUR AVERAGE			
$-0.106 \pm 0.050^{+0.036}_{-0.015}$	AUBERT	08AI BABR	$e^+ e^- \rightarrow \gamma(4S)$
$-0.31 \pm 0.25 \pm 0.08$	¹ AUBERT	06O BABR	$e^+ e^- \rightarrow \gamma(4S)$
$-0.077 \pm 0.065^{+0.046}_{-0.026}$	GARMASH	06 BELL	$e^+ e^- \rightarrow \gamma(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$0.088 \pm 0.095^{+0.097}_{-0.056}$	AUBERT,B	05N BABR	Repl. by AUBERT 08AI

¹ Measured in the $B^+ \rightarrow K^+ K^- K^+$ decay.

$A_{CP}(B^+ \rightarrow f_2(1270) K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.68^{+0.19}_{-0.17}$ OUR AVERAGE			
$-0.85 \pm 0.22^{+0.26}_{-0.13}$	AUBERT	08AI BABR	$e^+ e^- \rightarrow \gamma(4S)$
$-0.59 \pm 0.22 \pm 0.036$	GARMASH	06 BELL	$e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow f_X(1300) K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.28 \pm 0.26^{+0.15}_{-0.14}$	AUBERT	08AI BABR	$e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow \rho^0 K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.37 ± 0.10 OUR AVERAGE			
$0.44 \pm 0.10^{+0.06}_{-0.14}$	AUBERT	08AI BABR	$e^+ e^- \rightarrow \gamma(4S)$
$0.30 \pm 0.11^{+0.11}_{-0.04}$	GARMASH	06 BELL	$e^+ e^- \rightarrow \gamma(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$0.32 \pm 0.13^{+0.10}_{-0.08}$	AUBERT,B	05N BABR	Repl. by AUBERT 08AI

$A_{CP}(B^+ \rightarrow K_0^*(1430)^0 \pi^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.055 ± 0.033 OUR AVERAGE			
$0.032 \pm 0.035^{+0.034}_{-0.028}$	AUBERT	08AI	BABR $e^+ e^- \rightarrow \gamma(4S)$
$0.076 \pm 0.038^{+0.028}_{-0.022}$	GARMASH	06	BELL $e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$-0.064 \pm 0.032^{+0.023}_{-0.026}$	AUBERT,B	05N	BABR Repl. by AUBERT 08AI

$A_{CP}(B^+ \rightarrow K_2^*(1430)^0 \pi^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.05 \pm 0.23^{+0.18}_{-0.08}$	AUBERT	08AI	BABR $e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow K^0 \rho^+)$

<u>VALUE (units 10^{-6})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.12 \pm 0.17 \pm 0.02$	AUBERT	07Z	BABR $e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow K^{*+} \pi^+ \pi^-)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.07 \pm 0.07 \pm 0.04$	AUBERT,B	06U	BABR $e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow \rho \bar{\Lambda} \gamma)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$+0.17 \pm 0.16 \pm 0.05$	WANG	07C	BELL $e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow \rho \bar{\Lambda} \pi^0)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$+0.01 \pm 0.17 \pm 0.04$	WANG	07C	BELL $e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow \rho^0 K^*(892)^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.20^{+0.32}_{-0.29} \pm 0.04$	AUBERT	03V	BABR $e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow K^*(892)^+ f_0(980))$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.34 \pm 0.21 \pm 0.03$	AUBERT,B	06G	BABR $e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow a_1^+ K^0)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$+0.12 \pm 0.11 \pm 0.02$	AUBERT	08F	BABR $e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow b_1^+ K^0)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.03 \pm 0.15 \pm 0.02$	AUBERT	08AG	BABR $e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow K^*(892)^0 \rho^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.01 \pm 0.16 \pm 0.02$	AUBERT,B	06G	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

$A_{CP}(B^+ \rightarrow b_1^0 K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.46 \pm 0.20 \pm 0.02$	AUBERT	07BI	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

$A_{CP}(B^+ \rightarrow K^0 K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.12 ± 0.18 OUR AVERAGE			

$0.13^{+0.23}_{-0.24} \pm 0.02$ LIN 07 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

$0.10 \pm 0.26 \pm 0.03$ ¹ AUBERT,BE 06C BABR $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.15 \pm 0.33 \pm 0.03$ ² AUBERT,BE 05E BABR Repl. by AUBERT,BE 06C

¹ Corresponds to 90% confidence range $-0.31 < A_{CP} < 0.54$.

² Corresponds to 90% confidence range $-0.43 < A_{CP} < 0.68$.

$A_{CP}(B^+ \rightarrow K^+ K_S^0 K_S^0)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.04 \pm 0.11 \pm 0.02$	¹ AUBERT,B	04V	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Corresponds to 90% confidence range $-0.23 < A_{CP} < 0.15$.

$A_{CP}(B^+ \rightarrow K^+ K^- \pi^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.00 \pm 0.10 \pm 0.03$	AUBERT	07BB	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

$A_{CP}(B^+ \rightarrow K^+ K^- K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.017 \pm 0.026 \pm 0.015$	AUBERT	06O	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.02 \pm 0.07 \pm 0.03$ AUBERT 03M BABR Repl. by AUBERT 06O

$A_{CP}(B^+ \rightarrow \phi K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.01 ± 0.06 OUR AVERAGE			

$0.00 \pm 0.08 \pm 0.02$ AUBERT 06O BABR $e^+ e^- \rightarrow \Upsilon(4S)$

$-0.07 \pm 0.17^{+0.03}_{-0.02}$ ACOSTA 05J CDF $p\bar{p}$ at 1.96 TeV

$0.01 \pm 0.12 \pm 0.05$ ¹ CHEN 03B BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.04 \pm 0.09 \pm 0.01$ ² AUBERT 04A BABR Repl. by AUBERT 06O

$-0.05 \pm 0.20 \pm 0.03$ ³ AUBERT 02E BABR $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Corresponds to 90% confidence range $-0.20 < A_{CP} < 0.22$.

² Corresponds to 90% confidence range $-0.10 < A_{CP} < 0.18$.

³ Corresponds to 90% confidence range $-0.37 < A_{CP} < 0.28$.

$A_{CP}(B^+ \rightarrow X_0(1550)K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.04 \pm 0.07 \pm 0.02$	¹ AUBERT	06O BABR	$e^+e^- \rightarrow \gamma(4S)$

¹ Measured in the $B^+ \rightarrow K^+K^-K^+$ decay.

$A_{CP}(B^+ \rightarrow K^{*+}K^+K^-)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.11 \pm 0.08 \pm 0.03$	AUBERT,B	06U BABR	$e^+e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow \phi K^*(892)^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.01 ± 0.08 OUR AVERAGE			

$0.00 \pm 0.09 \pm 0.04$

AUBERT 07BA BABR $e^+e^- \rightarrow \gamma(4S)$

$-0.02 \pm 0.14 \pm 0.03$

¹ CHEN 05A BELL $e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.16 \pm 0.17 \pm 0.03$

AUBERT 03V BABR Repl. by AUBERT 07BA

$-0.13 \pm 0.29^{+0.08}_{-0.11}$

² CHEN 03B BELL Repl. by CHEN 05A

$-0.43^{+0.36}_{-0.30} \pm 0.06$

³ AUBERT 02E BABR Repl. by AUBERT 03V

¹ Corresponds to 90% confidence range $-0.25 < A_{CP} < 0.22$.

² Corresponds to 90% confidence range $-0.64 < A_{CP} < 0.36$.

³ Corresponds to 90% confidence range $-0.88 < A_{CP} < 0.18$.

$A_{CP}(B^+ \rightarrow \phi(K\pi)_0^{*+})$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.04 \pm 0.15 \pm 0.04$	AUBERT	08BI BABR	$e^+e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow \phi K_1(1270)^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.15 \pm 0.19 \pm 0.05$	AUBERT	08BI BABR	$e^+e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow \phi K_2^*(1430)^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.23 \pm 0.19 \pm 0.06$	AUBERT	08BI BABR	$e^+e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow \eta K^+ \gamma)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.12 ± 0.07 OUR AVERAGE			

$-0.09 \pm 0.10 \pm 0.01$

¹ AUBERT 09 BABR $e^+e^- \rightarrow \gamma(4S)$

$-0.16 \pm 0.09 \pm 0.06$

² NISHIDA 05 BELL $e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-0.09 \pm 0.12 \pm 0.01$

¹ AUBERT,B 06M BABR Repl. by AUBERT 09

¹ $m_{\eta K} < 3.25 \text{ GeV}/c^2$.

² $m_{\eta K} < 2.4 \text{ GeV}/c^2$

$A_{CP}(B^+ \rightarrow \phi K^+ \gamma)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.26 \pm 0.14 \pm 0.05$	AUBERT	07Q BABR	$e^+e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow \rho^+ \gamma)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.11 \pm 0.32 \pm 0.09$	TANIGUCHI	08	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

$A_{CP}(B^+ \rightarrow \pi^+ \pi^0)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.06 ± 0.05 OUR AVERAGE			

$0.07 \pm 0.06 \pm 0.01$	LIN	08	BELL $e^+ e^- \rightarrow \Upsilon(4S)$
$0.03 \pm 0.08 \pm 0.01$	AUBERT	07BC	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-0.01 \pm 0.10 \pm 0.02$	¹ AUBERT	05L	BABR Repl. by AUBERT 07BC
$0.00 \pm 0.10 \pm 0.02$	² CHAO	05A	BELL Repl. by CHAO 04B
$-0.02 \pm 0.10 \pm 0.01$	³ CHAO	04B	BELL Repl. by LIN 08
$-0.03^{+0.18}_{-0.17} \pm 0.02$	⁴ AUBERT	03L	BABR Repl. by AUBERT 05L
$0.30 \pm 0.30^{+0.06}_{-0.04}$	⁵ CASEY	02	BELL Repl. by CHAO 04B

¹ Corresponds to a 90% CL interval of $-0.19 < A_{CP} < 0.21$.

² Corresponds to a 90% CL interval of $-0.17 < A_{CP} < 0.16$.

³ This corresponds to 90% CL interval of $-0.18 < A_{CP} < 0.14$.

⁴ Corresponds to 90% confidence range $-0.32 < A_{CP} < 0.27$.

⁵ Corresponds to 90% confidence range $-0.23 < A_{CP} < +0.86$.

$A_{CP}(B^+ \rightarrow \pi^+ \pi^- \pi^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.007 \pm 0.077 \pm 0.025$	AUBERT,B	05G	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-0.39 \pm 0.33 \pm 0.12$	AUBERT	03M	BABR Repl. by AUBERT 05G
---------------------------	--------	-----	--------------------------

$A_{CP}(B^+ \rightarrow \rho^0 \pi^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.074 \pm 0.120^{+0.035}_{-0.055}$	AUBERT,B	05G	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-0.19 \pm 0.11 \pm 0.02$	AUBERT	04Z	BABR Repl. by AUBERT,B 05G
---------------------------	--------	-----	-------------------------------

$A_{CP}(B^+ \rightarrow f_2(1270) \pi^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.004 \pm 0.247^{+0.028}_{-0.032}$	AUBERT,B	05G	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

$A_{CP}(B^+ \rightarrow \rho^+ \pi^0)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.02 ± 0.11 OUR AVERAGE			

$-0.01 \pm 0.13 \pm 0.02$	AUBERT	07X	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
$0.06 \pm 0.17^{+0.04}_{-0.05}$	ZHANG	05A	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.24 \pm 0.16 \pm 0.06$	AUBERT	04Z	BABR Repl. by AUBERT 07X
--------------------------	--------	-----	--------------------------

$A_{CP}(B^+ \rightarrow \rho^+ \rho^0)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.05 ± 0.05 OUR AVERAGE			
$-0.054 \pm 0.055 \pm 0.010$	AUBERT	09G BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$0.00 \pm 0.22 \pm 0.03$	ZHANG	03B BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$-0.12 \pm 0.13 \pm 0.10$	AUBERT,BE	06G BABR	Repl. by AUBERT 09G
$-0.19 \pm 0.23 \pm 0.03$	AUBERT	03V BABR	Repl. by AUBERT,BE 06G

$A_{CP}(B^+ \rightarrow b_1^0 \pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
$+0.05 \pm 0.16 \pm 0.02$	AUBERT	07BI BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

$A_{CP}(B^+ \rightarrow \omega \pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.04 ± 0.06 OUR AVERAGE			
$-0.02 \pm 0.08 \pm 0.01$	AUBERT	07AE BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$-0.02 \pm 0.09 \pm 0.01$	JEN	06 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
-0.34 ± 0.25	¹ CHEN	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$-0.01 \pm 0.10 \pm 0.01$	AUBERT,B	06E BABR	Repl. by AUBERT 07AE
$0.03 \pm 0.16 \pm 0.01$	AUBERT	04H BABR	Repl. by AUBERT,B 06E
$0.50^{+0.23}_{-0.20} \pm 0.02$	² WANG	04A BELL	Repl. by JEN 06
$-0.01^{+0.29}_{-0.31} \pm 0.03$	³ AUBERT	02E BABR	Repl. by AUBERT 04H

¹ Corresponds to 90% confidence range $-0.75 < A_{CP} < 0.07$.

² Corresponds to 90% CL interval $-0.25 < A_{CP} < 0.41$

³ Corresponds to 90% confidence range $-0.50 < A_{CP} < 0.46$.

$A_{CP}(B^+ \rightarrow \omega \rho^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.20 \pm 0.09 \pm 0.02$	AUBERT	09H BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$0.04 \pm 0.18 \pm 0.02$	AUBERT,B	06T BABR	Repl. by AUBERT 09H
$0.05 \pm 0.26 \pm 0.02$	AUBERT	05O BABR	Repl. by AUBERT,B 06T

$A_{CP}(B^+ \rightarrow \eta \pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.16 ± 0.07 OUR AVERAGE	Error includes scale factor of 1.1.		
$-0.08 \pm 0.10 \pm 0.01$	AUBERT	07AE BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$-0.23 \pm 0.09 \pm 0.02$	CHANG	07B BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$-0.13 \pm 0.12 \pm 0.01$	AUBERT,B	05K BABR	Repl. by AUBERT 07AE
$0.07 \pm 0.15 \pm 0.03$	CHANG	05A BELL	Repl. by CHANG 07B
$-0.44 \pm 0.18 \pm 0.01$	AUBERT	04H BABR	Repl. by AUBERT,B 05K

$A_{CP}(B^+ \rightarrow \eta' \pi^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.21±0.15 OUR AVERAGE			
0.21±0.17±0.01	AUBERT	07AE BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
0.20 ^{+0.37} _{-0.36} ±0.04	SCHUEMANN	06 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.14±0.16±0.01	AUBERT,B	05K BABR	Repl. by AUBERT 07AE

$A_{CP}(B^+ \rightarrow \eta \rho^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.11±0.11 OUR AVERAGE			
0.13±0.11±0.02	AUBERT	08AH BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
-0.04 ^{+0.34} _{-0.32} ±0.01	WANG	07B BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.02±0.18±0.02	AUBERT,B	05K BABR	Repl. by AUBERT 08AH

$A_{CP}(B^+ \rightarrow \eta' \rho^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.04±0.28±0.02	¹ AUBERT	07E BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
¹ Reports A_{CP} with the opposite sign convention.			

$A_{CP}(B^+ \rightarrow \rho \bar{p} \pi^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.00±0.04 OUR AVERAGE			
-0.02±0.05±0.02	¹ WEI	08 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
+0.04±0.07±0.04	AUBERT	07AV BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
-0.16±0.22±0.01	WANG	04 BELL	Repl. by WEI 08
¹ Requires $m_{\rho \bar{p}} < 2.85 \text{ GeV}/c^2$.			

$A_{CP}(B^+ \rightarrow \rho \bar{p} K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.16±0.07 OUR AVERAGE			
-0.17±0.10±0.02	¹ WEI	08 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
-0.16 ^{+0.07} _{-0.08} ±0.04	¹ AUBERT,B	05L BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
-0.05±0.11±0.01	WANG	04 BELL	Repl. by WEI 08
¹ Requires $m_{\rho \bar{p}} < 2.85 \text{ GeV}/c^2$.			

$A_{CP}(B^+ \rightarrow \rho \bar{p} K^*(892)^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.21±0.16 OUR AVERAGE Error includes scale factor of 1.4.			
-0.01±0.19±0.02	CHEN	08C BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
+0.32±0.13±0.05	AUBERT	07AV BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

$A_{CP}(B^+ \rightarrow K^+ \ell^+ \ell^-)$

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.07 \pm 0.22 \pm 0.02$	AUBERT,B	06J	BABR $e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow K^{*+} \ell^+ \ell^-)$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.03 \pm 0.23 \pm 0.03$	AUBERT,B	06J	BABR $e^+ e^- \rightarrow \gamma(4S)$

$\gamma(B^+ \rightarrow D^{(*)} K^{(*)+})$

For angle $\gamma(\phi_3)$ of the CKM unitarity triangle, see the review on “CP Violation” in the Reviews section.

VALUE (°)	DOCUMENT ID	TECN	COMMENT
62 ± 15 OUR AVERAGE			

$76^{+22}_{-23} \pm 7.1$	¹ AUBERT	08AL	BABR $e^+ e^- \rightarrow \gamma(4S)$
$53^{+15}_{-18} \pm 10$	² POLUEKTOV	06	BELL $e^+ e^- \rightarrow \gamma(4S)$
$70 \pm 31^{+18}_{-15}$	³ AUBERT,B	05Y	BABR Repl. by AUBERT 08AL
$77^{+17}_{-19} \pm 17$	⁴ POLUEKTOV	04	BELL Repl. by POLUEKTOV 06

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Uses Dalitz plot analysis of $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ and $\bar{D}^0 \rightarrow K_S^0 K^+ K^-$ decays coming from $B^\pm \rightarrow D^{(*)} K^{(*)\pm}$ modes. The corresponding two standard deviation interval is $29^\circ < \gamma < 122^\circ$.

² Uses a Dalitz plot analysis of the $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ decays; Combines the DK^+ , $D^* K^+$ and DK^{*+} modes. The corresponding two standard deviations interval for gamma is $8^\circ < \gamma < 111^\circ$.

³ Uses a Dalitz plot analysis of neutral $D \rightarrow K_S^0 \pi^+ \pi^-$ decays coming from $B^\pm \rightarrow DK^\pm$ and $B^\pm \rightarrow D^{*0} K^\pm$ followed by $D^{*0} \rightarrow D\pi^0, D\gamma$. The corresponding two standard deviations interval for gamma is $12^\circ < \gamma < 137^\circ$. AUBERT,B 05Y also reports the amplitude ratios and the strong phases.

⁴ Uses a Dalitz plot analysis of the 3-body $D \rightarrow K_S^0 \pi^+ \pi^-$ decays coming from $B^\pm \rightarrow DK^\pm$ and $B^\pm \rightarrow D^* K^\pm$ followed by $D^* \rightarrow D\pi^0$; here we use D to denote that the neutral D meson produced in the decay is an admixture of D^0 and \bar{D}^0 . The corresponding two standard deviations interval for γ is $26^\circ < \gamma < 126^\circ$. POLUEKTOV 04 also reports the amplitude ratios and the strong phases.

B^\pm REFERENCES

AALTONEN	09B	PR D79 011104R	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AUBERT	09	PR D79 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09A	PR D79 012002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09F	PR D79 051102	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09G	PRL 102 141802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09H	PR D79 052005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09J	PRD 79 051101	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABAZOV	08O	PRL 100 211802	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ADACHI	08	PR D77 091101R	I. Adachi <i>et al.</i>	(BELLE Collab.)
AUBERT	08A	PR D77 011101R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08AA	PR D77 111102R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08AB	PR D78 012006	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08AD	PR D77 091104R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08AG	PR D78 011104R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08AH	PR D78 011107R	B. Aubert <i>et al.</i>	(BABAR Collab.)

AUBERT	08AI	PR D78 012004	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08AL	PR D78 034023	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08AT	PRL 100 231803	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08AV	PRL 101 081801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08B	PR D77 011102R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08BC	PR D78 072007	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08BD	PR D78 091101R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08BE	PR D78 091102R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08BF	PR D78 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08BH	PR D78 112001	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08BI	PRL 101 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08BK	PRL 101 201801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08BL	PRL 101 261802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08BN	PR D78 112003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08D	PR D77 011107R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08F	PRL 100 051803	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08G	PRL 100 171803	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08H	PR D77 031101R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08N	PRL 100 021801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08Q	PRL 100 151802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08W	PRL 101 082001	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08X	PRL 101 091801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08Y	PR D77 111101R	B. Aubert <i>et al.</i>	(BABAR Collab.)
BHARDWAJ	08	PR D78 051104R	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)
BRODZICKA	08	PRL 100 092001	J. Brodzicka <i>et al.</i>	(BELLE Collab.)
CHEN	08C	PRL 100 251801	J.-H. Chen <i>et al.</i>	(BELLE Collab.)
HORII	08	PR D78 071901	Y. Horii <i>et al.</i>	(BELLE Collab.)
IWABUCHI	08	PRL 101 041601	M. Iwabuchi <i>et al.</i>	(BELLE Collab.)
LIN	08	NAT 452 332	S.-W. Lin <i>et al.</i>	(BELLE Collab.)
LIVENTSEV	08	PR D77 091503R	D. Liventsev <i>et al.</i>	(BELLE Collab.)
TANIGUCHI	08	PRL 101 111801	N. Taniguchi <i>et al.</i>	(BELLE Collab.)
WEI	08	PL B659 80	J.-T. Wei <i>et al.</i>	(BELLE Collab.)
WEI	08A	PR D78 011101R	J.-T. Wei <i>et al.</i>	(BELLE Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
ADAM	07	PRL 99 041802	N.E. Adam <i>et al.</i>	(CLEO Collab.)
Also		PR D76 012007	D.M. Asner <i>et al.</i>	(CLEO Collab.)
AUBERT	07AC	PR D76 031101R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AE	PR D76 031103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AG	PRL 99 051801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AL	PR D76 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AN	PR D76 051101R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AR	PR D76 071103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AV	PR D76 092004	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AZ	PRL 99 201801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07BA	PRL 99 201802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07BB	PRL 99 221801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07BC	PR D76 091102R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07BI	PRL 99 241803	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07BJ	PRL 99 251801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07BL	PRL 99 261801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07BN	PR D76 111101R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07E	PRL 98 051802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07H	PR D75 031101R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07L	PRL 98 151802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07M	PRL 98 171801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07N	PR D75 072002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07Q	PR D75 051102R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07R	PRL 98 211804	B. Aubert <i>et al.</i>	(BABAR Collab.)
Also		PRL 100 189903E	B. Aubert <i>et al.</i>	(BABAR Collab.)
Also		PRL 100 199905E	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07X	PR D75 091103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07Z	PR D76 011103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
CHANG	07B	PR D75 071104R	P. Chang <i>et al.</i>	(BELLE Collab.)
CHEN	07D	PRL 99 221802	K.-F. Chen <i>et al.</i>	(BELLE Collab.)
HOKUUE	07	PL B648 139	T. Hokuue <i>et al.</i>	(BELLE Collab.)
LIN	07	PRL 98 181804	S.-W. Lin <i>et al.</i>	(BELLE Collab.)
LIN	07A	PRL 99 121601	S.-W. Lin <i>et al.</i>	(BELLE Collab.)
SATOYAMA	07	PL B647 67	N. Satoyama <i>et al.</i>	(BELLE Collab.)
SCHUEMANN	07	PR D75 092002	J. Schuemann <i>et al.</i>	(BELLE Collab.)
TSAI	07	PR D75 111101R	Y.-T. Tsai <i>et al.</i>	(BELLE Collab.)
URQUIJO	07	PR D75 032001	P. Urquijo <i>et al.</i>	(BELLE Collab.)

WANG	07B	PR D75 092005	C.H. Wang <i>et al.</i>	(BELLE Collab.)
WANG	07C	PR D76 052004	M.-Z. Wang <i>et al.</i>	(BELLE Collab.)
XIE	07	PR D75 017101	Q.L. Xie <i>et al.</i>	(BELLE Collab.)
ABE	06	PR D73 051106R	K. Abe <i>et al.</i>	(BELLE Collab.)
ABULENCIA	06J	PRL 96 191801	A. Abulencia <i>et al.</i>	(CDF Collab.)
ACOSTA	06	PRL 96 202001	D. Acosta <i>et al.</i>	(CDF Collab.)
AUBERT	06	PR D73 011101R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06F	PR D73 011103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06J	PR D73 051105R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06K	PR D73 057101	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06N	PR D74 031103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06O	PR D74 032003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06Z	PR D73 111104R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	06A	PR D73 112004	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	06C	PR D74 011102R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	06E	PR D74 011106R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	06G	PRL 97 201801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	06H	PRL 97 201802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	06J	PR D73 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	06M	PR D74 031102R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	06P	PR D74 031105R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	06T	PR D74 051102R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	06U	PR D74 051104R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	06Y	PR D74 091105R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06A	PR D74 099903 (errat.)	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06C	PRL 97 171805	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06G	PRL 97 261801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06H	PRL 97 261803	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06J	PR D74 111102R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06M	PR D74 071101R	B. Aubert <i>et al.</i>	(BABAR Collab.)
CHISTOV	06A	PR D74 111105R	R. Chistov <i>et al.</i>	(BELLE Collab.)
FANG	06	PR D74 012007	F. Fang <i>et al.</i>	(BELLE Collab.)
GABYSHEV	06	PRL 97 202003	N. Gabyshev <i>et al.</i>	(BELLE Collab.)
GABYSHEV	06A	PRL 97 242001	N. Gabyshev <i>et al.</i>	(BELLE Collab.)
GARMASH	06	PRL 96 251803	A. Garmash <i>et al.</i>	(BELLE Collab.)
GOKHROO	06	PRL 97 162002	G. Gokhroo <i>et al.</i>	(BELLE Collab.)
IKADO	06	PRL 97 251802	K. Ikado <i>et al.</i>	(BELLE Collab.)
JEN	06	PR D74 111101R	C.-M. Jen <i>et al.</i>	(BELLE Collab.)
KUMAR	06	PR D74 051103R	R. Kumar <i>et al.</i>	(BELLE Collab.)
MOHAPATRA	06	PRL 96 221601	D. Mohapatra <i>et al.</i>	(BELLE Collab.)
POLUEKTOV	06	PR D73 112009	A. Poluektov <i>et al.</i>	(BELLE Collab.)
SCHUEMANN	06	PRL 97 061802	J. Schuemann <i>et al.</i>	(BELLE Collab.)
SONI	06	PL B634 155	N. Soni <i>et al.</i>	(BELLE Collab.)
ABE	05A	PRL 94 221805	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	05B	PR D71 072003	K. Abe <i>et al.</i>	(BELLE Collab.)
Also		PR D71 079903 (errat.)	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	05G	PRL 95 231802	K. Abe <i>et al.</i>	(BELLE Collab.)
ACOSTA	05J	PRL 95 031801	D. Acosta <i>et al.</i>	(CDF Collab.)
AUBERT	05	PRL 94 011801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05B	PR D71 031501R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05G	PR D72 032004	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05H	PRL 94 101801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05J	PRL 94 141801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05K	PRL 94 171801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05L	PRL 94 181802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05M	PRL 94 191802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05N	PR D71 031102R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05O	PR D71 031103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05R	PR D71 071103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05U	PR D71 091103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05X	PR D71 111101	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05B	PRL 95 041804	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05E	PR D72 011102R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05G	PR D72 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05K	PRL 95 131803	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05L	PR D72 051101R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05N	PR D72 072003	B. Aubert <i>et al.</i>	(BABAR Collab.)
Also		PR D74 099903 (errat.)	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05O	PR D72 051102R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05T	PR D72 071102R	B. Aubert <i>et al.</i>	(BABAR Collab.)

AUBERT,B	05U	PR D72 071103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05V	PR D72 071104R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05Y	PRL 95 121802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	05E	PRL 95 221801	B. Aubert <i>et al.</i>	(BABAR Collab.)
CHANG	05	PR D71 072007	M.-C. Chang <i>et al.</i>	(BELLE Collab.)
CHANG	05A	PR D71 091106R	P. Chang <i>et al.</i>	(BELLE Collab.)
CHAO	05A	PR D71 031502R	Y. Chao <i>et al.</i>	(BELLE Collab.)
CHEN	05A	PRL 94 221804	K.-F. Chen <i>et al.</i>	(BELLE Collab.)
GARMASH	05	PR D71 092003	A. Garmash <i>et al.</i>	(BELLE Collab.)
ITOH	05	PRL 95 091601	R. Itoh <i>et al.</i>	(BELLE Collab.)
LEE	05	PRL 95 061802	Y.-J. Lee <i>et al.</i>	(BELLE Collab.)
LIVENTSEV	05	PR D72 051109R	D. Liventsev <i>et al.</i>	(BELLE Collab.)
MAJUMDER	05	PRL 95 041803	G. Majumder <i>et al.</i>	(BELLE Collab.)
MOHAPATRA	05	PR D72 011101R	D. Mohapatra <i>et al.</i>	(BELLE Collab.)
NISHIDA	05	PL B610 23	S. Nishida <i>et al.</i>	(BELLE Collab.)
OKABE	05	PL B614 27	T. Okabe <i>et al.</i>	(BELLE Collab.)
SAIGO	05	PRL 94 091601	M. Saigo <i>et al.</i>	(BELLE Collab.)
WANG	05A	PL B617 141	M.-Z. Wang <i>et al.</i>	(BELLE Collab.)
XIE	05	PR D72 051105R	Q.L. Xie <i>et al.</i>	(BELLE Collab.)
YANG	05	PRL 94 111802	H. Yang <i>et al.</i>	(BELLE Collab.)
ZHANG	05A	PRL 94 031801	J. Zhang <i>et al.</i>	(BELLE Collab.)
ZHANG	05B	PR D71 091107R	L.M. Zhang <i>et al.</i>	(BELLE Collab.)
ZHANG	05D	PRL 95 141801	J. Zhang <i>et al.</i>	(BELLE Collab.)
ABDALLAH	04E	EPJ C33 307	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ABE	04D	PR D69 112002	K. Abe <i>et al.</i>	(BELLE Collab.)
AUBERT	04A	PR D69 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04C	PRL 92 111801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	04H	PRL 92 061801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04K	PRL 92 141801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04M	PRL 92 201802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04N	PRL 92 202002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04O	PRL 92 221803	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04P	PRL 92 241802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04Q	PR D69 051101R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04T	PR D69 071103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04Y	PRL 93 041801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	04Z	PRL 93 051802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04B	PR D70 011101R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04D	PR D70 032006	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04L	PRL 93 131804	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04P	PR D70 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04S	PRL 93 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04U	PR D70 091105R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04V	PRL 93 181805	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	04	PR D70 111102R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	04A	PR D70 112006	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	04B	PR D70 091106	B. Aubert <i>et al.</i>	(BABAR Collab.)
CHAO	04	PR D69 111102R	Y. Chao <i>et al.</i>	(BELLE Collab.)
CHAO	04B	PRL 93 191802	Y. Chao <i>et al.</i>	(BELLE Collab.)
CHISTOV	04	PRL 93 051803	R. Chistov <i>et al.</i>	(BELLE Collab.)
DRUTSKOY	04	PRL 92 051801	A. Drutskoy <i>et al.</i>	(BELLE Collab.)
GARMASH	04	PR D69 012001	A. Garmash <i>et al.</i>	(BELLE Collab.)
LEE	04	PRL 93 211801	Y.-J. Lee <i>et al.</i>	(BELLE Collab.)
MAJUMDER	04	PR D70 111103R	G. Majumder <i>et al.</i>	(BELLE Collab.)
NAKAO	04	PR D69 112001	M. Nakao <i>et al.</i>	(BELLE Collab.)
POLUEKTOV	04	PR D70 072003	A. Poluektov <i>et al.</i>	(BELLE Collab.)
SCHWANDA	04	PRL 93 131803	C. Schwanda <i>et al.</i>	(BELLE Collab.)
WANG	04	PRL 92 131801	M.Z. Wang <i>et al.</i>	(BELLE Collab.)
WANG	04A	PR D70 012001	C.H. Wang <i>et al.</i>	(BELLE Collab.)
ZANG	04	PR D69 017101	S.L. Zang <i>et al.</i>	(BELLE Collab.)
ABE	03B	PR D67 032003	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	03D	PRL 90 131803	K. Abe <i>et al.</i>	(BELLE Collab.)
ADAM	03	PR D67 032001	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ADAM	03B	PR D68 012004	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ATHAR	03	PR D68 072003	S.B. Athar <i>et al.</i>	(CLEO Collab.)
AUBERT	03K	PRL 90 231801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	03L	PRL 91 021801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	03M	PRL 91 051801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	03O	PRL 91 071801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	03U	PRL 91 221802	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	03V	PRL 91 171802	B. Aubert <i>et al.</i>	(BaBar Collab.)

AUBERT	03W	PRL 91 161801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	03X	PR D68 092001	B. Aubert <i>et al.</i>	(BaBar Collab.)
BORNHEIM	03	PR D68 052002	A. Bornheim <i>et al.</i>	(CLEO Collab.)
CHEN	03B	PRL 91 201801	K.-F. Chen <i>et al.</i>	(BELLE Collab.)
CHOI	03	PRL 91 262001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
CSORNA	03	PR D67 112002	S.E. Csorna <i>et al.</i>	(CLEO Collab.)
EDWARDS	03	PR D68 011102R	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
FANG	03	PRL 90 071801	F. Fang <i>et al.</i>	(BELLE Collab.)
HUANG	03	PRL 91 241802	H.-C. Huang <i>et al.</i>	(BELLE Collab.)
ISHIKAWA	03	PRL 91 261601	A. Ishikawa <i>et al.</i>	(BELLE Collab.)
KROKOVNY	03B	PRL 91 262002	P. Krokovny <i>et al.</i>	(BELLE Collab.)
SWAIN	03	PR D68 051101R	S.K. Swain <i>et al.</i>	(BELLE Collab.)
UNNO	03	PR D68 011103R	Y. Unno <i>et al.</i>	(BELLE Collab.)
ZHANG	03B	PRL 91 221801	J. Zhang <i>et al.</i>	(BELLE Collab.)
ABE	02	PRL 88 021801	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02B	PRL 88 031802	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02H	PRL 88 171801	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02K	PRL 88 181803	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02N	PL B538 11	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02O	PR D65 091103R	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02W	PRL 89 151802	K. Abe <i>et al.</i>	(BELLE Collab.)
ACOSTA	02C	PR D65 092009	D. Acosta <i>et al.</i>	(CDF Collab.)
ACOSTA	02F	PR D66 052005	D. Acosta <i>et al.</i>	(CDF Collab.)
AHMED	02B	PR D66 031101R	S. Ahmed <i>et al.</i>	(CLEO Collab.)
AUBERT	02	PR D65 032001	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	02C	PRL 88 101805	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	02E	PR D65 051101R	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	02F	PR D65 091101R	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	02L	PRL 88 241801	B. Aubert <i>et al.</i>	(BaBar Collab.)
BRIERE	02	PRL 89 081803	R. Briere <i>et al.</i>	(CLEO Collab.)
CASEY	02	PR D66 092002	B.C.K. Casey <i>et al.</i>	(BELLE Collab.)
CHEN	02B	PL B546 196	K.-F. Chen <i>et al.</i>	(BELLE Collab.)
DRUTSKOY	02	PL B542 171	A. Drutskoy <i>et al.</i>	(BELLE Collab.)
DYTMAN	02	PR D66 091101R	S.A. Dytman <i>et al.</i>	(CLEO Collab.)
ECKHART	02	PRL 89 251801	E. Eckhart <i>et al.</i>	(CLEO Collab.)
EDWARDS	02B	PR D65 111102R	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
GABYSHEV	02	PR D66 091102R	N. Gabyshev <i>et al.</i>	(BELLE Collab.)
GARMASH	02	PR D65 092005	A. Garmash <i>et al.</i>	(BELLE Collab.)
GODANG	02	PRL 88 021802	R. Godang <i>et al.</i>	(CLEO Collab.)
GORDON	02	PL B542 183	A. Gordon <i>et al.</i>	(BELLE Collab.)
LU	02	PRL 89 191801	R.-S. Lu <i>et al.</i>	(BELLE Collab.)
MAHAPATRA	02	PRL 88 101803	R. Mahapatra <i>et al.</i>	(CLEO Collab.)
NISHIDA	02	PRL 89 231801	S. Nishida <i>et al.</i>	(BELLE Collab.)
ABE	01H	PRL 87 101801	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	01I	PRL 87 111801	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	01K	PR D64 071101	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	01L	PRL 87 161601	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	01M	PL B517 309	K. Abe <i>et al.</i>	(BELLE Collab.)
ALEXANDER	01B	PR D64 092001	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
AMMAR	01B	PRL 87 271801	R. Ammar <i>et al.</i>	(CLEO Collab.)
ANDERSON	01B	PRL 87 181803	S. Anderson <i>et al.</i>	(CLEO Collab.)
AUBERT	01D	PRL 87 151801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	01E	PRL 87 151802	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	01F	PRL 87 201803	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	01G	PRL 87 221802	B. Aubert <i>et al.</i>	(BaBar Collab.)
BARATE	01E	EPJ C19 213	R. Barate <i>et al.</i>	(ALEPH Collab.)
BRIERE	01	PRL 86 3718	R.A. Biere <i>et al.</i>	(CLEO Collab.)
BROWDER	01	PRL 86 2950	T.E. Browder <i>et al.</i>	(CLEO Collab.)
EDWARDS	01	PRL 86 30	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
GRITSAN	01	PR D64 077501	A. Gritsan <i>et al.</i>	(CLEO Collab.)
RICHICHI	01	PR D63 031103R	S.J. Richichi <i>et al.</i>	(CLEO Collab.)
ABBIENDI	00B	PL B476 233	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABE	00C	PR D62 071101R	K. Abe <i>et al.</i>	(SLD Collab.)
AHMED	00B	PR D62 112003	S. Ahmed <i>et al.</i>	(CLEO Collab.)
ANASTASSOV	00	PRL 84 1393	A. Anastassov <i>et al.</i>	(CLEO Collab.)
BARATE	00R	PL B492 275	R. Barate <i>et al.</i>	(ALEPH Collab.)
BEHRENS	00	PR D61 052001	B.H. Behrens <i>et al.</i>	(CLEO Collab.)
BONVICINI	00	PRL 84 5940	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
CHEN	00	PRL 85 525	S. Chen <i>et al.</i>	(CLEO Collab.)
COAN	00	PRL 84 5283	T.E. Coan <i>et al.</i>	(CLEO Collab.)
CRONIN-HEN...	00	PRL 85 515	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)

CSORNA	00	PR D61 111101	S.E. Csorna <i>et al.</i>	(CLEO Collab.)
JESSOP	00	PRL 85 2881	C.P. Jessop <i>et al.</i>	(CLEO Collab.)
RICHICHI	00	PRL 85 520	S.J. Richichi <i>et al.</i>	(CLEO Collab.)
ABBIENDI	99J	EPJ C12 609	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
AFFOLDER	99B	PRL 83 3378	T. Affolder <i>et al.</i>	(CDF Collab.)
BARTELT	99	PRL 82 3746	J. Bartelt <i>et al.</i>	(CLEO Collab.)
COAN	99	PR D59 111101	T.E. Coan <i>et al.</i>	(CLEO Collab.)
ABE	98B	PR D57 5382	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	98O	PR D58 072001	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	98Q	PR D58 092002	F. Abe <i>et al.</i>	(CDF Collab.)
ACCIARRI	98S	PL B438 417	M. Acciarri <i>et al.</i>	(L3 Collab.)
ANASTASSOV	98	PRL 80 4127	A. Anastassov <i>et al.</i>	(CLEO Collab.)
ATHANAS	98	PRL 80 5493	M. Athanas <i>et al.</i>	(CLEO Collab.)
BARATE	98Q	EPJ C4 387	R. Barate <i>et al.</i>	(ALEPH Collab.)
BEHRENS	98	PRL 80 3710	B.H. Behrens <i>et al.</i>	(CLEO Collab.)
BERGFELD	98	PRL 81 272	T. Bergfeld <i>et al.</i>	(CLEO Collab.)
BRANDENB...	98	PRL 80 2762	G. Brandenbrug <i>et al.</i>	(CLEO Collab.)
CAPRINI	98	NP B530 153	I. Caprini, L. Lellouch, M. Neubert	(BCIP, CERN)
GODANG	98	PRL 80 3456	R. Godang <i>et al.</i>	(CLEO Collab.)
ABE	97J	PRL 79 590	K. Abe <i>et al.</i>	(SLD Collab.)
ACCIARRI	97F	PL B396 327	M. Acciarri <i>et al.</i>	(L3 Collab.)
ARTUSO	97	PL B399 321	M. Artuso <i>et al.</i>	(CLEO Collab.)
ATHANAS	97	PRL 79 2208	M. Athanas <i>et al.</i>	(CLEO Collab.)
BROWDER	97	PR D56 11	T. Browder <i>et al.</i>	(CLEO Collab.)
FU	97	PRL 79 3125	X. Fu <i>et al.</i>	(CLEO Collab.)
JESSOP	97	PRL 79 4533	C.P. Jessop <i>et al.</i>	(CLEO Collab.)
ABE	96B	PR D53 3496	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	96C	PRL 76 4462	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	96H	PRL 76 2015	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	96L	PRL 76 4675	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	96Q	PR D54 6596	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	96R	PRL 77 5176	F. Abe <i>et al.</i>	(CDF Collab.)
ADAM	96D	ZPHY C72 207	W. Adam <i>et al.</i>	(DELPHI Collab.)
ALEXANDER	96T	PRL 77 5000	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
ASNER	96	PR D53 1039	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BARISH	96B	PRL 76 1570	B.C. Barish <i>et al.</i>	(CLEO Collab.)
BERGFELD	96B	PRL 77 4503	T. Bergfeld <i>et al.</i>	(CLEO Collab.)
BISHAI	96	PL B369 186	M. Bishai <i>et al.</i>	(CLEO Collab.)
BUSKULIC	96J	ZPHY C71 31	D. Buskalic <i>et al.</i>	(ALEPH Collab.)
GIBAUT	96	PR D53 4734	R. Gibaut <i>et al.</i>	(CLEO Collab.)
PDG	96	PR D54 1	R. M. Barnett <i>et al.</i>	
ABREU	95N	PL B357 255	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	95Q	ZPHY C68 13	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ADAM	95	ZPHY C68 363	W. Adam <i>et al.</i>	(DELPHI Collab.)
AKERS	95T	ZPHY C67 379	R. Akers <i>et al.</i>	(OPAL Collab.)
ALBRECHT	95D	PL B353 554	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	95	PL B341 435	J. Alexander <i>et al.</i>	(CLEO Collab.)
Also		PL B347 469 (erratum)	J. Alexander <i>et al.</i>	(CLEO Collab.)
ARTUSO	95	PRL 75 785	M. Artuso <i>et al.</i>	(CLEO Collab.)
BARISH	95	PR D51 1014	B.C. Barish <i>et al.</i>	(CLEO Collab.)
BUSKULIC	95	PL B343 444	D. Buskalic <i>et al.</i>	(ALEPH Collab.)
ABE	94D	PRL 72 3456	F. Abe <i>et al.</i>	(CDF Collab.)
ALAM	94	PR D50 43	M.S. Alam <i>et al.</i>	(CLEO Collab.)
ALBRECHT	94D	PL B335 526	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ATHANAS	94	PRL 73 3503	M. Athanas <i>et al.</i>	(CLEO Collab.)
Also		PRL 74 3090 (erratum)	M. Athanas <i>et al.</i>	(CLEO Collab.)
PDG	94	PR D50 1173	L. Montanet <i>et al.</i>	(CERN, LBL, BOST+)
STONE	94	HEPSY 93-11	S. Stone	
Published in B Decays, 2nd Edition, World Scientific, Singapore				
ABREU	93D	ZPHY C57 181	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	93G	PL B312 253	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACTON	93C	PL B307 247	P.D. Acton <i>et al.</i>	(OPAL Collab.)
ALBRECHT	93E	ZPHY C60 11	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	93B	PL B319 365	J. Alexander <i>et al.</i>	(CLEO Collab.)
AMMAR	93	PRL 71 674	R. Ammar <i>et al.</i>	(CLEO Collab.)
BEAN	93B	PRL 70 2681	A. Bean <i>et al.</i>	(CLEO Collab.)
BUSKULIC	93D	PL B307 194	D. Buskalic <i>et al.</i>	(ALEPH Collab.)
Also		PL B325 537 (erratum)	D. Buskalic <i>et al.</i>	(ALEPH Collab.)
SANGHERA	93	PR D47 791	S. Sanghera <i>et al.</i>	(CLEO Collab.)
ALBRECHT	92C	PL B275 195	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	92E	PL B277 209	H. Albrecht <i>et al.</i>	(ARGUS Collab.)

ALBRECHT	92G	ZPHY C54 1	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BORTOLETTO	92	PR D45 21	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
BUSKULIC	92G	PL B295 396	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ALBRECHT	91B	PL B254 288	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	91C	PL B255 297	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	91E	PL B262 148	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BERKELMAN	91	ARNPS 41 1	K. Berkelman, S. Stone	(CORN, SYRA)
"Decays of <i>B</i> Mesons"				
FULTON	91	PR D43 651	R. Fulton <i>et al.</i>	(CLEO Collab.)
ALBRECHT	90B	PL B241 278	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	90J	ZPHY C48 543	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ANTREASYAN	90B	ZPHY C48 553	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)
BORTOLETTO	90	PRL 64 2117	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
Also		PR D45 21	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
WEIR	90B	PR D41 1384	A.J. Weir <i>et al.</i>	(Mark II Collab.)
ALBRECHT	89G	PL B229 304	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
AVERY	89B	PL B223 470	P. Avery <i>et al.</i>	(CLEO Collab.)
BEBEK	89	PRL 62 8	C. Bebek <i>et al.</i>	(CLEO Collab.)
BORTOLETTO	89	PRL 62 2436	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
ALBRECHT	88F	PL B209 119	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	88K	PL B215 424	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	87C	PL B185 218	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	87D	PL B199 451	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
AVERY	87	PL B183 429	P. Avery <i>et al.</i>	(CLEO Collab.)
BEBEK	87	PR D36 1289	C. Bebek <i>et al.</i>	(CLEO Collab.)
ALAM	86	PR D34 3279	M.S. Alam <i>et al.</i>	(CLEO Collab.)
PDG	86	PL 170B 1	M. Aguilar-Benitez <i>et al.</i>	(CERN, CIT+)
GILES	84	PR D30 2279	R. Giles <i>et al.</i>	(CLEO Collab.)
