

$\Delta(1900)$ S_{31} $I(J^P) = \frac{3}{2}(\frac{1}{2}^-)$ Status: $\ast\ast$

OMITTED FROM SUMMARY TABLE

Some obsolete results published before 1980 were last included in our 2006 edition, Journal of Physics, G **33** 1 (2006). Some further obsolete results published before 1984 were last included in our 2006 edition, Journal of Physics, G **33** 1 (2006).

The latest GWU analysis (ARNDT 06) finds no evidence for this resonance.

 $\Delta(1900)$ BREIT-WIGNER MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1850 to 1950 (≈ 1900) OUR ESTIMATE			
1920 ± 24	MANLEY 92	IPWA	$\pi N \rightarrow \pi N & N\pi\pi$
1890 ± 50	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
1908 ± 30	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1802 ± 87	VRANA 00	DPWA	Multichannel
1918.5 ± 23.0	CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$

 $\Delta(1900)$ BREIT-WIGNER WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
140 to 240 (≈ 200) OUR ESTIMATE			
263 ± 39	MANLEY 92	IPWA	$\pi N \rightarrow \pi N & N\pi\pi$
170 ± 50	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
140 ± 40	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
48 ± 45	VRANA 00	DPWA	Multichannel
93.5 ± 54.0	CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$

 $\Delta(1900)$ POLE POSITION**REAL PART**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1780	¹ HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
1870 ± 40	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1795	VRANA 00	DPWA	Multichannel
not seen	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
2029 or 2025	² LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$

-2×IMAGINARY PART

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
180±50	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
58	VRANA 00	DPWA	Multichannel
not seen	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
164 or 163	² LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$

Δ(1900) ELASTIC POLE RESIDUE

MODULUS |r|

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
10±3	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ

<u>VALUE (°)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
+20±40	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

Δ(1900) DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 N\pi$	10–30 %
$\Gamma_2 \Sigma K$	
$\Gamma_3 N\pi\pi$	
$\Gamma_4 \Delta\pi$	
$\Gamma_5 \Delta(1232)\pi$, D-wave	
$\Gamma_6 N\rho$	
$\Gamma_7 N\rho$, S=1/2, S-wave	
$\Gamma_8 N\rho$, S=3/2, D-wave	
$\Gamma_9 N(1440)\pi$, S-wave	
$\Gamma_{10} N\gamma$, helicity=1/2	

Δ(1900) BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.1 to 0.3 OUR ESTIMATE			
0.41±0.04	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
0.10±0.03	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
0.08±0.04	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.33±0.10	VRANA 00	DPWA	Multichannel
0.28	CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1900) \rightarrow \Sigma K$	$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
<0.03	CANDLIN 84 DPWA $\pi^+ p \rightarrow \Sigma^+ K^+$
$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1900) \rightarrow \Delta(1232)\pi, D\text{-wave}$	$(\Gamma_1 \Gamma_5)^{1/2} / \Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
+0.25±0.07	MANLEY 92 IPWA $\pi N \rightarrow \pi N \& N\pi\pi$
$\Gamma(\Delta(1232)\pi, D\text{-wave}) / \Gamma_{\text{total}}$	Γ_5 / Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.28±0.01	VRANA 00 DPWA Multichannel
$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1900) \rightarrow N\rho, S=1/2, S\text{-wave}$	$(\Gamma_1 \Gamma_7)^{1/2} / \Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
-0.14±0.11	MANLEY 92 IPWA $\pi N \rightarrow \pi N \& N\pi\pi$
$\Gamma(N\rho, S=1/2, S\text{-wave}) / \Gamma_{\text{total}}$	Γ_7 / Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.30±0.02	VRANA 00 DPWA Multichannel
$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1900) \rightarrow N\rho, S=3/2, D\text{-wave}$	$(\Gamma_1 \Gamma_8)^{1/2} / \Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
-0.37±0.07	MANLEY 92 IPWA $\pi N \rightarrow \pi N \& N\pi\pi$
$\Gamma(N\rho, S=3/2, D\text{-wave}) / \Gamma_{\text{total}}$	Γ_8 / Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.05±0.01	VRANA 00 DPWA Multichannel
$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1900) \rightarrow N(1440)\pi, S\text{-wave}$	$(\Gamma_1 \Gamma_9)^{1/2} / \Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
-0.16±0.11	MANLEY 92 IPWA $\pi N \rightarrow \pi N \& N\pi\pi$
$\Gamma(N(1440)\pi, S\text{-wave}) / \Gamma_{\text{total}}$	Γ_9 / Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.04±0.01	VRANA 00 DPWA Multichannel

$\Delta(1900)$ PHOTON DECAY AMPLITUDES

Papers on γN amplitudes predating 1981 may be found in our 2006 edition,
Journal of Physics, G **33** 1 (2006).

$\Delta(1900) \rightarrow N\gamma$, helicity-1/2 amplitude $A_{1/2}$

<u>VALUE (GeV^{-1/2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.004±0.016	CRAWFORD 83	IPWA	$\gamma N \rightarrow \pi N$
0.029±0.008	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$

Δ(1900) FOOTNOTES

- ¹ See HOEHLER 93 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.
² LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.

Δ(1900) REFERENCES

For early references, see Physics Letters **111B** 1 (1982).

ARNDT	06	PR C74 045205	R.A. Arndt <i>et al.</i>	(GWU)
PDG	06	JPG 33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)
VRANA	00	PRPL 328 181	T.P. Vrana, S.A. Dytman,, T.-S.H. Lee	(PITT+)
HOEHLER	93	πN Newsletter 9 1	G. Hohler	(KARL)
MANLEY	92	PR D45 4002	D.M. Manley, E.M. Saleski	(KENT) IJP
Also		PR D30 904	D.M. Manley <i>et al.</i>	(VPI)
ARNDT	91	PR D43 2131	R.A. Arndt <i>et al.</i>	(VPI, TELE) IJP
CANDLIN	84	NP B238 477	D.J. Candlin <i>et al.</i>	(EDIN, RAL, LOWC)
CRAWFORD	83	NP B211 1	R.L. Crawford, W.T. Morton	(GLAS)
AWAJI	81	Bonn Conf. 352	N. Awaji, R. Kajikawa	(NAGO)
Also		NP B197 365	K. Fujii <i>et al.</i>	(NAGO)
CHEW	80	Toronto Conf. 123	D.M. Chew	(LBL) IJP
CUTKOSKY	80	Toronto Conf. 19	R.E. Cutkosky <i>et al.</i>	(CMU, LBL) IJP
Also		PR D20 2839	R.E. Cutkosky <i>et al.</i>	(CMU, LBL) IJP
HOEHLER	79	PDAT 12-1	G. Hohler <i>et al.</i>	(KARLT) IJP
Also		Toronto Conf. 3	R. Koch	(KARLT) IJP
LONGACRE	78	PR D17 1795	R.S. Longacre <i>et al.</i>	(LBL, SLAC)
