

$K(1630)$

$I(J^P) = \frac{1}{2}(? ?)$

OMMITTED FROM SUMMARY TABLE

Seen as a narrow peak, compatible with the experimental resolution, in the invariant mass of the $K_S^0\pi^+\pi^-$ system produced in $\pi^- p$ interactions at high momentum transfers.

$K(1630)$ MASS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1629 ± 7	~ 75	KARNAUKHOV98	BC	$16.0 \pi^- p \rightarrow (K_S^0\pi^+\pi^-) X^+\pi^- X^0$

$K(1630)$ WIDTH

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
16^{+19}_{-16}	~ 75	¹ KARNAUKHOV98	BC	$16.0 \pi^- p \rightarrow (K_S^0\pi^+\pi^-) X^+\pi^- X^0$

¹ Compatible with an experimental resolution of 14 ± 1 MeV.

$K(1630)$ DECAY MODES

Mode
$\Gamma_1 \quad K_S^0\pi^+\pi^-$

$K(1630)$ REFERENCES

KARNAUKHOV 98 PAN 61 203 V.M. Karnaughov, C. Coca, V.I. Moroz
Translated from YAF 61 252.

— OTHER RELATED PAPERS —

KARNAUKHOV 00 PAN 63 588 V.M. Karnaughov, C. Coca, V.I. Moroz
Translated from YAF 63 652.