

**$f_2(1565)$**

$I^G(J^{PC}) = 0^+(2^{++})$

## OMMITTED FROM SUMMARY TABLE

Seen in antinucleon-nucleon annihilation at rest. Needs confirmation.

---

### **$f_2(1565)$ MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b><math>1562 \pm 13</math> OUR AVERAGE</b>	Error includes scale factor of 2.1. See the ideogram below.		
1590 $\pm$ 10	<sup>1</sup> AMELIN 06	VES	$36 \pi^- p \rightarrow \omega\omega n$
1552 $\pm$ 13	<sup>2</sup> AMSLER 02	CBAR	$0.9 \bar{p}p \rightarrow \pi^0 \eta\eta, \pi^0 \pi^0 \pi^0$
1550 $\pm 10 \pm 20$	AMELIN 00	VES	$37 \pi^- p \rightarrow \eta\pi^+ \pi^- n$
1575 $\pm$ 18	BERTIN 98	OBLX	$0.05-0.405 \bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$
1507 $\pm$ 15	<sup>2</sup> BERTIN 97C	OBLX	$0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
1565 $\pm$ 20	MAY 90	ASTE	$0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>			
1598 $\pm 11 \pm 9$	BAKER 99B	SPEC	$0 \bar{p}p \rightarrow \omega\omega\pi^0$
1534 $\pm$ 20	<sup>3</sup> ABELE 96C	RVUE	Compilation
$\sim 1552$	<sup>4</sup> AMSLER 95D	CBAR	$0.0 \bar{p}p \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta\eta, \pi^0 \pi^0 \eta$
1598 $\pm 72$	BALOSHIN 95	SPEC	$40 \pi^- C \rightarrow K_S^0 K_S^0 X$
$1566^{+80}_{-50}$	<sup>5</sup> ANISOVICH 94	CBAR	$0.0 \bar{p}p \rightarrow 3\pi^0, \eta\eta\pi^0$
1502 $\pm 9$	ADAMO 93	OBLX	$\bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$
1488 $\pm 10$	<sup>6</sup> ARMSTRONG 93C	E760	$\bar{p}p \rightarrow \pi^0 \eta\eta \rightarrow 6\gamma$
1508 $\pm 10$	<sup>6</sup> ARMSTRONG 93D	E760	$\bar{p}p \rightarrow 3\pi^0 \rightarrow 6\gamma$
1525 $\pm 10$	<sup>6</sup> ARMSTRONG 93D	E760	$\bar{p}p \rightarrow \eta\pi^0 \pi^0 \rightarrow 6\gamma$
$\sim 1504$	<sup>7</sup> WEIDENAUER 93	ASTE	$0.0 \bar{p}N \rightarrow 3\pi^- 2\pi^+$
1540 $\pm 15$	<sup>6</sup> ADAMO 92	OBLX	$\bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$
1515 $\pm 10$	<sup>8</sup> AKER 91	CBAR	$0.0 \bar{p}p \rightarrow 3\pi^0$
1477 $\pm 5$	BRIDGES 86C	DBC	$0.0 \bar{p}N \rightarrow 3\pi^- 2\pi^+$

<sup>1</sup> Supersedes the  $\omega\omega$  state of BELADIDZE 92B earlier assigned to the  $f_2(1640)$ .

<sup>2</sup> T-matrix pole.

<sup>3</sup> T-matrix pole, large coupling to  $\rho\rho$  and  $\omega\omega$ , could be  $f_2(1640)$ .


<sup>4</sup> Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.

<sup>5</sup> From a simultaneous analysis of the annihilations  $\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta\eta$  including AKER 91 data.

<sup>6</sup>  $J^P$  not determined, could be partly  $f_0(1500)$ .

<sup>7</sup>  $J^P$  not determined.

<sup>8</sup> Superseded by AMSLER 95B.



### f<sub>2</sub>(1565) WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>134± 8 OUR AVERAGE</b>			
140± 11	<sup>9</sup> AMELIN	06 VES	$36 \pi^- p \rightarrow \omega\omega n$
113± 23	<sup>10</sup> AMSLER	02 CBAR	$0.9 \bar{p}p \rightarrow \pi^0 \eta\eta, \pi^0 \pi^0 \pi^0$
130± 20±40	AMELIN	00 VES	$37 \pi^- p \rightarrow \eta\pi^+ \pi^- n$
119± 24	BERTIN	98 OBLX	$0.05-0.405 \bar{p}p \rightarrow \pi^+ \pi^+ \pi^-$
130± 20	<sup>10</sup> BERTIN	97C OBLX	$0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
170± 40	MAY	90 ASTE	$0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
180± 60	<sup>11</sup> ABELE	96C RVUE	Compilation
~ 142	<sup>12</sup> AMSLER	95D CBAR	$0.0 \bar{p}p \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta\eta,$ $\pi^0 \pi^0 \eta$
263±101	BALOSHIN	95 SPEC	$40 \pi^- C \rightarrow K_S^0 K_S^0 X$
166 <sup>+ 80</sup> <sub>- 20</sub>	<sup>13</sup> ANISOVICH	94 CBAR	$0.0 \bar{p}p \rightarrow 3\pi^0, \eta\eta\pi^0$
130± 10	ADAMO	93 OBLX	$\bar{p}p \rightarrow \pi^+ \pi^+ \pi^-$
148± 27	<sup>15</sup> ARMSTRONG	E760	$\bar{p}p \rightarrow \pi^0 \eta\eta \rightarrow 6\gamma$
103± 15	<sup>15</sup> ARMSTRONG	E760	$\bar{p}p \rightarrow 3\pi^0 \rightarrow 6\gamma$
111± 10	<sup>15</sup> ARMSTRONG	E760	$\bar{p}p \rightarrow \eta\pi^0 \pi^0 \rightarrow 6\gamma$
~ 206	<sup>16</sup> WEIDENAUER	93 ASTE	$0.0 \bar{p}N \rightarrow 3\pi^- 2\pi^+$
132± 37	ADAMO	92 OBLX	$\bar{p}p \rightarrow \pi^+ \pi^+ \pi^-$
120± 10	<sup>17</sup> AKER	91 CBAR	$0.0 \bar{p}p \rightarrow 3\pi^0$
116± 9	BRIDGES	86C DBC	$0.0 \bar{p}N \rightarrow 3\pi^- 2\pi^+$

- <sup>9</sup> Supersedes the  $\omega\omega$  state of BELADIDZE 92B earlier assigned to the  $f_2(1640)$ .  
<sup>10</sup> T-matrix pole.  
<sup>11</sup> T-matrix pole, large coupling to  $\rho\rho$  and  $\omega\omega$ , could be  $f_2(1640)$ .  
<sup>12</sup> Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.  
<sup>13</sup> From a simultaneous analysis of the annihilations  $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta\eta$  including AKER 91 data.  
<sup>14</sup> Supersedes ADAMO 92.  
<sup>15</sup>  $J^P$  not determined, could be partly  $f_0(1500)$ .  
<sup>16</sup>  $J^P$  not determined.  
<sup>17</sup> Superseded by AMSLER 95B.
- 

## $f_2(1565)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \pi\pi$	seen
$\Gamma_2 \pi^+\pi^-$	seen
$\Gamma_3 \pi^0\pi^0$	seen
$\Gamma_4 \rho^0\rho^0$	seen
$\Gamma_5 2\pi^+ 2\pi^-$	seen
$\Gamma_6 \eta\eta$	seen
$\Gamma_7 a_2(1320)\pi$	
$\Gamma_8 \omega\omega$	seen
$\Gamma_9 K\bar{K}$	
$\Gamma_{10} \gamma\gamma$	

---

## $f_2(1565)$ PARTIAL WIDTHS

### $\Gamma(\eta\eta)$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_6$
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>					
$1.2 \pm 0.3$	870	<sup>18</sup> SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$	

### $\Gamma(K\bar{K})$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_9$
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>					
$2.0 \pm 1.0$	870	<sup>18</sup> SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$	

### $\Gamma(\gamma\gamma)$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{10}$
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>					
$0.70 \pm 0.14$	870	<sup>18</sup> SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$	

<sup>18</sup> From analysis of L3 data at 91 and 183–209 GeV, using  $f_2(1565)$  mass of 1570 MeV, width of 160 MeV,  $\Gamma(\pi\pi) = 25$  MeV, and SU(3) relations.

---

**$f_2(1565)$  BRANCHING RATIOS** **$\Gamma(\pi\pi)/\Gamma_{\text{total}}$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<b><math>\Gamma_1/\Gamma</math></b>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
seen	BAKER 99B	SPEC	$0 \bar{p}p \rightarrow \omega\omega\pi^0$	

 **$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<b><math>\Gamma_2/\Gamma</math></b>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
seen	BERTIN 98	OBLX	$0.05\text{--}0.405 \bar{p}p \rightarrow \pi^+\pi^+\pi^-$	
not seen	19 ANISOVICH 94B	RVUE	$\bar{p}p \rightarrow \pi^+\pi^-\pi^0$	
seen	MAY 89	ASTE	$\bar{p}p \rightarrow \pi^+\pi^-\pi^0$	

19 ANISOVICH 94B is from a reanalysis of MAY 90.

 **$\Gamma(\pi^0\pi^0)/\Gamma_{\text{total}}$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<b><math>\Gamma_3/\Gamma</math></b>
seen	AMSLER 95B	CBAR	$0.0 \bar{p}p \rightarrow 3\pi^0$	

 **$\Gamma(\pi^+\pi^-)/\Gamma(\rho^0\rho^0)$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<b><math>\Gamma_2/\Gamma_4</math></b>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
$0.042 \pm 0.013$	BRIDGES 86B	DBC	$\bar{p}N \rightarrow 3\pi^- 2\pi^+$	

 **$\Gamma(\eta\eta)/\Gamma(\pi^0\pi^0)$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<b><math>\Gamma_6/\Gamma_3</math></b>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
$0.024 \pm 0.005 \pm 0.012$	20 ARMSTRONG 93C	E760	$\bar{p}p \rightarrow \pi^0\eta\eta \rightarrow 6\gamma$	

20  $J^P$  not determined, could be partly  $f_0(1500)$ . **$\Gamma(\omega\omega)/\Gamma_{\text{total}}$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<b><math>\Gamma_8/\Gamma</math></b>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
seen	BAKER 99B	SPEC	$0 \bar{p}p \rightarrow \omega\omega\pi^0$	

 **$f_2(1565)$  REFERENCES**

AMELIN 06	PAN 69 690	D.V. Amelin <i>et al.</i>	(VES Collab.)
SCHEGELSKY 06A	Translated from YAF 69 715. EPJ A27 207	V.A. Schegelsky <i>et al.</i>	
AMSLER 02	EPJ C23 29	C. Amsler <i>et al.</i>	
AMELIN 00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)
BAKER 99B	PL B467 147	C.A. Baker <i>et al.</i>	
BERTIN 98	PR D57 55	A. Bertin <i>et al.</i>	(OBELIX Collab.)
BERTIN 97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)
ABELE 96C	NP A609 562	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER 95B	PL B342 433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER 95C	PL B353 571	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER 95D	PL B355 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
BALOSHIN 95	PAN 58 46	O.N. Baloshin <i>et al.</i>	(ITEP)
	Translated from YAF 58 50.		

AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ANISOVICH	94	PL B323 233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.)
ANISOVICH	94B	PR D50 1972	V.V. Anisovich <i>et al.</i>	(LOQM)
ADAMO	93	NP A558 13C	A. Adamo <i>et al.</i>	(OBELIX Collab.)
ARMSTRONG	93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
ARMSTRONG	93D	PL B307 399	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
WEIDENAUER	93	ZPHY C59 387	P. Weidenauer <i>et al.</i>	(ASTERIX Collab.)
ADAMO	92	PL B287 368	A. Adamo <i>et al.</i>	(OBELIX Collab.)
BELADIDZE	92B	ZPHY C54 367	G.M. Beladidze <i>et al.</i>	(VES Collab.)
AKER	91	PL B260 249	E. Aker <i>et al.</i>	(Crystal Barrel Collab.)
MAY	90	ZPHY C46 203	B. May <i>et al.</i>	(ASTERIX Collab.)
MAY	89	PL B225 450	B. May <i>et al.</i>	(ASTERIX Collab.) IJP
BRIDGES	86B	PRL 56 215	D.L. Bridges <i>et al.</i>	(SYRA, CASE)
BRIDGES	86C	PRL 57 1534	D.L. Bridges <i>et al.</i>	(SYRA)

## OTHER RELATED PAPERS

BUGG	07	EPJ C52 55	D. Bugg
ANISOVICH	05	JETPL 80 715	V.V. Anisovich

Translated from ZETFP 80 845.