

$\psi(4415)$

$$J^{PC} = 0^-(1^{--})$$

$\psi(4415)$ MASS

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|-----------------------|----------|--|
| 4421 ± 4 | OUR ESTIMATE | | |
| 4415.1 ± 7.9 | ¹ ABLIKIM | 08D BES2 | $e^+e^- \rightarrow$ hadrons |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 4411 ± 7 | ² PAKHLOVA | 08A BELL | $10.6 e^+e^- \rightarrow D^0 D^- \pi^+ \gamma$ |
| 4425 ± 6 | ³ SETH | 05A RVUE | $e^+e^- \rightarrow$ hadrons |
| 4429 ± 9 | ⁴ SETH | 05A RVUE | $e^+e^- \rightarrow$ hadrons |
| 4417 ± 10 | BRANDELIK | 78C DASP | e^+e^- |
| 4414 ± 7 | SIEGRIST | 76 MRK1 | e^+e^- |

¹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (234 \pm 88)^\circ$.

² Systematic uncertainties not estimated.

³ From a fit to Crystal Ball (OSTERHELD 86) data.

⁴ From a fit to BES (BAI 02C) data.

$\psi(4415)$ WIDTH

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|-----------------------|----------|--|
| 62 ± 20 | OUR ESTIMATE | | |
| 71.5 ± 19.0 | ⁵ ABLIKIM | 08D BES2 | $e^+e^- \rightarrow$ hadrons |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 77 ± 20 | ⁶ PAKHLOVA | 08A BELL | $10.6 e^+e^- \rightarrow D^0 D^- \pi^+ \gamma$ |
| 119 ± 16 | ⁷ SETH | 05A RVUE | $e^+e^- \rightarrow$ hadrons |
| 118 ± 35 | ⁸ SETH | 05A RVUE | $e^+e^- \rightarrow$ hadrons |
| 66 ± 15 | BRANDELIK | 78C DASP | e^+e^- |
| 33 ± 10 | SIEGRIST | 76 MRK1 | e^+e^- |

⁵ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (234 \pm 88)^\circ$.

⁶ Systematic uncertainties not estimated.

⁷ From a fit to Crystal Ball (OSTERHELD 86) data.

⁸ From a fit to BES (BAI 02C) data.

$\psi(4415)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) | Confidence level |
|---|--------------------------------|------------------|
| Γ_1 hadrons | dominant | |
| Γ_2 $D^0 D^- \pi^+$ | | |
| Γ_3 $(D^0 D^- \pi^+)_{non-res}$ | < 2.3 % | 90% |
| Γ_4 $D\bar{D}_2^*(2460) \rightarrow D^0 D^- \pi^+$ | (10 ± 4) % | |
| Γ_5 e^+e^- | (9.4 ± 3.2) × 10 ⁻⁶ | |

$\psi(4415)$ PARTIAL WIDTHS **$\Gamma(e^+e^-)$ Γ_5**

| <u>VALUE (keV)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------|--------------------|-------------|----------------|
|--------------------|--------------------|-------------|----------------|

0.58±0.07 OUR ESTIMATE**0.35±0.12** ⁹ ABLIKIM 08D BES2 $e^+e^- \rightarrow$ hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.72±0.11 ¹⁰ SETH 05A RVUE $e^+e^- \rightarrow$ hadrons0.64±0.23 ¹¹ SETH 05A RVUE $e^+e^- \rightarrow$ hadrons0.49±0.13 BRANDELIK 78C DASP e^+e^- 0.44±0.14 SIEGRIST 76 MRK1 e^+e^- ⁹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (234 \pm 88)^\circ$.¹⁰ From a fit to Crystal Ball (OSTERHELD 86) data.¹¹ From a fit to BES (BAI 02C) data. **$\psi(4415)$ BRANCHING RATIOS** **$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$ Γ_1/Γ**

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|--------------------|-------------|----------------|
|--------------|--------------------|-------------|----------------|

dominant SIEGRIST 76 MRK1 e^+e^- **$\Gamma(D\bar{D}_2^*(2460) \rightarrow D^0 D^- \pi^+)/\Gamma_{\text{total}}$ Γ_4/Γ**

| <u>VALUE (units 10^{-2})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|----------------|
|---|--------------------|-------------|----------------|

10.5±2.4±3.8 ¹² PAKHLOVA 08A BELL 10.6 $e^+e^- \rightarrow D^0 D^- \pi^+ \gamma$ ¹² Using 4421 ± 4 MeV for the mass and 62 ± 20 MeV for the width of $\psi(4415)$. **$\Gamma((D^0 D^- \pi^+)_{\text{non-res}})/\Gamma(D\bar{D}_2^*(2460) \rightarrow D^0 D^- \pi^+)$ Γ_3/Γ_4**

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|------------|--------------------|-------------|----------------|
|--------------|------------|--------------------|-------------|----------------|

<0.22 90 ¹³ PAKHLOVA 08A BELL 10.6 $e^+e^- \rightarrow D^0 D^- \pi^+ \gamma$ ¹³ Using 4421 ± 4 MeV for the mass and 62 ± 20 MeV for the width of $\psi(4415)$. **$\psi(4415)$ REFERENCES**

| | | | | |
|-----------|-----|----------------|-----------------------------|-----------------------------|
| ABLIKIM | 08D | PL B660 315 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| PAKHLOVA | 08A | PRL 100 062001 | G. Pakhlova <i>et al.</i> | (BELLE Collab.) |
| SETH | 05A | PR D72 017501 | K.K. Seth | |
| BAI | 02C | PRL 88 101802 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| OSTERHELD | 86 | SLAC-PUB-4160 | A. Osterheld <i>et al.</i> | (SLAC Crystal Ball Collab.) |
| BRANDELIK | 78C | PL 76B 361 | R. Brandelik <i>et al.</i> | (DASP Collab.) |
| SIEGRIST | 76 | PRL 36 700 | J.L. Siegrist <i>et al.</i> | (LBL, SLAC) |

OTHER RELATED PAPERS

| | | | | |
|-----------|----|----------------|----------------------------|---------------------|
| PAKHLOVA | 08 | PR D77 011103R | G. Pakhlova <i>et al.</i> | (BELLE Collab.) |
| PAKHLOVA | 07 | PRL 98 092001 | G. Pakhlova <i>et al.</i> | (BELLE Collab.) |
| BURMESTER | 77 | PL 66B 395 | J. Burmester <i>et al.</i> | (DESY, HAMB, SIEG+) |
| LUTH | 77 | PL 70B 120 | V. Luth <i>et al.</i> | (LBL, SLAC) |