

$\chi_{c2}(1P)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

See the Review on “ $\psi(2S)$ and χ_c branching ratios” before the $\chi_{c0}(1P)$ Listings.

$\chi_{c2}(1P)$ MASS

| <u>VALUE (MeV)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------------|-------------|--|
| 3556.20 ± 0.09 OUR AVERAGE | | | | |
| 3555.3 ± 0.6 ± 2.2 | 2.5k | UEHARA | 08 BELL | $\gamma\gamma \rightarrow$ hadrons |
| 3555.70 ± 0.59 ± 0.39 | | ABLIKIM | 05G BES2 | $\psi(2S) \rightarrow \gamma\chi_{c2}$ |
| 3556.173 ± 0.123 ± 0.020 | | ANDREOTTI | 05A E835 | $p\bar{p} \rightarrow e^+e^-\gamma$ |
| 3559.9 ± 2.9 | | EISENSTEIN | 01 CLE2 | $e^+e^- \rightarrow e^+e^-\chi_{c2}$ |
| 3556.4 ± 0.7 | | BAI | 99B BES | $\psi(2S) \rightarrow \gamma X$ |
| 3556.22 ± 0.131 ± 0.020 | 585 | ¹ ARMSTRONG | 92 E760 | $\bar{p}p \rightarrow e^+e^-\gamma$ |
| 3556.9 ± 0.4 ± 0.5 | 50 | BAGLIN | 86B SPEC | $\bar{p}p \rightarrow e^+e^- X$ |
| 3557.8 ± 0.2 ± 4 | | ² GAISER | 86 CBAL | $\psi(2S) \rightarrow \gamma X$ |
| 3553.4 ± 2.2 | 66 | ³ LEMOIGNE | 82 GOLI | $185 \pi^- \text{Be} \rightarrow \gamma\mu^+\mu^- A$ |
| 3555.9 ± 0.7 | | ⁴ OREGLIA | 82 CBAL | $e^+e^- \rightarrow J/\psi 2\gamma$ |
| 3557 ± 1.5 | 69 | ⁵ HIMEL | 80 MRK2 | $e^+e^- \rightarrow J/\psi 2\gamma$ |
| 3551 ± 11 | 15 | BRANDELIK | 79B DASP | $e^+e^- \rightarrow J/\psi 2\gamma$ |
| 3553 ± 4 | | ⁵ BARTEL | 78B CNTR | $e^+e^- \rightarrow J/\psi 2\gamma$ |
| 3553 ± 4 ± 4 | | ^{5,6} TANENBAUM | 78 MRK1 | e^+e^- |
| 3563 ± 7 | 360 | ⁵ BIDDICK | 77 CNTR | $e^+e^- \rightarrow \gamma X$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 3543 ± 10 | 4 | WHITAKER | 76 MRK1 | $e^+e^- \rightarrow J/\psi 2\gamma$ |

¹ Recalculated by ANDREOTTI 05A, using the value of $\psi(2S)$ mass from AULCHENKO 03.

² Using mass of $\psi(2S) = 3686.0$ MeV.

³ $J/\psi(1S)$ mass constrained to 3097 MeV.

⁴ Assuming $\psi(2S)$ mass = 3686 MeV and $J/\psi(1S)$ mass = 3097 MeV.

⁵ Mass value shifted by us by amount appropriate for $\psi(2S)$ mass = 3686 MeV and $J/\psi(1S)$ mass = 3097 MeV.

⁶ From a simultaneous fit to radiative and hadronic decay channels.

$\chi_{c2}(1P)$ WIDTH

| <u>VALUE (MeV)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------------|-------------|------------------------|-------------|-------------------------------------|
| 1.97 ± 0.11 OUR FIT | | | | |
| 1.95 ± 0.13 OUR AVERAGE | | | | |
| 1.915 ± 0.188 ± 0.013 | | ANDREOTTI | 05A E835 | $p\bar{p} \rightarrow e^+e^-\gamma$ |
| 1.96 ± 0.17 ± 0.07 | 585 | ⁷ ARMSTRONG | 92 E760 | $\bar{p}p \rightarrow e^+e^-\gamma$ |
| 2.6 ^{+1.4} / _{-1.0} | 50 | BAGLIN | 86B SPEC | $\bar{p}p \rightarrow e^+e^- X$ |
| 2.8 ^{+2.1} / _{-2.0} | | ⁸ GAISER | 86 CBAL | $\psi(2S) \rightarrow \gamma X$ |

⁷ Recalculated by ANDREOTTI 05A.

⁸ Errors correspond to 90% confidence level; authors give only width range.

$\chi_{c2}(1P)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) | Confidence level | |
|------------------------|--|---------------------------------|-----|
| Hadronic decays | | | |
| Γ_1 | $2(\pi^+\pi^-)$ | (1.11±0.11) % | |
| Γ_2 | $\rho\rho$ | | |
| Γ_3 | $\pi^+\pi^-\pi^0\pi^0$ | (2.00±0.26) % | |
| Γ_4 | $\rho^+\pi^-\pi^0 + \text{c.c.}$ | (2.4 ±0.4) % | |
| Γ_5 | $K^+K^-\pi^0\pi^0$ | (2.2 ±0.4) × 10 ⁻³ | |
| Γ_6 | $K^+\pi^-K^0\pi^0 + \text{c.c.}$ | (1.50±0.22) % | |
| Γ_7 | $\rho^+K^-K^0 + \text{c.c.}$ | (4.5 ±1.4) × 10 ⁻³ | |
| Γ_8 | $K^*(892)^0K^+\pi^- \rightarrow$ $K^+\pi^-K^0\pi^0 + \text{c.c.}$ | (3.2 ±0.9) × 10 ⁻³ | |
| Γ_9 | $K^*(892)^0K^0\pi^0 \rightarrow$ $K^+\pi^-K^0\pi^0 + \text{c.c.}$ | (4.2 ±0.9) × 10 ⁻³ | |
| Γ_{10} | $K^*(892)^-K^+\pi^0 \rightarrow$ $K^+\pi^-K^0\pi^0 + \text{c.c.}$ | (4.1 ±0.9) × 10 ⁻³ | |
| Γ_{11} | $K^*(892)^+K^0\pi^- \rightarrow$ $K^+\pi^-K^0\pi^0 + \text{c.c.}$ | (3.2 ±0.9) × 10 ⁻³ | |
| Γ_{12} | $K^+K^-\eta\pi^0$ | (1.4 ±0.5) × 10 ⁻³ | |
| Γ_{13} | $\pi^+\pi^-K^+K^-$ | (9.2 ±1.1) × 10 ⁻³ | |
| Γ_{14} | $K^+\bar{K}^*(892)^0\pi^- + \text{c.c.}$ | (2.3 ±1.2) × 10 ⁻³ | |
| Γ_{15} | $K^*(892)^0\bar{K}^*(892)^0$ | (2.5 ±0.5) × 10 ⁻³ | |
| Γ_{16} | $3(\pi^+\pi^-)$ | (8.6 ±1.8) × 10 ⁻³ | |
| Γ_{17} | $\phi\phi$ | (1.48±0.28) × 10 ⁻³ | |
| Γ_{18} | $\omega\omega$ | (1.9 ±0.6) × 10 ⁻³ | |
| Γ_{19} | $\pi\pi$ | (2.39±0.14) × 10 ⁻³ | |
| Γ_{20} | $\rho^0\pi^+\pi^-$ | (4.0 ±1.7) × 10 ⁻³ | |
| Γ_{21} | $\pi^+\pi^-\eta$ | (5.2 ±1.4) × 10 ⁻⁴ | |
| Γ_{22} | $\pi^+\pi^-\eta'$ | (5.4 ±2.0) × 10 ⁻⁴ | |
| Γ_{23} | $\eta\eta$ | (5.4 ±0.8) × 10 ⁻⁴ | |
| Γ_{24} | K^+K^- | (1.09±0.08) × 10 ⁻³ | |
| Γ_{25} | $K_S^0K_S^0$ | (5.8 ±0.5) × 10 ⁻⁴ | |
| Γ_{26} | $\bar{K}^0K^+\pi^- + \text{c.c.}$ | (1.32±0.20) × 10 ⁻³ | |
| Γ_{27} | $K^+K^-\pi^0$ | (3.3 ±0.8) × 10 ⁻⁴ | |
| Γ_{28} | $K^+K^-\eta$ | < 3.5 × 10 ⁻⁴ | 90% |
| Γ_{29} | $\eta\eta'$ | < 6 × 10 ⁻⁵ | 90% |
| Γ_{30} | $\eta'\eta'$ | < 1.1 × 10 ⁻⁴ | 90% |
| Γ_{31} | $\pi^+\pi^-K_S^0K_S^0$ | (2.4 ±0.6) × 10 ⁻³ | |
| Γ_{32} | $K^+K^-K_S^0K_S^0$ | < 4 × 10 ⁻⁴ | 90% |
| Γ_{33} | $K^+K^-K^+K^-$ | (1.78±0.22) × 10 ⁻³ | |
| Γ_{34} | $K^+K^-\phi$ | (1.55±0.32) × 10 ⁻³ | |
| Γ_{35} | $K_S^0K_S^0\rho\bar{\rho}$ | < 7.9 × 10 ⁻⁴ | 90% |
| Γ_{36} | $\rho\bar{\rho}$ | (7.2 ±0.4) × 10 ⁻⁵ | |

| | | | |
|---------------|-----------------------------------|----------------------------------|-----|
| Γ_{37} | $p\bar{p}\pi^0$ | $(4.7 \pm 1.0) \times 10^{-4}$ | |
| Γ_{38} | $p\bar{p}\eta$ | $(2.0 \pm 0.8) \times 10^{-4}$ | |
| Γ_{39} | $\pi^+\pi^-\rho\bar{\rho}$ | $(1.32 \pm 0.34) \times 10^{-3}$ | |
| Γ_{40} | $\pi^0\pi^0\rho\bar{\rho}$ | $(8.5 \pm 2.6) \times 10^{-4}$ | |
| Γ_{41} | $p\bar{n}\pi^-$ | $(1.1 \pm 0.4) \times 10^{-3}$ | |
| Γ_{42} | $\Lambda\bar{\Lambda}$ | $(1.86 \pm 0.27) \times 10^{-4}$ | |
| Γ_{43} | $\Lambda\bar{\Lambda}\pi^+\pi^-$ | $< 3.5 \times 10^{-3}$ | 90% |
| Γ_{44} | $K^+\bar{p}\Lambda + \text{c.c.}$ | $(9.1 \pm 1.8) \times 10^{-4}$ | |
| Γ_{45} | $\Sigma^0\bar{\Sigma}^0$ | $< 8 \times 10^{-5}$ | 90% |
| Γ_{46} | $\Sigma^+\bar{\Sigma}^-$ | $< 7 \times 10^{-5}$ | 90% |
| Γ_{47} | $\Xi^0\bar{\Xi}^0$ | $< 1.1 \times 10^{-4}$ | 90% |
| Γ_{48} | $\Xi^-\bar{\Xi}^+$ | $(1.55 \pm 0.35) \times 10^{-4}$ | |
| Γ_{49} | $J/\psi(1S)\pi^+\pi^-\pi^0$ | $< 1.5 \%$ | 90% |

Radiative decays

| | | | |
|---------------|---------------------|----------------------------------|-----|
| Γ_{50} | $\gamma J/\psi(1S)$ | $(19.5 \pm 0.8) \%$ | |
| Γ_{51} | $\gamma\rho^0$ | $< 5 \times 10^{-5}$ | 90% |
| Γ_{52} | $\gamma\omega$ | $< 6 \times 10^{-6}$ | 90% |
| Γ_{53} | $\gamma\phi$ | $< 1.2 \times 10^{-5}$ | 90% |
| Γ_{54} | $\gamma\gamma$ | $(2.56 \pm 0.16) \times 10^{-4}$ | |

CONSTRAINED FIT INFORMATION

A multiparticle fit to $\chi_{c1}(1P)$, $\chi_{c0}(1P)$, $\chi_{c2}(1P)$, and $\psi(2S)$ with 4 total widths, a partial width, 24 combinations of partial widths obtained from integrated cross section, and 82 branching ratios uses 213 measurements to determine 47 parameters. The overall fit has a $\chi^2 = 301.4$ for 166 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

| | | | | | | | | | | |
|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|
| x ₁₃ | 17 | | | | | | | | | |
| x ₁₄ | 4 | 22 | | | | | | | | |
| x ₁₅ | 10 | 8 | 2 | | | | | | | |
| x ₁₇ | 9 | 7 | 2 | 4 | | | | | | |
| x ₁₉ | 22 | 19 | 4 | 11 | 12 | | | | | |
| x ₂₀ | 20 | 4 | 1 | 2 | 2 | 5 | | | | |
| x ₂₄ | 18 | 16 | 3 | 9 | 10 | 36 | 4 | | | |
| x ₂₅ | 17 | 15 | 3 | 9 | 9 | 32 | 4 | 25 | | |
| x ₃₃ | 12 | 10 | 2 | 6 | 6 | 20 | 3 | 16 | 14 | |
| x ₃₆ | 7 | 6 | 1 | 4 | 2 | 1 | 2 | 1 | 2 | 2 |
| x ₄₂ | 8 | 7 | 2 | 4 | 5 | 18 | 2 | 14 | 13 | 8 |
| x ₅₀ | 28 | 24 | 5 | 14 | 15 | 55 | 6 | 44 | 39 | 25 |
| x ₅₄ | -18 | -15 | -3 | -9 | -5 | 2 | -5 | 0 | -2 | -3 |
| Γ | -25 | -21 | -5 | -13 | -12 | -33 | -6 | -27 | -25 | -17 |
| | x ₁ | x ₁₃ | x ₁₄ | x ₁₅ | x ₁₇ | x ₁₉ | x ₂₀ | x ₂₄ | x ₂₅ | x ₃₃ |
| x ₄₂ | 0 | | | | | | | | | |
| x ₅₀ | -11 | 22 | | | | | | | | |
| x ₅₄ | 26 | 2 | 9 | | | | | | | |
| Γ | -50 | -13 | -49 | -47 | | | | | | |
| | x ₃₆ | x ₄₂ | x ₅₀ | x ₅₄ | | | | | | |

$\chi_{c2}(1P)$ PARTIAL WIDTHS

———— $\chi_{c2}(1P) \Gamma(i) \Gamma(\gamma J/\psi(1S)) / \Gamma(\text{total})$ ————

| $\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S)) / \Gamma_{\text{total}}$ | | | | $\Gamma_{36} \Gamma_{50} / \Gamma$ |
|---|------------------------------|------|---------------------------------------|------------------------------------|
| VALUE (eV) | DOCUMENT ID | TECN | COMMENT | |
| 27.7 ± 1.4 OUR FIT | | | | |
| 27.5 ± 1.5 OUR AVERAGE | | | | |
| 27.0 ± 1.5 ± 1.1 | ⁹ ANDREOTTI 05A | E835 | $p\bar{p} \rightarrow e^+ e^- \gamma$ | |
| 27.7 ± 1.5 ± 2.0 | ^{9,10} ARMSTRONG 92 | E760 | $\bar{p}p \rightarrow e^+ e^- \gamma$ | |
| 36 ± 8 | ⁹ BAGLIN 86B | SPEC | $\bar{p}p \rightarrow e^+ e^- X$ | |

⁹ Calculated by us using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.

¹⁰ Recalculated by ANDREOTTI 05A.

| $\Gamma(\gamma\gamma) \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$ | | | | | $\Gamma_{54}\Gamma_{50}/\Gamma$ |
|---|------------|--------------------------------|----------|---|---------------------------------|
| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 98± 6 OUR FIT | | | | | |
| 117± 10 OUR AVERAGE | | | | | |
| 111± 12± 9 | 147 ± 15 | ¹¹ DOBBS | 06 CLE3 | 10.4 $e^+e^- \rightarrow e^+e^-\chi_{c2}$ | |
| 114± 11± 9 | 136 ± 13.3 | ^{11,12} ABE | 02T BELL | $e^+e^- \rightarrow e^+e^-\chi_{c2}$ | |
| 139± 55± 21 | | ^{11,13} ACCIARRI | 99E L3 | $e^+e^- \rightarrow e^+e^-\chi_{c2}$ | |
| 242± 65± 51 | | ^{11,14} ACKER...,K... | 98 OPAL | $e^+e^- \rightarrow e^+e^-\chi_{c2}$ | |
| 150± 42± 36 | | ^{11,15} DOMINICK | 94 CLE2 | $e^+e^- \rightarrow e^+e^-\chi_{c2}$ | |
| 470± 240± 120 | | ^{11,16} BAUER | 93 TPC | $e^+e^- \rightarrow e^+e^-\chi_{c2}$ | |

¹¹ Calculated by us using $B(J/\psi \rightarrow \ell^+\ell^-) = 0.1187 \pm 0.0008$.

¹² All systematic errors added in quadrature.

¹³ The value for $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$ reported in ACCIARRI 99E is derived using $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) \times B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.0162 \pm 0.0014$.

¹⁴ The value for $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$ reported in ACKERSTAFF,K 98 is derived using $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$ and $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1203 \pm 0.0038$.

¹⁵ The value for $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$ reported in DOMINICK 94 is derived using $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$, $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0627 \pm 0.0020$, and $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0597 \pm 0.0025$.

¹⁶ The value for $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$ reported in BAUER 93 is derived using $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$, $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0627 \pm 0.0020$, and $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0597 \pm 0.0025$.

———— $\chi_{c2}(1P) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ ————

| $\Gamma(2(\pi^+\pi^-)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ | | | | | $\Gamma_1\Gamma_{54}/\Gamma$ |
|---|------------|-------------|---------|--|------------------------------|
| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 5.6 ± 0.5 OUR FIT | | | | | |
| 5.2 ± 0.7 OUR AVERAGE | | | | | |
| 5.01± 0.44± 0.55 | 1597 ± 138 | UEHARA | 08 BELL | $\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(\pi^+\pi^-)$ | |
| 6.4 ± 1.8 ± 0.8 | | EISENSTEIN | 01 CLE2 | $e^+e^- \rightarrow e^+e^-\chi_{c2}$ | |

| $\Gamma(\rho^0\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ | | | | | $\Gamma_{20}\Gamma_{54}/\Gamma$ |
|--|-----------|-------------|---------|--|---------------------------------|
| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 2.0± 0.9 OUR FIT | | | | | |
| 3.2± 1.9± 0.5 | 986 ± 578 | UEHARA | 08 BELL | $\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(\pi^+\pi^-)$ | |

| $\Gamma(\rho\rho) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ | | | | | $\Gamma_2\Gamma_{54}/\Gamma$ |
|--|-----|------|-------------|------|------------------------------|
| VALUE (eV) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|------|----|------|--------|---------|--|
| <7.8 | 90 | <598 | UEHARA | 08 BELL | $\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(\pi^+\pi^-)$ |
|------|----|------|--------|---------|--|

| $\Gamma(\pi^+\pi^-K^+K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ | | | | | $\Gamma_{13}\Gamma_{54}/\Gamma$ |
|--|----------|-------------|---------|---|---------------------------------|
| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 4.6 ± 0.5 OUR FIT | | | | | |
| 4.42± 0.42± 0.53 | 780 ± 74 | UEHARA | 08 BELL | $\gamma\gamma \rightarrow \chi_{c2} \rightarrow K^+K^-\pi^+\pi^-$ | |

$$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{15}\Gamma_{54}/\Gamma$$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|----------|-------------|------|--|
| 1.26±0.24 OUR FIT | | | | |
| 0.8 ±0.17±0.27 | 151 ± 30 | UEHARA 08 | BELL | $\gamma\gamma \rightarrow \chi_{c2} \rightarrow K^+ K^- \pi^+ \pi^-$ |

$$\Gamma(K^+ K^- K^+ K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{33}\Gamma_{54}/\Gamma$$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|----------|-------------|------|---|
| 0.90±0.12 OUR FIT | | | | |
| 1.10±0.21±0.15 | 126 ± 24 | UEHARA 08 | BELL | $\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(K^+ K^-)$ |

$$\Gamma(\phi\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{17}\Gamma_{54}/\Gamma$$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------------|-------------|------|---|
| 0.75±0.14 OUR FIT | | | | |
| 0.58±0.18±0.16 | 26.5 ± 8.1 | UEHARA 08 | BELL | $\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(K^+ K^-)$ |

$$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{19}\Gamma_{54}/\Gamma$$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|---------|---------------------------|------|--|
| 1.21±0.09 OUR FIT | | | | |
| 1.18±0.25 OUR AVERAGE | | | | |
| 1.44±0.54±0.47 | 34 ± 13 | ¹⁷ UEHARA 09 | BELL | 10.6 $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$ |
| 1.14±0.21±0.17 | 54 ± 10 | ¹⁸ NAKAZAWA 05 | BELL | $e^+ e^- \rightarrow e^+ e^- \chi_{c2}$ |

¹⁷We multiplied the measurement by 3 to convert from $\pi^0 \pi^0$ to $\pi\pi$. Interference with the continuum included.

¹⁸We have multiplied $\pi^+ \pi^-$ measurement by 3/2 to obtain $\pi\pi$.

$$\Gamma(K^+ K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{24}\Gamma_{54}/\Gamma$$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|--------|-------------|------|---|
| 0.55±0.04 OUR FIT | | | | |
| 0.44±0.11±0.07 | 33 ± 8 | NAKAZAWA 05 | BELL | $e^+ e^- \rightarrow e^+ e^- \chi_{c2}$ |

$$\Gamma(K_S^0 K_S^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{25}\Gamma_{54}/\Gamma$$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|--------|-------------|------|---|
| 0.293±0.026 OUR FIT | | | | |
| 0.31 ±0.05 ±0.03 | 38 ± 7 | CHEN 07B | BELL | $e^+ e^- \rightarrow e^+ e^- \chi_{c2}$ |

$\chi_{c2}(1P)$ BRANCHING RATIOS

HADRONIC DECAYS

$$\Gamma(2(\pi^+ \pi^-))/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma$$

| VALUE | DOCUMENT ID |
|------------------------------|-------------|
| 0.0111±0.0011 OUR FIT | |

$$\Gamma(\rho^0 \pi^+ \pi^-)/\Gamma(2(\pi^+ \pi^-)) \quad \Gamma_{20}/\Gamma_1$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--------------------------|--------------|------|---|
| 0.36±0.15 OUR FIT | | | |
| 0.31±0.17 | TANENBAUM 78 | MRK1 | $\psi(2S) \rightarrow \gamma \chi_{c2}$ |

$\Gamma(\pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_3/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-------|------------------|----------|--|
| 2.00±0.25±0.08 | 903.5 | ¹⁹ HE | 08B CLEO | $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$ |

¹⁹ HE 08B reports $1.87 \pm 0.07 \pm 0.22 \pm 0.13$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\rho^+ \pi^- \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_4/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------|--------|---------------------|----------|--|
| 2.4±0.4±0.1 | 1031.9 | ^{20,21} HE | 08B CLEO | $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$ |

²⁰ HE 08B reports $2.23 \pm 0.11 \pm 0.32 \pm 0.16$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \rho^+ \pi^- \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²¹ Calculated by us. We have added the values from HE 08B for $\rho^+ \pi^- \pi^0$ and $\rho^- \pi^+ \pi^0$ decays assuming uncorrelated statistical and fully correlated systematic uncertainties.

$\Gamma(K^+ K^- \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_5/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|------------------|----------|--|
| 0.22±0.04±0.01 | 76.9 | ²² HE | 08B CLEO | $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$ |

²² HE 08B reports $0.21 \pm 0.03 \pm 0.03 \pm 0.01$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \pi^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_6/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-------|------------------|----------|--|
| 1.50±0.21±0.06 | 211.6 | ²³ HE | 08B CLEO | $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$ |

²³ HE 08B reports $1.41 \pm 0.11 \pm 0.16 \pm 0.10$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\rho^+ K^- K^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_7/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|------------------|----------|--|
| 0.45±0.13±0.02 | 62.9 | ²⁴ HE | 08B CLEO | $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$ |

²⁴ HE 08B reports $0.42 \pm 0.11 \pm 0.06 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \rho^+ K^- K^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^0 K^+ \pi^- \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_8/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|------------------|----------|--|
| 0.32±0.09±0.01 | 38.7 | ²⁵ HE | 08B CLEO | $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$ |

²⁵ HE 08B reports $0.30 \pm 0.07 \pm 0.04 \pm 0.02$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0 K^+ \pi^- \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^0 K^0 \pi^0 \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_9/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|------------------|----------|--|
| 0.42±0.09±0.02 | 63.0 | ²⁶ HE | 08B CLEO | $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$ |

²⁶ HE 08B reports $0.39 \pm 0.07 \pm 0.05 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0 K^0 \pi^0 \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{10}/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|------------------|----------|--|
| 0.41±0.09±0.02 | 51.1 | ²⁷ HE | 08B CLEO | $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$ |

²⁷ HE 08B reports $0.38 \pm 0.07 \pm 0.04 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^+ K^0 \pi^- \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{11}/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|------------------|----------|--|
| 0.32±0.09±0.01 | 39.3 | ²⁸ HE | 08B CLEO | $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$ |

²⁸ HE 08B reports $0.30 \pm 0.07 \pm 0.04 \pm 0.02$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^+ K^0 \pi^- \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ K^- \eta \pi^0)/\Gamma_{\text{total}}$ Γ_{12}/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|------------------|----------|--|
| 0.14±0.05±0.01 | 22.9 | ²⁹ HE | 08B CLEO | $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$ |

²⁹ HE 08B reports $0.13 \pm 0.04 \pm 0.02 \pm 0.01$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \eta \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\pi^+\pi^-K^+K^-)/\Gamma_{\text{total}} \quad \Gamma_{13}/\Gamma$$

VALUE (units 10^{-3}) DOCUMENT ID
9.2±1.1 OUR FIT

$$\Gamma(K^+\bar{K}^*(892)^0\pi^- + \text{c.c.})/\Gamma(\pi^+\pi^-K^+K^-) \quad \Gamma_{14}/\Gamma_{13}$$

VALUE DOCUMENT ID TECN COMMENT
0.25±0.13 OUR FIT
0.25±0.13 TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c2}$

$$\Gamma(K^+\bar{K}^*(892)^0\pi^- + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{14}/\Gamma$$

VALUE (units 10^{-4}) DOCUMENT ID
23±12 OUR FIT

$$\Gamma(K^*(892)^0\bar{K}^*(892)^0)/\Gamma_{\text{total}} \quad \Gamma_{15}/\Gamma$$

VALUE (units 10^{-3}) DOCUMENT ID
2.5±0.5 OUR FIT

$$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}} \quad \Gamma_{16}/\Gamma$$

VALUE (units 10^{-3}) DOCUMENT ID TECN COMMENT
8.6±1.8 OUR EVALUATION Treating systematic error as correlated.
8.6±1.8 OUR AVERAGE

8.6±0.9±1.6 ³⁰ BAI 99B BES $\psi(2S) \rightarrow \gamma\chi_{c2}$
 8.7±5.9±0.4 ³⁰ TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c2}$

³⁰ Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.3 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$. Multiplied by a factor of 2 to convert from $K_S^0 K^+\pi^-$ to $K^0 K^+\pi^-$ decay.

$$\Gamma(\phi\phi)/\Gamma_{\text{total}} \quad \Gamma_{17}/\Gamma$$

VALUE (units 10^{-3}) DOCUMENT ID
1.48±0.28 OUR FIT

$$\Gamma(\omega\omega)/\Gamma_{\text{total}} \quad \Gamma_{18}/\Gamma$$

VALUE (units 10^{-3}) EVTS DOCUMENT ID TECN COMMENT
1.9±0.6±0.1 27.7 ± 7.4 ³¹ ABLIKIM 05N BES2 $\psi(2S) \rightarrow \gamma\chi_{c2} \rightarrow \gamma\phi\pi$

³¹ ABLIKIM 05N reports $[\Gamma(\chi_{c2}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (0.165 \pm 0.044 \pm 0.032) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\pi\pi)/\Gamma_{\text{total}} \quad \Gamma_{19}/\Gamma$$

VALUE (units 10^{-3}) DOCUMENT ID
2.39±0.14 OUR FIT

$$\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}} \quad \Gamma_{20}/\Gamma$$

VALUE (units 10^{-4}) DOCUMENT ID
40±17 OUR FIT

$\Gamma(\pi^+\pi^-\eta)/\Gamma_{\text{total}}$ Γ_{21}/Γ

| VALUE (units 10^{-3}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------|
|--------------------------|-----|-------------|------|---------|

| | | | | |
|-----------------------|--|---------------------|----|--|
| 0.52±0.14±0.02 | | ³² ATHAR | 07 | CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$ |
|-----------------------|--|---------------------|----|--|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------|----|-----------------------|-----|--|
| <1.6 | 90 | ³³ ABLIKIM | 06R | BES2 $\psi(2S) \rightarrow \gamma \chi_{c2}$ |
|------|----|-----------------------|-----|--|

³² ATHAR 07 reports $(0.49 \pm 0.12 \pm 0.06) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³³ ABLIKIM 06R reports $< 1.7 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.74 \times 10^{-2}$.

$\Gamma(\pi^+\pi^-\eta')/\Gamma_{\text{total}}$ Γ_{22}/Γ

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

| | | | |
|-----------------------|---------------------|----|--|
| 0.54±0.20±0.02 | ³⁴ ATHAR | 07 | CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$ |
|-----------------------|---------------------|----|--|

³⁴ ATHAR 07 reports $(0.51 \pm 0.18 \pm 0.06) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$ Γ_{23}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|------|-------------|------|---------|
|--------------------------|-----|------|-------------|------|---------|

| | | | | | |
|--------------------|----------|--|---------------------|----|--|
| 5.4±0.7±0.2 | 156 ± 14 | | ³⁵ ASNER | 09 | CLEO $\psi(2S) \rightarrow \gamma \eta \eta$ |
|--------------------|----------|--|---------------------|----|--|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-----|----|--|---------------------|----|--|
| < 5 | 90 | | ³⁶ ADAMS | 07 | CLEO $\psi(2S) \rightarrow \gamma \chi_{c2}$ |
|-----|----|--|---------------------|----|--|

| | | | | | |
|-----|----|--|-------------------|-----|---|
| <12 | 90 | | ³⁷ BAI | 03C | BES $\psi(2S) \rightarrow \gamma \eta \eta \rightarrow 5\gamma$ |
|-----|----|--|-------------------|-----|---|

| | | | | | |
|-------------|--|--|-------------------|----|---|
| 7.9±4.1±2.4 | | | ³⁸ LEE | 85 | CBAL $\psi' \rightarrow \text{photons}$ |
|-------------|--|--|-------------------|----|---|

³⁵ ASNER 09 reports $(5.1 \pm 0.5 \pm 0.6) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³⁶ Superseded by ASNER 09. ADAMS 07 reports $< 4.7 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 0.0933 \pm 0.0014 \pm 0.0061$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.74 \times 10^{-2}$.

³⁷ Rescaled by us using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (8.3 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$. Multiplied by a factor of 2 to convert from $K_S^0 K^+\pi^-$ to $K^0 K^+\pi^-$ decay.

³⁸ Calculated using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 0.078 \pm 0.008$.

$\Gamma(K^+K^-)/\Gamma_{\text{total}}$ Γ_{24}/Γ

| VALUE (units 10^{-3}) | DOCUMENT ID |
|--------------------------|-------------|
|--------------------------|-------------|

| | |
|--------------------------|--|
| 1.09±0.08 OUR FIT | |
|--------------------------|--|

$\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{25}/Γ
VALUE (units 10^{-3}) DOCUMENT ID
0.58±0.05 OUR FIT

$\Gamma(K_S^0 K_S^0)/\Gamma(\pi\pi)$ Γ_{25}/Γ_{19}
VALUE DOCUMENT ID TECN COMMENT
0.242±0.020 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.27 ±0.07 ±0.04 ^{39,40} CHEN 07B BELL $e^+e^- \rightarrow e^+e^-\chi_{c2}$
³⁹ Using $\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ from the $\pi^+\pi^-$ measurement of NAKAZAWA 05 rescaled by 3/2 to convert to $\pi\pi$.
⁴⁰ Not independent from other measurements.

$\Gamma(K_S^0 K_S^0)/\Gamma(K^+K^-)$ Γ_{25}/Γ_{24}
VALUE DOCUMENT ID TECN COMMENT
0.53±0.05 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.70±0.21±0.12 ^{41,42} CHEN 07B BELL $e^+e^- \rightarrow e^+e^-\chi_{c2}$
⁴¹ Using $\Gamma(K^+K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ from NAKAZAWA 05.
⁴² Not independent from other measurements.

$\Gamma(\overline{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{26}/Γ
VALUE (units 10^{-3}) CL% EVTS DOCUMENT ID TECN COMMENT
1.32±0.20 OUR AVERAGE

1.39±0.22±0.05 ⁴³ ATHAR 07 CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$
1.11±0.41±0.04 28 ⁴⁴ ABLIKIM 06R BES2 $\psi(2S) \rightarrow \gamma\chi_{c2}$
• • • We do not use the following data for averages, fits, limits, etc. • • •
<2.0 90 ⁴⁵ BAI 99B BES $\psi(2S) \rightarrow \gamma\chi_{c2}$

⁴³ ATHAR 07 reports $(1.3 \pm 0.2 \pm 0.1) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \overline{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴⁴ We have multiplied the $K_S^0 K^+ \pi^-$ measurement by a factor of 2 to convert to $K^0 K^+ \pi^-$. ABLIKIM 06R reports $(1.2 \pm 0.4 \pm 0.2) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \overline{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.6) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴⁵ Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.3 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$. Multiplied by a factor of 2 to convert from $K_S^0 K^+ \pi^-$ to $K^0 K^+ \pi^-$ decay.

$\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{27}/Γ

| <u>VALUE (units 10^{-3})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|---------------------|-------------|--|
| $0.33 \pm 0.08 \pm 0.01$ | | ⁴⁶ ATHAR | 07 | CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$ |

⁴⁶ ATHAR 07 reports $(0.31 \pm 0.07 \pm 0.04) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ K^- \eta)/\Gamma_{\text{total}}$ Γ_{28}/Γ

| <u>VALUE (units 10^{-3})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|---------------------|-------------|--|
| <0.35 | 90 | ⁴⁷ ATHAR | 07 | CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$ |

⁴⁷ ATHAR 07 reports $< 0.33 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.74 \times 10^{-2}$.

$\Gamma(\eta\eta')/\Gamma_{\text{total}}$ Γ_{29}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|---------------|---------------------|-------------|---|
| <0.6 | 90 | 3.3 ± 8.0 | ⁴⁸ ASNER | 09 | CLEO $\psi(2S) \rightarrow \gamma \eta \eta'$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|------|----|--|---------------------|----|--|
| <2.5 | 90 | | ⁴⁹ ADAMS | 07 | CLEO $\psi(2S) \rightarrow \gamma \chi_{c2}$ |
|------|----|--|---------------------|----|--|

⁴⁸ ASNER 09 reports $< 0.6 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.74 \times 10^{-2}$.

⁴⁹ Superseded by ASNER 09. ADAMS 07 reports $< 2.3 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 0.0933 \pm 0.0014 \pm 0.0061$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.74 \times 10^{-2}$.

$\Gamma(\eta' \eta')/\Gamma_{\text{total}}$ Γ_{30}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|-------------|---------------------|-------------|--|
| <1.1 | 90 | 12 ± 7 | ⁵⁰ ASNER | 09 | CLEO $\psi(2S) \rightarrow \gamma \eta' \eta'$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|------|----|--|---------------------|----|--|
| <3.3 | 90 | | ⁵¹ ADAMS | 07 | CLEO $\psi(2S) \rightarrow \gamma \chi_{c2}$ |
|------|----|--|---------------------|----|--|

⁵⁰ ASNER 09 reports $< 1.0 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta' \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.74 \times 10^{-2}$.

⁵¹ Superseded by ASNER 09. ADAMS 07 reports $< 3.1 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta' \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 0.0933 \pm 0.0014 \pm 0.0061$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.74 \times 10^{-2}$.

$\Gamma(\pi^+\pi^-K_S^0K_S^0)/\Gamma_{\text{total}}$ Γ_{31}/Γ

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-------------|-----------------------|----------|--|
| $2.4 \pm 0.6 \pm 0.1$ | 57 ± 11 | ⁵² ABLIKIM | 050 BES2 | $\psi(2S) \rightarrow \gamma\chi_{c2}$ |

⁵² ABLIKIM 050 reports $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-K_S^0K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (0.207 \pm 0.039 \pm 0.033) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+K^-K_S^0K_S^0)/\Gamma_{\text{total}}$ Γ_{32}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------------|-----------------------|----------|--------------------------------------|
| <4 | 90 | 2.3 ± 2.2 | ⁵³ ABLIKIM | 050 BES2 | $e^+e^- \rightarrow \chi_{c2}\gamma$ |

⁵³ ABLIKIM 050 reports $[\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-K_S^0K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] < 3.5 \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 8.74 \times 10^{-2}$.

$\Gamma(K^+K^-K^+K^-)/\Gamma_{\text{total}}$ Γ_{33}/Γ

| VALUE (units 10^{-3}) | DOCUMENT ID |
|---|-------------|
| 1.78 ± 0.22 OUR FIT | |

$\Gamma(K^+K^-\phi)/\Gamma_{\text{total}}$ Γ_{34}/Γ

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-----------------------|----------|---|
| $1.55 \pm 0.32 \pm 0.06$ | 52 | ⁵⁴ ABLIKIM | 06T BES2 | $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$ |

⁵⁴ ABLIKIM 06T reports $(1.67 \pm 0.26 \pm 0.24) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K_S^0K_S^0\rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_{35}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-----------------------|----------|--|
| <7.9 | 90 | ⁵⁵ ABLIKIM | 06D BES2 | $\psi(2S) \rightarrow \chi_{c2}\gamma$ |

⁵⁵ Using $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (9.3 \pm 0.6)\%$.

$\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_{36}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID |
|---|-------------|
| 0.72 ± 0.04 OUR FIT | |

$\Gamma(\rho\bar{\rho}\pi^0)/\Gamma_{\text{total}}$ Γ_{37}/Γ

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|--|---------------------|---------|---|
| $0.47 \pm 0.10 \pm 0.02$ | ⁵⁶ ATHAR | 07 CLEO | $\psi(2S) \rightarrow \gamma h^+ h^- h^0$ |

⁵⁶ ATHAR 07 reports $(0.44 \pm 0.08 \pm 0.05) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \rho\bar{\rho}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\rho\bar{p}\eta)/\Gamma_{\text{total}}$ **Γ_{38}/Γ**

| <u>VALUE (units 10^{-3})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------------------|-------------|---|
| $0.20 \pm 0.08 \pm 0.01$ | ⁵⁷ ATHAR 07 | CLEO | $\psi(2S) \rightarrow \gamma h^+ h^- h^0$ |

⁵⁷ ATHAR 07 reports $(0.19 \pm 0.07 \pm 0.02) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \rho\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^+\pi^-\rho\bar{p})/\Gamma_{\text{total}}$ **Γ_{39}/Γ**

| <u>VALUE (units 10^{-3})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--|-------------|----------------|
| 1.32 ± 0.34 OUR EVALUATION | Treating systematic error as correlated. | | |
| 1.3 ± 0.4 OUR AVERAGE | Error includes scale factor of 1.3. | | |

| | | | |
|--------------------------|----------------------------|------|--|
| $1.17 \pm 0.19 \pm 0.30$ | ⁵⁸ BAI 99B | BES | $\psi(2S) \rightarrow \gamma\chi_{c2}$ |
| $2.64 \pm 1.03 \pm 0.14$ | ⁵⁸ TANENBAUM 78 | MRK1 | $\psi(2S) \rightarrow \gamma\chi_{c2}$ |

⁵⁸ Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.3 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$. Multiplied by a factor of 2 to convert from $K_S^0 K^+\pi^-$ to $K^0 K^+\pi^-$ decay.

$\Gamma(\pi^0\pi^0\rho\bar{p})/\Gamma_{\text{total}}$ **Γ_{40}/Γ**

| <u>VALUE (%)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|---|
| $0.085 \pm 0.026 \pm 0.003$ | 29.2 | ⁵⁹ HE 08B | CLEO | $e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$ |

⁵⁹ HE 08B reports $0.08 \pm 0.02 \pm 0.01 \pm 0.01\%$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^0\pi^0\rho\bar{p})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\rho\bar{n}\pi^-)/\Gamma_{\text{total}}$ **Γ_{41}/Γ**

| <u>VALUE (units 10^{-4})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|---------------------------|-------------|--|
| $11.1 \pm 3.8 \pm 0.4$ | ⁶⁰ ABLIKIM 06l | BES2 | $\psi(2S) \rightarrow \gamma p\pi^- X$ |

⁶⁰ ABLIKIM 06l reports $[\Gamma(\chi_{c2}(1P) \rightarrow \rho\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ = $(0.97 \pm 0.20 \pm 0.26) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$ **Γ_{42}/Γ**

| <u>VALUE (units 10^{-4})</u> | <u>DOCUMENT ID</u> |
|---|--------------------|
| 1.86 ± 0.27 OUR FIT | |

$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_{43}/Γ**

| <u>VALUE (units 10^{-3})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|---------------------------|-------------|--|
| < 3.5 | 90 | ⁶¹ ABLIKIM 06D | BES2 | $\psi(2S) \rightarrow \chi_{c2}\gamma$ |

⁶¹ Using $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (9.3 \pm 0.6)\%$.

$\Gamma(K^+ \bar{p} \Lambda + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{44}/Γ

| VALUE (units 10^{-3}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-----|------|---------------------|------|--|
| 0.91 ± 0.17 ± 0.04 | | | ⁶² ATHAR | 07 | CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$ |

⁶² ATHAR 07 reports $(0.85 \pm 0.14 \pm 0.10) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ \bar{p} \Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{45}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------------|--------------------|------|--|
| <0.8 | 90 | 7.5 ± 3.4 | ⁶³ NAIK | 08 | CLEO $\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$ |

⁶³ NAIK 08 reports $< 0.75 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.74 \times 10^{-2}$.

$\Gamma(\Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}$ Γ_{46}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------------|--------------------|------|--|
| <0.7 | 90 | 4.0 ± 3.5 | ⁶⁴ NAIK | 08 | CLEO $\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$ |

⁶⁴ NAIK 08 reports $< 0.67 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.74 \times 10^{-2}$.

$\Gamma(\Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}$ Γ_{47}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------------|--------------------|------|--|
| <1.1 | 90 | 2.9 ± 1.7 | ⁶⁵ NAIK | 08 | CLEO $\psi(2S) \rightarrow \gamma \Xi^0 \bar{\Xi}^0$ |

⁶⁵ NAIK 08 reports $< 1.06 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.74 \times 10^{-2}$.

$\Gamma(\Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}$ Γ_{48}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-----|------------|--------------------|------|--|
| 1.55 ± 0.34 ± 0.06 | | 29 ± 5 | ⁶⁶ NAIK | 08 | CLEO $\psi(2S) \rightarrow \gamma \Xi^+ \bar{\Xi}^-$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|---------|----|--|-----------------------|-----|--|
| < 3.7 | 90 | | ⁶⁷ ABLIKIM | 06D | BES2 $\psi(2S) \rightarrow \chi_{c2} \gamma$ |
|---------|----|--|-----------------------|-----|--|

⁶⁶ NAIK 08 reports $(1.45 \pm 0.30 \pm 0.15) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁶⁷ Using $B(\psi(2S) \rightarrow \chi_{c2} \gamma) = (9.3 \pm 0.6)\%$.

$\Gamma(J/\psi(1S) \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{49}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------|-----|-------------|------|--|
| <0.015 | 90 | BARATE | 81 | SPEC 190 GeV $\pi^- \text{Be} \rightarrow 2\pi 2\mu$ |

————— **RADIATIVE DECAYS** —————

$\Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$ Γ_{50}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|------------|-------------|--------------------|-------------|----------------|
|--------------|------------|-------------|--------------------|-------------|----------------|

0.195 ± 0.008 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|--|--|---------|----------|--|---|
| 0.199 ± 0.005 ± 0.012 | | 68 ADAM | 05A CLEO | | $e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$ |
| 68 Uses $B(\psi(2S) \rightarrow \gamma\chi_{c2} \rightarrow \gamma\gamma J/\psi)$ from ADAM 05A and $B(\psi(2S) \rightarrow \gamma\chi_{c2})$ from ATHAR 04. | | | | | |

$\Gamma(\gamma\rho^0)/\Gamma_{\text{total}}$ Γ_{51}/Γ

| <u>VALUE (units 10⁻⁶)</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------------------|------------|-------------|--------------------|-------------|----------------|
|--------------------------------------|------------|-------------|--------------------|-------------|----------------|

<50 90 17.2 ± 6.8 69 BENNETT 08A CLEO $\psi(2S) \rightarrow \gamma\gamma\rho^0$

69 BENNETT 08A reports $< 50 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 8.74 \times 10^{-2}$.

$\Gamma(\gamma\omega)/\Gamma_{\text{total}}$ Γ_{52}/Γ

| <u>VALUE (units 10⁻⁶)</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------------------|------------|-------------|--------------------|-------------|----------------|
|--------------------------------------|------------|-------------|--------------------|-------------|----------------|

<6 90 0.0 ± 1.8 70 BENNETT 08A CLEO $\psi(2S) \rightarrow \gamma\gamma\omega$

70 BENNETT 08A reports $< 7.0 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 8.74 \times 10^{-2}$.

$\Gamma(\gamma\phi)/\Gamma_{\text{total}}$ Γ_{53}/Γ

| <u>VALUE (units 10⁻⁶)</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------------------|------------|-------------|--------------------|-------------|----------------|
|--------------------------------------|------------|-------------|--------------------|-------------|----------------|

<12 90 1.3 ± 2.5 71 BENNETT 08A CLEO $\psi(2S) \rightarrow \gamma\gamma\phi$

71 BENNETT 08A reports $< 13 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 8.74 \times 10^{-2}$.

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{54}/Γ

| <u>VALUE (units 10⁻⁴)</u> | <u>DOCUMENT ID</u> |
|--------------------------------------|--------------------|
|--------------------------------------|--------------------|

2.56 ± 0.16 OUR FIT

$\Gamma(\gamma\gamma)/\Gamma(\gamma J/\psi(1S))$ Γ_{54}/Γ_{50}

| <u>VALUE (units 10⁻³)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------------------|--------------------|-------------|----------------|
|--------------------------------------|--------------------|-------------|----------------|

1.32 ± 0.09 OUR FIT

0.99 ± 0.18 72 AMBROGIANI 00B E835 $\bar{p}p \rightarrow \chi_{c2} \rightarrow \gamma\gamma, \gamma J/\psi$

72 Calculated by us using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}} \times \Gamma(p\bar{p})/\Gamma_{\text{total}}$ $\Gamma_{54}/\Gamma \times \Gamma_{36}/\Gamma$

| <u>VALUE (units 10⁻⁸)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------------------|--------------------|-------------|----------------|
|--------------------------------------|--------------------|-------------|----------------|

1.85 ± 0.18 OUR FIT

1.7 ± 0.4 OUR AVERAGE

| | | | |
|-------------|--------------|------|---------------------------------------|
| 1.60 ± 0.42 | ARMSTRONG 93 | E760 | $\bar{p}p \rightarrow \gamma\gamma X$ |
| 9.9 ± 4.5 | BAGLIN 87B | SPEC | $\bar{p}p \rightarrow \gamma\gamma X$ |

$\chi_{c2}(1P)$ CROSS-PARTICLE BRANCHING RATIOS

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \pi^+ \pi^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{13} / \Gamma \times \Gamma_{107}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-3}) DOCUMENT ID TECN COMMENT

2.39 ± 0.27 OUR FIT

2.5 ± 0.9 OUR AVERAGE Error includes scale factor of 2.3.

1.90 ± 0.14 ± 0.44

BAI

99B

BES

$\psi(2S) \rightarrow \gamma \chi_{c2}$

3.8 ± 0.67

⁷³ TANENBAUM

78

MRK1

$\psi(2S) \rightarrow \gamma \chi_{c2}$

⁷³ The reported value is derived using $B(\psi(2S) \rightarrow \pi^+ \pi^- J/\psi) \times B(J/\psi \rightarrow \ell^+ \ell^-) = (4.6 \pm 0.7)\%$. Calculated by us using $B(J/\psi \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0 \bar{K}^*(892)^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma_{15} / \Gamma \times \Gamma_{107}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$

VALUE (units 10^{-4}) DOCUMENT ID TECN COMMENT

2.2 ± 0.4 OUR FIT

3.11 ± 0.36 ± 0.48

ABLIKIM

04H

BES2

$\psi(2S) \rightarrow \gamma \chi_{c2}$

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow p \bar{p}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{36} / \Gamma \times \Gamma_{107}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-5}) DOCUMENT ID TECN COMMENT

1.88 ± 0.14 OUR FIT

1.4 ± 1.1

⁷⁴ BAI

98I

BES

$\psi(2S) \rightarrow \gamma \chi_{c2} \rightarrow \gamma \bar{p} p$

⁷⁴ Calculated by us. The value for $B(\chi_{c2} \rightarrow p \bar{p})$ reported in BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow p \bar{p}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma_{36} / \Gamma \times \Gamma_{107}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$

VALUE (units 10^{-6}) EVTS DOCUMENT ID TECN COMMENT

6.3 ± 0.5 OUR FIT

6.7 ± 1.1 OUR AVERAGE Error includes scale factor of 1.5.

7.2 ± 0.7 ± 0.4

121 ± 12

⁷⁵ NAIK

08

CLEO

$\psi(2S) \rightarrow \gamma p \bar{p}$

4.4^{+1.6}_{-1.4} ± 0.6

14.3^{+5.2}_{-4.7}

BAI

04F

BES

$\psi(2S) \rightarrow \gamma \chi_{c2}(1P) \rightarrow \gamma \bar{p} p$

⁷⁵ Calculated by us. NAIK 08 reports $B(\chi_{c2} \rightarrow p \bar{p}) = (7.7 \pm 0.8 \pm 0.4 \pm 0.5) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \Lambda \bar{\Lambda}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma_{42} / \Gamma \times \Gamma_{107}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$

VALUE (units 10^{-6}) EVTS DOCUMENT ID TECN COMMENT

16.3 ± 2.3 OUR FIT

15.9 ± 2.1 ± 1.0

71 ± 9

⁷⁶ NAIK

08

CLEO

$\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda}$

⁷⁶ Calculated by us. NAIK 08 reports $B(\chi_{c2} \rightarrow \Lambda \bar{\Lambda}) = (17.0 \pm 2.2 \pm 1.1 \pm 1.1) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \Lambda \bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{42}/\Gamma \times \Gamma_{107}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

4.8±0.7 OUR FIT

| | | | | |
|-----------------------------|---------------------|--------|---------|---|
| $7.1^{+3.1}_{-2.9} \pm 1.3$ | $8.3^{+3.7}_{-3.4}$ | 77 BAI | 03E BES | $\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda}$ |
|-----------------------------|---------------------|--------|---------|---|

⁷⁷ BAI 03E reports [$B(\chi_{c2} \rightarrow \Lambda \bar{\Lambda}) B(\psi(2S) \rightarrow \gamma \chi_{c2}) / B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-)$] × [$B^2(\Lambda \rightarrow \pi^- p) / B(J/\psi \rightarrow p \bar{p})$] = $(1.33^{+0.59}_{-0.55} \pm 0.25)\%$. We calculate from this measurement the presented value using $B(\Lambda \rightarrow \pi^- p) = (63.9 \pm 0.5)\%$ and $B(J/\psi \rightarrow p \bar{p}) = (2.17 \pm 0.07) \times 10^{-3}$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \pi\pi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{19}/\Gamma \times \Gamma_{107}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

2.09±0.09 OUR FIT

2.16±0.14 OUR AVERAGE Error includes scale factor of 1.3.

| | | | | |
|--------------------------|------|---------------------|---------|---|
| $2.23 \pm 0.06 \pm 0.10$ | 2.5k | ⁷⁸ ASNER | 09 CLEO | $\psi(2S) \rightarrow \gamma \pi^+ \pi^-$ |
| $1.90 \pm 0.08 \pm 0.20$ | 0.8k | ⁷⁹ ASNER | 09 CLEO | $\psi(2S) \rightarrow \gamma \pi^0 \pi^0$ |

⁷⁸ Calculated by us. ASNER 09 reports $B(\chi_{c2} \rightarrow \pi^+ \pi^-) = (1.59 \pm 0.04 \pm 0.07 \pm 0.10) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$. We have multiplied the $\pi^+ \pi^-$ measurement by 3/2 to obtain $\pi\pi$.

⁷⁹ Calculated by us. ASNER 09 reports $B(\chi_{c2} \rightarrow \pi^0 \pi^0) = (0.68 \pm 0.03 \pm 0.07 \pm 0.04) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$. We have multiplied the $\pi^0 \pi^0$ measurement by 3 to obtain $\pi\pi$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \pi\pi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{19}/\Gamma \times \Gamma_{107}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

0.623±0.028 OUR FIT

0.54 ±0.06 OUR AVERAGE

| | | | | |
|--------------------------|--------------|-------------------|---------|---|
| $0.66 \pm 0.18 \pm 0.37$ | 21 ± 6 | ⁸⁰ BAI | 03C BES | $\psi(2S) \rightarrow \gamma \pi^0 \pi^0$ |
| $0.54 \pm 0.05 \pm 0.04$ | 185 ± 16 | ⁸¹ BAI | 98I BES | $\psi(2S) \rightarrow \gamma \pi^+ \pi^-$ |

⁸⁰ We have multiplied $\pi^0 \pi^0$ measurement by 3 to obtain $\pi\pi$.

⁸¹ Calculated by us. The value for $B(\chi_{c2} \rightarrow \pi^+ \pi^-)$ reported by BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D]. We have multiplied $\pi^+ \pi^-$ measurement by 3/2 to obtain $\pi\pi$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{24}/\Gamma \times \Gamma_{107}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

9.5±0.6 OUR FIT

10.5±0.3±0.6 1.6k ⁸² ASNER 09 CLEO $\psi(2S) \rightarrow \gamma K^+ K^-$

⁸² Calculated by us. ASNER 09 reports $B(\chi_{c2} \rightarrow K^+ K^-) = (1.13 \pm 0.03 \pm 0.06 \pm 0.07) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{24} / \Gamma \times \Gamma_{107}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}}$$

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|----------|-------------|---------|---------------------------------------|
| 0.283 ± 0.017 OUR FIT | | | | |
| 0.190 ± 0.034 ± 0.019 | 115 ± 13 | 83 BAI | 98I BES | $\psi(2S) \rightarrow \gamma K^+ K^-$ |

⁸³ Calculated by us. The value for $B(\chi_{c2} \rightarrow K^+ K^-)$ reported by BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K_S^0 K_S^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma_{25} / \Gamma \times \Gamma_{107}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|----------|-------------|----------|---|
| 5.1 ± 0.4 OUR FIT | | | | |
| 5.0 ± 0.4 OUR AVERAGE | | | | |
| 4.9 ± 0.3 ± 0.3 | 373 ± 20 | 84 ASNER | 09 CLEO | $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$ |
| 5.72 ± 0.76 ± 0.63 | 65 | ABLIKIM | 05O BES2 | $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$ |

⁸⁴ Calculated by us. ASNER 09 reports $B(\chi_{c2} \rightarrow K_S^0 K_S^0) = (0.53 \pm 0.03 \pm 0.03 \pm 0.03) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K_S^0 K_S^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{25} / \Gamma \times \Gamma_{107}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}}$$

| VALUE (units 10^{-5}) | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-------------|---------|---|
| 15.1 ± 1.1 OUR FIT | | | |
| 14.7 ± 4.1 ± 3.3 | 85 BAI | 99B BES | $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$ |

⁸⁵ Calculated by us. The value of $B(\chi_{c2} \rightarrow K_S^0 K_S^0)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma_{50} / \Gamma \times \Gamma_{107}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|------|-------------|------|---|
| 1.70 ± 0.04 OUR FIT | | | | |
| 1.34 ± 0.14 OUR AVERAGE | | | | Error includes scale factor of 1.9. See the ideogram below. |

| | | | | |
|--------------------|------|--------------|----------|---|
| 1.62 ± 0.04 ± 0.12 | 5.8k | BAI | 04I BES2 | $\psi(2S) \rightarrow J/\psi \gamma \gamma$ |
| 0.99 ± 0.10 ± 0.08 | | GAISER | 86 CBAL | $\psi(2S) \rightarrow \gamma X$ |
| 1.47 ± 0.17 | | 86 OREGLIA | 82 CBAL | $\psi(2S) \rightarrow \gamma \chi_{c2}$ |
| 1.8 ± 0.5 | | 87 BRANDELIK | 79B DASP | $\psi(2S) \rightarrow \gamma \chi_{c2}$ |
| 1.2 ± 0.2 | | 87 BARTEL | 78B CNTR | $\psi(2S) \rightarrow \gamma \chi_{c2}$ |
| 2.2 ± 1.2 | | 88 BIDDICK | 77 CNTR | $e^+ e^- \rightarrow \gamma X$ |
| 1.2 ± 0.7 | | 86 WHITAKER | 76 MRK1 | $e^+ e^-$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------------------|-------|-----------|----------|---|
| 1.95 ± 0.02 ± 0.07 | 12.4k | 89 MENDEZ | 08 CLEO | $\psi(2S) \rightarrow \gamma \chi_{c2}$ |
| 1.85 ± 0.04 ± 0.07 | 1.9k | 90 ADAM | 05A CLEO | Repl. by MENDEZ 08 |


86 Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

87 Recalculated by us using $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$.

88 Assumes isotropic gamma distribution.

89 Not independent from other measurements of MENDEZ 08.

90 Not independent from other values reported by ADAM 05A.



$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}} \text{ (units } 10^{-2})$$

$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \text{ anything})$$

$$\Gamma_{50} / \Gamma \times \Gamma_{107}^{\psi(2S)} / \Gamma_9^{\psi(2S)}$$

$$\Gamma_{50} / \Gamma \times \Gamma_{107}^{\psi(2S)} / (\Gamma_{11}^{\psi(2S)} + \Gamma_{12}^{\psi(2S)} + \Gamma_{13}^{\psi(2S)}) + 0.344 \Gamma_{106}^{\psi(2S)} + 0.195 \Gamma_{107}^{\psi(2S)}$$

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
| 2.86±0.07 OUR FIT | | | | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|----------------|-------|----------------------|-----|------|---|
| 3.12±0.03±0.09 | 12.4k | ⁹¹ MENDEZ | 08 | CLEO | $\psi(2S) \rightarrow \gamma \chi_{c2}$ |
| 3.11±0.07±0.07 | 1.9k | ADAM | 05A | CLEO | Repl. by MENDEZ 08 |

⁹¹ Not independent from other measurements of MENDEZ 08.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{50} / \Gamma \times \Gamma_{107}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}}$$

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

5.07 ± 0.13 OUR FIT

5.53 ± 0.17 OUR AVERAGE

| | | | | |
|--------------------|-------|-----------------------|-----|--|
| 5.56 ± 0.05 ± 0.16 | 12.4k | MENDEZ | 08 | CLEO $\psi(2S) \rightarrow \gamma \chi_{c2}$ |
| 6.0 ± 2.8 | 1.3k | ⁹² ABLIKIM | 04B | BES $\psi(2S) \rightarrow J/\psi X$ |
| 3.9 ± 1.2 | | ⁹³ HIMEL | 80 | MRK2 $\psi(2S) \rightarrow \gamma \chi_{c2}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------------------|------|--------------------|-----|-------------------------|
| 5.52 ± 0.13 ± 0.13 | 1.9k | ⁹⁴ ADAM | 05A | CLEO Repl. by MENDEZ 08 |
|--------------------|------|--------------------|-----|-------------------------|

⁹² From a fit to the J/ψ recoil mass spectra.

⁹³ The value for $B(\psi(2S) \rightarrow \gamma \chi_{c2}) \times B(\chi_{c2} \rightarrow \gamma J/\psi(1S))$ reported in HIMEL 80 is derived using $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (33 \pm 3)\%$ and $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.138 \pm 0.018$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = (0.1181 \pm 0.0020)$.

⁹⁴ Not independent from other values reported by ADAM 05A.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \gamma \gamma) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma_{54} / \Gamma \times \Gamma_{107}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

2.24 ± 0.16 OUR FIT

2.73 ± 0.32 OUR AVERAGE

| | | | | |
|--------------------|----------|---------|-----|--|
| 2.68 ± 0.28 ± 0.15 | 333 ± 35 | ECKLUND | 08A | CLEO $\psi(2S) \rightarrow \gamma \chi_{c2} \rightarrow 3\gamma$ |
| 7.0 ± 2.1 ± 2.0 | | LEE | 85 | CBAL $\psi(2S) \rightarrow \gamma \chi_{c2}$ |

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow 2(\pi^+ \pi^-)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_1 / \Gamma \times \Gamma_{107}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}}$$

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

2.89 ± 0.27 OUR FIT

3.1 ± 1.0 OUR AVERAGE Error includes scale factor of 2.5.

| | | | |
|-----------------|-------------------------|-----|--|
| 2.3 ± 0.1 ± 0.5 | ⁹⁵ BAI | 99B | BES $\psi(2S) \rightarrow \gamma \chi_{c2}$ |
| 4.3 ± 0.6 | ⁹⁶ TANENBAUM | 78 | MRK1 $\psi(2S) \rightarrow \gamma \chi_{c2}$ |

⁹⁵ Calculated by us. The value for $B(\chi_{c2} \rightarrow 2\pi^+ 2\pi^-)$ reported in BAI 99B is derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

⁹⁶ The value for $B(\psi(2S) \rightarrow \gamma \chi_{c2}) \times B(\chi_{c2} \rightarrow 2\pi^+ \pi^-)$ reported in TANENBAUM 78 is derived using $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) \times B(J/\psi(1S) \ell^+ \ell^-) = (4.6 \pm 0.7)\%$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma_{33} / \Gamma \times \Gamma_{107}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

1.55 ± 0.19 OUR FIT

1.76 ± 0.16 ± 0.24 160 ⁹⁷ ABLIKIM 06T BES2 $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

⁹⁷ Calculated by us. The value of $B(\chi_{c2} \rightarrow 2K^+ 2K^-)$ reported by ABLIKIM 06T was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.1 \pm 0.4)\%$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{33} / \Gamma \times \Gamma_{107}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}}$$

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------------|---------|---|
| 4.6 ± 0.6 OUR FIT | | | |
| 3.6 ± 0.6 ± 0.6 | ⁹⁸ BAI | 99B BES | $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$ |

⁹⁸ Calculated by us. The value of $B(\chi_{c2} \rightarrow 2K^+ 2K^-)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \phi\phi) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma_{17} / \Gamma \times \Gamma_{107}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}}$$

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|------|-----------------------|----------|---|
| 1.29 ± 0.24 OUR FIT | | | | |
| 1.38 ± 0.24 ± 0.23 | 41 | ⁹⁹ ABLIKIM | 06T BES2 | $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$ |

⁹⁹ Calculated by us. The value of $B(\chi_{c2} \rightarrow \phi\phi)$ reported by ABLIKIM 06T was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.1 \pm 0.4)\%$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \phi\phi) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{17} / \Gamma \times \Gamma_{107}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}}$$

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|--------------------|---------|---|
| 3.8 ± 0.7 OUR FIT | | | |
| 4.8 ± 1.3 ± 1.3 | ¹⁰⁰ BAI | 99B BES | $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$ |

¹⁰⁰ Calculated by us. The value of $B(\chi_{c2} \rightarrow \phi\phi)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

MULTIPOLE AMPLITUDES IN $\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)$ RADIATIVE DECAY

$a_2 = M_2 / \sqrt{E_1^2 + M_2^2 + E_3^2}$ Magnetic quadrupole fractional transition amplitude

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-------|---------------------------|----------|--|
| -10.0 ± 1.5 OUR AVERAGE | | | | |
| - 9.3 ± 1.6 ± 0.3 | 19.8k | ¹⁰¹ ARTUSO | 09 CLEO | $\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$ |
| - 9.3 ⁺ ₋ 3.9 ⁺ _{4.1} ± 0.6 | 5.9k | ¹⁰² AMBROGIANI | 02 E835 | $p\bar{p} \rightarrow \chi_{c2} \rightarrow J/\psi \gamma$ |
| -14 ± 6 | 1.9k | ¹⁰² ARMSTRONG | 93E E760 | $p\bar{p} \rightarrow \chi_{c2} \rightarrow J/\psi \gamma$ |
| -33.3 ⁺ ₋ 11.6 ⁺ _{29.2} | 441 | ¹⁰² OREGLIA | 82 CBAL | $\psi(2S) \rightarrow \chi_{c1} \gamma \rightarrow J/\psi \gamma \gamma$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------------------|-------|-----------------------|---------|--|
| - 7.9 ± 1.9 ± 0.3 | 19.8k | ¹⁰³ ARTUSO | 09 CLEO | $\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$ |
|-------------------|-------|-----------------------|---------|--|

¹⁰¹ From a fit with floating M_2 amplitudes a_2 and b_2 , and fixed E_3 amplitudes $a_3 = b_3 = 0$.

¹⁰² Assuming $a_3 = 0$.

¹⁰³ From a fit with floating M_2 and E_3 amplitudes a_2 , b_2 , and a_3 , and b_3 .

$a_3 = E3/\sqrt{E1^2 + M2^2 + E3^2}$ Electric octupole fractional transition amplitude

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|-------|--------------------------|------|--|
| 1.6±1.3 OUR AVERAGE | | | | |
| 1.7±1.4±0.3 | 19.8k | ¹⁰⁴ ARTUSO 09 | CLEO | $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ |
| 2.0 ^{+5.5} _{-4.4} ±0.9 | 5908 | AMBROGIANI 02 | E835 | $\rho\bar{p} \rightarrow \chi_{c2} \rightarrow J/\psi\gamma$ |
| 0 ⁺⁶ ₋₅ | 1904 | ARMSTRONG 93E | E760 | $\rho\bar{p} \rightarrow \chi_{c2} \rightarrow J/\psi\gamma$ |
| ¹⁰⁴ From a fit with floating $M2$ and $E3$ amplitudes a_2 , b_2 , and a_3 , and b_3 . | | | | |

MULTIPOLE AMPLITUDES IN $\psi(2S) \rightarrow \gamma\chi_{c2}(1P)$ RADIATIVE DECAY

$b_2 = M2/\sqrt{E1^2 + M2^2 + E3^2}$ Magnetic quadrupole fractional transition amplitude

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-------|----------------------------|------|--|
| 1.0±1.4 OUR AVERAGE Error includes scale factor of 1.1. | | | | |
| 1.0±1.3±0.3 | 19.8k | ¹⁰⁵ ARTUSO 09 | CLEO | $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ |
| - 5.1 ^{+5.4} _{-3.6} | 721 | ¹⁰⁶ ABLIKIM 04I | BES2 | $\psi(2S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$ |
| 13.2 ^{+9.8} _{-7.5} | 441 | ¹⁰⁵ OREGLIA 82 | CBAL | $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.2±1.5±0.4 | 19.8k | ¹⁰⁷ ARTUSO 09 | CLEO | $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ |
| ¹⁰⁵ From a fit with floating $M2$ amplitudes a_2 and b_2 , and fixed $E3$ amplitudes $a_3=b_3=0$. | | | | |
| ¹⁰⁶ From a fit with floating $M2$ and $E3$ amplitudes a_2 and a_3 . | | | | |
| ¹⁰⁷ From a fit with floating $M2$ and $E3$ amplitudes a_2 , b_2 , and a_3 , and b_3 . | | | | |

$b_3 = E3/\sqrt{E1^2 + M2^2 + E3^2}$ Electric octupole fractional transition amplitude

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|-------|----------------------------|------|--|
| -1.0±1.1 OUR AVERAGE | | | | |
| -0.8±1.2±0.2 | 19.8k | ARTUSO 09 | CLEO | $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ |
| -2.7 ^{+4.3} _{-2.9} | 721 | ¹⁰⁸ ABLIKIM 04I | BES2 | $\psi(2S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$ |
| ¹⁰⁸ From a fit with floating $M2$ and $E3$ amplitudes a_2 and a_3 . | | | | |

MULTIPOLE AMPLITUDE RATIOS IN RADIATIVE DECAYS
 $\psi(2S) \rightarrow \gamma\chi_{c2}(1P)$ and $\chi_{c2} \rightarrow \gamma J/\psi(1S)$

b_2/a_2 Magnetic quadrupole transition amplitude ratio

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-------|--------------------------|------|---|
| -11 ⁺¹⁴ ₋₁₅ | 19.8k | ¹⁰⁹ ARTUSO 09 | CLEO | $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ |
| ¹⁰⁹ Statistical and systematic errors combined. From a fit with floating $M2$ amplitudes a_2 and b_2 , and fixed $E3$ amplitudes $a_3=b_3=0$. Not independent of values for $a_2(\chi_{c2}(1P))$ and $b_2(\chi_{c2}(1P))$ from ARTUSO 09. | | | | |

$\chi_{c2}(1P)$ REFERENCES

| | | | | |
|---------------|-----|----------------|---|---------------------------|
| ARTUSO | 09 | PR D80 112003 | M. Artuso <i>et al.</i> | (CLEO Collab.) |
| ASNER | 09 | PR D79 072007 | D.M. Asner <i>et al.</i> | (CLEO Collab.) |
| UEHARA | 09 | PR D79 052009 | S. Uehara <i>et al.</i> | (BELLE Collab.) |
| BENNETT | 08A | PRL 101 151801 | J.V. Bennett <i>et al.</i> | (CLEO Collab.) |
| ECKLUND | 08A | PR D78 091501R | K.M. Ecklund <i>et al.</i> | (CLEO Collab.) |
| HE | 08B | PR D78 092004 | Q. He <i>et al.</i> | (CLEO Collab.) |
| MENDEZ | 08 | PR D78 011102R | H. Mendez <i>et al.</i> | (CLEO Collab.) |
| NAIK | 08 | PR D78 031101R | P. Naik <i>et al.</i> | (CLEO Collab.) |
| UEHARA | 08 | EPJ C53 1 | S. Uehara <i>et al.</i> | (BELLE Collab.) |
| ADAMS | 07 | PR D75 071101R | G.S. Adams <i>et al.</i> | (CLEO Collab.) |
| ATHAR | 07 | PR D75 032002 | S.B. Athar <i>et al.</i> | (CLEO Collab.) |
| CHEN | 07B | PL B651 15 | W.T. Chen <i>et al.</i> | (BELLE Collab.) |
| ABLIKIM | 06D | PR D73 052006 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 06I | PR D74 012004 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 06R | PR D74 072001 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 06T | PL B642 197 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| DOBBS | 06 | PR D73 071101R | S. Dobbs <i>et al.</i> | (CLEO Collab.) |
| ABLIKIM | 05G | PR D71 092002 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 05N | PL B630 7 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 05O | PL B630 21 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ADAM | 05A | PRL 94 232002 | N.E. Adam <i>et al.</i> | (CLEO Collab.) |
| ANDREOTTI | 05A | NP B717 34 | M. Andreotti <i>et al.</i> | (FNAL E835 Collab.) |
| NAKAZAWA | 05 | PL B615 39 | H. Nakazawa <i>et al.</i> | (BELLE Collab.) |
| ABLIKIM | 04B | PR D70 012003 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 04H | PR D70 092003 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 04I | PR D70 092004 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ATHAR | 04 | PR D70 112002 | S.B. Athar <i>et al.</i> | (CLEO Collab.) |
| BAI | 04F | PR D69 092001 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 04I | PR D70 012006 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| AULCHENKO | 03 | PL B573 63 | V.M. Aulchenko <i>et al.</i> | (KEDR Collab.) |
| BAI | 03C | PR D67 032004 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 03E | PR D67 112001 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| ABE | 02T | PL B540 33 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| AMBROGIANI | 02 | PR D65 052002 | M. Ambrogiani <i>et al.</i> | (FNAL E835 Collab.) |
| EISENSTEIN | 01 | PRL 87 061801 | B.I. Eisenstein <i>et al.</i> | (CLEO Collab.) |
| AMBROGIANI | 00B | PR D62 052002 | M. Ambrogiani <i>et al.</i> | (FNAL E835 Collab.) |
| ACCIARRI | 99E | PL B453 73 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| BAI | 99B | PR D60 072001 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| ACKER...,K... | 98 | PL B439 197 | K. Ackerstaff <i>et al.</i> | (OPAL Collab.) |
| BAI | 98D | PR D58 092006 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 98I | PRL 81 3091 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| DOMINICK | 94 | PR D50 4265 | J. Dominick <i>et al.</i> | (CLEO Collab.) |
| ARMSTRONG | 93 | PRL 70 2988 | T.A. Armstrong <i>et al.</i> | (FNAL E760 Collab.) |
| ARMSTRONG | 93E | PR D48 3037 | T.A. Armstrong <i>et al.</i> | (FNAL-E760 Collab.) |
| BAUER | 93 | PL B302 345 | D.A. Bauer <i>et al.</i> | (TPC Collab.) |
| ARMSTRONG | 92 | NP B373 35 | T.A. Armstrong <i>et al.</i> | (FNAL, FERR, GENO+) |
| Also | | PRL 68 1468 | T.A. Armstrong <i>et al.</i> | (FNAL, FERR, GENO+) |
| BAGLIN | 87B | PL B187 191 | C. Baglin <i>et al.</i> | (R704 Collab.) |
| BAGLIN | 86B | PL B172 455 | C. Baglin (LAPP, CERN, GENO, LYON, OSLO+) | |
| GAISER | 86 | PR D34 711 | J. Gaiser <i>et al.</i> | (Crystal Ball Collab.) |
| LEE | 85 | SLAC 282 | R.A. Lee | (SLAC) |
| LEMOIGNE | 82 | PL 113B 509 | Y. Lemoigne <i>et al.</i> | (SACL, LOIC, SHMP+) |
| OREGLIA | 82 | PR D25 2259 | M.J. Oreglia <i>et al.</i> | (SLAC, CIT, HARV+) |
| Also | | Private Comm. | M.J. Oreglia | (EFI) |
| BARATE | 81 | PR D24 2994 | R. Barate <i>et al.</i> | (SACL, LOIC, SHMP, CERN+) |
| HIMEL | 80 | PRL 44 920 | T. Himel <i>et al.</i> | (LBL, SLAC) |
| Also | | Private Comm. | G. Trilling | (LBL, UCB) |
| BRANDELIK | 79B | NP B160 426 | R. Brandelik <i>et al.</i> | (DASP Collab.) |
| BARTEL | 78B | PL 79B 492 | W. Bartel <i>et al.</i> | (DESY, HEIDP) |
| TANENBAUM | 78 | PR D17 1731 | W.M. Tanenbaum <i>et al.</i> | (SLAC, LBL) |
| Also | | Private Comm. | G. Trilling | (LBL, UCB) |
| BIDDICK | 77 | PRL 38 1324 | C.J. Biddick <i>et al.</i> | (UCSD, UMD, PAVI+) |
| WHITAKER | 76 | PRL 37 1596 | J.S. Whitaker <i>et al.</i> | (SLAC, LBL) |