

$\Upsilon(1S)$

$$J^{PC} = 0^{--}(1^{--})$$

$\Upsilon(1S)$ MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
9460.30 ± 0.26 OUR AVERAGE	Error includes scale factor of 3.3.		
9460.51 ± 0.09 ± 0.05	¹ ARTAMONOV 00	MD1	$e^+e^- \rightarrow$ hadrons
9459.97 ± 0.11 ± 0.07	MACKAY 84	REDE	$e^+e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
9460.60 ± 0.09 ± 0.05	^{2,3} BARU	92B	REDE $e^+e^- \rightarrow$ hadrons
9460.59 ± 0.12	BARU 86	REDE	$e^+e^- \rightarrow$ hadrons
9460.6 ± 0.4	^{3,4} ARTAMONOV 84	REDE	$e^+e^- \rightarrow$ hadrons
¹ Reanalysis of BARU 92B and ARTAMONOV 84 using new electron mass (COHEN 87).			
² Superseding BARU 86.			
³ Superseded by ARTAMONOV 00.			
⁴ Value includes data of ARTAMONOV 82.			

$\Upsilon(1S)$ WIDTH

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>
54.02 ± 1.25 OUR EVALUATION	See the Note on "Width Determinations of the Υ States"

$\Upsilon(1S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $\tau^+\tau^-$	(2.60 ± 0.10) %	
Γ_2 e^+e^-	(2.48 ± 0.07) %	
Γ_3 $\mu^+\mu^-$	(2.48 ± 0.05) %	
Hadronic decays		
Γ_4 ggg	(81.7 ± 0.7) %	
Γ_5 γgg	(2.21 ± 0.22) %	
Γ_6 $\eta'(958)$ anything	(2.94 ± 0.24) %	
Γ_7 $J/\psi(1S)$ anything	(6.5 ± 0.7) × 10 ⁻⁴	
Γ_8 χ_{c0} anything	< 5 × 10 ⁻³	90%
Γ_9 χ_{c1} anything	(2.3 ± 0.7) × 10 ⁻⁴	
Γ_{10} χ_{c2} anything	(3.4 ± 1.0) × 10 ⁻⁴	
Γ_{11} $\psi(2S)$ anything	(2.7 ± 0.9) × 10 ⁻⁴	
Γ_{12} $\rho\pi$	< 2 × 10 ⁻⁴	90%
Γ_{13} $\pi^+\pi^-$	< 5 × 10 ⁻⁴	90%
Γ_{14} K^+K^-	< 5 × 10 ⁻⁴	90%
Γ_{15} $p\bar{p}$	< 5 × 10 ⁻⁴	90%
Γ_{16} $\pi^0\pi^+\pi^-$	< 1.84 × 10 ⁻⁵	90%
Γ_{17} $D^*(2010)^\pm$ anything	(2.52 ± 0.20) %	
Γ_{18} \bar{d} anything	(2.86 ± 0.28) × 10 ⁻⁵	

Radiative decays

Γ_{19}	$\gamma\pi^+\pi^-$		$(6.3 \pm 1.8) \times 10^{-5}$	
Γ_{20}	$\gamma\pi^0\pi^0$		$(1.7 \pm 0.7) \times 10^{-5}$	
Γ_{21}	$\gamma\pi^0\eta$		$< 2.4 \times 10^{-6}$	90%
Γ_{22}	$\gamma K^+ K^-$	[a]	$(1.14 \pm 0.13) \times 10^{-5}$	
Γ_{23}	$\gamma p\bar{p}$	[b]	$< 6 \times 10^{-6}$	90%
Γ_{24}	$\gamma 2h^+ 2h^-$		$(7.0 \pm 1.5) \times 10^{-4}$	
Γ_{25}	$\gamma 3h^+ 3h^-$		$(5.4 \pm 2.0) \times 10^{-4}$	
Γ_{26}	$\gamma 4h^+ 4h^-$		$(7.4 \pm 3.5) \times 10^{-4}$	
Γ_{27}	$\gamma\pi^+\pi^- K^+ K^-$		$(2.9 \pm 0.9) \times 10^{-4}$	
Γ_{28}	$\gamma 2\pi^+ 2\pi^-$		$(2.5 \pm 0.9) \times 10^{-4}$	
Γ_{29}	$\gamma 3\pi^+ 3\pi^-$		$(2.5 \pm 1.2) \times 10^{-4}$	
Γ_{30}	$\gamma 2\pi^+ 2\pi^- K^+ K^-$		$(2.4 \pm 1.2) \times 10^{-4}$	
Γ_{31}	$\gamma\pi^+\pi^- p\bar{p}$		$(1.5 \pm 0.6) \times 10^{-4}$	
Γ_{32}	$\gamma 2\pi^+ 2\pi^- p\bar{p}$		$(4 \pm 6) \times 10^{-5}$	
Γ_{33}	$\gamma 2K^+ 2K^-$		$(2.0 \pm 2.0) \times 10^{-5}$	
Γ_{34}	$\gamma\eta'(958)$		$< 1.9 \times 10^{-6}$	90%
Γ_{35}	$\gamma\eta$		$< 1.0 \times 10^{-6}$	90%
Γ_{36}	$\gamma f_0(980)$		$< 3 \times 10^{-5}$	90%
Γ_{37}	$\gamma f_2'(1525)$		$(3.7^{+1.2}_{-1.1}) \times 10^{-5}$	
Γ_{38}	$\gamma f_2(1270)$		$(1.01 \pm 0.09) \times 10^{-4}$	
Γ_{39}	$\gamma\eta(1405)$		$< 8.2 \times 10^{-5}$	90%
Γ_{40}	$\gamma f_0(1500)$		$< 1.5 \times 10^{-5}$	90%
Γ_{41}	$\gamma f_0(1710)$		$< 2.6 \times 10^{-4}$	90%
Γ_{42}	$\gamma f_0(1710) \rightarrow \gamma K^+ K^-$		$< 7 \times 10^{-6}$	90%
Γ_{43}	$\gamma f_0(1710) \rightarrow \gamma\pi^0\pi^0$		$< 1.4 \times 10^{-6}$	90%
Γ_{44}	$\gamma f_0(1710) \rightarrow \gamma\eta\eta$		$< 1.8 \times 10^{-6}$	90%
Γ_{45}	$\gamma f_4(2050)$		$< 5.3 \times 10^{-5}$	90%
Γ_{46}	$\gamma f_0(2200) \rightarrow \gamma K^+ K^-$		$< 2 \times 10^{-4}$	90%
Γ_{47}	$\gamma f_J(2220) \rightarrow \gamma K^+ K^-$		$< 8 \times 10^{-7}$	90%
Γ_{48}	$\gamma f_J(2220) \rightarrow \gamma\pi^+\pi^-$		$< 6 \times 10^{-7}$	90%
Γ_{49}	$\gamma f_J(2220) \rightarrow \gamma p\bar{p}$		$< 1.1 \times 10^{-6}$	90%
Γ_{50}	$\gamma\eta(2225) \rightarrow \gamma\phi\phi$		$< 3 \times 10^{-3}$	90%
Γ_{51}	γX	[c]	$< 3 \times 10^{-5}$	90%
Γ_{52}	$\gamma X\bar{X}$	[d]	$< 1 \times 10^{-3}$	90%
Γ_{53}	$\gamma X \rightarrow \gamma + \geq 4$ prongs	[e]	$< 1.78 \times 10^{-4}$	95%
Γ_{54}	$\gamma a_1^0 \rightarrow \gamma\mu^+\mu^-$	[f]	$< 9 \times 10^{-6}$	90%
Γ_{55}	$\gamma a_1^0 \rightarrow \gamma\tau^+\tau^-$	[a]	$< 5.0 \times 10^{-5}$	90%

Lepton Flavor (LF) violating or Invisible decays

Γ_{56}	$\mu^\pm \tau^\mp$	LF	$< 6.0 \times 10^{-6}$	95%
Γ_{57}	invisible		$< 3.0 \times 10^{-4}$	90%

- [a] $2m_\tau < M(\tau^+\tau^-) < 7500$ MeV.
- [b] $2 < m_{K^+K^-} < 3$ GeV.
- [c] X = pseudoscalar with $m < 7.2$ GeV
- [d] $X\bar{X}$ = vectors with $m < 3.1$ GeV
- [e] 1.5 GeV $< m_X < 5.0$ GeV
- [f] $201 < M(\mu^+\mu^-) < 3565$ MeV.

$\Upsilon(1S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

$\Gamma(e^+e^-) \times \Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$				$\Gamma_2\Gamma_3/\Gamma$
<u>VALUE (eV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
31.2±1.6±1.7	KOBEL	92	CBAL	$e^+e^- \rightarrow \mu^+\mu^-$

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_0\Gamma_2/\Gamma$
<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
1.240±0.016 OUR AVERAGE				
1.252±0.004±0.019	⁵ ROSNER	06	CLEO	$9.5 e^+e^- \rightarrow \text{hadrons}$
1.187±0.023±0.031	⁵ BARU	92B	MD1	$e^+e^- \rightarrow \text{hadrons}$
1.23 ±0.02 ±0.05	⁵ JAKUBOWSKI	88	CBAL	$e^+e^- \rightarrow \text{hadrons}$
1.37 ±0.06 ±0.09	⁶ GILES	84B	CLEO	$e^+e^- \rightarrow \text{hadrons}$
1.23 ±0.08 ±0.04	⁶ ALBRECHT	82	DASP	$e^+e^- \rightarrow \text{hadrons}$
1.13 ±0.07 ±0.11	⁶ NICZYPORUK	82	LENA	$e^+e^- \rightarrow \text{hadrons}$
1.09 ±0.25	⁶ BOCK	80	CNTR	$e^+e^- \rightarrow \text{hadrons}$
1.35 ±0.14	⁷ BERGER	79	PLUT	$e^+e^- \rightarrow \text{hadrons}$

⁵ Radiative corrections evaluated following KURAEV 85.

⁶ Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

⁷ Radiative corrections reevaluated by ALEXANDER 89 using $B(\mu\mu) = 0.026$.

$\Upsilon(1S)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$				Γ_2
<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>			
1.340±0.018 OUR EVALUATION				

$\Upsilon(1S)$ BRANCHING RATIOS

$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$				Γ_1/Γ
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.60±0.10 OUR AVERAGE				
2.53±0.13±0.05	60k	⁸ BESSON	07	CLEO $e^+e^- \rightarrow \Upsilon(1S) \rightarrow \tau^+\tau^-$
2.61±0.12 ^{+0.09} _{-0.13}	25k	CINABRO	94B	CLE2 $e^+e^- \rightarrow \tau^+\tau^-$
2.7 ±0.4 ±0.2		⁹ ALBRECHT	85C	ARG $\Upsilon(2S) \rightarrow \pi^+\pi^-\tau^+\tau^-$
3.4 ±0.4 ±0.4		GILES	83	CLEO $e^+e^- \rightarrow \tau^+\tau^-$

⁸ BESSON 07 reports $[\Gamma(\Upsilon(1S) \rightarrow \tau^+ \tau^-) / \Gamma_{\text{total}}] / [B(\Upsilon(1S) \rightarrow \mu^+ \mu^-)] = 1.02 \pm 0.02 \pm 0.05$ which we multiply by our best value $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁹ Using $B(\Upsilon(1S) \rightarrow ee) = B(\Upsilon(1S) \rightarrow \mu\mu) = 0.0256$; not used for width evaluations.

$\Gamma(e^+ e^-) / \Gamma_{\text{total}}$ **Γ_2 / Γ**

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.38 ± 0.11 OUR AVERAGE				
2.29 ± 0.08 ± 0.11		ALEXANDER 98	CLE2	$\Upsilon(2S) \rightarrow \pi^+ \pi^- e^+ e^-$
2.42 ± 0.14 ± 0.14	307	ALBRECHT 87	ARG	$\Upsilon(2S) \rightarrow \pi^+ \pi^- e^+ e^-$
2.8 ± 0.3 ± 0.2	826	BESSON 84	CLEO	$\Upsilon(2S) \rightarrow \pi^+ \pi^- e^+ e^-$
5.1 ± 3.0		BERGER 80C	PLUT	$e^+ e^- \rightarrow e^+ e^-$

$\Gamma(\mu^+ \mu^-) / \Gamma_{\text{total}}$ **Γ_3 / Γ**

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0248 ± 0.0005 OUR AVERAGE				
0.0249 ± 0.0002 ± 0.0007	345k	ADAMS 05	CLEO	$e^+ e^- \rightarrow \mu^+ \mu^-$
0.0249 ± 0.0008 ± 0.0013		ALEXANDER 98	CLE2	$\Upsilon(2S) \rightarrow \pi^+ \pi^- \mu^+ \mu^-$
0.0212 ± 0.0020 ± 0.0010		¹⁰ BARU 92	MD1	$e^+ e^- \rightarrow \mu^+ \mu^-$
0.0231 ± 0.0012 ± 0.0010		¹⁰ KOBEL 92	CBAL	$e^+ e^- \rightarrow \mu^+ \mu^-$
0.0252 ± 0.0007 ± 0.0007		CHEN 89B	CLEO	$e^+ e^- \rightarrow \mu^+ \mu^-$
0.0261 ± 0.0009 ± 0.0011		KAARSBERG 89	CSB2	$e^+ e^- \rightarrow \mu^+ \mu^-$
0.0230 ± 0.0025 ± 0.0013	86	ALBRECHT 87	ARG	$\Upsilon(2S) \rightarrow \pi^+ \pi^- \mu^+ \mu^-$
0.029 ± 0.003 ± 0.002	864	BESSON 84	CLEO	$\Upsilon(2S) \rightarrow \pi^+ \pi^- \mu^+ \mu^-$
0.027 ± 0.003 ± 0.003		ANDREWS 83	CLEO	$e^+ e^- \rightarrow \mu^+ \mu^-$
0.032 ± 0.013 ± 0.003		ALBRECHT 82	DASP	$e^+ e^- \rightarrow \mu^+ \mu^-$
0.038 ± 0.015 ± 0.002		NICZYPORUK 82	LENA	$e^+ e^- \rightarrow \mu^+ \mu^-$
0.014 ^{+0.034} _{-0.014}		BOCK 80	CNTR	$e^+ e^- \rightarrow \mu^+ \mu^-$
0.022 ± 0.020		BERGER 79	PLUT	$e^+ e^- \rightarrow \mu^+ \mu^-$

¹⁰ Taking into account interference between the resonance and continuum.

$\Gamma(\tau^+ \tau^-) / \Gamma(\mu^+ \mu^-)$ **Γ_1 / Γ_3**

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.02 ± 0.02 ± 0.05	60k	BESSON 07	CLEO	$e^+ e^- \rightarrow \Upsilon(1S)$

$\Gamma(g g g) / \Gamma_{\text{total}}$ **Γ_4 / Γ**

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
81.7 ± 0.7	20M	¹¹ BESSON 06A	CLEO	$\Upsilon(1S) \rightarrow \text{hadrons}$

¹¹ Calculated using the value $\Gamma(\gamma g g) / \Gamma(g g g) = (2.70 \pm 0.01 \pm 0.13 \pm 0.24)\%$ from BESSON 06A and PDG 08 values of $B(\mu^+ \mu^-) = (2.48 \pm 0.05)\%$ and $R_{\text{hadrons}} = 3.51$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(\gamma g g) / \Gamma_{\text{total}}$ measurement of BESSON 06A.

$\Gamma(\gamma g g)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.20±0.60	400k	¹² BESSON	06A CLEO	$\Upsilon(1S) \rightarrow \gamma + \text{hadrons}$

¹² Calculated using BESSON 06A values of $\Gamma(\gamma g g)/\Gamma(g g g) = (2.70 \pm 0.01 \pm 0.13 \pm 0.24)\%$ and $\Gamma(g g g)/\Gamma_{\text{total}}$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(g g g)/\Gamma_{\text{total}}$ measurement of BESSON 06A.

$\Gamma(\gamma g g)/\Gamma(g g g)$ Γ_5/Γ_4

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.70±0.01±0.27	20M	BESSON	06A CLEO	$\Upsilon(1S) \rightarrow (\gamma +) \text{hadrons}$

$\Gamma(\eta'(958) \text{ anything})/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0294±0.0024 OUR AVERAGE			
0.030 ±0.002 ±0.002	AQUINES	06A CLE3	$\Upsilon(1S) \rightarrow \eta' \text{ anything}$
0.028 ±0.004 ±0.002	ARTUSO	03 CLE2	$\Upsilon(1S) \rightarrow \eta' \text{ anything}$

$\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.65±0.07 OUR AVERAGE					
0.64±0.04±0.06		730 ± 40	BRIERE	04 CLEO	$e^+e^- \rightarrow J/\psi X$
1.1 ±0.4 ±0.2			¹³ FULTON	89 CLEO	$e^+e^- \rightarrow \mu^+\mu^- X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.68	90	ALBRECHT	92J ARG	$e^+e^- \rightarrow e^+e^- X, \mu^+\mu^- X$
<1.7	90	MASCHMANN	90 CBAL	$e^+e^- \rightarrow \text{hadrons}$
<20	90	NICZYPORUK	83 LENA	

¹³ Using $B((J/\psi) \rightarrow \mu^+\mu^-) = (6.9 \pm 0.9)\%$.

$\Gamma(\chi_{c0} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$ Γ_8/Γ_7

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<7.4	90	BRIERE	04 CLEO	$e^+e^- \rightarrow J/\psi X$

$\Gamma(\chi_{c1} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$ Γ_9/Γ_7

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.35±0.08±0.06	52 ± 12	BRIERE	04 CLEO	$e^+e^- \rightarrow J/\psi X$

$\Gamma(\chi_{c2} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$ Γ_{10}/Γ_7

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.52±0.12±0.09	47 ± 11	BRIERE	04 CLEO	$e^+e^- \rightarrow J/\psi X$

$\Gamma(\psi(2S) \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$ Γ_{11}/Γ_7

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.41±0.11±0.08	42 ± 11	BRIERE	04 CLEO	$e^+e^- \rightarrow J/\psi \pi^+ \pi^- X$

$\Gamma(\rho\pi)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 2	90	FULTON	90B	$\Upsilon(1S) \rightarrow \rho^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<10	90	BLINOV	90	MD1 $\Upsilon(1S) \rightarrow \rho^0 \pi^0$
<21	90	NICZYPORUK	83	LENA $\Upsilon(1S) \rightarrow \rho^0 \pi^0$

$\Gamma(\pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	BARU	92	MD1 $\Upsilon(1S) \rightarrow \pi^+ \pi^-$

$\Gamma(K^+ K^-)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	BARU	92	MD1 $\Upsilon(1S) \rightarrow K^+ K^-$

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	¹⁴ BARU	96	MD1 $\Upsilon(1S) \rightarrow p\bar{p}$

¹⁴Supersedes BARU 92 in this node.

$\Gamma(\pi^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.84	90	ANASTASSOV	99	CLE2 $e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(D^*(2010)^\pm \text{ anything})/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
25.2±1.3±1.5	$\approx 2k$	¹⁵	AUBERT	10c	BABR $\Upsilon(2S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<19	90	¹⁶	ALBRECHT	92J	ARG $e^+ e^- \rightarrow D^0 \pi^\pm X$
-----	----	---------------	----------	-----	---

¹⁵For $x_p > 0.1$.

¹⁶For $x_p > 0.2$.

$\Gamma(\bar{d} \text{ anything})/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
2.86±0.19±0.21	455	ASNER	07	CLEO $e^+ e^- \rightarrow \bar{d} X$

$\Gamma(ggg, gg\gamma \rightarrow \bar{d} \text{ anything})/\Gamma(ggg, gg\gamma \rightarrow \text{anything})$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.36±0.23±0.25	455	ASNER	07	CLEO $e^+ e^- \rightarrow \bar{d} X$

$\Gamma(\gamma\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
6.3±1.2±1.3	¹⁷ ANASTASSOV	99	CLE2 $e^+ e^- \rightarrow \text{hadrons}$

¹⁷For $m_{\pi\pi} > 1 \text{ GeV}$.

$\Gamma(\gamma\pi^0\pi^0)/\Gamma_{\text{total}}$					Γ_{20}/Γ
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$1.7 \pm 0.6 \pm 0.3$		¹⁸ ANASTASSOV 99	CLE2	$e^+e^- \rightarrow$ hadrons	
¹⁸ For $m_{\pi\pi} > 1$ GeV.					
$\Gamma(\gamma\pi^0\eta)/\Gamma_{\text{total}}$					Γ_{21}/Γ
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 2.4	90	¹⁹ BESSON 07A	CLEO	$e^+e^- \rightarrow \Upsilon(1S)$	
¹⁹ BESSON 07A obtained this limit for $0.7 < m_{\pi^0\eta} < 3$ GeV.					
$\Gamma(\gamma K^+ K^-)/\Gamma_{\text{total}}$ ($2 < m_{K^+ K^-} < 3$ GeV)					Γ_{22}/Γ
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$1.14 \pm 0.08 \pm 0.10$	90	ATHAR 06	CLE3	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$	
$\Gamma(\gamma p\bar{p})/\Gamma_{\text{total}}$ ($2 < m_{p\bar{p}} < 3$ GeV)					Γ_{23}/Γ
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 0.6	90	ATHAR 06	CLE3	$\Upsilon(1S) \rightarrow \gamma p\bar{p}$	
$\Gamma(\gamma 2h^+ 2h^-)/\Gamma_{\text{total}}$					Γ_{24}/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$7.0 \pm 1.1 \pm 1.0$	80 ± 12	FULTON 90B	CLEO	$e^+e^- \rightarrow$ hadrons	
$\Gamma(\gamma 3h^+ 3h^-)/\Gamma_{\text{total}}$					Γ_{25}/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$5.4 \pm 1.5 \pm 1.3$	39 ± 11	FULTON 90B	CLEO	$e^+e^- \rightarrow$ hadrons	
$\Gamma(\gamma 4h^+ 4h^-)/\Gamma_{\text{total}}$					Γ_{26}/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$7.4 \pm 2.5 \pm 2.5$	36 ± 12	FULTON 90B	CLEO	$e^+e^- \rightarrow$ hadrons	
$\Gamma(\gamma\pi^+\pi^-\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{27}/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$2.9 \pm 0.7 \pm 0.6$	29 ± 8	FULTON 90B	CLEO	$e^+e^- \rightarrow$ hadrons	
$\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$					Γ_{28}/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$2.5 \pm 0.7 \pm 0.5$	26 ± 7	FULTON 90B	CLEO	$e^+e^- \rightarrow$ hadrons	
$\Gamma(\gamma 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$					Γ_{29}/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$2.5 \pm 0.9 \pm 0.8$	17 ± 5	FULTON 90B	CLEO	$e^+e^- \rightarrow$ hadrons	

$\Gamma(\gamma 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{30}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.4 \pm 0.9 \pm 0.8$	18 ± 7	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma \pi^+ \pi^- \rho \bar{\rho})/\Gamma_{\text{total}}$ Γ_{31}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.5 \pm 0.5 \pm 0.3$	22 ± 6	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma 2\pi^+ 2\pi^- \rho \bar{\rho})/\Gamma_{\text{total}}$ Γ_{32}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.4 \pm 0.4 \pm 0.4$	7 ± 6	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma 2K^+ 2K^-)/\Gamma_{\text{total}}$ Γ_{33}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.2 ± 0.2	2 ± 2	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma \eta'(958))/\Gamma_{\text{total}}$ Γ_{34}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 1.9	90	ATHAR 07A	CLEO	$\Upsilon(1S) \rightarrow \gamma \eta' \rightarrow \gamma \pi^+ \pi^- \eta, \gamma \rho$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 16	90	RICHICHI 01B	CLE2	$\Upsilon(1S) \rightarrow \gamma \eta' \rightarrow \gamma \eta \pi^+ \pi^-$

$\Gamma(\gamma \eta)/\Gamma_{\text{total}}$ Γ_{35}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 1.0	90	ATHAR 07A	CLEO	$\Upsilon(1S) \rightarrow \gamma \eta \rightarrow \gamma \gamma \gamma, \gamma \pi^+ \pi^- \pi^0, \gamma 3\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 21	90	MASEK 02	CLEO	$\Upsilon(1S) \rightarrow \gamma \eta$

$\Gamma(\gamma f_0(980))/\Gamma_{\text{total}}$ Γ_{36}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 3	90	²⁰ ATHAR 06	CLE3	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
²⁰ Assuming $B(f_0(980) \rightarrow \pi \pi) = 1$.				

$\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$ Γ_{37}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.7^{+0.9}_{-0.7} \pm 0.8$		ATHAR 06	CLE3	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 14	90	²¹ FULTON 90B	CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
< 19.4	90	²¹ ALBRECHT 89	ARG	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
²¹ Assuming $B(f'_2(1525) \rightarrow K \bar{K}) = 0.71$.				

$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
10.1 ± 0.9 OUR AVERAGE				
$10.5 \pm 1.6^{+1.9}_{-1.8}$		22 BESSON	07A CLE3	$\Upsilon(1S) \rightarrow \gamma \pi^0 \pi^0$
$10.2 \pm 0.8 \pm 0.7$		ATHAR	06 CLE3	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
$8.1 \pm 2.3^{+2.9}_{-2.7}$		23 ANASTASSOV 99	CLE2	$e^+ e^- \rightarrow \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<21	90	23 FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
<13	90	23 ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
<81	90	SCHMITT	88 CBAL	$\Upsilon(1S) \rightarrow \gamma X$

²² Using $B(f_2(1270) \rightarrow \pi^0 \pi^0) = B(f_2(1270) \rightarrow \pi \pi)/3$ and $B(f_2(1270) \rightarrow \pi \pi) = (0.845^{+0.025}_{-0.012})\%$.

²³ Using $B(f_2(1270) \rightarrow \pi \pi) = 0.84$.

$\Gamma(\gamma \eta(1405))/\Gamma_{\text{total}}$ Γ_{39}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<8.2	90	24 FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K^\pm \pi^\mp K_S^0$

²⁴ Includes unknown branching ratio of $\eta(1405) \rightarrow K^\pm \pi^\mp K_S^0$.

$\Gamma(\gamma f_0(1500))/\Gamma_{\text{total}}$ Γ_{40}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.5	90	25 BESSON	07A CLEO	$e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \pi^0 \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.1	90	26 BESSON	07A CLEO	$e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \eta \eta$
------	----	-----------	----------	---

²⁵ Using $B(f_0(1500) \rightarrow \pi^0 \pi^0) = B(f_0(1500) \rightarrow \pi \pi)/3$ and $B(f_0(1500) \rightarrow \pi \pi) = (0.349 \pm 0.023)\%$.

²⁶ Calculated by us using $B(f_0(1500) \rightarrow \eta \eta) = (5.1 \pm 0.9)\%$.

$\Gamma(\gamma f_0(1710))/\Gamma_{\text{total}}$ Γ_{41}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.6	90	27 ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 6.3	90	27 FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
<19	90	27 FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K_S^0 K_S^0$
< 8	90	28 ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
<24	90	29 SCHMITT	88 CBAL	$\Upsilon(1S) \rightarrow \gamma X$

²⁷ Assuming $B(f_0(1710) \rightarrow K \bar{K}) = 0.38$.

²⁸ Assuming $B(f_0(1710) \rightarrow \pi \pi) = 0.04$.

²⁹ Assuming $B(f_0(1710) \rightarrow \eta \eta) = 0.18$.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma K^+ K^-)/\Gamma_{\text{total}}$ Γ_{42}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<0.7	90	ATHAR	06 CLEO	$e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma K^+ K^-$

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi^0 \pi^0) / \Gamma_{\text{total}}$ Γ_{43} / Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<1.4	90	BESSON	07A CLEO	$e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \pi^0 \pi^0$

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{44} / Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<1.8	90	BESSON	07A CLEO	$e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \eta \eta$

$\Gamma(\gamma f_4(2050)) / \Gamma_{\text{total}}$ Γ_{45} / Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<5.3	90	³⁰ ATHAR	06 CLE3	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$

³⁰ Assuming $B(f_4(2050) \rightarrow \pi \pi) = 0.17$.

$\Gamma(\gamma f_0(2200) \rightarrow \gamma K^+ K^-) / \Gamma_{\text{total}}$ Γ_{46} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0002	90	BARU	89 MD1	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$

$\Gamma(\gamma f_J(2220) \rightarrow \gamma K^+ K^-) / \Gamma_{\text{total}}$ Γ_{47} / Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
< 8	90	ATHAR	06 CLE3	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 160	90	MASEK	02 CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
< 150	90	FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
< 290	90	ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
<2000	90	BARU	89 MD1	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$

$\Gamma(\gamma f_J(2220) \rightarrow \gamma \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{48} / Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
< 6	90	ATHAR	06 CLE3	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<120	90	MASEK	02 CLEO	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
------	----	-------	---------	---

$\Gamma(\gamma f_J(2220) \rightarrow \gamma p \bar{p}) / \Gamma_{\text{total}}$ Γ_{49} / Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
< 11	90	ATHAR	06 CLE3	$\Upsilon(1S) \rightarrow \gamma p \bar{p}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<160	90	MASEK	02 CLEO	$\Upsilon(1S) \rightarrow \gamma p \bar{p}$
------	----	-------	---------	---

$\Gamma(\gamma \eta(2225) \rightarrow \gamma \phi \phi) / \Gamma_{\text{total}}$ Γ_{50} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.003	90	BARU	89 MD1	$\Upsilon(1S) \rightarrow \gamma K^+ K^- K^+ K^-$

$\Gamma(\gamma X) / \Gamma_{\text{total}}$ Γ_{51} / Γ
 (X = pseudoscalar with $m < 7.2$ GeV)

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<3	90	³¹ BALEST	95 CLEO	$e^+ e^- \rightarrow \gamma + X$

³¹ For a noninteracting pseudoscalar X with mass < 7.2 GeV.

$\Gamma(\gamma X \bar{X})/\Gamma_{\text{total}}$ **Γ_{52}/Γ**

($X \bar{X}$ = vectors with $m < 3.1$ GeV)

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1	90	³² BALEST	95	CLEO $e^+e^- \rightarrow \gamma + X \bar{X}$

³²For a noninteracting vector X with mass < 3.1 GeV.

$\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$ **Γ_{53}/Γ**

($1.5 \text{ GeV} < m_X < 5.0 \text{ GeV}$)

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.78	95	ROSNER	07A	CLEO $e^+e^- \rightarrow \gamma X$

$\Gamma(\gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{54}/Γ**

($201 < M(\mu^+ \mu^-) < 3565 \text{ MeV}$)

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<9	90	³³ LOVE	08	CLEO $e^+e^- \rightarrow \gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-$

³³For a narrow scalar or pseudoscalar a_1^0 with $201 < M(\mu^+ \mu^-) < 3565 \text{ MeV}$, excluding J/ψ . Measured 90% CL limits as a function of $M(\mu^+ \mu^-)$ range from $1-9 \times 10^{-6}$.

$\Gamma(\gamma a_1^0 \rightarrow \gamma \tau^+ \tau^-)/\Gamma_{\text{total}}$ **Γ_{55}/Γ**

($2m_\tau < M(\tau^+ \tau^-) < 7500 \text{ MeV}$)

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<50	90	³⁴ LOVE	08	CLEO $e^+e^- \rightarrow \gamma a_1^0 \rightarrow \gamma \tau^+ \tau^-$

³⁴For a narrow scalar or pseudoscalar a_1^0 with $2m_\tau < M(\tau^+ \tau^-) < 7500 \text{ MeV}$. Measured 90% CL limits as a function of $M(\tau^+ \tau^-)$ range from $1-5 \times 10^{-5}$.

$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$ **Γ_{56}/Γ**

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<6.0	95	LOVE	08A	CLEO $e^+e^- \rightarrow \mu^\pm \tau^\mp$

$\Gamma(\text{invisible})/\Gamma_{\text{total}}$ **Γ_{57}/Γ**

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.0	90	AUBERT	09AX	BABR $\Upsilon(3S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<39	90	RUBIN	07	CLEO $\Upsilon(2S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$
<25	90	TAJIMA	07	BELL $\Upsilon(3S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$

$\Upsilon(1S)$ REFERENCES

AUBERT	10C	PR D81 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09AX	PRL 103 251801	B. Aubert <i>et al.</i>	(BABAR Collab.)
LOVE	08	PRL 101 151802	W. Love <i>et al.</i>	(CLEO Collab.)
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
ASNER	07	PR D75 012009	D.M. Asner <i>et al.</i>	(CLEO Collab.)
ATHAR	07A	PR D76 072003	S.B. Athar <i>et al.</i>	(CLEO Collab.)
BESSION	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)
BESSION	07A	PR D75 072001	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
RUBIN	07	PR D75 031104	P. Rubin <i>et al.</i>	(CLEO Collab.)

TAJIMA	07	PRL 98 132001	O. Tajima <i>et al.</i>	(BELLE Collab.)
AQUINES	06A	PR D74 092006	O. Aquines <i>et al.</i>	(CLEO Collab.)
ATHAR	06	PR D73 032001	S.B. Athar <i>et al.</i>	(CLEO Collab.)
BESSON	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)
BRIERE	04	PR D70 072001	R.A. Briere <i>et al.</i>	(CLEO Collab.)
ARTUSO	03	PR D67 052003	M. Artuso <i>et al.</i>	(CLEO Collab.)
MASEK	02	PR D65 072002	G. Masek <i>et al.</i>	(CLEO Collab.)
RICHICHI	01B	PRL 87 141801	S.J. Richichi <i>et al.</i>	(CLEO Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
ANASTASSOV	99	PRL 82 286	A. Anastassov <i>et al.</i>	(CLEO Collab.)
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
BARU	96	PRPL 267 71	S.E. Baru <i>et al.</i>	(NOVO)
BALEST	95	PR D51 2053	R. Balest <i>et al.</i>	(CLEO Collab.)
CINABRO	94B	PL B340 129	D. Cinabro <i>et al.</i>	(CLEO Collab.)
ALBRECHT	92J	ZPHY C55 25	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BARU	92	ZPHY C54 229	S.E. Baru <i>et al.</i>	(NOVO)
BARU	92B	ZPHY C56 547	S.E. Baru <i>et al.</i>	(NOVO)
KOBEL	92	ZPHY C53 193	M. Kobel <i>et al.</i>	(Crystal Ball Collab.)
BLINOV	90	PL B245 311	A.E. Blinov <i>et al.</i>	(NOVO)
FULTON	90B	PR D41 1401	R. Fulton <i>et al.</i>	(CLEO Collab.)
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	89	ZPHY C42 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)
BARU	89	ZPHY C42 505	S.E. Baru <i>et al.</i>	(NOVO)
CHEN	89B	PR D39 3528	W.Y. Chen <i>et al.</i>	(CLEO Collab.)
FULTON	89	PL B224 445	R. Fulton <i>et al.</i>	(CLEO Collab.)
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)
BUCHMUEL...	88	HE e^+e^- Physics 412	W. Buchmueller, S. Cooper	(HANN, DESY, MIT)
Editors: A. Ali and P. Soeding, World Scientific, Singapore				
JAKUBOWSKI	88	ZPHY C40 49	Z. Jakubowski <i>et al.</i>	(Crystal Ball Collab.) IGJPC
SCHMITT	88	ZPHY C40 199	P. Schmitt <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	87	ZPHY C35 283	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
BARU	86	ZPHY C30 551	S.E. Baru <i>et al.</i>	(NOVO)
ALBRECHT	85C	PL 154B 452	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
Translated from YAF 41 733.				
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)
BESSON	84	PR D30 1433	D. Besson <i>et al.</i>	(CLEO Collab.)
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)
MACKAY	84	PR D29 2483	W.W. MacKay <i>et al.</i>	(CUSB Collab.)
ANDREWS	83	PRL 50 807	D.E. Andrews <i>et al.</i>	(CLEO Collab.)
GILES	83	PRL 50 877	R. Giles <i>et al.</i>	(HARV, OSU, ROCH, RUTG+)
NICZYPORUK	83	ZPHY C17 197	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
ALBRECHT	82	PL 116B 383	H. Albrecht <i>et al.</i>	(DESY, DORT, HEIDH+)
ARTAMONOV	82	PL 118B 225	A.S. Artamonov <i>et al.</i>	(NOVO)
NICZYPORUK	82	ZPHY C15 299	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
BERGER	80C	PL 93B 497	C. Berger <i>et al.</i>	(PLUTO Collab.)
BOCK	80	ZPHY C6 125	P. Bock <i>et al.</i>	(HEIDP, MPIM, DESY, HAMB)
BERGER	79	ZPHY C1 343	C. Berger <i>et al.</i>	(PLUTO Collab.)