

# LIGHT UNFLAVORED MESONS

## ( $S = C = B = 0$ )

For  $I = 1$  ( $\pi, \rho, \omega$ ):  $u\bar{d}, (u\bar{u}-d\bar{d})/\sqrt{2}, d\bar{u}$ ;  
 for  $I = 0$  ( $\eta, \eta', h, h', \omega, \phi, f, f'$ ):  $c_1(u\bar{u} + d\bar{d}) + c_2(s\bar{s})$

$\pi^\pm$

$$I^G(J^P) = 1^-(0^-)$$

Mass  $m = 139.57018 \pm 0.00035$  MeV ( $S = 1.2$ )  
 Mean life  $\tau = (2.6033 \pm 0.0005) \times 10^{-8}$  s ( $S = 1.2$ )  
 $c\tau = 7.8045$  m

$\pi^\pm \rightarrow \ell^\pm \nu \gamma$  form factors [a]

$F_V = 0.0254 \pm 0.0017$   
 $F_A = 0.0119 \pm 0.0001$   
 $F_V$  slope parameter  $a = 0.10 \pm 0.06$   
 $R = 0.059^{+0.009}_{-0.008}$

$\pi^-$  modes are charge conjugates of the modes below.

For decay limits to particles which are not established, see the section on Searches for Axions and Other Very Light Bosons.

| $\pi^+$ DECAY MODES       | Fraction ( $\Gamma_i/\Gamma$ )           | Confidence level       | $p$<br>(MeV/c) |
|---------------------------|--|------------------------|----------------|
| $\mu^+ \nu_\mu$           | [b] (99.98770 ± 0.00004) %               |                        | 30             |
| $\mu^+ \nu_\mu \gamma$    | [c] ( 2.00 ± 0.25 ) × 10 <sup>-4</sup>   |                        | 30             |
| $e^+ \nu_e$               | [b] ( 1.230 ± 0.004 ) × 10 <sup>-4</sup> |                        | 70             |
| $e^+ \nu_e \gamma$        | [c] ( 7.39 ± 0.05 ) × 10 <sup>-7</sup>   |                        | 70             |
| $e^+ \nu_e \pi^0$         | ( 1.036 ± 0.006 ) × 10 <sup>-8</sup>     |                        | 4              |
| $e^+ \nu_e e^+ e^-$       | ( 3.2 ± 0.5 ) × 10 <sup>-9</sup>         |                        | 70             |
| $e^+ \nu_e \nu \bar{\nu}$ | < 5                                      | × 10 <sup>-6</sup> 90% | 70             |

### Lepton Family number (LF) or Lepton number (L) violating modes

|                     |    |           |                        |    |
|---------------------|----|-----------|------------------------|----|
| $\mu^+ \bar{\nu}_e$ | L  | [d] < 1.5 | × 10 <sup>-3</sup> 90% | 30 |
| $\mu^+ \nu_e$       | LF | [d] < 8.0 | × 10 <sup>-3</sup> 90% | 30 |
| $\mu^- e^+ e^+ \nu$ | LF | < 1.6     | × 10 <sup>-6</sup> 90% | 30 |

$\pi^0$

$$I^G(J^{PC}) = 1^-(0^{-+})$$

Mass  $m = 134.9766 \pm 0.0006$  MeV ( $S = 1.1$ )  
 $m_{\pi^\pm} - m_{\pi^0} = 4.5936 \pm 0.0005$  MeV  
 Mean life  $\tau = (8.4 \pm 0.5) \times 10^{-17}$  s ( $S = 2.6$ )  
 $c\tau = 25.1$  nm

For decay limits to particles which are not established, see the appropriate Search sections ( $A^0$  (axion) and Other Light Boson ( $X^0$ ) Searches, etc.).

| $\pi^0$ DECAY MODES  | Fraction ( $\Gamma_i/\Gamma$ )   | Scale factor/<br>Confidence level | $p$<br>(MeV/c)           |    |
|--|----------------------------------|-----------------------------------|--------------------------|----|
| $2\gamma$  | $(98.823 \pm 0.034) \%$          | S=1.5                             | 67                       |    |
| $e^+ e^- \gamma$   | $(1.174 \pm 0.035) \%$           | S=1.5                             | 67                       |    |
| $\gamma$ positronium   | $(1.82 \pm 0.29) \times 10^{-9}$ |                                   | 67                       |    |
| $e^+ e^+ e^- e^-$  | $(3.34 \pm 0.16) \times 10^{-5}$ |                                   | 67                       |    |
| $e^+ e^-$  | $(6.46 \pm 0.33) \times 10^{-8}$ |                                   | 67                       |    |
| $4\gamma$  | $< 2$                            | $\times 10^{-8}$ CL=90%           | 67                       |    |
| $\nu \bar{\nu}$  | [e] $< 2.7$                      | $\times 10^{-7}$ CL=90%           | 67                       |    |
| $\nu_e \bar{\nu}_e$  | $< 1.7$                          | $\times 10^{-6}$ CL=90%           | 67                       |    |
| $\nu_\mu \bar{\nu}_\mu$  | $< 1.6$                          | $\times 10^{-6}$ CL=90%           | 67                       |    |
| $\nu_\tau \bar{\nu}_\tau$  | $< 2.1$                          | $\times 10^{-6}$ CL=90%           | 67                       |    |
| $\gamma \nu \bar{\nu}$   | $< 6$                            | $\times 10^{-4}$ CL=90%           | 67                       |    |
| <b>Charge conjugation (C) or Lepton Family number (LF) violating modes</b> |                                  |                                   |                          |    |
| $3\gamma$  | C                                | $< 3.1$                           | $\times 10^{-8}$ CL=90%  | 67 |
| $\mu^+ e^-$  | LF                               | $< 3.8$                           | $\times 10^{-10}$ CL=90% | 26 |
| $\mu^- e^+$  | LF                               | $< 3.4$                           | $\times 10^{-9}$ CL=90%  | 26 |
| $\mu^+ e^- + \mu^- e^+$  | LF                               | $< 3.6$                           | $\times 10^{-10}$ CL=90% | 26 |

**$\eta$**

$$J^{PC} = 0^+(0^-+)$$

Mass  $m = 547.853 \pm 0.024$  MeV

Full width  $\Gamma = 1.30 \pm 0.07$  keV

**C-nonconserving decay parameters**

$$\begin{aligned} \pi^+ \pi^- \pi^0 \text{ left-right asymmetry} &= (0.09_{-0.12}^{+0.11}) \times 10^{-2} \\ \pi^+ \pi^- \pi^0 \text{ sextant asymmetry} &= (0.12_{-0.11}^{+0.10}) \times 10^{-2} \\ \pi^+ \pi^- \pi^0 \text{ quadrant asymmetry} &= (-0.09 \pm 0.09) \times 10^{-2} \\ \pi^+ \pi^- \gamma \text{ left-right asymmetry} &= (0.9 \pm 0.4) \times 10^{-2} \\ \pi^+ \pi^- \gamma \beta \text{ (D-wave)} &= -0.02 \pm 0.07 \quad (S = 1.3) \end{aligned}$$

**CP-nonconserving decay parameters**

$$\pi^+ \pi^- e^+ e^- \text{ decay-plane asymmetry } A_\phi = (-0.6 \pm 3.1) \times 10^{-2}$$

**Dalitz plot parameter**

$$\pi^0 \pi^0 \pi^0 \quad \alpha = -0.0317 \pm 0.0016$$

| $\eta$ DECAY MODES  | Fraction ( $\Gamma_i/\Gamma$ )  | Scale factor/<br>Confidence level | $p$<br>(MeV/c) |     |
|---|---------------------------------|-----------------------------------|----------------|-----|
| <b>Neutral modes</b>  |                                 |                                   |                |     |
| neutral modes   | (71.90±0.34) %                  | S=1.2                             | –              |     |
| 2 $\gamma$  | (39.31±0.20) %                  | S=1.1                             | 274            |     |
| 3 $\pi^0$   | (32.57±0.23) %                  | S=1.1                             | 179            |     |
| $\pi^0 2\gamma$   | ( 2.7 ±0.5 ) × 10 <sup>-4</sup> | S=1.1                             | 257            |     |
| 2 $\pi^0 2\gamma$   | < 1.2 × 10 <sup>-3</sup>        | CL=90%                            | 238            |     |
| 4 $\gamma$  | < 2.8 × 10 <sup>-4</sup>        | CL=90%                            | 274            |     |
| invisible   | < 6 × 10 <sup>-4</sup>          | CL=90%                            | –              |     |
| <b>Charged modes</b>  |                                 |                                   |                |     |
| charged modes   | (28.10±0.34) %                  | S=1.2                             | –              |     |
| $\pi^+ \pi^- \pi^0$   | (22.74±0.28) %                  | S=1.2                             | 174            |     |
| $\pi^+ \pi^- \gamma$  | ( 4.60±0.16) %                  | S=2.1                             | 236            |     |
| $e^+ e^- \gamma$  | ( 7.0 ±0.7 ) × 10 <sup>-3</sup> | S=1.5                             | 274            |     |
| $\mu^+ \mu^- \gamma$  | ( 3.1 ±0.4 ) × 10 <sup>-4</sup> |                                   | 253            |     |
| $e^+ e^-$   | < 2.7 × 10 <sup>-5</sup>        | CL=90%                            | 274            |     |
| $\mu^+ \mu^-$   | ( 5.8 ±0.8 ) × 10 <sup>-6</sup> |                                   | 253            |     |
| 2 $e^+ 2e^-$  | < 6.9 × 10 <sup>-5</sup>        | CL=90%                            | 274            |     |
| $\pi^+ \pi^- e^+ e^- (\gamma)$  | ( 2.68±0.11) × 10 <sup>-4</sup> |                                   | 235            |     |
| $e^+ e^- \mu^+ \mu^-$   | < 1.6 × 10 <sup>-4</sup>        | CL=90%                            | 253            |     |
| 2 $\mu^+ 2\mu^-$  | < 3.6 × 10 <sup>-4</sup>        | CL=90%                            | 161            |     |
| $\mu^+ \mu^- \pi^+ \pi^-$   | < 3.6 × 10 <sup>-4</sup>        | CL=90%                            | 113            |     |
| $\pi^+ \pi^- 2\gamma$   | < 2.0 × 10 <sup>-3</sup>        |                                   | 236            |     |
| $\pi^+ \pi^- \pi^0 \gamma$  | < 5 × 10 <sup>-4</sup>          | CL=90%                            | 174            |     |
| $\pi^0 \mu^+ \mu^- \gamma$  | < 3 × 10 <sup>-6</sup>          | CL=90%                            | 210            |     |
| <b>Charge conjugation (C), Parity (P),<br/>Charge conjugation × Parity (CP), or<br/>Lepton Family number (LF) violating modes</b> |                                 |                                   |                |     |
| $\pi^0 \gamma$  | C                               | < 9 × 10 <sup>-5</sup>            | CL=90%         | 257 |
| $\pi^+ \pi^-$   | P,CP                            | < 1.3 × 10 <sup>-5</sup>          | CL=90%         | 236 |
| 2 $\pi^0$   | P,CP                            | < 3.5 × 10 <sup>-4</sup>          | CL=90%         | 238 |
| 2 $\pi^0 \gamma$  | C                               | < 5 × 10 <sup>-4</sup>            | CL=90%         | 238 |
| 3 $\pi^0 \gamma$  | C                               | < 6 × 10 <sup>-5</sup>            | CL=90%         | 179 |
| 3 $\gamma$  | C                               | < 1.6 × 10 <sup>-5</sup>          | CL=90%         | 274 |
| 4 $\pi^0$   | P,CP                            | < 6.9 × 10 <sup>-7</sup>          | CL=90%         | 40  |
| $\pi^0 e^+ e^-$   | C                               | [f] < 4 × 10 <sup>-5</sup>        | CL=90%         | 257 |
| $\pi^0 \mu^+ \mu^-$   | C                               | [f] < 5 × 10 <sup>-6</sup>        | CL=90%         | 210 |
| $\mu^+ e^- + \mu^- e^+$   | LF                              | < 6 × 10 <sup>-6</sup>            | CL=90%         | 264 |

**$f_0(600)$  [g]**  
or  $\sigma$

$$I^G(J^{PC}) = 0^+(0^{++})$$

Mass  $m = (400\text{--}1200)$  MeV

Full width  $\Gamma = (600\text{--}1000)$  MeV

| <b><math>f_0(600)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|--|--------------------------------|-------------|
| $\pi\pi$                                 | dominant                       | —           |
| $\gamma\gamma$                           | seen                           | —           |

**$\rho(770)$  [h]**

$$I^G(J^{PC}) = 1^+(1^{--})$$

Mass  $m = 775.49 \pm 0.34$  MeV

Full width  $\Gamma = 149.1 \pm 0.8$  MeV

$\Gamma_{ee} = 7.04 \pm 0.06$  keV

| <b><math>\rho(770)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ )                       | Scale factor/<br>Confidence level | $p$<br>(MeV/c) |
|---|--|-----------------------------------|----------------|
| $\pi\pi$                                  | $\sim 100$ %   |                                   | 363            |
| <b><math>\rho(770)^\pm</math> decays</b>  |  |                                   |                |
| $\pi^\pm\gamma$                           | ( $4.5 \pm 0.5$ ) $\times 10^{-4}$                   | S=2.2                             | 375            |
| $\pi^\pm\eta$                             | < 6 $\times 10^{-3}$                                 | CL=84%                            | 153            |
| $\pi^\pm\pi^+\pi^-\pi^0$                  | < 2.0 $\times 10^{-3}$                               | CL=84%                            | 254            |
| <b><math>\rho(770)^0</math> decays</b>    |  |                                   |                |
| $\pi^+\pi^-\gamma$                        | ( $9.9 \pm 1.6$ ) $\times 10^{-3}$                   |                                   | 362            |
| $\pi^0\gamma$                             | ( $6.0 \pm 0.8$ ) $\times 10^{-4}$                   |                                   | 376            |
| $\eta\gamma$                              | ( $3.00 \pm 0.20$ ) $\times 10^{-4}$                 |                                   | 194            |
| $\pi^0\pi^0\gamma$                        | ( $4.5 \pm 0.8$ ) $\times 10^{-5}$                   |                                   | 363            |
| $\mu^+\mu^-$                              | [i] ( $4.55 \pm 0.28$ ) $\times 10^{-5}$             |                                   | 373            |
| $e^+e^-$                                  | [i] ( $4.72 \pm 0.05$ ) $\times 10^{-5}$             |                                   | 388            |
| $\pi^+\pi^-\pi^0$                         | ( $1.01^{+0.54}_{-0.36} \pm 0.34$ ) $\times 10^{-4}$ |                                   | 323            |
| $\pi^+\pi^-\pi^+\pi^-$                    | ( $1.8 \pm 0.9$ ) $\times 10^{-5}$                   |                                   | 251            |
| $\pi^+\pi^-\pi^0\pi^0$                    | ( $1.6 \pm 0.8$ ) $\times 10^{-5}$                   |                                   | 257            |
| $\pi^0e^+e^-$                             | < 1.2 $\times 10^{-5}$                               | CL=90%                            | 376            |

**$\omega(782)$**

$$I^G(J^{PC}) = 0^-(1^{--})$$

Mass  $m = 782.65 \pm 0.12$  MeV (S = 1.9)

Full width  $\Gamma = 8.49 \pm 0.08$  MeV

$\Gamma_{ee} = 0.60 \pm 0.02$  keV

| $\omega(782)$ DECAY MODES           | Fraction ( $\Gamma_i/\Gamma$ )      | Scale factor/<br>Confidence level | $p$<br>(MeV/c) |
|-------------------------------------|-------------------------------------|-----------------------------------|----------------|
| $\pi^+\pi^-\pi^0$                   | (89.2 $\pm$ 0.7 ) %                 |                                   | 327            |
| $\pi^0\gamma$                       | ( 8.28 $\pm$ 0.28) %                | S=2.1                             | 380            |
| $\pi^+\pi^-$                        | ( 1.53 $^{+0.11}_{-0.13}$ ) %       | S=1.2                             | 366            |
| neutrals (excluding $\pi^0\gamma$ ) | ( 8 $^{+8}_{-5}$ ) $\times 10^{-3}$ | S=1.1                             | –              |
| $\eta\gamma$                        | ( 4.6 $\pm$ 0.4 ) $\times 10^{-4}$  | S=1.1                             | 200            |
| $\pi^0e^+e^-$                       | ( 7.7 $\pm$ 0.6 ) $\times 10^{-4}$  |                                   | 380            |
| $\pi^0\mu^+\mu^-$                   | ( 1.3 $\pm$ 0.4 ) $\times 10^{-4}$  | S=2.1                             | 349            |
| $e^+e^-$                            | ( 7.28 $\pm$ 0.14) $\times 10^{-5}$ | S=1.3                             | 391            |
| $\pi^+\pi^-\pi^0\pi^0$              | < 2 $\times 10^{-4}$                | CL=90%                            | 262            |
| $\pi^+\pi^-\gamma$                  | < 3.6 $\times 10^{-3}$              | CL=95%                            | 366            |
| $\pi^+\pi^-\pi^+\pi^-$              | < 1 $\times 10^{-3}$                | CL=90%                            | 256            |
| $\pi^0\pi^0\gamma$                  | ( 6.6 $\pm$ 1.1 ) $\times 10^{-5}$  |                                   | 367            |
| $\eta\pi^0\gamma$                   | < 3.3 $\times 10^{-5}$              | CL=90%                            | 162            |
| $\mu^+\mu^-$                        | ( 9.0 $\pm$ 3.1 ) $\times 10^{-5}$  |                                   | 377            |
| $3\gamma$                           | < 1.9 $\times 10^{-4}$              | CL=95%                            | 391            |

**Charge conjugation (C) violating modes**

|             |   |                        |        |     |
|-------------|---|------------------------|--------|-----|
| $\eta\pi^0$ | C | < 2.1 $\times 10^{-4}$ | CL=90% | 162 |
| $2\pi^0$    | C | < 2.1 $\times 10^{-4}$ | CL=90% | 367 |
| $3\pi^0$    | C | < 2.3 $\times 10^{-4}$ | CL=90% | 330 |

**$\eta'(958)$**

$$I^G(J^{PC}) = 0^+(0^{-+})$$

Mass  $m = 957.78 \pm 0.06$  MeV

Full width  $\Gamma = 0.194 \pm 0.009$  MeV

| $\eta'(958)$ DECAY MODES                                       | Fraction ( $\Gamma_i/\Gamma$ )            | Confidence level | $p$<br>(MeV/c) |
|--|---|------------------|----------------|
| $\pi^+\pi^-\eta$   | (43.2 $\pm$ 0.7 ) %                       |                  | 232            |
| $\rho^0\gamma$ (including non-resonant<br>$\pi^+\pi^-\gamma$ ) | (29.3 $\pm$ 0.5 ) %                       |                  | 165            |
| $\pi^0\pi^0\eta$   | (21.7 $\pm$ 0.8 ) %                       |                  | 239            |
| $\omega\gamma$   | ( 2.75 $\pm$ 0.22) %                      |                  | 159            |
| $\gamma\gamma$   | ( 2.22 $\pm$ 0.08) %                      |                  | 479            |
| $3\pi^0$   | ( 1.68 $\pm$ 0.22) $\times 10^{-3}$       |                  | 430            |
| $\mu^+\mu^-\gamma$   | ( 1.09 $\pm$ 0.27) $\times 10^{-4}$       |                  | 467            |
| $\pi^+\pi^-\mu^+\mu^-$   | < 2.2 $\times 10^{-4}$                    | 90%              | 401            |
| $\pi^+\pi^-\pi^0$  | ( 3.6 $^{+1.1}_{-0.9}$ ) $\times 10^{-3}$ |                  | 428            |
| $\pi^0\rho^0$  | < 4 %                                     | 90%              | 111            |
| $2(\pi^+\pi^-)$  | < 2.4 $\times 10^{-4}$                    | 90%              | 372            |

|                           |                                      |                  |     |     |
|---------------------------|--------------------------------------|------------------|-----|-----|
| $\pi^+ \pi^- 2\pi^0$      | < 2.5                                | $\times 10^{-3}$ | 90% | 376 |
| $2(\pi^+ \pi^-)$ neutrals | < 1                                  | %                | 95% | —   |
| $2(\pi^+ \pi^-)\pi^0$     | < 1.9                                | $\times 10^{-3}$ | 90% | 298 |
| $2(\pi^+ \pi^-)2\pi^0$    | < 1                                  | %                | 95% | 197 |
| $3(\pi^+ \pi^-)$          | < 5                                  | $\times 10^{-4}$ | 90% | 189 |
| $\pi^+ \pi^- e^+ e^-$     | $(2.4^{+1.3}_{-1.0}) \times 10^{-3}$ |                  |     | 458 |
| $\gamma e^+ e^-$          | < 9                                  | $\times 10^{-4}$ | 90% | 479 |
| $\pi^0 \gamma \gamma$     | < 8                                  | $\times 10^{-4}$ | 90% | 469 |
| $4\pi^0$                  | < 5                                  | $\times 10^{-4}$ | 90% | 380 |
| $e^+ e^-$                 | < 2.1                                | $\times 10^{-7}$ | 90% | 479 |
| invisible                 | < 9                                  | $\times 10^{-4}$ | 90% | —   |

**Charge conjugation (C), Parity (P),  
Lepton family number (LF) violating modes**

|                     |              |                    |                  |     |     |
|---------------------|--------------|--------------------|------------------|-----|-----|
| $\pi^+ \pi^-$       | <i>P, CP</i> | < 2.9              | $\times 10^{-3}$ | 90% | 458 |
| $\pi^0 \pi^0$       | <i>P, CP</i> | < 1.0              | $\times 10^{-3}$ | 90% | 459 |
| $\pi^0 e^+ e^-$     | <i>C</i>     | [ <i>f</i> ] < 1.4 | $\times 10^{-3}$ | 90% | 469 |
| $\eta e^+ e^-$      | <i>C</i>     | [ <i>f</i> ] < 2.4 | $\times 10^{-3}$ | 90% | 322 |
| $3\gamma$           | <i>C</i>     | < 1.0              | $\times 10^{-4}$ | 90% | 479 |
| $\mu^+ \mu^- \pi^0$ | <i>C</i>     | [ <i>f</i> ] < 6.0 | $\times 10^{-5}$ | 90% | 445 |
| $\mu^+ \mu^- \eta$  | <i>C</i>     | [ <i>f</i> ] < 1.5 | $\times 10^{-5}$ | 90% | 273 |
| $e\mu$              | <i>LF</i>    | < 4.7              | $\times 10^{-4}$ | 90% | 473 |

**$f_0(980)$**  [*J*]

$$I^G(J^{PC}) = 0^+(0^{++})$$

Mass  $m = 980 \pm 10$  MeV

Full width  $\Gamma = 40$  to  $100$  MeV

| <b><math>f_0(980)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | <i>p</i> (MeV/c) |
|--|--------------------------------|------------------|
| $\pi \pi$                                | dominant                       | 471              |
| $K \bar{K}$                              | seen                           | †                |
| $\gamma \gamma$                          | seen                           | 490              |

**$a_0(980)$**  [*J*]

$$I^G(J^{PC}) = 1^-(0^{++})$$

Mass  $m = 980 \pm 20$  MeV

Full width  $\Gamma = 50$  to  $100$  MeV

| <b><math>a_0(980)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|--|--------------------------------|-------------|
| $\eta\pi$                                | dominant                       | 319         |
| $K\bar{K}$                               | seen                           | †           |
| $\gamma\gamma$                           | seen                           | 490         |

**$\phi(1020)$**

$$J^{PC} = 0^-(1^--)$$

Mass  $m = 1019.455 \pm 0.020$  MeV (S = 1.1)

Full width  $\Gamma = 4.26 \pm 0.04$  MeV (S = 1.4)

| <b><math>\phi(1020)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ )                               | Scale factor/<br>Confidence level | $p$<br>(MeV/c) |
|--|--|-----------------------------------|----------------|
| $K^+K^-$                                   | (48.9 ± 0.5 ) %  | S=1.1                             | 127            |
| $K_L^0 K_S^0$                              | (34.2 ± 0.4 ) %  | S=1.1                             | 110            |
| $\rho\pi + \pi^+\pi^-\pi^0$                | (15.32 ± 0.32 ) %  | S=1.1                             | —              |
| $\eta\gamma$                               | ( 1.309 ± 0.024 ) %  | S=1.2                             | 363            |
| $\pi^0\gamma$                              | ( 1.27 ± 0.06 ) × 10 <sup>-3</sup>                           |                                   | 501            |
| $\ell^+\ell^-$                             | —  |                                   | 510            |
| $e^+e^-$                                   | ( 2.954 ± 0.030 ) × 10 <sup>-4</sup>                         | S=1.1                             | 510            |
| $\mu^+\mu^-$                               | ( 2.87 ± 0.19 ) × 10 <sup>-4</sup>                           |                                   | 499            |
| $\eta e^+e^-$                              | ( 1.15 ± 0.10 ) × 10 <sup>-4</sup>                           |                                   | 363            |
| $\pi^+\pi^-$                               | ( 7.4 ± 1.3 ) × 10 <sup>-5</sup>                             |                                   | 490            |
| $\omega\pi^0$                              | ( 4.7 ± 0.5 ) × 10 <sup>-5</sup>                             |                                   | 171            |
| $\omega\gamma$                             | < 5 %  | CL=84%                            | 209            |
| $\rho\gamma$                               | < 1.2 × 10 <sup>-5</sup>                                     | CL=90%                            | 215            |
| $\pi^+\pi^-\gamma$                         | ( 4.1 ± 1.3 ) × 10 <sup>-5</sup>                             |                                   | 490            |
| $f_0(980)\gamma$                           | ( 3.22 ± 0.19 ) × 10 <sup>-4</sup>                           | S=1.1                             | 39             |
| $\pi^0\pi^0\gamma$                         | ( 1.13 ± 0.06 ) × 10 <sup>-4</sup>                           |                                   | 492            |
| $\pi^+\pi^-\pi^+\pi^-$                     | ( 4.0 <sup>+2.8</sup> / <sub>-2.2</sub> ) × 10 <sup>-6</sup> |                                   | 410            |
| $\pi^+\pi^+\pi^-\pi^-\pi^0$                | < 4.6 × 10 <sup>-6</sup>                                     | CL=90%                            | 342            |
| $\pi^0 e^+ e^-$                            | ( 1.12 ± 0.28 ) × 10 <sup>-5</sup>                           |                                   | 501            |
| $\pi^0\eta\gamma$                          | ( 7.27 ± 0.30 ) × 10 <sup>-5</sup>                           | S=1.5                             | 346            |
| $a_0(980)\gamma$                           | ( 7.6 ± 0.6 ) × 10 <sup>-5</sup>                             |                                   | 39             |
| $K^0\bar{K}^0\gamma$                       | < 1.9 × 10 <sup>-8</sup>                                     | CL=90%                            | 110            |
| $\eta'(958)\gamma$                         | ( 6.25 ± 0.21 ) × 10 <sup>-5</sup>                           |                                   | 60             |
| $\eta\pi^0\pi^0\gamma$                     | < 2 × 10 <sup>-5</sup>                                       | CL=90%                            | 293            |
| $\mu^+\mu^-\gamma$                         | ( 1.4 ± 0.5 ) × 10 <sup>-5</sup>                             |                                   | 499            |
| $\rho\gamma\gamma$                         | < 1.2 × 10 <sup>-4</sup>                                     | CL=90%                            | 215            |
| $\eta\pi^+\pi^-$                           | < 1.8 × 10 <sup>-5</sup>                                     | CL=90%                            | 288            |
| $\eta\mu^+\mu^-$                           | < 9.4 × 10 <sup>-6</sup>                                     | CL=90%                            | 321            |

**$h_1(1170)$**

$$I^G(J^{PC}) = 0^-(1^{+-})$$

Mass  $m = 1170 \pm 20$  MeV

Full width  $\Gamma = 360 \pm 40$  MeV

| <b><math>h_1(1170)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---|--------------------------------|-------------|
| $\rho\pi$                                 | seen                           | 307         |

**$b_1(1235)$**

$$I^G(J^{PC}) = 1^+(1^{+-})$$

Mass  $m = 1229.5 \pm 3.2$  MeV ( $S = 1.6$ )

Full width  $\Gamma = 142 \pm 9$  MeV ( $S = 1.2$ )

| <b><math>b_1(1235)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | Confidence level | $p$ (MeV/c) |
|---|--------------------------------|------------------|-------------|
| $\omega\pi$                               | dominant                       |                  | 348         |
| $\pi^\pm\gamma$                           | $(1.6 \pm 0.4) \times 10^{-3}$ |                  | 607         |
| $\eta\rho$                                | seen                           |                  | †           |
| $\pi^+\pi^+\pi^-\pi^0$                    | < 50 %                         | 84%              | 535         |
| $(K\bar{K})^\pm\pi^0$                     | < 8 %                          | 90%              | 248         |
| $K_S^0 K_L^0 \pi^\pm$                     | < 6 %                          | 90%              | 235         |
| $K_S^0 K_S^0 \pi^\pm$                     | < 2 %                          | 90%              | 235         |
| $\phi\pi$                                 | < 1.5 %                        | 84%              | 147         |

**$a_1(1260)$  <sup>[k]</sup>**

$$I^G(J^{PC}) = 1^-(1^{++})$$

Mass  $m = 1230 \pm 40$  MeV <sup>[l]</sup>

Full width  $\Gamma = 250$  to  $600$  MeV

| <b><math>a_1(1260)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---|--------------------------------|-------------|
| $(\rho\pi)_{S\text{-wave}}$               | seen                           | 353         |
| $(\rho\pi)_{D\text{-wave}}$               | seen                           | 353         |
| $(\rho(1450)\pi)_{S\text{-wave}}$         | seen                           | †           |
| $(\rho(1450)\pi)_{D\text{-wave}}$         | seen                           | †           |
| $\sigma\pi$                               | seen                           | –           |
| $f_0(980)\pi$                             | not seen                       | 189         |
| $f_0(1370)\pi$                            | seen                           | †           |
| $f_2(1270)\pi$                            | seen                           | †           |
| $K K^*(892) + \text{c.c.}$                | seen                           | †           |
| $\pi\gamma$                               | seen                           | 608         |

**$f_2(1270)$** 

$$I^G(J^{PC}) = 0^+(2^{++})$$

 Mass  $m = 1275.1 \pm 1.2$  MeV (S = 1.1)

 Full width  $\Gamma = 185.1^{+2.9}_{-2.4}$  MeV (S = 1.5)

| <b><math>f_2(1270)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ )       | Scale factor/<br>Confidence level | $\rho$<br>(MeV/c) |
|---|--------------------------------------|-----------------------------------|-------------------|
| $\pi\pi$                                  | (84.8 $^{+2.4}_{-1.2}$ ) %           | S=1.2                             | 623               |
| $\pi^+\pi^-2\pi^0$                        | ( 7.1 $^{+1.4}_{-2.7}$ ) %           | S=1.3                             | 562               |
| $K\bar{K}$                                | ( 4.6 $\pm 0.4$ ) %                  | S=2.8                             | 403               |
| $2\pi^+2\pi^-$                            | ( 2.8 $\pm 0.4$ ) %                  | S=1.2                             | 559               |
| $\eta\eta$                                | ( 4.0 $\pm 0.8$ ) $\times 10^{-3}$   | S=2.1                             | 326               |
| $4\pi^0$                                  | ( 3.0 $\pm 1.0$ ) $\times 10^{-3}$   |                                   | 564               |
| $\gamma\gamma$                            | ( 1.64 $\pm 0.19$ ) $\times 10^{-5}$ | S=1.9                             | 638               |
| $\eta\pi\pi$                              | < 8 $\times 10^{-3}$                 | CL=95%                            | 477               |
| $K^0K^-\pi^+ + \text{c.c.}$               | < 3.4 $\times 10^{-3}$               | CL=95%                            | 293               |
| $e^+e^-$                                  | < 6 $\times 10^{-10}$                | CL=90%                            | 638               |

 **$f_1(1285)$** 

$$I^G(J^{PC}) = 0^+(1^{++})$$

 Mass  $m = 1281.8 \pm 0.6$  MeV (S = 1.6)

 Full width  $\Gamma = 24.3 \pm 1.1$  MeV (S = 1.4)

| <b><math>f_1(1285)</math> DECAY MODES</b>                 | Fraction ( $\Gamma_i/\Gamma$ )     | Scale factor/<br>Confidence level | $\rho$<br>(MeV/c) |
|---|------------------------------------|-----------------------------------|-------------------|
| $4\pi$  | (33.1 $^{+2.1}_{-1.8}$ ) %         | S=1.3                             | 568               |
| $\pi^0\pi^0\pi^+\pi^-$                                    | (22.0 $^{+1.4}_{-1.2}$ ) %         | S=1.3                             | 566               |
| $2\pi^+2\pi^-$  | (11.0 $^{+0.7}_{-0.6}$ ) %         | S=1.3                             | 563               |
| $\rho^0\pi^+\pi^-$  | (11.0 $^{+0.7}_{-0.6}$ ) %         | S=1.3                             | 336               |
| $\rho^0\rho^0$  | seen                               |                                   | †                 |
| $4\pi^0$  | < 7 $\times 10^{-4}$               | CL=90%                            | 568               |
| $\eta\pi\pi$  | (52 $\pm 5$ ) %                    |                                   | 482               |
| $a_0(980)\pi$ [ignoring $a_0(980) \rightarrow K\bar{K}$ ] | (36 $\pm 7$ ) %                    |                                   | 238               |
| $\eta\pi\pi$ [excluding $a_0(980)\pi$ ]                   | (16 $\pm 7$ ) %                    |                                   | 482               |
| $K\bar{K}\pi$   | ( 9.0 $\pm 0.4$ ) %                | S=1.1                             | 308               |
| $K\bar{K}^*(892)$   | not seen                           |                                   | †                 |
| $\gamma\rho^0$  | ( 5.5 $\pm 1.3$ ) %                | S=2.8                             | 406               |
| $\phi\gamma$  | ( 7.4 $\pm 2.6$ ) $\times 10^{-4}$ |                                   | 236               |

**$\eta(1295)$**

$$I^G(J^{PC}) = 0^+(0^-+)$$

Mass  $m = 1294 \pm 4$  MeV (S = 1.6)

Full width  $\Gamma = 55 \pm 5$  MeV

| $\eta(1295)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|--------------------------|--------------------------------|-------------|
| $\eta\pi^+\pi^-$         | seen                           | 487         |
| $a_0(980)\pi$            | seen                           | 248         |
| $\eta\pi^0\pi^0$         | seen                           | 490         |
| $\eta(\pi\pi)$ S-wave    | seen                           | —           |

**$\pi(1300)$**

$$I^G(J^{PC}) = 1^-(0^-+)$$

Mass  $m = 1300 \pm 100$  MeV [1]

Full width  $\Gamma = 200$  to 600 MeV

| $\pi(1300)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-------------------------|--------------------------------|-------------|
| $\rho\pi$               | seen                           | 404         |
| $\pi(\pi\pi)$ S-wave    | seen                           | —           |

**$a_2(1320)$**

$$I^G(J^{PC}) = 1^-(2^{++})$$

Mass  $m = 1318.3 \pm 0.6$  MeV (S = 1.2)

Full width  $\Gamma = 107 \pm 5$  MeV [1]

| $a_2(1320)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ )       | Scale factor/<br>Confidence level | $p$<br>(MeV/c) |
|-------------------------|--------------------------------------|-----------------------------------|----------------|
| $3\pi$                  | (70.1 $\pm$ 2.7 ) %                  | S=1.2                             | 624            |
| $\eta\pi$               | (14.5 $\pm$ 1.2 ) %                  |                                   | 535            |
| $\omega\pi\pi$          | (10.6 $\pm$ 3.2 ) %                  | S=1.3                             | 366            |
| $K\bar{K}$              | ( 4.9 $\pm$ 0.8 ) %                  |                                   | 437            |
| $\eta'(958)\pi$         | ( 5.3 $\pm$ 0.9 ) $\times 10^{-3}$   |                                   | 288            |
| $\pi^\pm\gamma$         | ( 2.68 $\pm$ 0.31 ) $\times 10^{-3}$ |                                   | 652            |
| $\gamma\gamma$          | ( 9.4 $\pm$ 0.7 ) $\times 10^{-6}$   |                                   | 659            |
| $e^+e^-$                | < 5 $\times 10^{-9}$                 | CL=90%                            | 659            |

**$f_0(1370)$  [1]**

$$I^G(J^{PC}) = 0^+(0^{++})$$

Mass  $m = 1200$  to 1500 MeV

Full width  $\Gamma = 200$  to 500 MeV

| <b><math>f_0(1370)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---|--------------------------------|-------------|
| $\pi\pi$                                  | seen                           | 672         |
| $4\pi$                                    | seen                           | 617         |
| $4\pi^0$                                  | seen                           | 617         |
| $2\pi^+2\pi^-$                            | seen                           | 612         |
| $\pi^+\pi^-2\pi^0$                        | seen                           | 615         |
| $\rho\rho$                                | dominant                       | †           |
| $2(\pi\pi)_S\text{-wave}$                 | seen                           | —           |
| $\pi(1300)\pi$                            | seen                           | †           |
| $a_1(1260)\pi$                            | seen                           | 35          |
| $\eta\eta$                                | seen                           | 411         |
| $K\bar{K}$                                | seen                           | 475         |
| $K\bar{K}n\pi$                            | not seen                       | †           |
| $6\pi$                                    | not seen                       | 508         |
| $\omega\omega$                            | not seen                       | †           |
| $\gamma\gamma$                            | seen                           | 685         |
| $e^+e^-$                                  | not seen                       | 685         |

**$\pi_1(1400)$  [ $m$ ]**

$$I^G(J^{PC}) = 1^-(1^-+)$$

Mass  $m = 1354 \pm 25$  MeV ( $S = 1.8$ )

Full width  $\Gamma = 330 \pm 35$  MeV

| <b><math>\pi_1(1400)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---|--------------------------------|-------------|
| $\eta\pi^0$                                 | seen                           | 557         |
| $\eta\pi^-$                                 | seen                           | 556         |

**$\eta(1405)$  [ $n$ ]**

$$I^G(J^{PC}) = 0^+(0^-+)$$

Mass  $m = 1409.8 \pm 2.5$  MeV [ $l$ ] ( $S = 2.2$ )

Full width  $\Gamma = 51.1 \pm 3.4$  MeV [ $l$ ] ( $S = 2.0$ )

| <b><math>\eta(1405)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | Confidence level | $p$ (MeV/c) |
|--|--------------------------------|------------------|-------------|
| $K\bar{K}\pi$                              | seen                           |                  | 425         |
| $\eta\pi\pi$                               | seen                           |                  | 563         |
| $a_0(980)\pi$                              | seen                           |                  | 345         |
| $\eta(\pi\pi)_S\text{-wave}$               | seen                           |                  | —           |

|                |       |        |     |
|----------------|-------|--------|-----|
| $f_0(980)\eta$ | seen  |        | †   |
| $4\pi$         | seen  |        | 639 |
| $\rho\rho$     | <58 % | 99.85% | †   |
| $\rho^0\gamma$ | seen  |        | 492 |
| $K^*(892)K$    | seen  |        | 125 |

**$f_1(1420)$**  [0]

$$I^G(J^{PC}) = 0^+(1^{++})$$

Mass  $m = 1426.4 \pm 0.9$  MeV (S = 1.1)

Full width  $\Gamma = 54.9 \pm 2.6$  MeV

| <b><math>f_1(1420)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---|--------------------------------|-------------|
| $K\bar{K}\pi$                             | dominant                       | 438         |
| $K\bar{K}^*(892) + \text{c.c.}$           | dominant                       | 163         |
| $\eta\pi\pi$                              | possibly seen                  | 573         |
| $\phi\gamma$                              | seen                           | 349         |

**$\omega(1420)$**  [ρ]

$$I^G(J^{PC}) = 0^-(1^{--})$$

Mass  $m$  (1400–1450) MeV

Full width  $\Gamma$  (180–250) MeV

| <b><math>\omega(1420)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|--|--------------------------------|-------------|
| $\rho\pi$                                    | dominant                       | 486         |
| $\omega\pi\pi$                               | seen                           | 444         |
| $b_1(1235)\pi$                               | seen                           | 125         |
| $e^+e^-$                                     | seen                           | 710         |

**$a_0(1450)$**  [j]

$$I^G(J^{PC}) = 1^-(0^{++})$$

Mass  $m = 1474 \pm 19$  MeV

Full width  $\Gamma = 265 \pm 13$  MeV

| <b><math>a_0(1450)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---|--------------------------------|-------------|
| $\pi\eta$                                 | seen                           | 627         |
| $\pi\eta'(958)$                           | seen                           | 410         |
| $K\bar{K}$                                | seen                           | 547         |
| $\omega\pi\pi$                            | seen                           | 484         |
| $a_0(980)\pi\pi$                          | seen                           | 342         |
| $\gamma\gamma$                            | seen                           | 737         |

**$\rho(1450)$**  [*q*]

$$J^{PC} = 1^{+}(1^{-}-)$$

Mass  $m = 1465 \pm 25$  MeV [*l*]

Full width  $\Gamma = 400 \pm 60$  MeV [*l*]

| <b><math>\rho(1450)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|--|--------------------------------|-------------|
| $\pi\pi$                                   | seen                           | 720         |
| $4\pi$                                     | seen                           | 669         |
| $e^+e^-$                                   | seen                           | 732         |
| $\eta\rho$                                 | possibly seen                  | 310         |
| $a_2(1320)\pi$                             | not seen                       | 55          |
| $K\bar{K}$                                 | not seen                       | 541         |
| $K\bar{K}^*(892) + \text{c.c.}$            | possibly seen                  | 229         |
| $\eta\gamma$                               | possibly seen                  | 630         |

**$\eta(1475)$**  [*n*]

$$J^{PC} = 0^{+}(0^{-}+)$$

Mass  $m = 1476 \pm 4$  MeV ( $S = 1.3$ )

Full width  $\Gamma = 85 \pm 9$  MeV ( $S = 1.5$ )

| <b><math>\eta(1475)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|--|--------------------------------|-------------|
| $K\bar{K}\pi$                              | dominant                       | 477         |
| $K\bar{K}^*(892) + \text{c.c.}$            | seen                           | 245         |
| $a_0(980)\pi$                              | seen                           | 396         |
| $\gamma\gamma$                             | seen                           | 738         |

**$f_0(1500)$**  [*m*]

$$J^{PC} = 0^{+}(0^{++})$$

Mass  $m = 1505 \pm 6$  MeV ( $S = 1.3$ )

Full width  $\Gamma = 109 \pm 7$  MeV

| <b><math>f_0(1500)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | Scale factor | $p$ (MeV/c) |
|---|--------------------------------|--------------|-------------|
| $\pi\pi$                                  | $(34.9 \pm 2.3) \%$            | 1.2          | 741         |
| $\pi^+\pi^-$                              | seen                           |              | 740         |
| $2\pi^0$                                  | seen                           |              | 741         |
| $4\pi$                                    | $(49.5 \pm 3.3) \%$            | 1.2          | 691         |
| $4\pi^0$                                  | seen                           |              | 691         |
| $2\pi^+2\pi^-$                            | seen                           |              | 687         |
| $2(\pi\pi)_{S\text{-wave}}$               | seen                           |              | —           |

|                  |               |     |     |
|------------------|---------------|-----|-----|
| $\rho\rho$       | seen          |     | †   |
| $\pi(1300)\pi$   | seen          |     | 144 |
| $a_1(1260)\pi$   | seen          |     | 218 |
| $\eta\eta$       | ( 5.1±0.9 ) % | 1.4 | 516 |
| $\eta\eta'(958)$ | ( 1.9±0.8 ) % | 1.7 | †   |
| $K\bar{K}$       | ( 8.6±1.0 ) % | 1.1 | 568 |
| $\gamma\gamma$   | not seen      |     | 753 |

### $f'_2(1525)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

Mass  $m = 1525 \pm 5$  MeV [l]

Full width  $\Gamma = 73^{+6}_{-5}$  MeV [l]

| $f'_2(1525)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ )   | $p$ (MeV/c) |
|--------------------------|----------------------------------|-------------|
| $K\bar{K}$               | (88.7 ± 2.2 ) %                  | 581         |
| $\eta\eta$               | (10.4 ± 2.2 ) %                  | 530         |
| $\pi\pi$                 | ( 8.2 ± 1.5 ) × 10 <sup>-3</sup> | 750         |
| $\gamma\gamma$           | ( 1.11±0.14 ) × 10 <sup>-6</sup> | 763         |

### $\pi_1(1600)$ [m]

$$I^G(J^{PC}) = 1^-(1^{-+})$$

Mass  $m = 1662^{+15}_{-11}$  MeV (S = 1.2)

Full width  $\Gamma = 234 \pm 50$  MeV (S = 1.7)

| $\pi_1(1600)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---------------------------|--------------------------------|-------------|
| $\pi\pi\pi$               | not seen                       | 803         |
| $\rho^0\pi^-$             | not seen                       | 641         |
| $f_2(1270)\pi^-$          | not seen                       | 319         |
| $b_1(1235)\pi$            | seen                           | 357         |
| $\eta'(958)\pi^-$         | seen                           | 543         |
| $f_1(1285)\pi$            | seen                           | 315         |

### $\eta_2(1645)$

$$I^G(J^{PC}) = 0^+(2^{-+})$$

Mass  $m = 1617 \pm 5$  MeV

Full width  $\Gamma = 181 \pm 11$  MeV

| $\eta_2(1645)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|----------------------------|--------------------------------|-------------|
| $a_2(1320)\pi$             | seen                           | 242         |
| $K\bar{K}\pi$              | seen                           | 580         |
| $K^*\bar{K}$               | seen                           | 404         |
| $\eta\pi^+\pi^-$           | seen                           | 685         |
| $a_0(980)\pi$              | seen                           | 499         |
| $f_2(1270)\eta$            | not seen                       | †           |

### $\omega(1650)$ [r]

$$I^G(J^{PC}) = 0^-(1^{--})$$

Mass  $m = 1670 \pm 30$  MeV

Full width  $\Gamma = 315 \pm 35$  MeV

| $\omega(1650)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|----------------------------|--------------------------------|-------------|
| $\rho\pi$                  | seen                           | 646         |
| $\omega\pi\pi$             | seen                           | 617         |
| $\omega\eta$               | seen                           | 500         |
| $e^+e^-$                   | seen                           | 835         |

### $\omega_3(1670)$

$$I^G(J^{PC}) = 0^-(3^{--})$$

Mass  $m = 1667 \pm 4$  MeV

Full width  $\Gamma = 168 \pm 10$  MeV [l]

| $\omega_3(1670)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|------------------------------|--------------------------------|-------------|
| $\rho\pi$                    | seen                           | 645         |
| $\omega\pi\pi$               | seen                           | 615         |
| $b_1(1235)\pi$               | possibly seen                  | 361         |

### $\pi_2(1670)$

$$I^G(J^{PC}) = 1^-(2^{-+})$$

Mass  $m = 1672.4 \pm 3.2$  MeV [l] (S = 1.4)

Full width  $\Gamma = 259 \pm 9$  MeV [l] (S = 1.3)

| $\pi_2(1670)$ DECAY MODES     | Fraction ( $\Gamma_i/\Gamma$ ) | Confidence level | $p$ (MeV/c) |
|-------------------------------|--------------------------------|------------------|-------------|
| $3\pi$                        | (95.8±1.4) %                   |                  | 809         |
| $f_2(1270)\pi$                | (56.3±3.2) %                   |                  | 329         |
| $\rho\pi$                     | (31 ±4 ) %                     |                  | 648         |
| $\sigma\pi$                   | (10.9±3.4) %                   |                  | —           |
| $(\pi\pi)_{S\text{-wave}}$    | ( 8.7±3.4) %                   |                  | —           |
| $K\bar{K}^*(892)+\text{c.c.}$ | ( 4.2±1.4) %                   |                  | 455         |
| $\omega\rho$                  | ( 2.7±1.1) %                   |                  | 304         |
| $\gamma\gamma$                | < 2.8 × 10 <sup>-7</sup>       | 90%              | 836         |
| $\rho(1450)\pi$               | < 3.6 × 10 <sup>-3</sup>       | 97.7%            | 148         |
| $b_1(1235)\pi$                | < 1.9 × 10 <sup>-3</sup>       | 97.7%            | 366         |
| $f_1(1285)\pi$                | possibly seen                  |                  | 323         |
| $a_2(1320)\pi$                | not seen                       |                  | 292         |

### $\phi(1680)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

Mass  $m = 1680 \pm 20$  MeV [1]

Full width  $\Gamma = 150 \pm 50$  MeV [1]

| $\phi(1680)$ DECAY MODES      | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-------------------------------|--------------------------------|-------------|
| $K\bar{K}^*(892)+\text{c.c.}$ | dominant                       | 462         |
| $K_S^0 K\pi$                  | seen                           | 621         |
| $K\bar{K}$                    | seen                           | 680         |
| $e^+e^-$                      | seen                           | 840         |
| $\omega\pi\pi$                | not seen                       | 623         |
| $K^+K^-\pi^+\pi^-$            | seen                           | 544         |

### $\rho_3(1690)$

$$I^G(J^{PC}) = 1^+(3^{--})$$

Mass  $m = 1688.8 \pm 2.1$  MeV [1]

Full width  $\Gamma = 161 \pm 10$  MeV [1] (S = 1.5)

| $\rho_3(1690)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | Scale factor | $p$ (MeV/c) |
|----------------------------|--------------------------------|--------------|-------------|
| $4\pi$                     | (71.1 ± 1.9 ) %                |              | 790         |
| $\pi^\pm\pi^+\pi^-\pi^0$   | (67 ±22 ) %                    |              | 787         |
| $\omega\pi$                | (16 ± 6 ) %                    |              | 655         |
| $\pi\pi$                   | (23.6 ± 1.3 ) %                |              | 834         |
| $K\bar{K}\pi$              | ( 3.8 ± 1.2 ) %                |              | 629         |

|  |                  |     |     |
|--|------------------|-----|-----|
| $K\bar{K}$                             | ( 1.58 ± 0.26) % | 1.2 | 685 |
| $\eta\pi^+\pi^-$                       | seen             |     | 727 |
| $\rho(770)\eta$                        | seen             |     | 520 |
| $\pi\pi\rho$                           | seen             |     | 633 |
| Excluding $2\rho$ and $a_2(1320)\pi$ . |                  |     |     |
| $a_2(1320)\pi$                         | seen             |     | 307 |
| $\rho\rho$                             | seen             |     | 334 |

**$\rho(1700)$  [q]**

$$I^G(J^{PC}) = 1^+(1^- -)$$

Mass  $m = 1720 \pm 20$  MeV [l] ( $\eta\rho^0$  and  $\pi^+\pi^-$  modes)

Full width  $\Gamma = 250 \pm 100$  MeV [l] ( $\eta\rho^0$  and  $\pi^+\pi^-$  modes)

| <b><math>\rho(1700)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|--|--------------------------------|-------------|
| $2(\pi^+\pi^-)$                            | large                          | 803         |
| $\rho\pi\pi$                               | dominant                       | 653         |
| $\rho^0\pi^+\pi^-$                         | large                          | 650         |
| $\rho^\pm\pi^\mp\pi^0$                     | large                          | 652         |
| $a_1(1260)\pi$                             | seen                           | 404         |
| $h_1(1170)\pi$                             | seen                           | 447         |
| $\pi(1300)\pi$                             | seen                           | 349         |
| $\rho\rho$                                 | seen                           | 372         |
| $\pi^+\pi^-$                               | seen                           | 849         |
| $\pi\pi$                                   | seen                           | 849         |
| $K\bar{K}^*(892) + \text{c.c.}$            | seen                           | 496         |
| $\eta\rho$                                 | seen                           | 545         |
| $a_2(1320)\pi$                             | not seen                       | 334         |
| $K\bar{K}$                                 | seen                           | 704         |
| $e^+e^-$                                   | seen                           | 860         |
| $\pi^0\omega$                              | seen                           | 674         |

**$f_0(1710)$  [s]**

$$I^G(J^{PC}) = 0^+(0^{++})$$

Mass  $m = 1720 \pm 6$  MeV (S = 1.6)

Full width  $\Gamma = 135 \pm 8$  MeV (S = 1.1)

| <b><math>f_0(1710)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---|--------------------------------|-------------|
| $K \bar{K}$                               | seen                           | 704         |
| $\eta\eta$                                | seen                           | 663         |
| $\pi\pi$                                  | seen                           | 849         |
| $\omega\omega$                            | seen                           | 357         |

**$\pi(1800)$**

$$I^G(J^{PC}) = 1^-(0^-+)$$

Mass  $m = 1816 \pm 14$  MeV ( $S = 2.3$ )

Full width  $\Gamma = 208 \pm 12$  MeV

| <b><math>\pi(1800)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---|--------------------------------|-------------|
| $\pi^+\pi^-\pi^-$                         | seen                           | 881         |
| $f_0(600)\pi^-$                           | seen                           | —           |
| $f_0(980)\pi^-$                           | seen                           | 634         |
| $f_0(1370)\pi^-$                          | seen                           | 371         |
| $f_0(1500)\pi^-$                          | not seen                       | 254         |
| $\rho\pi^-$                               | not seen                       | 735         |
| $\eta\eta\pi^-$                           | seen                           | 664         |
| $a_0(980)\eta$                            | seen                           | 477         |
| $a_2(1320)\eta$                           | not seen                       | †           |
| $f_2(1270)\pi$                            | not seen                       | 445         |
| $f_0(1370)\pi^-$                          | not seen                       | 371         |
| $f_0(1500)\pi^-$                          | seen                           | 254         |
| $\eta\eta'(958)\pi^-$                     | seen                           | 380         |
| $K_0^*(1430)K^-$                          | seen                           | †           |
| $K^*(892)K^-$                             | not seen                       | 573         |

**$\phi_3(1850)$**

$$I^G(J^{PC}) = 0^-(3^{--})$$

Mass  $m = 1854 \pm 7$  MeV

Full width  $\Gamma = 87^{+28}_{-23}$  MeV ( $S = 1.2$ )

| <b><math>\phi_3(1850)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|--|--------------------------------|-------------|
| $K \bar{K}$                                  | seen                           | 785         |
| $K \bar{K}^*(892) + \text{c.c.}$             | seen                           | 602         |

**$\pi_2(1880)$**

$$I^G(J^{PC}) = 1^-(2^-+)$$

Mass  $m = 1895 \pm 16$  MeV

Full width  $\Gamma = 235 \pm 34$  MeV

**$f_2(1950)$**

$$I^G(J^{PC}) = 0^+(2^{++})$$

Mass  $m = 1944 \pm 12$  MeV (S = 1.5)

Full width  $\Gamma = 472 \pm 18$  MeV

| <b><math>f_2(1950)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---|--------------------------------|-------------|
| $K^*(892)\bar{K}^*(892)$                  | seen                           | 387         |
| $\pi^+\pi^-$                              | seen                           | 962         |
| $\pi^0\pi^0$                              | seen                           | 963         |
| $4\pi$                                    | seen                           | 925         |
| $\eta\eta$                                | seen                           | 803         |
| $K\bar{K}$                                | seen                           | 837         |
| $\gamma\gamma$                            | seen                           | 972         |

**$f_2(2010)$**

$$I^G(J^{PC}) = 0^+(2^{++})$$

Mass  $m = 2011^{+60}_{-80}$  MeV

Full width  $\Gamma = 202 \pm 60$  MeV

| <b><math>f_2(2010)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---|--------------------------------|-------------|
| $\phi\phi$                                | seen                           | †           |
| $K\bar{K}$                                | seen                           | 876         |

**$a_4(2040)$**

$$I^G(J^{PC}) = 1^-(4^{++})$$

Mass  $m = 2001 \pm 10$  MeV

Full width  $\Gamma = 235 \pm 29$  MeV (S = 1.3)

| <b><math>a_4(2040)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---|--------------------------------|-------------|
| $K\bar{K}$                                | seen                           | 870         |
| $\pi^+\pi^-\pi^0$                         | seen                           | 976         |
| $\rho\pi$                                 | seen                           | 844         |
| $f_2(1270)\pi$                            | seen                           | 583         |

|                      |      |     |
|----------------------|------|-----|
| $\omega \pi^- \pi^0$ | seen | 821 |
| $\omega \rho$        | seen | 627 |
| $\eta \pi^0$         | seen | 920 |
| $\eta'(958) \pi$     | seen | 764 |

**$f_4(2050)$**

$$I^G(J^{PC}) = 0^+(4^{++})$$

Mass  $m = 2018 \pm 11$  MeV (S = 2.1)

Full width  $\Gamma = 237 \pm 18$  MeV (S = 1.9)

| <b><math>f_4(2050)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ )       | $p$ (MeV/c) |
|---|--------------------------------------|-------------|
| $\omega \omega$                           | seen                                 | 637         |
| $\pi \pi$                                 | $(17.0 \pm 1.5) \%$                  | 1000        |
| $K \bar{K}$                               | $(6.8^{+3.4}_{-1.8}) \times 10^{-3}$ | 880         |
| $\eta \eta$                               | $(2.1 \pm 0.8) \times 10^{-3}$       | 848         |
| $4\pi^0$                                  | $< 1.2 \%$                           | 964         |
| $a_2(1320) \pi$                           | seen                                 | 567         |

**$\phi(2170)$**

$$I^G(J^{PC}) = 0^-(1^{--})$$

Mass  $m = 2175 \pm 15$  MeV (S = 1.6)

Full width  $\Gamma = 61 \pm 18$  MeV

| <b><math>\phi(2170)</math> DECAY MODES</b>              | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---|--------------------------------|-------------|
| $e^+ e^-$   | seen                           | 1087        |
| $\phi f_0(980)$   | seen                           | 427         |
| $K^+ K^- f_0(980) \rightarrow$<br>$K^+ K^- \pi^+ \pi^-$ | seen                           | —           |
| $K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^0 \pi^0$      | seen                           | —           |
| $K^{*0} K^\pm \pi^\mp$                                  | not seen                       | 770         |

**$f_2(2300)$**

$$I^G(J^{PC}) = 0^+(2^{++})$$

Mass  $m = 2297 \pm 28$  MeV

Full width  $\Gamma = 149 \pm 40$  MeV

| $f_2(2300)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-------------------------|--------------------------------|-------------|
| $\phi\phi$              | seen                           | 529         |
| $K\bar{K}$              | seen                           | 1037        |
| $\gamma\gamma$          | seen                           | 1149        |

**$f_2(2340)$**

$$I^G(J^{PC}) = 0^+(2^{++})$$

Mass  $m = 2339 \pm 60$  MeV

Full width  $\Gamma = 319^{+80}_{-70}$  MeV

| $f_2(2340)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-------------------------|--------------------------------|-------------|
| $\phi\phi$              | seen                           | 573         |
| $\eta\eta$              | seen                           | 1033        |

### NOTES

- [a] See the “Note on  $\pi^\pm \rightarrow \ell^\pm \nu \gamma$  and  $K^\pm \rightarrow \ell^\pm \nu \gamma$  Form Factors” in the  $\pi^\pm$  Particle Listings for definitions and details.
- [b] Measurements of  $\Gamma(e^+ \nu_e)/\Gamma(\mu^+ \nu_\mu)$  always include decays with  $\gamma$ 's, and measurements of  $\Gamma(e^+ \nu_e \gamma)$  and  $\Gamma(\mu^+ \nu_\mu \gamma)$  never include low-energy  $\gamma$ 's. Therefore, since no clean separation is possible, we consider the modes with  $\gamma$ 's to be subreactions of the modes without them, and let  $[\Gamma(e^+ \nu_e) + \Gamma(\mu^+ \nu_\mu)]/\Gamma_{\text{total}} = 100\%$ .
- [c] See the  $\pi^\pm$  Particle Listings for the energy limits used in this measurement; low-energy  $\gamma$ 's are not included.
- [d] Derived from an analysis of neutrino-oscillation experiments.
- [e] Astrophysical and cosmological arguments give limits of order  $10^{-13}$ ; see the  $\pi^0$  Particle Listings.
- [f] C parity forbids this to occur as a single-photon process.
- [g] See the “Note on scalar mesons” in the  $f_0(1370)$  Particle Listings . The interpretation of this entry as a particle is controversial.
- [h] See the “Note on  $\rho(770)$ ” in the  $\rho(770)$  Particle Listings .
- [i] The  $\omega\rho$  interference is then due to  $\omega\rho$  mixing only, and is expected to be small. If  $e\mu$  universality holds,  $\Gamma(\rho^0 \rightarrow \mu^+ \mu^-) = \Gamma(\rho^0 \rightarrow e^+ e^-) \times 0.99785$ .
- [j] See the “Note on scalar mesons” in the  $f_0(1370)$  Particle Listings .
- [k] See the “Note on  $a_1(1260)$ ” in the  $a_1(1260)$  Particle Listings in PDG 06, Journal of Physics, G **33** 1 (2006).

- [l] This is only an educated guess; the error given is larger than the error on the average of the published values. See the Particle Listings for details.
- [m] See the “Note on non- $q\bar{q}$  mesons” in the Particle Listings in PDG 06, Journal of Physics, G **33** 1 (2006).
- [n] See the “Note on the  $\eta(1405)$ ” in the  $\eta(1405)$  Particle Listings.
- [o] See the “Note on the  $f_1(1420)$ ” in the  $\eta(1405)$  Particle Listings.
- [p] See also the  $\omega(1650)$  Particle Listings.
- [q] See the “Note on the  $\rho(1450)$  and the  $\rho(1700)$ ” in the  $\rho(1700)$  Particle Listings.
- [r] See also the  $\omega(1420)$  Particle Listings.
- [s] See the “Note on  $f_0(1710)$ ” in the  $f_0(1710)$  Particle Listings in 2004 edition of *Review of Particle Physics*.