

$\chi_{c2}(1P)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

See the Review on “ $\psi(2S)$ and χ_c branching ratios” before the $\chi_{c0}(1P)$ Listings.

$\chi_{c2}(1P)$ MASS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3556.20 ± 0.09	OUR AVERAGE			
3555.3 ± 0.6 ± 2.2	2.5k	UEHARA	08 BELL	$\gamma\gamma \rightarrow$ hadrons
3555.70 ± 0.59 ± 0.39		ABLIKIM	05G BES2	$\psi(2S) \rightarrow \gamma\chi_{c2}$
3556.173 ± 0.123 ± 0.020		ANDREOTTI	05A E835	$p\bar{p} \rightarrow e^+e^-\gamma$
3559.9 ± 2.9		EISENSTEIN	01 CLE2	$e^+e^- \rightarrow e^+e^-\chi_{c2}$
3556.4 ± 0.7		BAI	99B BES	$\psi(2S) \rightarrow \gamma X$
3556.22 ± 0.131 ± 0.020	585	¹ ARMSTRONG	92 E760	$\bar{p}p \rightarrow e^+e^-\gamma$
3556.9 ± 0.4 ± 0.5	50	BAGLIN	86B SPEC	$\bar{p}p \rightarrow e^+e^- X$
3557.8 ± 0.2 ± 4		² GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$
3553.4 ± 2.2	66	³ LEMOIGNE	82 GOLI	$185 \pi^- \text{Be} \rightarrow \gamma\mu^+\mu^- A$
3555.9 ± 0.7		⁴ OREGLIA	82 CBAL	$e^+e^- \rightarrow J/\psi 2\gamma$
3557 ± 1.5	69	⁵ HIMEL	80 MRK2	$e^+e^- \rightarrow J/\psi 2\gamma$
3551 ± 11	15	BRANDELIK	79B DASP	$e^+e^- \rightarrow J/\psi 2\gamma$
3553 ± 4		⁵ BARTEL	78B CNTR	$e^+e^- \rightarrow J/\psi 2\gamma$
3553 ± 4 ± 4		^{5,6} TANENBAUM	78 MRK1	e^+e^-
3563 ± 7	360	⁵ BIDDICK	77 CNTR	$e^+e^- \rightarrow \gamma X$
• • •				We do not use the following data for averages, fits, limits, etc. • • •
3543 ± 10	4	WHITAKER	76 MRK1	$e^+e^- \rightarrow J/\psi 2\gamma$

¹ Recalculated by ANDREOTTI 05A, using the value of $\psi(2S)$ mass from AULCHENKO 03.

² Using mass of $\psi(2S) = 3686.0$ MeV.

³ $J/\psi(1S)$ mass constrained to 3097 MeV.

⁴ Assuming $\psi(2S)$ mass = 3686 MeV and $J/\psi(1S)$ mass = 3097 MeV.

⁵ Mass value shifted by us by amount appropriate for $\psi(2S)$ mass = 3686 MeV and $J/\psi(1S)$ mass = 3097 MeV.

⁶ From a simultaneous fit to radiative and hadronic decay channels.

$\chi_{c2}(1P)$ WIDTH

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.98 ± 0.11	OUR FIT			
1.95 ± 0.13	OUR AVERAGE			
1.915 ± 0.188 ± 0.013		ANDREOTTI	05A E835	$p\bar{p} \rightarrow e^+e^-\gamma$
1.96 ± 0.17 ± 0.07	585	⁷ ARMSTRONG	92 E760	$\bar{p}p \rightarrow e^+e^-\gamma$
2.6 ^{+1.4} _{-1.0}	50	BAGLIN	86B SPEC	$\bar{p}p \rightarrow e^+e^- X$
2.8 ^{+2.1} _{-2.0}		⁸ GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$

⁷ Recalculated by ANDREOTTI 05A.

⁸ Errors correspond to 90% confidence level; authors give only width range.

$\chi_{c2}(1P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Hadronic decays		
Γ_1 $2(\pi^+\pi^-)$	(1.10±0.11) %	
Γ_2 $\rho\rho$		
Γ_3 $\pi^+\pi^-\pi^0\pi^0$	(2.00±0.26) %	
Γ_4 $\rho^+\pi^-\pi^0 + \text{c.c.}$	(2.4 ±0.4) %	
Γ_5 $4\pi^0$	(1.21±0.17) × 10 ⁻³	
Γ_6 $K^+K^-\pi^0\pi^0$	(2.2 ±0.5) × 10 ⁻³	
Γ_7 $K^+\pi^-K^0\pi^0 + \text{c.c.}$	(1.51±0.22) %	
Γ_8 $\rho^+K^-K^0 + \text{c.c.}$	(4.5 ±1.4) × 10 ⁻³	
Γ_9 $K^*(892)^0K^+\pi^- \rightarrow$ $K^+\pi^-K^0\pi^0 + \text{c.c.}$	(3.2 ±0.9) × 10 ⁻³	
Γ_{10} $K^*(892)^0K^0\pi^0 \rightarrow$ $K^+\pi^-K^0\pi^0 + \text{c.c.}$	(4.2 ±0.9) × 10 ⁻³	
Γ_{11} $K^*(892)^-K^+\pi^0 \rightarrow$ $K^+\pi^-K^0\pi^0 + \text{c.c.}$	(4.1 ±0.9) × 10 ⁻³	
Γ_{12} $K^*(892)^+K^0\pi^- \rightarrow$ $K^+\pi^-K^0\pi^0 + \text{c.c.}$	(3.2 ±0.9) × 10 ⁻³	
Γ_{13} $K^+K^-\eta\pi^0$	(1.4 ±0.5) × 10 ⁻³	
Γ_{14} $K^+K^-\pi^+\pi^-$	(9.1 ±1.1) × 10 ⁻³	
Γ_{15} $K^+K^-\pi^+\pi^-\pi^0$	(1.3 ±0.4) %	
Γ_{16} $K^+\bar{K}^*(892)^0\pi^- + \text{c.c.}$	(2.3 ±1.2) × 10 ⁻³	
Γ_{17} $K^*(892)^0\bar{K}^*(892)^0$	(2.5 ±0.5) × 10 ⁻³	
Γ_{18} $3(\pi^+\pi^-)$	(8.6 ±1.8) × 10 ⁻³	
Γ_{19} $\phi\phi$	(1.14±0.12) × 10 ⁻³	
Γ_{20} $\omega\omega$	(9.2 ±1.1) × 10 ⁻⁴	
Γ_{21} $\omega\phi$		
Γ_{22} $\pi\pi$	(2.43±0.13) × 10 ⁻³	
Γ_{23} $\rho^0\pi^+\pi^-$	(4.0 ±1.7) × 10 ⁻³	
Γ_{24} $\pi^+\pi^-\eta$	(5.2 ±1.4) × 10 ⁻⁴	
Γ_{25} $\pi^+\pi^-\eta'$	(5.5 ±2.0) × 10 ⁻⁴	
Γ_{26} $\eta\eta$	(5.9 ±0.5) × 10 ⁻⁴	
Γ_{27} K^+K^-	(1.09±0.08) × 10 ⁻³	
Γ_{28} $K_S^0K_S^0$	(5.8 ±0.5) × 10 ⁻⁴	
Γ_{29} $\bar{K}^0K^+\pi^- + \text{c.c.}$	(1.40±0.20) × 10 ⁻³	
Γ_{30} $K^+K^-\pi^0$	(3.3 ±0.8) × 10 ⁻⁴	
Γ_{31} $K^+K^-\eta$	< 3.5 × 10 ⁻⁴	90%
Γ_{32} $\eta\eta'$	< 6 × 10 ⁻⁵	90%
Γ_{33} $\eta'\eta'$	< 1.1 × 10 ⁻⁴	90%
Γ_{34} $\pi^+\pi^-K_S^0K_S^0$	(2.4 ±0.6) × 10 ⁻³	
Γ_{35} $K^+K^-K_S^0K_S^0$	< 4 × 10 ⁻⁴	90%
Γ_{36} $K^+K^-K^+K^-$	(1.78±0.22) × 10 ⁻³	

Γ_{37}	$K^+ K^- \phi$	$(1.55 \pm 0.33) \times 10^{-3}$	
Γ_{38}	$\rho \bar{\rho}$	$(7.2 \pm 0.4) \times 10^{-5}$	
Γ_{39}	$\rho \bar{\rho} \pi^0$	$(5.1 \pm 0.5) \times 10^{-4}$	
Γ_{40}	$\rho \bar{\rho} \eta$	$(1.90 \pm 0.28) \times 10^{-4}$	
Γ_{41}	$\rho \bar{\rho} \omega$	$(3.9 \pm 0.5) \times 10^{-4}$	
Γ_{42}	$\rho \bar{\rho} \phi$	$(3.0 \pm 1.0) \times 10^{-5}$	
Γ_{43}	$\rho \bar{\rho} \pi^+ \pi^-$	$(1.32 \pm 0.34) \times 10^{-3}$	
Γ_{44}	$\rho \bar{\rho} \pi^0 \pi^0$	$(8.6 \pm 2.6) \times 10^{-4}$	
Γ_{45}	$\rho \bar{\rho} K^+ K^-$ (non-resonant)	$(2.1 \pm 0.4) \times 10^{-4}$	
Γ_{46}	$\rho \bar{\rho} K_S^0 K_S^0$	$< 7.9 \times 10^{-4}$	90%
Γ_{47}	$\rho \bar{\eta} \pi^-$	$(1.1 \pm 0.4) \times 10^{-3}$	
Γ_{48}	$\Lambda \bar{\Lambda}$	$(1.86 \pm 0.27) \times 10^{-4}$	
Γ_{49}	$\Lambda \bar{\Lambda} \pi^+ \pi^-$	$< 3.5 \times 10^{-3}$	90%
Γ_{50}	$K^+ \bar{p} \Lambda + \text{c.c.}$	$(9.1 \pm 1.8) \times 10^{-4}$	
Γ_{51}	$K^+ p \Lambda(1520) + \text{c.c.}$	$(3.1 \pm 0.7) \times 10^{-4}$	
Γ_{52}	$\Lambda(1520) \bar{\Lambda}(1520)$	$(5.1 \pm 1.6) \times 10^{-4}$	
Γ_{53}	$\Sigma^0 \bar{\Sigma}^0$	$< 8 \times 10^{-5}$	90%
Γ_{54}	$\Sigma^+ \bar{\Sigma}^-$	$< 7 \times 10^{-5}$	90%
Γ_{55}	$\Xi^0 \bar{\Xi}^0$	$< 1.1 \times 10^{-4}$	90%
Γ_{56}	$\Xi^- \bar{\Xi}^+$	$(1.55 \pm 0.35) \times 10^{-4}$	
Γ_{57}	$J/\psi(1S) \pi^+ \pi^- \pi^0$	$< 1.5 \%$	90%

Radiative decays

Γ_{58}	$\gamma J/\psi(1S)$	$(19.5 \pm 0.8) \%$	
Γ_{59}	$\gamma \rho^0$	$< 2.1 \times 10^{-5}$	90%
Γ_{60}	$\gamma \omega$	$< 6 \times 10^{-6}$	90%
Γ_{61}	$\gamma \phi$	$< 8 \times 10^{-6}$	90%
Γ_{62}	$\gamma \gamma$	$(2.59 \pm 0.16) \times 10^{-4}$	

CONSTRAINED FIT INFORMATION

A multiparticle fit to $\chi_{c1}(1P)$, $\chi_{c0}(1P)$, $\chi_{c2}(1P)$, and $\psi(2S)$ with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 84 branching ratios uses 223 measurements to determine 49 parameters. The overall fit has a $\chi^2 = 312.2$ for 174 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

x ₁₄	17									
x ₁₆	4	22								
x ₁₇	10	8	2							
x ₁₉	12	11	2	6						
x ₂₂	23	20	4	12	27					
x ₂₃	20	4	1	2	3	5				
x ₂₆	13	12	3	7	17	32	3			
x ₂₇	18	16	3	9	20	39	4	24		
x ₂₈	17	15	3	9	18	34	4	21	25	
x ₂₉	9	8	2	5	10	19	2	12	14	13
x ₃₆	12	10	2	6	11	22	3	13	16	14
x ₃₈	7	6	1	4	1	1	2	0	1	2
x ₄₈	8	7	2	4	10	20	2	12	14	13
x ₅₈	28	24	5	14	30	59	6	36	44	39
x ₆₂	-18	-15	-3	-9	1	3	-5	3	0	-2
Γ	-25	-21	-5	-13	-19	-36	-6	-21	-27	-25
	x ₁	x ₁₄	x ₁₆	x ₁₇	x ₁₉	x ₂₂	x ₂₃	x ₂₆	x ₂₇	x ₂₈

x ₃₆	8									
x ₃₈	1	2								
x ₄₈	7	8	0							
x ₅₈	21	25	-11	22						
x ₆₂	0	-3	27	2	9					
Γ	-13	-17	-50	-13	-49	-48				
	x ₂₉	x ₃₆	x ₃₈	x ₄₈	x ₅₈	x ₆₂				

χ_{c2}(1P) PARTIAL WIDTHS

———— χ_{c2}(1P) Γ(i)Γ(γJ/ψ(1S))/Γ(total) ————

Γ(p̄p̄) × Γ(γJ/ψ(1S))/Γ_{total}

Γ₃₈Γ₅₈/Γ

VALUE (eV) DOCUMENT ID TECN COMMENT

27.7±1.4 OUR FIT

27.5±1.5 OUR AVERAGE

27.0±1.5±1.1

⁹ ANDREOTTI 05A E835 p̄p̄ → e⁺e⁻γ

27.7±1.5±2.0

^{9,10} ARMSTRONG 92 E760 p̄p̄ → e⁺e⁻γ

36 ±8

⁹ BAGLIN 86B SPEC p̄p̄ → e⁺e⁻X

⁹ Calculated by us using B(J/ψ(1S) → e⁺e⁻) = 0.0593 ± 0.0010.

¹⁰ Recalculated by ANDREOTTI 05A.

$\Gamma(\gamma\gamma) \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$ $\Gamma_{62}\Gamma_{58}/\Gamma$

<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
100± 6 OUR FIT				
117± 10 OUR AVERAGE				
111± 12± 9	147 ± 15	11 DOBBS	06 CLE3	10.4 $e^+e^- \rightarrow e^+e^-\chi_{c2}$
114± 11± 9	136 ± 13.3	11,12 ABE	02T BELL	$e^+e^- \rightarrow e^+e^-\chi_{c2}$
139± 55± 21		11,13 ACCIARRI	99E L3	$e^+e^- \rightarrow e^+e^-\chi_{c2}$
242± 65± 51		11,14 ACKER...,K...	98 OPAL	$e^+e^- \rightarrow e^+e^-\chi_{c2}$
150± 42± 36		11,15 DOMINICK	94 CLE2	$e^+e^- \rightarrow e^+e^-\chi_{c2}$
470±240±120		11,16 BAUER	93 TPC	$e^+e^- \rightarrow e^+e^-\chi_{c2}$

¹¹ Calculated by us using $B(J/\psi \rightarrow \ell^+\ell^-) = 0.1187 \pm 0.0008$.

¹² All systematic errors added in quadrature.

¹³ The value for $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$ reported in ACCIARRI 99E is derived using $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) \times B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.0162 \pm 0.0014$.

¹⁴ The value for $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$ reported in ACKERSTAFF,K 98 is derived using $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$ and $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1203 \pm 0.0038$.

¹⁵ The value for $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$ reported in DOMINICK 94 is derived using $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$, $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0627 \pm 0.0020$, and $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0597 \pm 0.0025$.

¹⁶ The value for $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$ reported in BAUER 93 is derived using $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$, $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0627 \pm 0.0020$, and $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0597 \pm 0.0025$.

———— **$\chi_{c2}(1P) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$** ————

$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{22}\Gamma_{62}/\Gamma$

<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.24±0.08 OUR FIT				
1.18±0.25 OUR AVERAGE				
1.44±0.54±0.47	34 ± 13	17 UEHARA	09 BELL	10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
1.14±0.21±0.17	54 ± 10	18 NAKAZAWA	05 BELL	10.6 $e^+e^- \rightarrow e^+e^-\pi^+\pi^-$

¹⁷ We multiplied the measurement by 3 to convert from $\pi^0\pi^0$ to $\pi\pi$. Interference with the continuum included.

¹⁸ We have multiplied $\pi^+\pi^-$ measurement by 3/2 to obtain $\pi\pi$.

$\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{26}\Gamma_{62}/\Gamma$

<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.53±0.22±0.09				
	8	19 UEHARA	10A BELL	10.6 $e^+e^- \rightarrow e^+e^-\eta\eta$

¹⁹ Interference with the continuum not included.

$\Gamma(K^+K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{27}\Gamma_{62}/\Gamma$

<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.56±0.05 OUR FIT				
0.44±0.11±0.07	33 ± 8	NAKAZAWA	05 BELL	10.6 $e^+e^- \rightarrow e^+e^-K^+K^-$

$\Gamma(K_S^0K_S^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{28}\Gamma_{62}/\Gamma$

<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.297±0.026 OUR FIT				
0.31 ±0.05 ±0.03	38 ± 7	CHEN	07B BELL	$e^+e^- \rightarrow e^+e^-\chi_{c2}$

$\Gamma(2(\pi^+\pi^-)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_1\Gamma_{62}/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
5.6 ± 0.5 OUR FIT					
5.2 ± 0.7 OUR AVERAGE					
5.01 ± 0.44 ± 0.55	1597 ± 138	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(\pi^+\pi^-)$	
6.4 ± 1.8 ± 0.8		EISENSTEIN	01 CLE2	$e^+e^- \rightarrow e^+e^-\chi_{c2}$	
$\Gamma(\rho^0\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{23}\Gamma_{62}/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
2.0 ± 0.9 OUR FIT					
3.2 ± 1.9 ± 0.5	986 ± 578	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(\pi^+\pi^-)$	
$\Gamma(\rho\rho) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_2\Gamma_{62}/\Gamma$
VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<7.8	90	<598	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(\pi^+\pi^-)$
$\Gamma(K^+K^-\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{14}\Gamma_{62}/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
4.7 ± 0.5 OUR FIT					
4.42 ± 0.42 ± 0.53	780 ± 74	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c2} \rightarrow K^+K^-\pi^+\pi^-$	
$\Gamma(\bar{K}^0K^+\pi^- + \text{c.c.}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{29}\Gamma_{62}/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.72 ± 0.11 OUR FIT					
1.20 ± 0.33 ± 0.13	126	²⁰ DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$	
²⁰ We have multiplied $\bar{K}K\pi$ by 2/3 to obtain $\bar{K}^0K^+\pi^- + \text{c.c.}$					
$\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{15}\Gamma_{62}/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
6.5 ± 0.9 ± 1.5	1250	DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$	
$\Gamma(K^*(892)^0\bar{K}^*(892)^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{17}\Gamma_{62}/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
1.26 ± 0.24 OUR FIT					
0.8 ± 0.17 ± 0.27	151 ± 30	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c2} \rightarrow K^+K^-\pi^+\pi^-$	
$\Gamma(K^+K^-K^+K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{36}\Gamma_{62}/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.91 ± 0.12 OUR FIT					
1.10 ± 0.21 ± 0.15	126 ± 24	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(K^+K^-)$	
$\Gamma(\phi\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{19}\Gamma_{62}/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.58 ± 0.06 OUR FIT					
0.58 ± 0.18 ± 0.16	26.5 ± 8.1	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(K^+K^-)$	

$\chi_{c2}(1P)$ BRANCHING RATIOS

HADRONIC DECAYS

$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$	Γ_1/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>
0.0110±0.0011 OUR FIT	

$\Gamma(\rho^0\pi^+\pi^-)/\Gamma(2(\pi^+\pi^-))$	Γ_{23}/Γ_1
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.36±0.15 OUR FIT	
0.31±0.17	TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c2}$

$\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}$	Γ_3/Γ
<u>VALUE (%)</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
2.00±0.25±0.08	903.5 21 HE 08B CLEO $e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

²¹ HE 08B reports $1.87 \pm 0.07 \pm 0.22 \pm 0.13$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\rho^+\pi^-\pi^0 + \text{c.c.})/\Gamma_{\text{total}}$	Γ_4/Γ
<u>VALUE (%)</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
2.4±0.4±0.1	1031.9 22,23 HE 08B CLEO $e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

²² HE 08B reports $2.23 \pm 0.11 \pm 0.32 \pm 0.16$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \rho^+\pi^-\pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²³ Calculated by us. We have added the values from HE 08B for $\rho^+\pi^-\pi^0$ and $\rho^-\pi^+\pi^0$ decays assuming uncorrelated statistical and fully correlated systematic uncertainties.

$\Gamma(4\pi^0)/\Gamma_{\text{total}}$	Γ_5/Γ
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
1.21±0.16±0.05	1164 24 ABLIKIM 11A BES3 $e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$

²⁴ ABLIKIM 11A reports $(1.21 \pm 0.05 \pm 0.16) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow 4\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}$	Γ_6/Γ
<u>VALUE (%)</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.22±0.04±0.01	76.9 25 HE 08B CLEO $e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

²⁵ HE 08B reports $0.21 \pm 0.03 \pm 0.03 \pm 0.01$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.51±0.21±0.06	211.6	²⁶ HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

²⁶ HE 08B reports $1.41 \pm 0.11 \pm 0.16 \pm 0.10$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\rho^+ K^- K^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.45±0.13±0.02	62.9	²⁷ HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

²⁷ HE 08B reports $0.42 \pm 0.11 \pm 0.06 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \rho^+ K^- K^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^0 K^+ \pi^- \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.32±0.09±0.01	38.7	²⁸ HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

²⁸ HE 08B reports $0.30 \pm 0.07 \pm 0.04 \pm 0.02$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0 K^+ \pi^- \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^0 K^0 \pi^0 \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.42±0.09±0.02	63.0	²⁹ HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

²⁹ HE 08B reports $0.39 \pm 0.07 \pm 0.05 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0 K^0 \pi^0 \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.41±0.09±0.02	51.1	³⁰ HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

³⁰ HE 08B reports $0.38 \pm 0.07 \pm 0.04 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^+ K^0 \pi^- \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_{12}/Γ**

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
------------------	-------------	--------------------	-------------	----------------

0.32±0.09±0.01	39.3	³¹ HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$
-----------------------	------	------------------	----------	--

³¹ HE 08B reports $0.30 \pm 0.07 \pm 0.04 \pm 0.02$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^+ K^0 \pi^- \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ K^- \eta \pi^0)/\Gamma_{\text{total}}$ **Γ_{13}/Γ**

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
------------------	-------------	--------------------	-------------	----------------

0.14±0.05±0.01	22.9	³² HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$
-----------------------	------	------------------	----------	--

³² HE 08B reports $0.13 \pm 0.04 \pm 0.02 \pm 0.01$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \eta \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ **Γ_{14}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>
---	--------------------

9.1±1.1 OUR FIT

$\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma(K^+ K^- \pi^+ \pi^-)$ **Γ_{16}/Γ_{14}**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

0.25±0.13 OUR FIT

0.25±0.13	TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma \chi_{c2}$
------------------	--------------	------	---

$\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_{16}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>
---	--------------------

23±12 OUR FIT

$\Gamma(K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ **Γ_{17}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>
---	--------------------

2.5±0.5 OUR FIT

$\Gamma(3(\pi^+ \pi^-))/\Gamma_{\text{total}}$ **Γ_{18}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

8.6±1.8 OUR EVALUATION Treating systematic error as correlated.

8.6±1.8 OUR AVERAGE

8.6±0.9±1.6	³³ BAI	99B BES	$\psi(2S) \rightarrow \gamma \chi_{c2}$
-------------	-------------------	---------	---

8.7±5.9±0.4	³³ TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma \chi_{c2}$
-------------	----------------------------	------	---

³³ Rescaled by us using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (8.3 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.6 \pm 0.5)\%$. Multiplied by a factor of 2 to convert from $K_S^0 K^+ \pi^-$ to $K^0 K^+ \pi^-$ decay.

$\Gamma(\phi\phi)/\Gamma_{\text{total}}$ **Γ_{19}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>
---	--------------------

1.14±0.12 OUR FIT

$\Gamma(\omega\omega)/\Gamma_{\text{total}}$ **Γ_{20}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

0.92±0.11 OUR AVERAGE

0.89±0.11±0.04	762	³⁴ ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma \text{ hadrons}$
1.9 ±0.6 ±0.1	27.7 ±7.4	³⁵ ABLIKIM	05N BES2	$\psi(2S) \rightarrow \gamma\chi_{c2} \rightarrow \gamma 6\pi$

³⁴ ABLIKIM 11K reports $(8.9 \pm 0.3 \pm 1.1) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³⁵ ABLIKIM 05N reports $[\Gamma(\chi_{c2}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (0.165 \pm 0.044 \pm 0.032) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\omega\phi)/\Gamma_{\text{total}}$ **Γ_{21}/Γ**

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<2.0	90	³⁶ ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma \text{ hadrons}$
------	----	-----------------------	----------	---

³⁶ ABLIKIM 11K reports $< 2 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 8.72 \times 10^{-2}$.

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$ **Γ_{22}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>
---	--------------------

2.43±0.13 OUR FIT

$\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_{23}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>
---	--------------------

40±17 OUR FIT

$\Gamma(\pi^+\pi^-\eta)/\Gamma_{\text{total}}$ **Γ_{24}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

0.52±0.14±0.02		³⁷ ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$
-----------------------	--	---------------------	---------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.6	90	³⁸ ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma\chi_{c2}$
------	----	-----------------------	----------	--

³⁷ ATHAR 07 reports $(0.49 \pm 0.12 \pm 0.06) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³⁸ ABLIKIM 06R reports $< 1.7 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 8.72 \times 10^{-2}$.

$\Gamma(\pi^+ \pi^- \eta')/\Gamma_{\text{total}}$ Γ_{25}/Γ

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.55±0.20±0.02	³⁹ ATHAR	07	CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

³⁹ ATHAR 07 reports $(0.51 \pm 0.18 \pm 0.06) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+ \pi^- \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$ Γ_{26}/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>
5.9±0.5 OUR FIT	

$\Gamma(K^+ K^-)/\Gamma_{\text{total}}$ Γ_{27}/Γ

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>
1.09±0.08 OUR FIT	

$\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{28}/Γ

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>
0.58±0.05 OUR FIT	

$\Gamma(K_S^0 K_S^0)/\Gamma(\pi\pi)$ Γ_{28}/Γ_{22}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.239±0.019 OUR FIT			

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.27 \pm 0.07 \pm 0.04$	^{40,41} CHEN	07B	BELL $e^+ e^- \rightarrow e^+ e^- \chi_{c2}$
--------------------------	-----------------------	-----	--

⁴⁰ Using $\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ from the $\pi^+ \pi^-$ measurement of NAKAZAWA 05 rescaled by 3/2 to convert to $\pi\pi$.

⁴¹ Not independent from other measurements.

$\Gamma(K_S^0 K_S^0)/\Gamma(K^+ K^-)$ Γ_{28}/Γ_{27}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.53±0.05 OUR FIT			

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.70 \pm 0.21 \pm 0.12$	^{42,43} CHEN	07B	BELL $e^+ e^- \rightarrow e^+ e^- \chi_{c2}$
--------------------------	-----------------------	-----	--

⁴² Using $\Gamma(K^+ K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ from NAKAZAWA 05.

⁴³ Not independent from other measurements.

$\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{30}/Γ

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.33±0.08±0.01	⁴⁴ ATHAR	07	CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

⁴⁴ ATHAR 07 reports $(0.31 \pm 0.07 \pm 0.04) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ K^- \eta)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.35	90	⁴⁵ ATHAR 07	CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$
⁴⁵ ATHAR 07 reports $< 0.33 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.72 \times 10^{-2}$.				

$\Gamma(\eta\eta')/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.6	90	3.3 ± 8.0	⁴⁶ ASNER 09	CLEO	$\psi(2S) \rightarrow \gamma \eta \eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<2.5	90		⁴⁷ ADAMS 07	CLEO	$\psi(2S) \rightarrow \gamma \chi_{c2}$
⁴⁶ ASNER 09 reports $< 0.6 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.72 \times 10^{-2}$.					
⁴⁷ Superseded by ASNER 09. ADAMS 07 reports $< 2.3 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 0.0933 \pm 0.0014 \pm 0.0061$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.72 \times 10^{-2}$.					

$\Gamma(\eta' \eta')/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<1.1	90	12 ± 7	⁴⁸ ASNER 09	CLEO	$\psi(2S) \rightarrow \gamma \eta' \eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<3.3	90		⁴⁹ ADAMS 07	CLEO	$\psi(2S) \rightarrow \gamma \chi_{c2}$
⁴⁸ ASNER 09 reports $< 1.0 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta' \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.72 \times 10^{-2}$.					
⁴⁹ Superseded by ASNER 09. ADAMS 07 reports $< 3.1 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta' \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 0.0933 \pm 0.0014 \pm 0.0061$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.72 \times 10^{-2}$.					

$\Gamma(\pi^+ \pi^- K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$2.4 \pm 0.6 \pm 0.1$		57 ± 11	⁵⁰ ABLIKIM 050	BES2	$\psi(2S) \rightarrow \gamma \chi_{c2}$
⁵⁰ ABLIKIM 050 reports $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+ \pi^- K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ = $(0.207 \pm 0.039 \pm 0.033) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

$\Gamma(K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<4	90	2.3 ± 2.2	⁵¹ ABLIKIM	050 BES2	$e^+ e^- \rightarrow \chi_{c2} \gamma$

⁵¹ ABLIKIM 050 reports $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] < 3.5 \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.72 \times 10^{-2}$.

$\Gamma(K^+ K^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{36}/Γ

VALUE (units 10^{-3})	DOCUMENT ID
1.78 ± 0.22 OUR FIT	

$\Gamma(K^+ K^- \phi)/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.55 ± 0.32 ± 0.06	52	⁵² ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

⁵² ABLIKIM 06T reports $(1.67 \pm 0.26 \pm 0.24) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE (units 10^{-4})	DOCUMENT ID
0.72 ± 0.04 OUR FIT	

$\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$ Γ_{39}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.51 ± 0.05 OUR AVERAGE			
$0.52 \pm 0.04 \pm 0.02$	⁵³ ONYISI	10 CLE3	$\psi(2S) \rightarrow \gamma p\bar{p}X$
$0.47 \pm 0.10 \pm 0.02$	⁵⁴ ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

⁵³ ONYISI 10 reports $(4.83 \pm 0.25 \pm 0.35 \pm 0.31) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵⁴ ATHAR 07 reports $(0.44 \pm 0.08 \pm 0.05) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$ Γ_{40}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.190 ± 0.028 OUR AVERAGE			
$0.188 \pm 0.028 \pm 0.007$	⁵⁵ ONYISI	10 CLE3	$\psi(2S) \rightarrow \gamma p\bar{p}X$
$0.20 \pm 0.08 \pm 0.01$	⁵⁶ ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

⁵⁵ ONYISI 10 reports $(1.76 \pm 0.23 \pm 0.14 \pm 0.11) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵⁶ ATHAR 07 reports $(0.19 \pm 0.07 \pm 0.02) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$ **Γ_{41}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.39±0.05±0.02	⁵⁷ ONYISI 10	CLE3	$\psi(2S) \rightarrow \gamma p\bar{p}X$

⁵⁷ ONYISI 10 reports $(3.68 \pm 0.35 \pm 0.26 \pm 0.24) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$ **Γ_{42}/Γ**

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.0±0.9±0.1	24 ± 7	⁵⁸ ABLIKIM 11F	BES3	$\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$

⁵⁸ ABLIKIM 11F reports $(3.04 \pm 0.85 \pm 0.43) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_{43}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.32±0.34 OUR EVALUATION	Treating systematic error as correlated.		
1.3 ± 0.4 OUR AVERAGE	Error includes scale factor of 1.3.		

1.17±0.19±0.30 ⁵⁹ BAI 99B BES $\psi(2S) \rightarrow \gamma\chi_{c2}$

2.64±1.03±0.14 ⁵⁹ TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c2}$

⁵⁹ Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.3 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$. Multiplied by a factor of 2 to convert from $K_S^0 K^+\pi^-$ to $K^0 K^+\pi^-$ decay.

$\Gamma(p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}$ **Γ_{44}/Γ**

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.086±0.026±0.003	29.2	⁶⁰ HE 08B	CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

⁶⁰ HE 08B reports $0.08 \pm 0.02 \pm 0.01 \pm 0.01 \%$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\rho\bar{p}K^+K^- \text{ (non-resonant)})/\Gamma_{\text{total}}$ **Γ_{45}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.09 \pm 0.35 \pm 0.08$	131 ± 12	⁶¹ ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma\rho\bar{p}K^+K^-$

⁶¹ ABLIKIM 11F reports $(2.08 \pm 0.19 \pm 0.30) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \rho\bar{p}K^+K^- \text{ (non-resonant)})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\rho\bar{p}K_S^0K_S^0)/\Gamma_{\text{total}}$ **Γ_{46}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<7.9	90	⁶² ABLIKIM	06D BES2	$\psi(2S) \rightarrow \chi_{c2}\gamma$

⁶² Using $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (9.3 \pm 0.6)\%$.

$\Gamma(\rho\bar{n}\pi^-)/\Gamma_{\text{total}}$ **Γ_{47}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$11.1 \pm 3.8 \pm 0.4$	⁶³ ABLIKIM	06i BES2	$\psi(2S) \rightarrow \gamma\rho\pi^-X$

⁶³ ABLIKIM 06i reports $[\Gamma(\chi_{c2}(1P) \rightarrow \rho\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ = $(0.97 \pm 0.20 \pm 0.26) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$ **Γ_{48}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>
1.86 ± 0.27 OUR FIT	

$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_{49}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3.5	90	⁶⁴ ABLIKIM	06D BES2	$\psi(2S) \rightarrow \chi_{c2}\gamma$

⁶⁴ Using $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (9.3 \pm 0.6)\%$.

$\Gamma(K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_{50}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.91 \pm 0.17 \pm 0.04$	⁶⁵ ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

⁶⁵ ATHAR 07 reports $(0.85 \pm 0.14 \pm 0.10) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+\rho\Lambda(1520) + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_{51}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.1 \pm 0.7 \pm 0.1$	79 ± 13	⁶⁶ ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma\rho\bar{p}K^+K^-$

⁶⁶ ABLIKIM 11F reports $(3.06 \pm 0.50 \pm 0.54) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ p \Lambda(1520) + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Lambda(1520)\bar{\Lambda}(1520))/\Gamma_{\text{total}}$ **Γ_{52}/Γ**

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$5.1 \pm 1.6 \pm 0.2$	29 ± 7	⁶⁷ ABLIKIM	11F	BES3 $\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$

⁶⁷ ABLIKIM 11F reports $(5.05 \pm 1.29 \pm 0.93) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Lambda(1520)\bar{\Lambda}(1520))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}$ **Γ_{53}/Γ**

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 0.8	90	7.5 ± 3.4	⁶⁸ NAIK	08	CLEO $\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$

⁶⁸ NAIK 08 reports $< 0.75 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.72 \times 10^{-2}$.

$\Gamma(\Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}$ **Γ_{54}/Γ**

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 0.7	90	4.0 ± 3.5	⁶⁹ NAIK	08	CLEO $\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$

⁶⁹ NAIK 08 reports $< 0.67 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.72 \times 10^{-2}$.

$\Gamma(\Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}$ **Γ_{55}/Γ**

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 1.1	90	2.9 ± 1.7	⁷⁰ NAIK	08	CLEO $\psi(2S) \rightarrow \gamma \Xi^0 \bar{\Xi}^0$

⁷⁰ NAIK 08 reports $< 1.06 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.72 \times 10^{-2}$.

$\Gamma(\Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}$ **Γ_{56}/Γ**

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$1.55 \pm 0.34 \pm 0.06$		29 ± 5	⁷¹ NAIK	08	CLEO $\psi(2S) \rightarrow \gamma \Xi^+ \bar{\Xi}^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.7	90		⁷² ABLIKIM	06D	BES2 $\psi(2S) \rightarrow \chi_{c2} \gamma$
---------	----	--	-----------------------	-----	--

⁷¹ NAIK 08 reports $(1.45 \pm 0.30 \pm 0.15) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁷² Using $B(\psi(2S) \rightarrow \chi_{c2} \gamma) = (9.3 \pm 0.6)\%$.

$\Gamma(J/\psi(1S)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{57}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.015	90	BARATE	81	SPEC 190 GeV π^- Be $\rightarrow 2\pi 2\mu$

————— **RADIATIVE DECAYS** —————

$\Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$ Γ_{58}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.195 ± 0.008 OUR FIT			

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.199 ± 0.005 ± 0.012	⁷³ ADAM	05A	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$
⁷³ Uses $B(\psi(2S) \rightarrow \gamma\chi_{c2} \rightarrow \gamma\gamma J/\psi)$ from ADAM 05A and $B(\psi(2S) \rightarrow \gamma\chi_{c2})$ from ATHAR 04.			

$\Gamma(\gamma\rho^0)/\Gamma_{\text{total}}$ Γ_{59}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<21	90	13 ± 11	⁷⁴ ABLIKIM	11E	BES3 $\psi(2S) \rightarrow \gamma\gamma\rho^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<50	90	17.2 ± 6.8	⁷⁵ BENNETT	08A	CLEO $\psi(2S) \rightarrow \gamma\gamma\rho^0$
⁷⁴ ABLIKIM 11E reports $< 20.8 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 8.72 \times 10^{-2}$.					
⁷⁵ BENNETT 08A reports $< 50 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 8.72 \times 10^{-2}$.					

$\Gamma(\gamma\omega)/\Gamma_{\text{total}}$ Γ_{60}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<6	90	1 ± 6	⁷⁶ ABLIKIM	11E	BES3 $\psi(2S) \rightarrow \gamma\gamma\omega$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<7	90	0.0 ± 1.8	⁷⁷ BENNETT	08A	CLEO $\psi(2S) \rightarrow \gamma\gamma\omega$
⁷⁶ ABLIKIM 11E reports $< 6.1 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 8.72 \times 10^{-2}$.					
⁷⁷ BENNETT 08A reports $< 7.0 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 8.72 \times 10^{-2}$.					

$\Gamma(\gamma\phi)/\Gamma_{\text{total}}$ Γ_{61}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 8	90	5 ± 5	⁷⁸ ABLIKIM	11E	BES3 $\psi(2S) \rightarrow \gamma\gamma\phi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<12	90	1.3 ± 2.5	⁷⁹ BENNETT	08A	CLEO $\psi(2S) \rightarrow \gamma\gamma\phi$
⁷⁸ ABLIKIM 11E reports $< 8.1 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 8.72 \times 10^{-2}$.					
⁷⁹ BENNETT 08A reports $< 13 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 8.72 \times 10^{-2}$.					

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	Γ_{62}/Γ
<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>
2.59 ± 0.16 OUR FIT	

$\Gamma(\gamma\gamma)/\Gamma(\gamma J/\psi(1S))$	Γ_{62}/Γ_{58}
<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
1.33 ± 0.09 OUR FIT	
0.99 ± 0.18	⁸⁰ AMBROGIANI 00B E835 $\bar{p}p \rightarrow \chi_{c2} \rightarrow \gamma\gamma, \gamma J/\psi$

⁸⁰ Calculated by us using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}} \times \Gamma(\rho\bar{\rho})/\Gamma_{\text{total}}$	$\Gamma_{62}/\Gamma \times \Gamma_{38}/\Gamma$
<u>VALUE (units 10^{-8})</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
1.86 ± 0.18 OUR FIT	
1.7 ± 0.4 OUR AVERAGE	
1.60 ± 0.42	ARMSTRONG 93 E760 $\bar{p}p \rightarrow \gamma\gamma X$
9.9 ± 4.5	BAGLIN 87B SPEC $\bar{p}p \rightarrow \gamma\gamma X$

$\chi_{c2}(1P)$ CROSS-PARTICLE BRANCHING RATIOS

$\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-\pi^+\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$	$\Gamma_{14}/\Gamma \times \Gamma_{111}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$
<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
2.36 ± 0.27 OUR FIT	
2.5 ± 0.9 OUR AVERAGE	Error includes scale factor of 2.3.

1.90 ± 0.14 ± 0.44	BAI 99B BES $\psi(2S) \rightarrow \gamma\chi_{c2}$
3.8 ± 0.67	⁸¹ TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c2}$

⁸¹ The reported value is derived using $B(\psi(2S) \rightarrow \pi^+\pi^- J/\psi) \times B(J/\psi \rightarrow \ell^+\ell^-) = (4.6 \pm 0.7)\%$. Calculated by us using $B(J/\psi \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.

$\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0\bar{K}^*(892)^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}$	$\Gamma_{17}/\Gamma \times \Gamma_{111}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$
<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
2.2 ± 0.4 OUR FIT	
3.11 ± 0.36 ± 0.48	ABLIKIM 04H BES2 $\psi(2S) \rightarrow \gamma\chi_{c2}$

$\Gamma(\chi_{c2}(1P) \rightarrow \rho\bar{\rho})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$	$\Gamma_{38}/\Gamma \times \Gamma_{111}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
1.86 ± 0.14 OUR FIT	
1.4 ± 1.1	⁸² BAI 98I BES $\psi(2S) \rightarrow \gamma\chi_{c2} \rightarrow \gamma\bar{p}p$

⁸² Calculated by us. The value for $B(\chi_{c2} \rightarrow \rho\bar{\rho})$ reported in BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}} \quad \Gamma_{38}/\Gamma \times \Gamma_{111}^{\psi(2S)}/\Gamma_{\psi(2S)}$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

6.3±0.5 OUR FIT

6.7±1.1 OUR AVERAGE Error includes scale factor of 1.5.

7.2±0.7±0.4	121 ± 12	⁸³ NAIK	08	CLEO $\psi(2S) \rightarrow \gamma p\bar{p}$
4.4 ^{+1.6} _{-1.4} ±0.6	14.3 ^{+5.2} _{-4.7}	BAI	04F	BES $\psi(2S) \rightarrow \gamma\chi_{c2}(1P) \rightarrow \gamma p\bar{p}$

⁸³ Calculated by us. NAIK 08 reports $B(\chi_{c2} \rightarrow p\bar{p}) = (7.7 \pm 0.8 \pm 0.4 \pm 0.5) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$.

$$\Gamma(\chi_{c2}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}} \quad \Gamma_{48}/\Gamma \times \Gamma_{111}^{\psi(2S)}/\Gamma_{\psi(2S)}$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

16.3±2.3 OUR FIT

15.9±2.1±1.0 71 ± 9 ⁸⁴ NAIK 08 CLEO $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$

⁸⁴ Calculated by us. NAIK 08 reports $B(\chi_{c2} \rightarrow \Lambda\bar{\Lambda}) = (17.0 \pm 2.2 \pm 1.1 \pm 1.1) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$.

$$\Gamma(\chi_{c2}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \quad \Gamma_{48}/\Gamma \times \Gamma_{111}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

4.8±0.7 OUR FIT

7.1^{+3.1}_{-2.9}±1.3 8.3^{+3.7}_{-3.4} ⁸⁵ BAI 03E BES $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$

⁸⁵ BAI 03E reports $[B(\chi_{c2} \rightarrow \Lambda\bar{\Lambda}) B(\psi(2S) \rightarrow \gamma\chi_{c2}) / B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-)] \times [B^2(\Lambda \rightarrow \pi^-p) / B(J/\psi \rightarrow p\bar{p})] = (1.33^{+0.59}_{-0.55} \pm 0.25)\%$. We calculate from this measurement the presented value using $B(\Lambda \rightarrow \pi^-p) = (63.9 \pm 0.5)\%$ and $B(J/\psi \rightarrow p\bar{p}) = (2.17 \pm 0.07) \times 10^{-3}$.

$$\Gamma(\chi_{c2}(1P) \rightarrow \pi\pi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}} \quad \Gamma_{22}/\Gamma \times \Gamma_{111}^{\psi(2S)}/\Gamma_{\psi(2S)}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.11±0.08 OUR FIT

2.17±0.09 OUR AVERAGE

2.19±0.05±0.15	4.5k	⁸⁶ ABLIKIM	10A	BES3 $e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$
2.23±0.06±0.10	2.5k	⁸⁷ ASNER	09	CLEO $\psi(2S) \rightarrow \gamma\pi^+\pi^-$
1.90±0.08±0.20	0.8k	⁸⁸ ASNER	09	CLEO $\psi(2S) \rightarrow \gamma\pi^0\pi^0$

⁸⁶ Calculated by us. ABLIKIM 10A reports $B(\chi_{c2} \rightarrow \pi^0\pi^0) = (0.88 \pm 0.02 \pm 0.06 \pm 0.04) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.3 \pm 0.4)\%$. We have multiplied the $\pi^0\pi^0$ measurement by 3 to obtain $\pi\pi$.

⁸⁷ Calculated by us. ASNER 09 reports $B(\chi_{c2} \rightarrow \pi^+\pi^-) = (1.59 \pm 0.04 \pm 0.07 \pm 0.10) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$. We have multiplied the $\pi^+\pi^-$ measurement by 3/2 to obtain $\pi\pi$.

⁸⁸ Calculated by us. ASNER 09 reports $B(\chi_{c2} \rightarrow \pi^0\pi^0) = (0.68 \pm 0.03 \pm 0.07 \pm 0.04) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$. We have multiplied the $\pi^0\pi^0$ measurement by 3 to obtain $\pi\pi$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \pi\pi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{22}/\Gamma \times \Gamma_{111}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10⁻³) EVTS DOCUMENT ID TECN COMMENT

0.629±0.024 OUR FIT

0.54 ±0.06 OUR AVERAGE

0.66 ±0.18 ±0.37	21 ± 6	89 BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi^0\pi^0$
0.54 ±0.05 ±0.04	185 ± 16	90 BAI	98I BES	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$

⁸⁹ We have multiplied $\pi^0\pi^0$ measurement by 3 to obtain $\pi\pi$.

⁹⁰ Calculated by us. The value for $B(\chi_{c2} \rightarrow \pi^+\pi^-)$ reported by BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D]. We have multiplied $\pi^+\pi^-$ measurement by 3/2 to obtain $\pi\pi$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \eta\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{26}/\Gamma \times \Gamma_{111}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10⁻⁴) CL% EVTS DOCUMENT ID TECN COMMENT

0.52±0.04 OUR FIT

0.52±0.04 OUR AVERAGE

0.54±0.03±0.04	386	⁹¹ ABLIKIM	10A BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$
0.47±0.05±0.05	156 ± 14	ASNER	09 CLEO	$\psi(2S) \rightarrow \gamma\eta\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.44	90	⁹² ADAMS	07 CLEO	$\psi(2S) \rightarrow \gamma\chi_{c2}$
< 3	90	BAI	03C BES	$\psi(2S) \rightarrow \gamma\eta\eta \rightarrow 5\gamma$
0.62±0.31±0.19		LEE	85 CBAL	$\psi(2S) \rightarrow \text{photons}$

⁹¹ Calculated by us. ABLIKIM 10A reports $B(\chi_{c2} \rightarrow \eta\eta) = (0.65 \pm 0.04 \pm 0.05 \pm 0.03) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.3 \pm 0.4)\%$.

⁹² Superseded by ASNER 09.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{27}/\Gamma \times \Gamma_{111}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10⁻⁵) EVTS DOCUMENT ID TECN COMMENT

9.5±0.6 OUR FIT

10.5±0.3±0.6 1.6k ⁹³ ASNER 09 CLEO $\psi(2S) \rightarrow \gamma K^+K^-$

⁹³ Calculated by us. ASNER 09 reports $B(\chi_{c2} \rightarrow K^+K^-) = (1.13 \pm 0.03 \pm 0.06 \pm 0.07) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{27}/\Gamma \times \Gamma_{111}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10⁻³) EVTS DOCUMENT ID TECN COMMENT

0.283±0.017 OUR FIT

0.190±0.034±0.019 115 ± 13 ⁹⁴ BAI 98I BES $\psi(2S) \rightarrow \gamma K^+K^-$

⁹⁴ Calculated by us. The value for $B(\chi_{c2} \rightarrow K^+K^-)$ reported by BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\Gamma(\chi_{c2}(1P) \rightarrow K_S^0 K_S^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}} \times \Gamma_{28} / \Gamma \times \Gamma_{111}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

5.1 ± 0.4 OUR FIT

5.0 ± 0.4 OUR AVERAGE

4.9 ± 0.3 ± 0.3	373 ± 20	⁹⁵ ASNER	09 CLEO	$\psi(2S) \rightarrow \gamma K_S^0 K_S^0$
5.72 ± 0.76 ± 0.63	65	ABLIKIM	050 BES2	$\psi(2S) \rightarrow \gamma K_S^0 K_S^0$

⁹⁵ Calculated by us. ASNER 09 reports $B(\chi_{c2} \rightarrow K_S^0 K_S^0) = (0.53 \pm 0.03 \pm 0.03 \pm 0.03) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$.

$$\Gamma(\chi_{c2}(1P) \rightarrow K_S^0 K_S^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) \times \Gamma_{28} / \Gamma \times \Gamma_{111}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

15.0 ± 1.1 OUR FIT

14.7 ± 4.1 ± 3.3

⁹⁶ BAI	99B BES	$\psi(2S) \rightarrow \gamma K_S^0 K_S^0$
-------------------	---------	---

⁹⁶ Calculated by us. The value of $B(\chi_{c2} \rightarrow K_S^0 K_S^0)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\Gamma(\chi_{c2}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}} \times \Gamma_{29} / \Gamma \times \Gamma_{111}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.22 ± 0.17 OUR FIT

1.15 ± 0.18 OUR AVERAGE

1.21 ± 0.19 ± 0.09	37	⁹⁷ ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
0.97 ± 0.32 ± 0.13	28	⁹⁸ ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

⁹⁷ Calculated by us. ATHAR 07 reports $B(\chi_{c2} \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) = (1.3 \pm 0.2 \pm 0.1 \pm 0.1) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$.

⁹⁸ Calculated by us. ABLIKIM 06R reports $B(\chi_{c2} \rightarrow K_S^0 K^\pm \pi^\mp) = (0.6 \pm 0.2 \pm 0.1) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (8.1 \pm 0.6)\%$. We have multiplied by 2 to obtain $\bar{K}^0 K^+ \pi^- + \text{c.c.}$ from $K_S^0 K^\pm \pi^\mp$.

$$\Gamma(\chi_{c2}(1P) \rightarrow 2(\pi^+ \pi^-)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) \times \Gamma_1 / \Gamma \times \Gamma_{111}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

2.86 ± 0.27 OUR FIT

3.1 ± 1.0 OUR AVERAGE Error includes scale factor of 2.5.

2.3 ± 0.1 ± 0.5	⁹⁹ BAI	99B BES	$\psi(2S) \rightarrow \gamma \chi_{c2}$
4.3 ± 0.6	¹⁰⁰ TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c2}$

⁹⁹ Calculated by us. The value for $B(\chi_{c2} \rightarrow 2\pi^+ 2\pi^-)$ reported in BAI 99B is derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

¹⁰⁰ The value for $B(\psi(2S) \rightarrow \gamma \chi_{c2}) \times B(\chi_{c2} \rightarrow 2\pi^+ \pi^-)$ reported in TANENBAUM 78 is derived using $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) \times B(J/\psi(1S) \ell^+ \ell^-) = (4.6 \pm 0.7)\%$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma_{36} / \Gamma \times \Gamma_{111}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.55±0.19 OUR FIT				
1.76±0.16±0.24	160	¹⁰¹ ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

¹⁰¹ Calculated by us. The value of $B(\chi_{c2} \rightarrow 2K^+ 2K^-)$ reported by ABLIKIM 06T was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.1 \pm 0.4)\%$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) \times \Gamma_{36} / \Gamma \times \Gamma_{111}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
4.6±0.6 OUR FIT			
3.6±0.6±0.6	¹⁰² BAI	99B BES	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

¹⁰² Calculated by us. The value of $B(\chi_{c2} \rightarrow 2K^+ 2K^-)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \phi\phi) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma_{19} / \Gamma \times \Gamma_{111}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.00±0.10 OUR FIT				
0.98±0.13 OUR AVERAGE				Error includes scale factor of 1.3.
0.94±0.03±0.10	849	¹⁰³ ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma$ hadrons
1.38±0.24±0.23	41	¹⁰⁴ ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

¹⁰³ Calculated by us. The value of $B(\chi_{c2} \rightarrow \phi\phi)$ reported by ABLIKIM 11K was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35)\%$.

¹⁰⁴ Calculated by us. The value of $B(\chi_{c2} \rightarrow \phi\phi)$ reported by ABLIKIM 06T was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.1 \pm 0.4)\%$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \phi\phi) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) \times \Gamma_{19} / \Gamma \times \Gamma_{111}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$


VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
2.96±0.29 OUR FIT			
4.8 ± 1.3 ± 1.3	¹⁰⁵ BAI	99B BES	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

¹⁰⁵ Calculated by us. The value of $B(\chi_{c2} \rightarrow \phi\phi)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma_{58} / \Gamma \times \Gamma_{111}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.70±0.04 OUR FIT				
1.34±0.14 OUR AVERAGE				Error includes scale factor of 1.9. See the ideogram below.
1.62±0.04±0.12	5.8k	BAI	04I BES2	$\psi(2S) \rightarrow J/\psi \gamma \gamma$
0.99±0.10±0.08		GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$
1.47±0.17		¹⁰⁶ OREGLIA	82 CBAL	$\psi(2S) \rightarrow \gamma \chi_{c2}$

- 1.8 ± 0.5 107 BRANDELIK 79B DASP $\psi(2S) \rightarrow \gamma\chi_{c2}$
- 1.2 ± 0.2 107 BARTEL 78B CNTR $\psi(2S) \rightarrow \gamma\chi_{c2}$
- 2.2 ± 1.2 108 BIDDICK 77 CNTR $e^+e^- \rightarrow \gamma X$
- 1.2 ± 0.7 106 WHITAKER 76 MRK1 e^+e^-
- • • We do not use the following data for averages, fits, limits, etc. • • •
- 1.95±0.02±0.07 12.4k 109 MENDEZ 08 CLEO $\psi(2S) \rightarrow \gamma\chi_{c2}$
- 1.85±0.04±0.07 1.9k 110 ADAM 05A CLEO Repl. by MENDEZ 08
- 106 Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.
- 107 Recalculated by us using $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$.
- 108 Assumes isotropic gamma distribution.
- 109 Not independent from other measurements of MENDEZ 08.
- 110 Not independent from other values reported by ADAM 05A.



$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) / \Gamma_{\text{total}} \text{ (units } 10^{-2}\text{)}$$

$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \text{ anything})$$

$$\Gamma_{58}/\Gamma \times \Gamma_{111}^{\psi(2S)} / \Gamma_9^{\psi(2S)} = \Gamma_{58}/\Gamma \times \Gamma_{111}^{\psi(2S)} / (\Gamma_{11}^{\psi(2S)} + \Gamma_{12}^{\psi(2S)} + \Gamma_{13}^{\psi(2S)}) + 0.344 \Gamma_{110}^{\psi(2S)} + 0.195 \Gamma_{111}^{\psi(2S)}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.86±0.07 OUR FIT				

- • • We do not use the following data for averages, fits, limits, etc. • • •
- 3.12±0.03±0.09 12.4k 111 MENDEZ 08 CLEO $\psi(2S) \rightarrow \gamma\chi_{c2}$
- 3.11±0.07±0.07 1.9k ADAM 05A CLEO Repl. by MENDEZ 08
- 111 Not independent from other measurements of MENDEZ 08.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))}{\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)} \times \frac{\Gamma_{58}/\Gamma \times \Gamma_{111}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}{\Gamma_{111}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

5.07±0.13 OUR FIT

5.53±0.17 OUR AVERAGE

5.56±0.05±0.16	12.4k	MENDEZ	08	CLEO $\psi(2S) \rightarrow \gamma \chi_{c2}$
6.0 ±2.8	1.3k	¹¹² ABLIKIM	04B	BES $\psi(2S) \rightarrow J/\psi X$
3.9 ±1.2		¹¹³ HIMEL	80	MRK2 $\psi(2S) \rightarrow \gamma \chi_{c2}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.52±0.13±0.13	1.9k	¹¹⁴ ADAM	05A	CLEO Repl. by MENDEZ 08
----------------	------	---------------------	-----	-------------------------

¹¹² From a fit to the J/ψ recoil mass spectra.

¹¹³ The value for $B(\psi(2S) \rightarrow \gamma \chi_{c2}) \times B(\chi_{c2} \rightarrow \gamma J/\psi(1S))$ reported in HIMEL 80 is derived using $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (33 \pm 3)\%$ and $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.138 \pm 0.018$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = (0.1181 \pm 0.0020)$.

¹¹⁴ Not independent from other values reported by ADAM 05A.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \gamma \gamma)}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))}{\Gamma_{\text{total}}} \times \frac{\Gamma_{62}/\Gamma \times \Gamma_{111}^{\psi(2S)}/\Gamma_{\psi(2S)}}{\Gamma_{111}^{\psi(2S)}/\Gamma_{\psi(2S)}}$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.26±0.16 OUR FIT

2.73±0.32 OUR AVERAGE

2.68±0.28±0.15	333 ± 35	ECKLUND	08A	CLEO $\psi(2S) \rightarrow \gamma \chi_{c2} \rightarrow 3\gamma$
7.0 ±2.1 ±2.0		LEE	85	CBAL $\psi(2S) \rightarrow \gamma \chi_{c2}$

MULTIPOLE AMPLITUDES IN $\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)$ RADIATIVE DECAY

$a_2 = M2/\sqrt{E1^2 + M2^2 + E3^2}$ Magnetic quadrupole fractional transition amplitude

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

-10.0± 1.5 OUR AVERAGE

- 9.3± 1.6±0.3	19.8k	¹¹⁵ ARTUSO	09	CLEO $\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
- 9.3 ⁺ ₋ 3.9 ⁺ _{4.1} ±0.6	5.9k	¹¹⁶ AMBROGIANI	02	E835 $p\bar{p} \rightarrow \chi_{c2} \rightarrow J/\psi \gamma$
-14 ± 6	1.9k	¹¹⁶ ARMSTRONG	93E	E760 $p\bar{p} \rightarrow \chi_{c2} \rightarrow J/\psi \gamma$
-33.3 ⁺ ₋ 11.6 _{29.2}	441	¹¹⁶ OREGLIA	82	CBAL $\psi(2S) \rightarrow \chi_{c1} \gamma \rightarrow J/\psi \gamma \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

- 7.9± 1.9±0.3	19.8k	¹¹⁷ ARTUSO	09	CLEO $\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
----------------	-------	-----------------------	----	---

¹¹⁵ From a fit with floating $M2$ amplitudes a_2 and b_2 , and fixed $E3$ amplitudes $a_3=b_3=0$.

¹¹⁶ Assuming $a_3=0$.

¹¹⁷ From a fit with floating $M2$ and $E3$ amplitudes a_2 , b_2 , and a_3 , and b_3 .

$a_3 = E3/\sqrt{E1^2 + M2^2 + E3^2}$ Electric octupole fractional transition amplitude

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.6±1.3 OUR AVERAGE				
1.7±1.4±0.3	19.8k	¹¹⁸ ARTUSO 09	CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
2.0 ^{+5.5} _{-4.4} ±0.9	5908	AMBROGIANI 02	E835	$\rho\bar{p} \rightarrow \chi_{c2} \rightarrow J/\psi\gamma$
0 ⁺⁶ ₋₅	1904	ARMSTRONG 93E	E760	$\rho\bar{p} \rightarrow \chi_{c2} \rightarrow J/\psi\gamma$

¹¹⁸ From a fit with floating $M2$ and $E3$ amplitudes a_2 , b_2 , and a_3 , and b_3 .

MULTIPOLE AMPLITUDES IN $\psi(2S) \rightarrow \gamma\chi_{c2}(1P)$ RADIATIVE DECAY

$b_2 = M2/\sqrt{E1^2 + M2^2 + E3^2}$ Magnetic quadrupole fractional transition amplitude

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.2±1.8 OUR AVERAGE Error includes scale factor of 1.7. See the ideogram below.				
4.6±1.0±1.3	13.8k	¹¹⁹ ABLIKIM 11I	BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$
0.2±1.5±0.4	19.8k	¹²⁰ ARTUSO 09	CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
- 5.1 ^{+5.4} _{-3.6}	721	¹¹⁹ ABLIKIM 04I	BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$
13.2 ^{+9.8} _{-7.5}	441	¹²¹ OREGLIA 82	CBAL	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$


• • • We do not use the following data for averages, fits, limits, etc. • • •

1.0±1.3±0.3 19.8k ¹²¹ ARTUSO 09 CLEO $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

¹¹⁹ From a fit with floating $M2$ and $E3$ amplitudes b_2 and b_3 .

¹²⁰ From a fit with floating $M2$ and $E3$ amplitudes a_2 , b_2 , and a_3 , and b_3 .

¹²¹ From a fit with floating $M2$ amplitudes a_2 and b_2 , and fixed $E3$ amplitudes $a_3=b_3=0$.



$b_2 = M2/\sqrt{E1^2 + M2^2 + E3^2}$ Magnetic quadrupole fractional transition amplitude (units 10^{-2})

$b_3 = E3/\sqrt{E1^2 + M2^2 + E3^2}$ Electric octupole fractional transition amplitude

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
-0.3±1.0 OUR AVERAGE				
1.5±0.8±1.8	13.8k	122 ABLIKIM	11I BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$
-0.8±1.2±0.2	19.8k	ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
-2.7 ^{+4.3} / _{-2.9}	721	122 ABLIKIM	04I BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$

¹²² From a fit with floating $M2$ and $E3$ amplitudes b_2 and b_3 .

MULTIPOLE AMPLITUDE RATIOS IN RADIATIVE DECAYS $\psi(2S) \rightarrow \gamma\chi_{c2}(1P)$ and $\chi_{c2} \rightarrow \gamma J/\psi(1S)$

b_2/a_2 Magnetic quadrupole transition amplitude ratio

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
-11⁺¹⁴/₋₁₅	19.8k	¹²³ ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

¹²³ Statistical and systematic errors combined. From a fit with floating $M2$ amplitudes a_2 and b_2 , and fixed $E3$ amplitudes $a_3=b_3=0$. Not independent of values for $a_2(\chi_{c2}(1P))$ and $b_2(\chi_{c2}(1P))$ from ARTUSO 09.

$\chi_{c2}(1P)$ REFERENCES

ABLIKIM	11A	PR D83 012006	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	11E	PR D83 112005	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	11F	PR D83 112009	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	11I	PR D84 092006	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	11K	PRL 107 092001	M. Ablikim <i>et al.</i>	(BES III Collab.)
DEL-AMO-SA...	11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
ABLIKIM	10A	PR D81 052005	M. Ablikim <i>et al.</i>	(BES III Collab.)
ONYISI	10	PR D82 011103R	P.U.E. Onyisi <i>et al.</i>	(CLEO Collab.)
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)
ARTUSO	09	PR D80 112003	M. Artuso <i>et al.</i>	(CLEO Collab.)
ASNER	09	PR D79 072007	D.M. Asner <i>et al.</i>	(CLEO Collab.)
UEHARA	09	PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.)
BENNETT	08A	PRL 101 151801	J.V. Bennett <i>et al.</i>	(CLEO Collab.)
ECKLUND	08A	PR D78 091501R	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)
HE	08B	PR D78 092004	Q. He <i>et al.</i>	(CLEO Collab.)
MENDEZ	08	PR D78 011102R	H. Mendez <i>et al.</i>	(CLEO Collab.)
NAIK	08	PR D78 031101R	P. Naik <i>et al.</i>	(CLEO Collab.)
UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)
ADAMS	07	PR D75 071101R	G.S. Adams <i>et al.</i>	(CLEO Collab.)
ATHAR	07	PR D75 032002	S.B. Athar <i>et al.</i>	(CLEO Collab.)
CHEN	07B	PL B651 15	W.T. Chen <i>et al.</i>	(BELLE Collab.)
ABLIKIM	06D	PR D73 052006	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06I	PR D74 012004	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06T	PL B642 197	M. Ablikim <i>et al.</i>	(BES Collab.)
DOBBS	06	PR D73 071101R	S. Dobbs <i>et al.</i>	(CLEO Collab.)
ABLIKIM	05G	PR D71 092002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05N	PL B630 7	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAM	05A	PRL 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ANDREOTTI	05A	NP B717 34	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)
NAKAZAWA	05	PL B615 39	H. Nakazawa <i>et al.</i>	(BELLE Collab.)
ABLIKIM	04B	PR D70 012003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04H	PR D70 092003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04I	PR D70 092004	M. Ablikim <i>et al.</i>	(BES Collab.)
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)
BAI	04F	PR D69 092001	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04I	PR D70 012006	J.Z. Bai <i>et al.</i>	(BES Collab.)
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
BAI	03C	PR D67 032004	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	03E	PR D67 112001	J.Z. Bai <i>et al.</i>	(BES Collab.)
ABE	02T	PL B540 33	K. Abe <i>et al.</i>	(BELLE Collab.)
AMBROGIANI	02	PR D65 052002	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
EISENSTEIN	01	PRL 87 061801	B.I. Eisenstein <i>et al.</i>	(CLEO Collab.)
AMBROGIANI	00B	PR D62 052002	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
ACCIARRI	99E	PL B453 73	M. Acciarri <i>et al.</i>	(L3 Collab.)
BAI	99B	PR D60 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)
ACKER...,K...	98	PL B439 197	K. Akerstaff <i>et al.</i>	(OPAL Collab.)
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98I	PRL 81 3091	J.Z. Bai <i>et al.</i>	(BES Collab.)
DOMINICK	94	PR D50 4265	J. Dominick <i>et al.</i>	(CLEO Collab.)
ARMSTRONG	93	PRL 70 2988	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)
ARMSTRONG	93E	PR D48 3037	T.A. Armstrong <i>et al.</i>	(FNAL-E760 Collab.)
BAUER	93	PL B302 345	D.A. Bauer <i>et al.</i>	(TPC Collab.)
ARMSTRONG	92	NP B373 35	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
Also		PRL 68 1468	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
BAGLIN	87B	PL B187 191	C. Baglin <i>et al.</i>	(R704 Collab.)
BAGLIN	86B	PL B172 455	C. Baglin	(LAPP, CERN, GENO, LYON, OSLO+)
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)
LEE	85	SLAC 282	R.A. Lee	(SLAC)
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)
OREGLIA	82	PR D25 2259	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)
Also		Private Comm.	M.J. Oreglia	(EFI)
BARATE	81	PR D24 2994	R. Barate <i>et al.</i>	(SACL, LOIC, SHMP, CERN+)
HIMEL	80	PRL 44 920	T. Himel <i>et al.</i>	(LBL, SLAC)
Also		Private Comm.	G. Trilling	(LBL, UCB)
BRANDELIK	79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)

TANENBAUM	78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)
Also		Private Comm.	G. Trilling	(LBL, UCB)
BIDDICK	77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)
WHITAKER	76	PRL 37 1596	J.S. Whitaker <i>et al.</i>	(SLAC, LBL)
