



$$I(J^P) = ?(??) \quad \text{Status: } *$$

OMITTED FROM SUMMARY TABLE

This would presumably be an isospin-1/2 particle, a $ccu \Xi_{cc}^{++}$ and a $ccd \Xi_{cc}^+$. However, opposed to the evidence cited below, the BABAR experiment has found no evidence for a Ξ_{cc}^+ in a search in $\Lambda_c^+ K^- \pi^+$ and $\Xi_c^0 \pi^+$ modes, and no evidence of a Ξ_{cc}^{++} in $\Lambda_c^+ K^- \pi^+ \pi^+$ and $\Xi_c^0 \pi^+ \pi^+$ modes (AUBERT,B 06D). Nor has the BELLE experiment found any evidence for a Ξ_{cc}^+ in the $\Lambda_c^+ K^- \pi^+$ mode (CHISTOV 06).

Ξ_{cc}^+ MASS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3518.9 ± 0.9 OUR AVERAGE				
3518 ± 3	6	¹ OCHERASHVI.05	SELX	Σ^- nucleus \approx 600 GeV
3519 ± 1	16	² MATTSON 02	SELX	Σ^- nucleus \approx 600 GeV

¹ OCHERASHVILI 05 claims "an excess of 5.62 events over ... 1.38 ± 0.13 events" for a significance of 4.8σ in $pD^+ K^-$ events.

² MATTSON 02 claims "an excess of 15.9 events over an expected background of 6.1 ± 0.5 events, a statistical significance of 6.3σ " in the $\Lambda_c^+ K^- \pi^+$ invariant-mass spectrum.

The probability that the peak is a fluctuation increases from 1.0×10^{-6} to 1.1×10^{-4} when the number of bins searched is considered.

Ξ_{cc}^+ MEAN LIFE

<u>VALUE (10^{-15} s)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<33	90	MATTSON 02	SELX	Σ^- nucleus, \approx 600 GeV

Ξ_{cc}^+ DECAY MODES

Mode		
Γ_1	$\Lambda_c^+ K^- \pi^+$	
Γ_2	$pD^+ K^-$	

$\Gamma(pD^+ K^-)/\Gamma(\Lambda_c^+ K^- \pi^+)$			Γ_2/Γ_1	
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.36 ± 0.21	6	OCHERASHVI.05	SELX	$\Sigma^- \approx$ 600 GeV

 **REFERENCES**

AUBERT,B	06D	PR D74 011103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
CHISTOV	06	PRL 97 162001	R. Chistov <i>et al.</i>	(BELLE Collab.)
OCHERASHVI...	05	PL B628 18	A. Ocherashvili <i>et al.</i>	(FNAL SELEX Collab.)
MATTSON	02	PRL 89 112001	M. Mattson <i>et al.</i>	(FNAL SELEX Collab.)
