

$\chi_{c1}(1P)$ $I^G(J^{PC}) = 0^+(1^{++})$

See the Review on “ $\psi(2S)$ and χ_c branching ratios” before the
 $\chi_{c0}(1P)$ Listings.

 $\chi_{c1}(1P)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
3510.66 ± 0.07 OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.			
3510.30 ± 0.14 ± 0.16		ABLIKIM 05G	BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$	
3510.719 ± 0.051 ± 0.019		ANDREOTTI 05A	E835	$p\bar{p} \rightarrow e^+ e^- \gamma$	
3509.4 ± 0.9		BAI 99B	BES	$\psi(2S) \rightarrow \gamma X$	
3510.60 ± 0.087 ± 0.019	513	¹ ARMSTRONG 92	E760	$\bar{p}p \rightarrow e^+ e^- \gamma$	
3511.3 ± 0.4 ± 0.4	30	BAGLIN 86B	SPEC	$\bar{p}p \rightarrow e^+ e^- X$	
3512.3 ± 0.3 ± 4.0		² GAISER 86	CBAL	$\psi(2S) \rightarrow \gamma X$	
3507.4 ± 1.7	91	³ LEMOIGNE 82	GOLI	$185 \pi^- Be \rightarrow \gamma \mu^+ \mu^- A$	
3510.4 ± 0.6		OREGLIA 82	CBAL	$e^+ e^- \rightarrow J/\psi 2\gamma$	
3510.1 ± 1.1	254	⁴ HIMEL 80	MRK2	$e^+ e^- \rightarrow J/\psi 2\gamma$	
3509 ± 11	21	BRANDELIK 79B	DASP	$e^+ e^- \rightarrow J/\psi 2\gamma$	
3507 ± 3		⁴ BARTEL 78B	CNTR	$e^+ e^- \rightarrow J/\psi 2\gamma$	
3505.0 ± 4 ± 4		^{4,5} TANENBAUM 78	MRK1	$e^+ e^-$	
3513 ± 7	367	⁴ BIDDICK 77	CNTR	$\psi(2S) \rightarrow \gamma X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
3500 ± 10	40	TANENBAUM 75	MRK1	Hadrons γ	


¹ Recalculated by ANDREOTTI 05A, using the value of $\psi(2S)$ mass from AULCHENKO 03.

² Using mass of $\psi(2S) = 3686.0$ MeV.

³ $J/\psi(1S)$ mass constrained to 3097 MeV.

⁴ Mass value shifted by us by amount appropriate for $\psi(2S)$ mass = 3686 MeV and $J/\psi(1S)$ mass = 3097 MeV.

⁵ From a simultaneous fit to radiative and hadronic decay channels.



$\chi_{c1}(1P)$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.86 ±0.05 OUR FIT					
0.88 ±0.05 OUR AVERAGE					
1.39 +0.40 -0.38	+0.26 -0.77		ABLIKIM	05G BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$
0.876±0.045±0.026			ANDREOTTI	05A E835	$p\bar{p} \rightarrow e^+ e^- \gamma$
0.87 ±0.11 ±0.08		513	⁶ ARMSTRONG	92 E760	$\bar{p}p \rightarrow e^+ e^- \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.3	95		BAGLIN	86B SPEC	$\bar{p}p \rightarrow e^+ e^- X$
<3.8	90		GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$

⁶ Recalculated by ANDREOTTI 05A.

$\chi_{c1}(1P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
------	--------------------------------	-----------------------------------

Hadronic decays

Γ_1	$3(\pi^+\pi^-)$	$(5.8 \pm 1.4) \times 10^{-3}$	S=1.2
Γ_2	$2(\pi^+\pi^-)$	$(7.6 \pm 2.6) \times 10^{-3}$	
Γ_3	$\pi^+\pi^-\pi^0\pi^0$	$(1.25 \pm 0.17)\%$	
Γ_4	$\rho^+\pi^-\pi^0 + \text{c.c.}$	$(1.53 \pm 0.26)\%$	
Γ_5	$\rho^0\pi^+\pi^-$	$(3.9 \pm 3.5) \times 10^{-3}$	
Γ_6	$4\pi^0$	$(5.7 \pm 0.8) \times 10^{-4}$	
Γ_7	$\pi^+\pi^-K^+K^-$	$(4.5 \pm 1.0) \times 10^{-3}$	
Γ_8	$K^+K^-\pi^0\pi^0$	$(1.17 \pm 0.29) \times 10^{-3}$	
Γ_9	$K^+\pi^-K^0\pi^0 + \text{c.c.}$	$(9.0 \pm 1.4) \times 10^{-3}$	
Γ_{10}	$\rho^+K^-K^0 + \text{c.c.}$	$(5.3 \pm 1.3) \times 10^{-3}$	
Γ_{11}	$K^*(892)^0K^0\pi^0 \rightarrow K^+\pi^-K^0\pi^0 + \text{c.c.}$	$(2.4 \pm 0.7) \times 10^{-3}$	
Γ_{12}	$K^+K^-\eta\pi^0$	$(1.2 \pm 0.4) \times 10^{-3}$	
Γ_{13}	$\pi^+\pi^-K_S^0K_S^0$	$(7.2 \pm 3.1) \times 10^{-4}$	
Γ_{14}	$K^+K^-\eta$	$(3.3 \pm 1.0) \times 10^{-4}$	
Γ_{15}	$K^0K^+\pi^- + \text{c.c.}$	$(7.3 \pm 0.6) \times 10^{-3}$	
Γ_{16}	$K^*(892)^0\bar{K}^0 + \text{c.c.}$	$(1.0 \pm 0.4) \times 10^{-3}$	
Γ_{17}	$K^*(892)^+K^- + \text{c.c.}$	$(1.5 \pm 0.7) \times 10^{-3}$	
Γ_{18}	$K_J^*(1430)^0\bar{K}^0 + \text{c.c.} \rightarrow K_S^0K^+\pi^- + \text{c.c.}$	$< 8 \times 10^{-4}$	CL=90%
Γ_{19}	$K_J^*(1430)^+K^- + \text{c.c.} \rightarrow K_S^0K^+\pi^- + \text{c.c.}$	$< 2.3 \times 10^{-3}$	CL=90%
Γ_{20}	$K^+K^-\pi^0$	$(1.91 \pm 0.26) \times 10^{-3}$	
Γ_{21}	$\eta\pi^+\pi^-$	$(5.0 \pm 0.5) \times 10^{-3}$	
Γ_{22}	$a_0(980)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	$(1.9 \pm 0.7) \times 10^{-3}$	
Γ_{23}	$f_2(1270)\eta$	$(2.8 \pm 0.8) \times 10^{-3}$	
Γ_{24}	$\pi^+\pi^-\eta'$	$(2.3 \pm 0.5) \times 10^{-3}$	
Γ_{25}	$\pi^0f_0(980) \rightarrow \pi^0\pi^+\pi^-$	$< 6 \times 10^{-6}$	CL=90%
Γ_{26}	$K^+\bar{K}^*(892)^0\pi^- + \text{c.c.}$	$(3.2 \pm 2.1) \times 10^{-3}$	
Γ_{27}	$K^*(892)^0\bar{K}^*(892)^0$	$(1.5 \pm 0.4) \times 10^{-3}$	
Γ_{28}	$K^+K^-K_S^0K_S^0$	$< 5 \times 10^{-4}$	CL=90%
Γ_{29}	$K^+K^-K^+K^-$	$(5.6 \pm 1.2) \times 10^{-4}$	
Γ_{30}	$K^+K^-\phi$	$(4.3 \pm 1.6) \times 10^{-4}$	
Γ_{31}	$\omega\omega$	$(6.0 \pm 0.7) \times 10^{-4}$	
Γ_{32}	$\omega\phi$	$(2.2 \pm 0.6) \times 10^{-5}$	
Γ_{33}	$\phi\phi$	$(4.4 \pm 0.6) \times 10^{-4}$	
Γ_{34}	$p\bar{p}$	$(7.3 \pm 0.4) \times 10^{-5}$	
Γ_{35}	$p\bar{p}\pi^0$	$(1.63 \pm 0.20) \times 10^{-4}$	
Γ_{36}	$p\bar{p}\eta$	$(1.53 \pm 0.26) \times 10^{-4}$	
Γ_{37}	$p\bar{p}\omega$	$(2.23 \pm 0.33) \times 10^{-4}$	
Γ_{38}	$p\bar{p}\phi$	$< 1.8 \times 10^{-5}$	CL=90%
Γ_{39}	$p\bar{p}\pi^+\pi^-$	$(5.0 \pm 1.9) \times 10^{-4}$	
Γ_{40}	$p\bar{p}\pi^0\pi^0$		

Γ_{41}	$p\bar{p}K^+K^-$ (non-resonant)	$(1.34 \pm 0.24) \times 10^{-4}$	
Γ_{42}	$p\bar{p}K_S^0K_S^0$	$< 4.5 \times 10^{-4}$	CL=90%
Γ_{43}	$\Lambda\bar{\Lambda}$	$(1.18 \pm 0.19) \times 10^{-4}$	
Γ_{44}	$\Lambda\bar{\Lambda}\pi^+\pi^-$	$< 1.5 \times 10^{-3}$	CL=90%
Γ_{45}	$K^+\bar{p}\Lambda$	$(4.3 \pm 0.4) \times 10^{-4}$	S=1.1
Γ_{46}	$K^+p\Lambda(1520) + \text{c.c.}$	$(1.8 \pm 0.5) \times 10^{-4}$	
Γ_{47}	$\Lambda(1520)\bar{\Lambda}(1520)$	$< 1.0 \times 10^{-4}$	CL=90%
Γ_{48}	$\Sigma^0\bar{\Sigma}^0$	$< 4 \times 10^{-5}$	CL=90%
Γ_{49}	$\Sigma^+\bar{\Sigma}^-$	$< 6 \times 10^{-5}$	CL=90%
Γ_{50}	$\Xi^0\bar{\Xi}^0$	$< 6 \times 10^{-5}$	CL=90%
Γ_{51}	$\Xi^-\bar{\Xi}^+$	$(8.4 \pm 2.3) \times 10^{-5}$	
Γ_{52}	$\pi^+\pi^- + K^+K^-$	$< 2.1 \times 10^{-3}$	
Γ_{53}	$K_S^0K_S^0$	$< 6 \times 10^{-5}$	CL=90%

Radiative decays

Γ_{54}	$\gamma J/\psi(1S)$	$(34.8 \pm 1.5) \%$
Γ_{55}	$\gamma\rho^0$	$(2.27 \pm 0.19) \times 10^{-4}$
Γ_{56}	$\gamma\omega$	$(7.1 \pm 0.9) \times 10^{-5}$
Γ_{57}	$\gamma\phi$	$(2.6 \pm 0.6) \times 10^{-5}$
Γ_{58}	$\gamma\gamma$	

CONSTRAINED FIT INFORMATION

A multiparticle fit to $\chi_{c1}(1P)$, $\chi_{c0}(1P)$, $\chi_{c2}(1P)$, and $\psi(2S)$ with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 84 branching ratios uses 227 measurements to determine 49 parameters. The overall fit has a $\chi^2 = 325.4$ for 178 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

x_{29}	8				
x_{34}	-9	-4			
x_{43}	11	5	-5		
x_{54}	38	16	-31	21	
Γ	-13	-5	-60	-7	-29
	x_{15}	x_{29}	x_{34}	x_{43}	x_{54}

$\chi_{c1}(1P)$ PARTIAL WIDTHS

— $\chi_{c1}(1P) \Gamma(i)\Gamma(\gamma J/\psi(1S))/\Gamma(\text{total})$ —

$\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$	$\Gamma_{34}\Gamma_{54}/\Gamma$		
VALUE (eV)	DOCUMENT ID	TECN	COMMENT
21.7±0.8 OUR FIT			
21.4±0.9 OUR AVERAGE			
21.5±0.5±0.8	⁷ ANDREOTTI 05A	E835	$p\bar{p} \rightarrow e^+ e^- \gamma$
21.4±1.5±2.2	^{7,8} ARMSTRONG 92	E760	$\bar{p}p \rightarrow e^+ e^- \gamma$
19.9 ^{+4.4} _{-4.0}	7 BAGLIN	86B SPEC	$\bar{p}p \rightarrow e^+ e^- X$
⁷ Calculated by us using $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$.			
⁸ Recalculated by ANDREOTTI 05A.			

$\chi_{c1}(1P)$ BRANCHING RATIOS

— HADRONIC DECAYS —

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$	Γ_1/Γ		
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
5.8±1.4 OUR EVALUATION			
Error includes scale factor of 1.2. Treating systematic error as correlated.			
5.8±1.1 OUR AVERAGE			
5.4±0.7±0.9	⁹ BAI	99B BES	$\psi(2S) \rightarrow \gamma\chi_{c1}$
16.0±5.9±0.8	⁹ TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma\chi_{c1}$
⁹ Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (8.8 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$.			

$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$	Γ_2/Γ		
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
7.6±2.6 OUR EVALUATION			
Treating systematic error as correlated.			
8 ±4 OUR AVERAGE			
Error includes scale factor of 1.5.			
4.6±2.1±2.6	¹⁰ BAI	99B BES	$\psi(2S) \rightarrow \gamma\chi_{c1}$
12.5±4.2±0.6	¹⁰ TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma\chi_{c1}$
¹⁰ Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (8.8 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$.			

$\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}$	Γ_3/Γ			
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.25±0.16±0.05	604.7	¹¹ HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$
¹¹ HE 08B reports $1.28 \pm 0.06 \pm 0.15 \pm 0.08\%$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(\rho^+\pi^-\pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.53±0.25±0.06	712.3	12,13 HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+h^-h^0h^0$

¹² HE 08B reports $1.56 \pm 0.13 \pm 0.22 \pm 0.10$ % from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \rho^+\pi^-\pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹³ Calculated by us. We have added the values from HE 08B for $\rho^+\pi^-\pi^0$ and $\rho^-\pi^+\pi^0$ decays assuming uncorrelated statistical and fully correlated systematic uncertainties.

 $\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
39±35		14 TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma\chi_{c1}$

¹⁴ Estimated using $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.087$. The errors do not contain the uncertainty in the $\psi(2S)$ decay.

 $\Gamma(4\pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.57±0.08±0.02	608	15 ABLIKIM	11A BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c1}$

¹⁵ ABLIKIM 11A reports $(0.57 \pm 0.03 \pm 0.08) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow 4\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.5±1.0 OUR EVALUATION		Treating systematic error as correlated.		

4.5±0.9 OUR AVERAGE

4.2±0.4±0.9	16 BAI	99B BES	$\psi(2S) \rightarrow \gamma\chi_{c1}$
7.3±3.0±0.4	16 TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma\chi_{c1}$

¹⁶ Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (8.8 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$.

 $\Gamma(K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.117±0.028±0.005	45.1	17 HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+h^-h^0h^0$

¹⁷ HE 08B reports $0.12 \pm 0.02 \pm 0.02 \pm 0.01$ % from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.90±0.14±0.03	141.3	18 HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

18 HE 08B reports $0.92 \pm 0.09 \pm 0.11 \pm 0.06$ % from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\rho^+ K^- K^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.53±0.13±0.02	141.3	19 HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

19 HE 08B reports $0.54 \pm 0.11 \pm 0.07 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \rho^+ K^- K^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^*(892)^0 K^0 \pi^0 \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.24±0.07±0.01	141.3	20 HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

20 HE 08B reports $0.25 \pm 0.06 \pm 0.03 \pm 0.02$ % from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^0 K^0 \pi^0 \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^+ K^- \eta \pi^0)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.117±0.036±0.005	141.3	21 HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

21 HE 08B reports $0.12 \pm 0.03 \pm 0.02 \pm 0.01$ % from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- \eta \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\pi^+ \pi^- K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.2±3.0±0.3	19.8 ± 7.7	22 ABLIKIM	050 BES2	$\psi(2S) \rightarrow \chi_{c1} \gamma$

22 ABLIKIM 050 reports $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^+ \pi^- K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (0.67 \pm 0.26 \pm 0.11) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ K^- \eta)/\Gamma_{\text{total}}$ Γ_{14}/Γ

<u>VALUE</u> (units 10^{-3})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.33 \pm 0.10 \pm 0.01$	23 ATTHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

²³ ATTHAR 07 reports $(0.34 \pm 0.10 \pm 0.04) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{15}/Γ

<u>VALUE</u> (units 10^{-3})	<u>DOCUMENT ID</u>
7.3 ± 0.6 OUR FIT	

$\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{16}/Γ

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.03 \pm 0.38 \pm 0.04$	22	24 ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$

²⁴ ABLIKIM 06R reports $(1.1 \pm 0.4 \pm 0.1) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{17}/Γ

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.5 \pm 0.7 \pm 0.1$	27	25 ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$

²⁵ ABLIKIM 06R reports $(1.6 \pm 0.7 \pm 0.2) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K_J^*(1430)^0 \bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{18}/Γ

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.8	90	26 ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$

²⁶ ABLIKIM 06R reports $< 0.9 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K_J^*(1430)^0 \bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.3 \times 10^{-2}$.

$\Gamma(K_J^*(1430)^+ K^- + \text{c.c.} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{19}/Γ

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.3	90	27 ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$

²⁷ ABLIKIM 06R reports $< 2.4 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K_J^*(1430)^+ K^- + \text{c.c.} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.3 \times 10^{-2}$.

$\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{20}/Γ

<u>VALUE</u> (units 10^{-3})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.91±0.25±0.07	28 ATHAR	07	CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$
28 ATHAR 07 reports $(1.95 \pm 0.16 \pm 0.23) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			

 $\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{21}/Γ

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.0±0.5 OUR AVERAGE				
4.9±0.5±0.2		29 ATHAR	07	CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$
5.5±1.0±0.2	222	30 ABLIKIM	06R	BES2 $\psi(2S) \rightarrow \gamma \chi_{c1}$
29 ATHAR 07 reports $(5.0 \pm 0.3 \pm 0.5) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
30 ABLIKIM 06R reports $(5.9 \pm 0.7 \pm 0.8) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(a_0(980)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{22}/Γ

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.9±0.7±0.1	58	31 ABLIKIM	06R	BES2 $\psi(2S) \rightarrow \gamma \chi_{c1}$
31 ABLIKIM 06R reports $(2.0 \pm 0.5 \pm 0.5) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow a_0(980)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(f_2(1270)\eta)/\Gamma_{\text{total}}$ Γ_{23}/Γ

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.8±0.8±0.1	53	32 ABLIKIM	06R	BES2 $\psi(2S) \rightarrow \gamma \chi_{c1}$
32 ABLIKIM 06R reports $(3.0 \pm 0.7 \pm 0.5) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow f_2(1270)\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(\pi^+\pi^-\eta')/\Gamma_{\text{total}}$ Γ_{24}/Γ

<u>VALUE</u> (units 10^{-3})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.3±0.5±0.1	33 ATHAR	07	CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

³³ ATHAR 07 reports $(2.4 \pm 0.4 \pm 0.3) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^+ \pi^- \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^0 f_0(980) \rightarrow \pi^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<6 × 10⁻⁶	90	34 ABLIKIM	11D BES3	$\psi(2S) \rightarrow \gamma \pi^0 \pi^+ \pi^-$
³⁴ ABLIKIM 11D reports $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^0 f_0(980) \rightarrow \pi^0 \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] < 6.0 \times 10^{-7}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.3 \times 10^{-2}$.				

$\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE (units 10 ⁻⁴)	DOCUMENT ID	TECN	COMMENT
32±21	35 TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c1}$

³⁵ Estimated using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.087$. The errors do not contain the uncertainty in the $\psi(2S)$ decay.

$\Gamma(K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
1.5±0.4±0.1	28.4 ± 5.5	36,37 ABLIKIM	04H BES	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$
³⁶ ABLIKIM 04H reports $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (1.40 \pm 0.27 \pm 0.22) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

³⁷ Assumes $B(K^*(892)^0 \rightarrow K^- \pi^+) = 2/3$.

$\Gamma(K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (units 10 ⁻⁴)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<5	90	3.2 ± 2.4	38 ABLIKIM	050 BES2	$\psi(2S) \rightarrow \chi_{c1} \gamma$
³⁸ ABLIKIM 050 reports $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] < 4.2 \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.3 \times 10^{-2}$.					

$\Gamma(K^+ K^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE (units 10 ⁻³)	DOCUMENT ID
0.56±0.12 OUR FIT	

$\Gamma(K^+ K^- \phi)/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
0.43±0.16±0.02	17	39 ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$
³⁹ ABLIKIM 06T reports $(0.46 \pm 0.16 \pm 0.06) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(\omega\omega)/\Gamma_{\text{total}}$	Γ_{31}/Γ			
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.0±0.7±0.2	597	40	ABLIKIM	11K BES3 $\psi(2S) \rightarrow \gamma$ hadrons

40 ABLIKIM 11K reports $(6.0 \pm 0.3 \pm 0.7) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\omega\phi)/\Gamma_{\text{total}}$	Γ_{32}/Γ			
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.22±0.06±0.01	15	41	ABLIKIM	11K BES3 $\psi(2S) \rightarrow \gamma$ hadrons

41 ABLIKIM 11K reports $(0.22 \pm 0.06 \pm 0.02) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi\phi)/\Gamma_{\text{total}}$	Γ_{33}/Γ			
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.4±0.5±0.2	366	42	ABLIKIM	11K BES3 $\psi(2S) \rightarrow \gamma$ hadrons

42 ABLIKIM 11K reports $(4.4 \pm 0.3 \pm 0.5) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \phi\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$	Γ_{34}/Γ
<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>
0.73±0.04 OUR FIT	

$\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$	Γ_{35}/Γ			
<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.163±0.020 OUR AVERAGE				
0.171±0.020±0.007	43	ONYISI	10	CLE3 $\psi(2S) \rightarrow \gamma p\bar{p}X$
0.117±0.049±0.005	44	ATHAR	07	CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

43 ONYISI 10 reports $(1.75 \pm 0.16 \pm 0.13 \pm 0.11) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

44 ATHAR 07 reports $(1.2 \pm 0.5 \pm 0.1) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$ **Γ_{36}/Γ**

<i>VALUE</i> (units 10^{-3})	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
$0.153 \pm 0.026 \pm 0.006$		45 ONYISI	10 CLE3	$\psi(2S) \rightarrow \gamma p\bar{p}X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.16	90	46 ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$
45	ONYISI 10 reports $(1.56 \pm 0.22 \pm 0.14 \pm 0.10) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			
46	ATHAR 07 reports $< 0.16 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.3 \times 10^{-2}$.			

$\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$ **Γ_{37}/Γ**

<i>VALUE</i> (units 10^{-3})	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
$0.223 \pm 0.032 \pm 0.009$	47 ONYISI	10 CLE3	$\psi(2S) \rightarrow \gamma p\bar{p}X$
47 ONYISI 10 reports $(2.28 \pm 0.28 \pm 0.16 \pm 0.14) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			

$\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$ **Γ_{38}/Γ**

<i>VALUE</i> (units 10^{-5})	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
<1.8	90	48 ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p\bar{p}K^+ K^-$
48 ABLIKIM 11F reports $< 1.82 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.3 \times 10^{-2}$.				

$\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_{39}/Γ**

<i>VALUE</i> (units 10^{-3})	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
0.50 ± 0.19 OUR EVALUATION			Treating systematic error as correlated.
0.50 ± 0.19 OUR AVERAGE			
0.46 $\pm 0.12 \pm 0.15$			
0.46 $\pm 0.12 \pm 0.15$	49 BAI	99B BES	$\psi(2S) \rightarrow \gamma\chi_{c1}$
1.08 $\pm 0.77 \pm 0.05$	49 TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma\chi_{c1}$
49 Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (8.8 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$.			

$\Gamma(p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}$ **Γ_{40}/Γ**

<i>VALUE</i> (%)	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
<0.05	90	50 HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$
50 HE 08B reports $< 0.05 \%$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.3 \times 10^{-2}$.				

$\Gamma(p\bar{p}K^+K^- \text{(non-resonant)})/\Gamma_{\text{total}}$ Γ_{41}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.34 \pm 0.23 \pm 0.05$	82 ± 9	51 ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$
⁵¹ ABLIKIM 11F reports $(1.35 \pm 0.15 \pm 0.19) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}K^+K^- \text{(non-resonant)})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(p\bar{p}K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{42}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<4.5	90	52 ABLIKIM	06D BES2	$\psi(2S) \rightarrow \gamma\chi_{c1}$
⁵² Using $B(\psi(2S) \rightarrow \chi_{c1}\gamma) (9.1 \pm 0.6)\%$.				

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{43}/Γ

VALUE (units 10^{-4})	DOCUMENT ID
$1.18 \pm 0.19 \text{ OUR FIT}$	

$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{44}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1.5	90	53 ABLIKIM	06D BES2	$\psi(2S) \rightarrow \gamma\chi_{c1}$
⁵³ Using $B(\psi(2S) \rightarrow \chi_{c1}\gamma) (9.1 \pm 0.6)\%$.				

$\Gamma(K^+\bar{p}\Lambda)/\Gamma_{\text{total}}$ Γ_{45}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.3 \pm 0.4 \text{ OUR AVERAGE}$		Error includes scale factor of 1.1.		
^{4.5 $\pm 0.4 \pm 0.2$} ABLIKIM 3k 54,55 $\psi(2S) \rightarrow \gamma\Lambda\bar{p}K^+$				
^{3.2 $\pm 0.9 \pm 0.1$} ATHAR 56 CLEO $\psi(2S) \rightarrow \gamma h^+h^-h^0$				
⁵⁴ ABLIKIM 13D reports $(4.5 \pm 0.2 \pm 0.4) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+\bar{p}\Lambda)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
⁵⁵ Using $B(\Lambda \rightarrow p\pi^-) = 63.9\%$.				
⁵⁶ ATHAR 07 reports $(3.3 \pm 0.9 \pm 0.4) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+\bar{p}\Lambda)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(K^+p\Lambda(1520)+\text{c.c.})/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.8 \pm 0.5 \pm 0.1$	48 ± 10	57 ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$
⁵⁷ ABLIKIM 11F reports $(1.81 \pm 0.38 \pm 0.28) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+p\Lambda(1520)+\text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(\Lambda(1520)\bar{\Lambda}(1520))/\Gamma_{\text{total}}$ Γ_{47}/Γ

<i>VALUE</i> (units 10^{-4})	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
<1.0	90	58 ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p\bar{p} K^+ K^-$
58 ABLIKIM 11F reports $< 1.00 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Lambda(1520)\bar{\Lambda}(1520))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.3 \times 10^{-2}$.				

 $\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{48}/Γ

<i>VALUE</i> (units 10^{-4})	<i>CL%</i>	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
<0.4	90	3.8 ± 2.5	59 NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma\Sigma^0\bar{\Sigma}^0$
59 NAIK 08 reports $< 0.44 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.3 \times 10^{-2}$.					

 $\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$ Γ_{49}/Γ

<i>VALUE</i> (units 10^{-4})	<i>CL%</i>	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
<0.6	90	4.3 ± 2.3	60 NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma\Sigma^+\bar{\Sigma}^-$
60 NAIK 08 reports $< 0.65 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.3 \times 10^{-2}$.					

 $\Gamma(\Xi^0\bar{\Xi}^0)/\Gamma_{\text{total}}$ Γ_{50}/Γ

<i>VALUE</i> (units 10^{-4})	<i>CL%</i>	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
<0.6	90	1.7 ± 2.4	61 NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma\Xi^0\bar{\Xi}^0$
61 NAIK 08 reports $< 0.60 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Xi^0\bar{\Xi}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.3 \times 10^{-2}$.					

 $\Gamma(\Xi^-\bar{\Xi}^+)/\Gamma_{\text{total}}$ Γ_{51}/Γ

<i>VALUE</i> (units 10^{-4})	<i>CL%</i>	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
0.84 ± 0.22 ± 0.03		16.4 ± 4.3	62 NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma\Xi^-\bar{\Xi}^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 3.4 90 63 ABLIKIM 06D BES2 $\psi(2S) \rightarrow \gamma\chi_{c1}$					
62 NAIK 08 reports $(0.86 \pm 0.22 \pm 0.08) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Xi^-\bar{\Xi}+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
63 Using $B(\psi(2S) \rightarrow \chi_{c1}\gamma)$ $(9.1 \pm 0.6)\%$.					

$\Gamma(\pi^+\pi^-) + \Gamma(K^+K^-)]/\Gamma_{\text{total}}$ Γ_{52}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<21		64 FELDMAN	77 MRK1	$\psi(2S) \rightarrow \gamma\chi_{c1}$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<38	90	64 BRANDELIK	79B DASP	$\psi(2S) \rightarrow \gamma\chi_{c1}$
64 Estimated using $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.087$. The errors do not contain the uncertainty in the $\psi(2S)$ decay.				

$\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{53}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.6	90	65 ABLIKIM	050 BES2	$\psi(2S) \rightarrow \chi_{c1}\gamma$
65 ABLIKIM 050 reports $[\Gamma(\chi_{c1}(1P) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ $< 0.6 \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.3 \times 10^{-2}$.				

RADIATIVE DECAYS

$\Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$ Γ_{54}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.348 ± 0.015 OUR FIT			
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.379 ± 0.008 ± 0.021	66 ADAM	05A CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c1}$
66 Uses $B(\psi(2S) \rightarrow \gamma\chi_{c1} \rightarrow \gamma\gamma J/\psi)$ from ADAM 05A and $B(\psi(2S) \rightarrow \gamma\chi_{c1})$ from ATHAR 04.			

$\Gamma(\gamma\rho^0)/\Gamma_{\text{total}}$ Γ_{55}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
227 ± 19 OUR AVERAGE				
226 ± 23 ± 9	432 ± 25	67 ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\rho^0$
228 ± 25 ± 9	186 ± 15	68 BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\rho^0$
67 ABLIKIM 11E reports $(228 \pm 13 \pm 22) \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
68 BENNETT 08A reports $(243 \pm 19 \pm 22) \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(\gamma\omega)/\Gamma_{\text{total}}$ Γ_{56}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
71 ± 9 OUR AVERAGE				
69 ± 9 ± 3	136 ± 14	69 ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\omega$
78 ± 18 ± 3	39 ± 7	70 BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\omega$

⁶⁹ ABLIKIM 11E reports $(69.7 \pm 7.2 \pm 6.6) \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁷⁰ BENNETT 08A reports $(83 \pm 15 \pm 12) \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\gamma\phi)/\Gamma_{\text{total}}$

Γ_{57}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
26±6±1		43 ± 9	71 ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\phi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<24 90 5.2 ± 3.1 72 BENNETT 08A CLEO $\psi(2S) \rightarrow \gamma\gamma\phi$

⁷¹ ABLIKIM 11E reports $(25.8 \pm 5.2 \pm 2.3) \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.3 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁷² BENNETT 08A reports $< 26 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.3 \times 10^{-2}$.

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

Γ_{58}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				

< 3.5 90 ECKLUND 08A CLEO $\psi(2S) \rightarrow \gamma\chi_{c1} \rightarrow 3\gamma$
<150 90 73 YAMADA 77 DASP $e^+ e^- \rightarrow 3\gamma$

⁷³ Estimated using $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.087$. The errors do not contain the uncertainty in the $\psi(2S)$ decay.

$\chi_{c1}(1P)$ CROSS-PARTICLE BRANCHING RATIOS

$$\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \quad \Gamma_{34}/\Gamma \times \Gamma_{119}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
1.98±0.16 OUR FIT			

1.1 ±1.0 74 BAI 98I BES $\psi(2S) \rightarrow \gamma\chi_{c1} \rightarrow \gamma\bar{p}p$

⁷⁴ Calculated by us. The value for $B(\chi_{c1} \rightarrow p\bar{p})$ reported in BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (8.7 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma_{\text{total}}}{\Gamma_{43}/\Gamma \times \Gamma_{119}^{\psi(2S)}/\Gamma^{\psi(2S)}}$$

<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
10.9 ± 1.7 OUR FIT				
$10.5 \pm 1.6 \pm 0.6$	46 ± 7	⁷⁵ NAIK	08	CLEO $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$
⁷⁵ Calculated by us. NAIK 08 reports $B(\chi_{c1} \rightarrow \Lambda\bar{\Lambda}) = (11.6 \pm 1.8 \pm 0.7 \pm 0.7) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (9.07 \pm 0.11 \pm 0.54)\%$.				

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{43}/\Gamma \times \Gamma_{119}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.2 ± 0.5 OUR FIT				

$7.1^{+2.8}_{-2.4} \pm 1.3$	$9.0^{+3.5}_{-3.1}$	⁷⁶ BAI	03E	BES $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$
---	---------------------	-------------------	-----	---

⁷⁶ BAI 03E reports [$B(\chi_{c1} \rightarrow \Lambda\bar{\Lambda}) B(\psi(2S) \rightarrow \gamma\chi_{c1}) / B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-)$] \times $[B^2(\Lambda \rightarrow \pi^- p) / B(J/\psi \rightarrow p\bar{p})] = (1.33^{+0.52}_{-0.46} \pm 0.25)\%$. We calculate from this measurement the presented value using $B(\Lambda \rightarrow \pi^- p) = (63.9 \pm 0.5)\%$ and $B(J/\psi \rightarrow p\bar{p}) = (2.17 \pm 0.07) \times 10^{-3}$.

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma_{\text{total}}}{\Gamma_{54}/\Gamma \times \Gamma_{119}^{\psi(2S)}/\Gamma^{\psi(2S)}}$$

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.23 ± 0.07 OUR FIT				

2.93 ± 0.15 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.

$3.377 \pm 0.009 \pm 0.183$	142k	ABLIKIM	120	BES3 $\psi(2S) \rightarrow \gamma\chi_{c1}$
$2.81 \pm 0.05 \pm 0.23$	13k	BAI	04I	BES2 $\psi(2S) \rightarrow J/\psi\gamma\gamma$
$2.56 \pm 0.12 \pm 0.20$		GAISER	86	CBAL $\psi(2S) \rightarrow \gamma X$
2.78 ± 0.30		77 OREGLIA	82	CBAL $\psi(2S) \rightarrow \gamma\chi_{c1}$
2.2 ± 0.5		78 BRANDELIK	79B	DASP $\psi(2S) \rightarrow \gamma\chi_{c1}$
2.9 ± 0.5		78 BARTEL	78B	CNTR $\psi(2S) \rightarrow \gamma\chi_{c1}$
5.0 ± 1.5		79 BIDDICK	77	CNTR $e^+e^- \rightarrow \gamma X$
2.8 ± 0.9		77 WHITAKER	76	MRK1 e^+e^-

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.56 \pm 0.03 \pm 0.12$	24.9k	⁸⁰ MENDEZ	08	CLEO $\psi(2S) \rightarrow \gamma\chi_{c1}$
$3.44 \pm 0.06 \pm 0.13$	3.7k	⁸¹ ADAM	05A	CLEO Repl. by MENDEZ 08


⁷⁷ Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.

⁷⁸ Recalculated by us using $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$.

⁷⁹ Assumes isotropic gamma distribution.

⁸⁰ Not independent from other measurements of MENDEZ 08.

⁸¹ Not independent from other values reported by ADAM 05A.



$$\Gamma(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{\text{total}} (\text{units } 10^{-2})$$

$$\begin{aligned} & \Gamma(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma(\psi(2S) \rightarrow \\ & J/\psi(1S) \text{anything}) \quad \Gamma_{54}/\Gamma \times \Gamma_{119}^{\psi(2S)} / \Gamma_9^{\psi(2S)} \\ & \Gamma_{54}/\Gamma \times \Gamma_{119}^{\psi(2S)} / \Gamma_9^{\psi(2S)} = \Gamma_{54}/\Gamma \times \Gamma_{119}^{\psi(2S)} / (\Gamma_{11}^{\psi(2S)} + \Gamma_{12}^{\psi(2S)} + \Gamma_{13}^{\psi(2S)} + \\ & 0.348\Gamma_{119}^{\psi(2S)} + 0.198\Gamma_{120}^{\psi(2S)}) \end{aligned}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
5.36±0.12 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.70 \pm 0.04 \pm 0.15$	24.9k	82 MENDEZ	08 CLEO	$\psi(2S) \rightarrow \gamma \chi_{c1}$
$5.77 \pm 0.10 \pm 0.12$	3.7k	ADAM	05A CLEO	Repl. by MENDEZ 08

⁸² Not independent from other measurements of MENDEZ 08.

$$\begin{aligned} & \Gamma(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma(\psi(2S) \rightarrow \\ & J/\psi(1S) \pi^+ \pi^-) \quad \Gamma_{54}/\Gamma \times \Gamma_{119}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)} \end{aligned}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
9.49±0.21 OUR FIT				

10.15±0.28 OUR AVERAGE

$10.17 \pm 0.07 \pm 0.27$	24.9k	MENDEZ	08 CLEO	$\psi(2S) \rightarrow \gamma \chi_{c1}$
$12.6 \pm 0.3 \pm 3.8$	3k	83 ABLIKIM	04B BES	$\psi(2S) \rightarrow J/\psi X$
8.5 ± 2.1		84 HIMEL	80 MRK2	$\psi(2S) \rightarrow \gamma \chi_{c1}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$10.24 \pm 0.17 \pm 0.23$	3.7k	85 ADAM	05A CLEO	Repl. by MENDEZ 08
---------------------------	------	---------	----------	--------------------

⁸³ From a fit to the J/ψ recoil mass spectra.

⁸⁴ The value for $B(\psi(2S) \rightarrow \gamma\chi_{c1}) \times B(\chi_{c1} \rightarrow \gamma J/\psi(1S))$ quoted in HIMEI 80 is derived using $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (33 \pm 3)\%$ and $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.138 \pm 0.018$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.

⁸⁵ Not independent from other values reported by ADAM 05A.

$$\Gamma(\chi_{c1}(1P) \rightarrow K^0 K^+ \pi^- + c.c.) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) / \Gamma_{\text{total}}$$

$$\Gamma_{15}/\Gamma \times \Gamma_{119}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

6.8±0.5 OUR FIT

7.2±0.6 OUR AVERAGE

$7.3 \pm 0.5 \pm 0.5$ ⁸⁶ ATHAR 07 CLEO $\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^-$

$7.0 \pm 0.5 \pm 0.9$ ⁸⁷ ABLIKIM 06R BES2 $\psi(2S) \rightarrow \gamma\chi_{c1}$

⁸⁶ Calculated by us. The value of $B(\chi_{c1} \rightarrow K^0 K^+ \pi^- + c.c.)$ reported by ATHAR 07 was derived using $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54)\%$.

⁸⁷ Calculated by us. ABLIKIM 06R reports $B(\chi_{c1} \rightarrow K_S^0 K^+ \pi^-) = (4.0 \pm 0.3 \pm 0.5) \times 10^{-3}$. We use $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (8.7 \pm 0.4) \times 10^{-2}$.

$$\Gamma(\chi_{c1}(1P) \rightarrow K^0 K^+ \pi^- + c.c.) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) / \Gamma_{\text{total}}$$

$$\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) / \Gamma_{15}/\Gamma \times \Gamma_{119}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

19.9±1.6 OUR FIT

13.2±2.4±3.2 ⁸⁸ BAI 99B BES $\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^-$

⁸⁸ Calculated by us. The value of $B(\chi_{c1} \rightarrow K_S^0 K^+ \pi^-)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) / \Gamma_{\text{total}}$$

$$\Gamma_{29}/\Gamma \times \Gamma_{119}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.52±0.11 OUR FIT

0.61±0.11±0.08 54 ⁸⁹ ABLIKIM 06T BES2 $\psi(2S) \rightarrow \gamma K^+ K^+ K^- K^-$

⁸⁹ Calculated by us. The value of $B(\chi_{c1} \rightarrow 2K^+ 2K^-)$ reported by ABLIKIM 06T was derived using $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.8)\%$.

$$\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) / \Gamma_{\text{total}}$$

$$\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) / \Gamma_{29}/\Gamma \times \Gamma_{119}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

1.53±0.31 OUR FIT

1.13±0.40±0.29 ⁹⁰ BAI 99B BES $\psi(2S) \rightarrow \gamma K^+ K^+ K^- K^-$

⁹⁰ Calculated by us. The value of $B(\chi_{c1} \rightarrow 2K^+ 2K^-)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma_{\text{total}}}{\Gamma_{34}/\Gamma \times \Gamma_{119}^{\psi(2S)}/\Gamma^{\psi(2S)}}$$

<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.7 ± 0.5 OUR FIT				
7.5 ± 1.4 OUR AVERAGE	Error includes scale factor of 2.0.			
$8.2 \pm 0.7 \pm 0.4$	141 ± 13	91 NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$
$4.8^{+1.4}_{-1.3} \pm 0.6$	$18.2^{+5.5}_{-4.9}$	BAI	04F BES	$\psi(2S) \rightarrow \gamma\chi_{c1}(1P) \rightarrow \gamma\bar{p}p$


⁹¹ Calculated by us. NAIK 08 reports $B(\chi_{c1} \rightarrow p\bar{p}) = (9.0 \pm 0.8 \pm 0.4 \pm 0.5) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (9.07 \pm 0.11 \pm 0.54)\%$.

MULTIPOLE AMPLITUDES IN $\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)$

$$a_2 = M2/\sqrt{E1^2 + M2^2} \text{ Magnetic quadrupole fractional transition amplitude}$$

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-5.4^{+1.2}_{-1.5}$ OUR AVERAGE Error includes scale factor of 2.4. See the ideogram below.				
$-6.26 \pm 0.63 \pm 0.24$	39k	ARTUSO 09	CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
$0.2 \pm 3.2 \pm 0.4$	2090	AMBROGIANI 02	E835	$p\bar{p} \rightarrow \chi_{c1} \rightarrow J/\psi\gamma$
$-0.2^{+0.8}_{-2.0}$	921	OREGLIA 82	CBAL	$\psi(2S) \rightarrow \chi_{c1}\gamma \rightarrow J/\psi\gamma\gamma$

WEIGHTED AVERAGE
 $-5.4+1.2-1.5$ (Error scaled by 2.4)



$$a_2 = M2/\sqrt{E1^2 + M2^2} \text{ (units } 10^{-2})$$

MULTIPOLE AMPLITUDES IN $\psi(2S) \rightarrow \gamma\chi_{c1}(1S)$ RADIATIVE DECAY

$b_2 = M2/\sqrt{E1^2 + M2^2}$ Magnetic quadrupole fractional transition amplitude

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.9 \pm 0.8 OUR AVERAGE				
$2.76 \pm 0.73 \pm 0.23$	39k	ARTUSO	09	CLEO $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
$7.7 \begin{array}{l} +5.0 \\ -4.5 \end{array}$	921	OREGLIA	82	CBAL $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

MULTIPOLE AMPLITUDE RATIOS IN RADIATIVE DECAYS

$\psi(2S) \rightarrow \gamma\chi_{c1}(1S)$ and $\chi_{c1} \rightarrow \gamma J/\psi(1S)$

a_2/b_2 Magnetic quadrupole transition amplitude ratio

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$-2.27 \begin{array}{l} +0.57 \\ -0.99 \end{array}$	39k	92	ARTUSO	09 CLEO $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

⁹² Statistical and systematic errors combined. Not independent of $a_2(\chi_{c1})$ and $b_2(\chi_{c1})$ values from ARTUSO 09.

$\chi_{c1}(1P)$ REFERENCES

ABLIKIM	13D	PR D87 012007	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12O	PRL 109 172002	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	11A	PR D83 012006	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	11D	PR D83 032003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	11E	PR D83 112005	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	11F	PR D83 112009	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	11K	PRL 107 092001	M. Ablikim <i>et al.</i>	(BES III Collab.)
ONYISI	10	PR D82 011103	P.U.E. Onyisi <i>et al.</i>	(CLEO Collab.)
ARTUSO	09	PR D80 112003	M. Artuso <i>et al.</i>	(CLEO Collab.)
BENNETT	08A	PRL 101 151801	J.V. Bennett <i>et al.</i>	(CLEO Collab.)
ECKLUND	08A	PR D78 091501	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)
HE	08B	PR D78 092004	Q. He <i>et al.</i>	(CLEO Collab.)
MENDEZ	08	PR D78 011102	H. Mendez <i>et al.</i>	(CLEO Collab.)
NAIK	08	PR D78 031101	P. Naik <i>et al.</i>	(CLEO Collab.)
ATHAR	07	PR D75 032002	S.B. Athar <i>et al.</i>	(CLEO Collab.)
ABLIKIM	06D	PR D73 052006	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06T	PL B642 197	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05G	PR D71 092002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAM	05A	PRL 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ANDREOTTI	05A	NP B717 34	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)
ABLIKIM	04B	PR D70 012003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04H	PR D70 092003	M. Ablikim <i>et al.</i>	(BES Collab.)
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)
BAI	04F	PR D69 092001	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04I	PR D70 012006	J.Z. Bai <i>et al.</i>	(BES Collab.)
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
BAI	03E	PR D67 112001	J.Z. Bai <i>et al.</i>	(BES Collab.)
AMBROGIANI	02	PR D65 052002	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
BAI	99B	PR D60 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98I	PRL 81 3091	J.Z. Bai <i>et al.</i>	(BES Collab.)
ARMSTRONG	92	NP B373 35	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
Also		PRL 68 1468	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
BAGLIN	86B	PL B172 455	C. Baglin	(LAPP, CERN, GENO, LYON, OSLO+)
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)
OREGLIA	82	PR D25 2259	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)
Also		Private Comm.	M.J. Oreglia	(EFI)

HIMEL	80	PRL 44 920	T. Himel <i>et al.</i>	(LBL, SLAC)
Also		Private Comm.	G. Trilling	(LBL, UCB)
BRANDELIK	79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)
TANENBAUM	78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)
Also		Private Comm.	G. Trilling	(LBL, UCB)
BIDDICK	77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)
YAMADA	77	Hamburg Conf. 69	S. Yamada	(DASP Collab.)
WHITAKER	76	PRL 37 1596	J.S. Whitaker <i>et al.</i>	(SLAC, LBL)
TANENBAUM	75	PRL 35 1323	W.M. Tanenbaum <i>et al.</i>	(LBL, SLAC)
