

**$\eta_b(2S)$**

$I^G(J^{PC}) = 0^+(0^-+)$

## OMMITTED FROM SUMMARY TABLE

Quantum numbers shown are quark-model predictions.

### **$\eta_b(2S)$ MASS**

| VALUE (MeV)  | EVTS   | DOCUMENT ID          | TECN | COMMENT   |
|--|--------|----------------------|------|---|
| <b>9999.0±3.5<sup>+2.8</sup><sub>-1.9</sub></b>  | 26k    | <sup>1</sup> MIZUK   | 12   | BELL $e^+ e^- \rightarrow \gamma \pi^+ \pi^- +$ hadrons |
| • • • We do not use the following data for averages, fits, limits, etc. • • •  |        |                      |      |   |
| 9974.6±2.3±2.1   | 11 ± 4 | <sup>2,3</sup> DOBBS | 12   | $\Upsilon(2S) \rightarrow \gamma$ hadrons               |
| <sup>1</sup> Assuming $\Gamma_{\eta_b}(2S) = 4.9$ MeV. Not independent of the corresponding mass difference measurement. |        |                      |      |   |
| <sup>2</sup> Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.                             |        |                      |      |   |
| <sup>3</sup> Assuming $\Gamma_{\eta_b}(2S) = 5$ MeV. Not independent of the corresponding mass difference measurement.   |        |                      |      |   |

### **$m_{\Upsilon(2S)} - m_{\eta_b(2S)}$**

| VALUE (MeV)   | EVTS   | DOCUMENT ID          | TECN | COMMENT   |
|---|--------|----------------------|------|---|
| <b>24.3±3.5<sup>+2.8</sup><sub>-1.9</sub></b>   | 26k    | <sup>4</sup> MIZUK   | 12   | BELL $e^+ e^- \rightarrow \gamma \pi^+ \pi^- +$ hadrons |
| • • • We do not use the following data for averages, fits, limits, etc. • • •                                 |        |                      |      |   |
| 48.7±2.3±2.1  | 11 ± 4 | <sup>5,6</sup> DOBBS | 12   | $\Upsilon(2S) \rightarrow \gamma$ hadrons               |
| <sup>4</sup> Assuming $\Gamma_{\eta_b}(2S) = 4.9$ MeV. Not independent of the corresponding mass measurement. |        |                      |      |   |
| <sup>5</sup> Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.                  |        |                      |      |   |
| <sup>6</sup> Assuming $\Gamma_{\eta_b}(2S) = 5$ MeV. Not independent of the corresponding mass measurement.   |        |                      |      |   |

### **$\eta_b(2S)$ WIDTH**

| VALUE (MeV) | CL% | DOCUMENT ID | TECN | COMMENT   |
|-------------|-----|-------------|------|---|
| <24         | 90  | MIZUK       | 12   | BELL $e^+ e^- \rightarrow \gamma \pi^+ \pi^-$ hadrons |

### **$\eta_b(2S)$ DECAY MODES**

| Mode               | Fraction ( $\Gamma_i/\Gamma$ ) |
|--------------------|--------------------------------|
| $\Gamma_1$ hadrons | seen                           |

## $\eta_b(2S)$ BRANCHING RATIOS

| $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$  | $\Gamma_1/\Gamma$                                |
|---|--|
| <i>VALUE</i>  | <i>EVTS</i>                                      |
| <b>seen</b>   | 26k  |
| MIZUK   | MIZUK  |
| 12  | 12   |
| BELL  | $e^+ e^- \rightarrow \gamma \pi^+ \pi^-$ hadrons |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ |  |
| seen  | 7 DOBBS  |
|   | 12   |
|   | $\gamma(2S) \rightarrow \gamma$ hadrons          |
| 7 Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.                                     |  |

## $\eta_b(2S)$ REFERENCES

|       |    |                |                        |                 |
|-------|----|----------------|------------------------|-----------------|
| DOBBS | 12 | PRL 109 082001 | S. Dobbs <i>et al.</i> |                 |
| MIZUK | 12 | PRL 109 232002 | R. Mizuk <i>et al.</i> | (BELLE Collab.) |