

$\Upsilon(2S)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

$\Upsilon(2S)$ MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
10023.26 ± 0.31 OUR AVERAGE			
10023.5 ± 0.5	¹ ARTAMONOV 00	MD1	$e^+ e^- \rightarrow$ hadrons
10023.1 ± 0.4	BARBER 84	REDE	$e^+ e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
10023.6 ± 0.5	^{2,3} BARU	86B REDE	$e^+ e^- \rightarrow$ hadrons
¹ Reanalysis of BARU 86B using new electron mass (COHEN 87).			
² Reanalysis of ARTAMONOV 84.			
³ Superseded by ARTAMONOV 00.			

$m\Upsilon(3S) - m\Upsilon(2S)$

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
331.50 ± 0.02 ± 0.13	LEES	11C BABR	$e^+ e^- \rightarrow \pi^+ \pi^- X$

$\Upsilon(2S)$ WIDTH

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>
31.98 ± 2.63 OUR EVALUATION	See the Note on "Width Determinations of the Υ States"

$\Upsilon(2S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $\Upsilon(1S)\pi^+\pi^-$	(17.85 ± 0.26) %	
Γ_2 $\Upsilon(1S)\pi^0\pi^0$	(8.6 ± 0.4) %	
Γ_3 $\tau^+\tau^-$	(2.00 ± 0.21) %	
Γ_4 $\mu^+\mu^-$	(1.93 ± 0.17) %	S=2.2
Γ_5 e^+e^-	(1.91 ± 0.16) %	
Γ_6 $\Upsilon(1S)\pi^0$	< 4 × 10 ⁻⁵	CL=90%
Γ_7 $\Upsilon(1S)\eta$	(2.9 ± 0.4) × 10 ⁻⁴	S=2.0
Γ_8 $J/\psi(1S)$ anything	< 6 × 10 ⁻³	CL=90%
Γ_9 \bar{d} anything	(3.4 ± 0.6) × 10 ⁻⁵	
Γ_{10} hadrons	(94 ± 11) %	
Γ_{11} ggg	(58.8 ± 1.2) %	
Γ_{12} γgg	(8.8 ± 1.1) %	
Γ_{13} Sum of 100 exclusive modes	(2.90 ± 0.30) × 10 ⁻³	

Radiative decays

Γ_{14}	$\gamma \chi_{b1}(1P)$	$(6.9 \pm 0.4) \%$			
Γ_{15}	$\gamma \chi_{b2}(1P)$	$(7.15 \pm 0.35) \%$			
Γ_{16}	$\gamma \chi_{b0}(1P)$	$(3.8 \pm 0.4) \%$			
Γ_{17}	$\gamma f_0(1710)$	< 5.9	$\times 10^{-4}$	CL=90%	
Γ_{18}	$\gamma f_2'(1525)$	< 5.3	$\times 10^{-4}$	CL=90%	
Γ_{19}	$\gamma f_2(1270)$	< 2.41	$\times 10^{-4}$	CL=90%	
Γ_{20}	$\gamma f_J(2220)$				
Γ_{21}	$\gamma \eta_c(1S)$	< 2.7	$\times 10^{-5}$	CL=90%	
Γ_{22}	$\gamma \chi_{c0}$	< 1.0	$\times 10^{-4}$	CL=90%	
Γ_{23}	$\gamma \chi_{c1}$	< 3.6	$\times 10^{-6}$	CL=90%	
Γ_{24}	$\gamma \chi_{c2}$	< 1.5	$\times 10^{-5}$	CL=90%	
Γ_{25}	$\gamma X(3872) \rightarrow \pi^+ \pi^- J/\psi$	< 8	$\times 10^{-7}$	CL=90%	
Γ_{26}	$\gamma X(3872) \rightarrow \pi^+ \pi^- \pi^0 J/\psi$	< 2.4	$\times 10^{-6}$	CL=90%	
Γ_{27}	$\gamma X(3915) \rightarrow \omega J/\psi$	< 2.8	$\times 10^{-6}$	CL=90%	
Γ_{28}	$\gamma X(4140) \rightarrow \phi J/\psi$	< 1.2	$\times 10^{-6}$	CL=90%	
Γ_{29}	$\gamma X(4350) \rightarrow \phi J/\psi$	< 1.3	$\times 10^{-6}$	CL=90%	
Γ_{30}	$\gamma \eta_b(1S)$	(3.9 ± 1.5)	$\times 10^{-4}$		
Γ_{31}	$\gamma \eta_b(2S)$				
Γ_{32}	$\gamma X \rightarrow \gamma + \geq 4$ prongs	[a]	< 1.95	$\times 10^{-4}$	CL=95%
Γ_{33}	$\gamma A^0 \rightarrow \gamma$ hadrons		< 8	$\times 10^{-5}$	CL=90%
Γ_{34}	$\gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-$		< 8.3	$\times 10^{-6}$	CL=90%

Lepton Family number (*LF*) violating modes

Γ_{35}	$e^\pm \tau^\mp$	<i>LF</i>	< 3.2	$\times 10^{-6}$	CL=90%
Γ_{36}	$\mu^\pm \tau^\mp$	<i>LF</i>	< 3.3	$\times 10^{-6}$	CL=90%

[a] $1.5 \text{ GeV} < m_X < 5.0 \text{ GeV}$

CONSTRAINED FIT INFORMATION

An overall fit to 3 branching ratios uses 13 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 11.8$ for 11 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$x_7 \left| \begin{array}{c} \underline{\quad} \\ 2 \\ x_1 \end{array} \right.$$

$\Upsilon(2S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

$\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_4\Gamma_5/\Gamma$
<u>VALUE (eV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
6.5±1.5±1.0	KOBEL	92	CBAL	$e^+e^- \rightarrow \mu^+\mu^-$

$\Gamma(\Upsilon(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_1\Gamma_5/\Gamma$
<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
105.4±1.0±4.2	11.8K	¹ AUBERT	08BP BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$
¹ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.				

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_{10}\Gamma_5/\Gamma$
<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.577±0.009 OUR AVERAGE				
0.581±0.004±0.009	¹ ROSNER	06	CLEO	10.0 $e^+e^- \rightarrow \text{hadrons}$
0.552±0.031±0.017	¹ BARU	96	MD1	$e^+e^- \rightarrow \text{hadrons}$
0.54 ±0.04 ±0.02	¹ JAKUBOWSKI	88	CBAL	$e^+e^- \rightarrow \text{hadrons}$
0.58 ±0.03 ±0.04	² GILES	84B	CLEO	$e^+e^- \rightarrow \text{hadrons}$
0.60 ±0.12 ±0.07	² ALBRECHT	82	DASP	$e^+e^- \rightarrow \text{hadrons}$
0.54 ±0.07 ^{+0.09} _{-0.05}	² NICZYPORUK	81C	LENA	$e^+e^- \rightarrow \text{hadrons}$
0.41 ±0.18	² BOCK	80	CNTR	$e^+e^- \rightarrow \text{hadrons}$

¹ Radiative corrections evaluated following KURAEV 85.

² Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

$\Upsilon(2S)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$	Γ_5
<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>
0.612±0.011 OUR EVALUATION	

$\Upsilon(2S)$ BRANCHING RATIOS

$\Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$				Γ_1/Γ
Abbreviation MM in the <i>COMMENT</i> field below stands for missing mass.				
<u>VALUE (units 10⁻²)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
17.85±0.26 OUR FIT				
17.92±0.26 OUR AVERAGE				
16.8 ±1.1 ±1.3	906k	¹ LEES	11C BABR	$e^+e^- \rightarrow \pi^+\pi^-X$
17.80±0.05±0.37	170k	² LEES	11L BABR	$\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$
18.02±0.02±0.61	851k	³ BHARI	09 CLEO	$e^+e^- \rightarrow \pi^+\pi^- \text{MM}$
17.22±0.17±0.75	11.8K	⁴ AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$
19.2 ±0.2 ±1.0	52.6k	⁵ ALEXANDER	98 CLE2	$\pi^+\pi^-\ell^+\ell^-$, $\pi^+\pi^- \text{MM}$
18.1 ±0.5 ±1.0	11.6k	ALBRECHT	87 ARG	$e^+e^- \rightarrow \pi^+\pi^- \text{MM}$
16.9 ±4.0		GELPHMAN	85 CBAL	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
19.1 ±1.2 ±0.6		BESSON	84 CLEO	$\pi^+\pi^- \text{MM}$
18.9 ±2.6		FONSECA	84 CUSB	$e^+e^- \rightarrow \ell^+\ell^-\pi^+\pi^-$
21 ±7	7	NICZYPORUK	81B LENA	$e^+e^- \rightarrow \ell^+\ell^-\pi^+\pi^-$

¹ LEES 11C reports $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \Upsilon(2S)\text{anything})] = (1.78 \pm 0.02 \pm 0.11) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \Upsilon(2S)\text{anything}) = (10.6 \pm 0.8) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

³ A weighted average of the inclusive and exclusive results.

⁴ Using $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$, $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$ and, $\Gamma_{ee}(\Upsilon(2S)) = 0.612 \pm 0.011$ keV.

⁵ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.52 \pm 0.17)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.07)\%$.

$\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_2/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
8.6 ± 0.4 OUR AVERAGE				
8.43 ± 0.16 ± 0.42	38k	¹ BHARI 09	CLEO	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
9.2 ± 0.6 ± 0.8	275	² ALEXANDER 98	CLE2	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
9.5 ± 1.9 ± 1.9	25	ALBRECHT 87	ARG	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
8.0 ± 1.5		GELPHMAN 85	CBAL	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
10.3 ± 2.3		FONSECA 84	CUSB	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$

¹ Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.

² Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.52 \pm 0.17)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.07)\%$.

$\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_2/Γ_1

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.462 ± 0.037	¹ BHARI 09	CLEO	$e^+e^- \rightarrow \Upsilon(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Not independent of other values reported by BHARI 09.

$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$ Γ_3/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.00 ± 0.21 OUR AVERAGE				
2.00 ± 0.12 ± 0.18	22k	¹ BESSON 07	CLEO	$e^+e^- \rightarrow \Upsilon(2S) \rightarrow \tau^+\tau^-$
1.7 ± 1.5 ± 0.6		HAAS 84B	CLEO	$e^+e^- \rightarrow \tau^+\tau^-$


¹ BESSON 07 reports $[\Gamma(\Upsilon(2S) \rightarrow \tau^+\tau^-)/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \mu^+\mu^-)] = 1.04 \pm 0.04 \pm 0.05$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_4/Γ

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0193 ± 0.0017 OUR AVERAGE Error includes scale factor of 2.2. See the ideogram below.					
0.0203 ± 0.0003 ± 0.0008		120k	ADAMS 05	CLEO	$e^+e^- \rightarrow \mu^+\mu^-$
0.0122 ± 0.0028 ± 0.0019			¹ KOBEL 92	CBAL	$e^+e^- \rightarrow \mu^+\mu^-$
0.0138 ± 0.0025 ± 0.0015			KAARSBERG 89	CSB2	$e^+e^- \rightarrow \mu^+\mu^-$
0.009 ± 0.006 ± 0.006			² ALBRECHT 85	ARG	$e^+e^- \rightarrow \mu^+\mu^-$
0.018 ± 0.008 ± 0.005			HAAS 84B	CLEO	$e^+e^- \rightarrow \mu^+\mu^-$
< 0.038		90	NICZYPORUK 81C	LENA	$e^+e^- \rightarrow \mu^+\mu^-$

¹ Taking into account interference between the resonance and continuum.

² Re-evaluated using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 0.026$.



$\Gamma(\tau^+ \tau^-) / \Gamma(\mu^+ \mu^-)$ **Γ_3 / Γ_4**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.04±0.04±0.05	22k	BESSION 07	CLEO	$e^+ e^- \rightarrow \Upsilon(2S)$

$\Gamma(\Upsilon(1S)\pi^0) / \Gamma_{\text{total}}$ **Γ_6 / Γ**

VALUE (units 10 ⁻⁵)	CL%	DOCUMENT ID	TECN	COMMENT
---------------------------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4	90	¹ TAMPONI 13	BELL	$e^+ e^- \rightarrow \Upsilon(1S)\pi^0$
< 18	90	² HE 08A	CLEO	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
< 110	90	ALEXANDER 98	CLE2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
< 800	90	LURZ 87	CBAL	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$

¹ TAMPONI 13 reports $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0) / \Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+ \pi^-)] < 2.3 \times 10^{-4}$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+ \pi^-) = 17.85 \times 10^{-2}$.

² Authors assume $B(\Upsilon(1S) \rightarrow e^+ e^-) + B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 4.96\%$.

$\Gamma(\Upsilon(1S)\pi^0) / \Gamma(\Upsilon(1S)\pi^+ \pi^-)$ **Γ_6 / Γ_1**

VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT
< 2.3	90	TAMPONI 13	BELL	$e^+ e^- \rightarrow \Upsilon(1S)\pi^0$

$\Gamma(\Upsilon(1S)\eta)/\Gamma_{\text{total}}$

Γ_7/Γ

VALUE (units 10^{-4}) CL% EVTS DOCUMENT ID TECN COMMENT

2.9 ± 0.4 OUR FIT Error includes scale factor of 2.0.
2.9 ± 0.4 OUR AVERAGE Error includes scale factor of 1.9. See the ideogram below.

2.39 ± 0.31 ± 0.14	112	¹ LEES	11L	BABR	$\Upsilon(2S) \rightarrow \ell^+ \ell^- \eta$
2.1 $^{+0.7}_{-0.6}$ ± 0.3	14	² HE	08A	CLEO	$e^+ e^- \rightarrow \ell^+ \ell^- \eta$

• • • We use the following data for averages but not for fits. • • •

3.55 ± 0.32 ± 0.05	241	³ TAMPONI	13	BELL	$e^+ e^- \rightarrow \Upsilon(1S)\eta$
--------------------	-----	----------------------	----	------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 9	90	^{1,4} AUBERT	08BP	BABR	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- \pi^0 \ell^+ \ell^-$
< 28	90	ALEXANDER	98	CLE2	$e^+ e^- \rightarrow \ell^+ \ell^- \eta$
< 50	90	ALBRECHT	87	ARG	$e^+ e^- \rightarrow \pi^+ \pi^- \ell^+ \ell^- \text{MM}$
< 70	90	LURZ	87	CBAL	$e^+ e^- \rightarrow \ell^+ \ell^- (\gamma\gamma, 3\pi^0)$
< 100	90	BESSON	84	CLEO	$e^+ e^- \rightarrow \pi^+ \pi^- \ell^+ \ell^- \text{MM}$
< 20	90	FONSECA	84	CUSB	$e^+ e^- \rightarrow \ell^+ \ell^- (\gamma\gamma, \pi^+ \pi^- \pi^0)$


¹ Using $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$.

² Authors assume $B(\Upsilon(1S) \rightarrow e^+ e^-) + B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 4.96\%$.

³ TAMPONI 13 reports $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\eta)/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+ \pi^-)] = (1.99 \pm 0.14 \pm 0.11) \times 10^{-3}$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+ \pi^-) = (17.85 \pm 0.26) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ Using $\Gamma_{ee}(\Upsilon(2S)) = 0.612 \pm 0.011$ keV.

WEIGHTED AVERAGE
 2.9±0.4 (Error scaled by 1.9)



$\Gamma(\Upsilon(1S)\eta)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_7/Γ_1

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

1.64±0.25 OUR FIT Error includes scale factor of 2.0.

1.99±0.14±0.11 241 TAMPONI 13 BELL $e^+e^- \rightarrow \Upsilon(1S)\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.35±0.17±0.08 ¹ LEES 11L BABR $\Upsilon(2S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\mu^+\mu^-$

< 5.2 90 ² AUBERT 08BP BABR $e^+e^- \rightarrow \gamma\pi^+\pi^-(\pi^0)\ell^+\ell^-$

¹ Not independent of other values reported by LEES 11L.

² Not independent of other values reported by AUBERT 08BP.

$\Gamma(\Upsilon(1S)\pi^0)/\Gamma(\Upsilon(1S)\eta)$ Γ_6/Γ_7

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.13 90 TAMPONI 13 BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^0$

$\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.006 90 MASCHMANN 90 CBAL $e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\bar{d} \text{ anything})/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

3.37±0.50±0.25 58 ASNER 07 CLEO $e^+e^- \rightarrow \bar{d}X$

$\Gamma(ggg)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

58.8±1.2 6M ¹ BESSON 06A CLEO $\Upsilon(2S) \rightarrow \text{hadrons}$

¹ Calculated using the value $\Gamma(\gamma gg)/\Gamma(ggg) = (3.18 \pm 0.04 \pm 0.22 \pm 0.41)\%$ from BESSON 06A and PDG 08 values of $B(\pi^+\pi^-\Upsilon(1S)) = (18.1 \pm 0.4)\%$, $B(\pi^0\pi^0\Upsilon(1S)) = (8.6 \pm 0.4)\%$, $B(\mu^+\mu^-) = (1.93 \pm 0.17)\%$, and $R_{\text{hadrons}} = 3.51$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ measurement of BESSON 06A.

$\Gamma(\gamma gg)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

8.79±1.05 100k ¹ BESSON 06A CLEO $\Upsilon(2S) \rightarrow \gamma + \text{hadrons}$

¹ Calculated using BESSON 06A values of $\Gamma(\gamma gg)/\Gamma(ggg) = (3.18 \pm 0.04 \pm 0.22 \pm 0.41)\%$ and $\Gamma(ggg)/\Gamma_{\text{total}}$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(ggg)/\Gamma_{\text{total}}$ measurement of BESSON 06A.

$\Gamma(\text{Sum of 100 exclusive modes})/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10^{-2})	DOCUMENT ID	COMMENT
--------------------------	-------------	---------

0.29±0.03 1,2 DOBBS 12A $\Upsilon(2S) \rightarrow \text{hadrons}$

¹ DOBBS 12A presents individual exclusive branching fractions or upper limits for 100 modes of four to ten pions, kaons, or protons.

² Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

$\Gamma(\gamma g g)/\Gamma(g g g)$

Γ_{12}/Γ_{11}

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.18±0.04±0.47	6M	BESSON	06A CLEO	$\Upsilon(2S) \rightarrow (\gamma +)$ hadrons

$\Gamma(\gamma \chi_{b1}(1P))/\Gamma_{\text{total}}$

Γ_{14}/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.069 ±0.004 OUR AVERAGE				
0.0693±0.0012±0.0041	407k	ARTUSO	05 CLEO	$e^+e^- \rightarrow \gamma X$
0.069 ±0.005 ±0.009		EDWARDS	99 CLE2	$\Upsilon(2S) \rightarrow \gamma \chi(1P)$
0.091 ±0.018 ±0.022		ALBRECHT	85E ARG	$e^+e^- \rightarrow \gamma \text{conv. } X$
0.065 ±0.007 ±0.012		NERNST	85 CBAL	$e^+e^- \rightarrow \gamma X$
0.080 ±0.017 ±0.016		HAAS	84 CLEO	$e^+e^- \rightarrow \gamma \text{conv. } X$
0.059 ±0.014		KLOPFEN...	83 CUSB	$e^+e^- \rightarrow \gamma X$

$\Gamma(\gamma \chi_{b2}(1P))/\Gamma_{\text{total}}$

Γ_{15}/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0715±0.0035 OUR AVERAGE				
0.0724±0.0011±0.0040	410k	ARTUSO	05 CLEO	$e^+e^- \rightarrow \gamma X$
0.074 ±0.005 ±0.008		EDWARDS	99 CLE2	$\Upsilon(2S) \rightarrow \gamma \chi(1P)$
0.098 ±0.021 ±0.024		ALBRECHT	85E ARG	$e^+e^- \rightarrow \gamma \text{conv. } X$
0.058 ±0.007 ±0.010		NERNST	85 CBAL	$e^+e^- \rightarrow \gamma X$
0.102 ±0.018 ±0.021		HAAS	84 CLEO	$e^+e^- \rightarrow \gamma \text{conv. } X$
0.061 ±0.014		KLOPFEN...	83 CUSB	$e^+e^- \rightarrow \gamma X$

$\Gamma(\gamma \chi_{b0}(1P))/\Gamma_{\text{total}}$

Γ_{16}/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.038 ±0.004 OUR AVERAGE				
0.0375±0.0012±0.0047	198k	ARTUSO	05 CLEO	$e^+e^- \rightarrow \gamma X$
0.034 ±0.005 ±0.006		EDWARDS	99 CLE2	$\Upsilon(2S) \rightarrow \gamma \chi(1P)$
0.064 ±0.014 ±0.016		ALBRECHT	85E ARG	$e^+e^- \rightarrow \gamma \text{conv. } X$
0.036 ±0.008 ±0.009		NERNST	85 CBAL	$e^+e^- \rightarrow \gamma X$
0.044 ±0.023 ±0.009		HAAS	84 CLEO	$e^+e^- \rightarrow \gamma \text{conv. } X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.035 ±0.014		KLOPFEN...	83 CUSB	$e^+e^- \rightarrow \gamma X$

$\Gamma(\gamma f_0(1710))/\Gamma_{\text{total}}$

Γ_{17}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<59	90	¹ ALBRECHT	89 ARG	$\Upsilon(2S) \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 5.9	90	² ALBRECHT	89 ARG	$\Upsilon(2S) \rightarrow \gamma \pi^+ \pi^-$
¹ Re-evaluated assuming $B(f_0(1710) \rightarrow K^+ K^-) = 0.19$.				
² Includes unknown branching ratio of $f_0(1710) \rightarrow \pi^+ \pi^-$.				

$\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$

Γ_{18}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<53	90	¹ ALBRECHT	89 ARG	$\Upsilon(2S) \rightarrow \gamma K^+ K^-$
¹ Re-evaluated assuming $B(f'_2(1525) \rightarrow K \bar{K}) = 0.71$.				

$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$						Γ_{19}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		
<24.1	90	¹ ALBRECHT 89	ARG	$\Upsilon(2S) \rightarrow \gamma \pi^+ \pi^-$		
¹ Using $B(f_2(1270) \rightarrow \pi \pi) = 0.84$.						
$\Gamma(\gamma f_J(2220))/\Gamma_{\text{total}}$						Γ_{20}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<6.8	90	¹ ALBRECHT 89	ARG	$\Upsilon(2S) \rightarrow \gamma K^+ K^-$		
¹ Includes unknown branching ratio of $f_J(2220) \rightarrow K^+ K^-$.						
$\Gamma(\gamma \eta_c(1S))/\Gamma_{\text{total}}$						Γ_{21}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<2.7 $\times 10^{-5}$	90	WANG 11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$		
$\Gamma(\gamma \chi_{c0})/\Gamma_{\text{total}}$						Γ_{22}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<1.0 $\times 10^{-4}$	90	WANG 11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$		
$\Gamma(\gamma \chi_{c1})/\Gamma_{\text{total}}$						Γ_{23}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<3.6 $\times 10^{-6}$	90	WANG 11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$		
$\Gamma(\gamma \chi_{c2})/\Gamma_{\text{total}}$						Γ_{24}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<1.5 $\times 10^{-5}$	90	WANG 11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$		
$\Gamma(\gamma X(3872) \rightarrow \pi^+ \pi^- J/\psi)/\Gamma_{\text{total}}$						Γ_{25}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.8 $\times 10^{-6}$	90	WANG 11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$		
$\Gamma(\gamma X(3872) \rightarrow \pi^+ \pi^- \pi^0 J/\psi)/\Gamma_{\text{total}}$						Γ_{26}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<2.4 $\times 10^{-6}$	90	WANG 11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$		
$\Gamma(\gamma X(3915) \rightarrow \omega J/\psi)/\Gamma_{\text{total}}$						Γ_{27}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<2.8 $\times 10^{-6}$	90	WANG 11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$		
$\Gamma(\gamma X(4140) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$						Γ_{28}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<1.2 $\times 10^{-6}$	90	WANG 11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$		
$\Gamma(\gamma X(4350) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$						Γ_{29}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<1.3 $\times 10^{-6}$	90	WANG 11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$		

$\Gamma(\gamma\eta_b(1S))/\Gamma_{\text{total}}$ **Γ_{30}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.9 \pm 1.1^{+1.1}_{-0.9}$		$13 \pm 5k$	¹ AUBERT	09AQ BABR	$\Upsilon(2S) \rightarrow \gamma X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<21	90		LEES	11J BABR	$\Upsilon(2S) \rightarrow X\gamma$
< 8.4	90		¹ BONVICINI	10 CLEO	$\Upsilon(2S) \rightarrow \gamma X$
< 5.1	90		² ARTUSO	05 CLEO	$e^+e^- \rightarrow \gamma X$

¹ Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV.

² Superseded by BONVICINI 10.

$\Gamma(\gamma\eta_b(2S))/\Gamma_{\text{total}}$ **Γ_{31}/Γ**

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
--------------	-------------	--------------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	11 ± 4	¹ DOBBS	12 $\Upsilon(2S) \rightarrow \gamma$ hadrons
------	------------	--------------------	--

¹ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

$\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$ **Γ_{32}/Γ**
($1.5 \text{ GeV} < m_X < 5.0 \text{ GeV}$)

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.95	95	ROSNER	07A CLEO	$e^+e^- \rightarrow \gamma X$

$\Gamma(\gamma A^0 \rightarrow \gamma \text{ hadrons})/\Gamma_{\text{total}}$ **Γ_{33}/Γ**
($0.3 \text{ GeV} < m_{A^0} < 7 \text{ GeV}$)

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<8×10^{-5}	90	¹ LEES	11H BABR	$\Upsilon(2S) \rightarrow \gamma$ hadrons

¹ For a narrow scalar or pseudoscalar A^0 , excluding known resonances, with mass in the range 0.3–7 GeV. Measured 90% CL limits as a function of m_{A^0} range from 1×10^{-6} to 8×10^{-5} .

$\Gamma(\gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{34}/Γ**

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<8.3	90	¹ AUBERT	09Z BABR	$e^+e^- \rightarrow \gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-$

¹ For a narrow scalar or pseudoscalar a_1^0 with mass in the range 212–9300 MeV, excluding J/ψ and $\psi(2S)$. Measured 90% CL limits as a function of $m_{a_1^0}$ range from 0.26– 8.3×10^{-6} .

———— LEPTON FAMILY NUMBER (LF) VIOLATING MODES ————

$\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$ **Γ_{35}/Γ**

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3.2	90	LEES	10B BABR	$e^+e^- \rightarrow e^\pm \tau^\mp$

$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$					Γ_{36}/Γ
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	
< 3.3	90	LEES	10B	BABR $e^+ e^- \rightarrow \mu^\pm \tau^\mp$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<14.4	95	LOVE	08A	CLEO $e^+ e^- \rightarrow \mu^\pm \tau^\mp$	

$\Upsilon(2S)$ Cross-Particle Branching Ratios

$B(\Upsilon(2S) \rightarrow \pi^+ \pi^-) \times B(\Upsilon(3S) \rightarrow \Upsilon(2S) X)$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.78 ± 0.02 ± 0.11	906k	LEES	11C	BABR $e^+ e^- \rightarrow \pi^+ \pi^- X$

$\Upsilon(2S)$ REFERENCES

TAMPONI	13	PR D87 011104	U. Tamponi <i>et al.</i>	(BELLE Collab.)
DOBBS	12	PRL 109 082001	S. Dobbs <i>et al.</i>	
DOBBS	12A	PR D86 052003	S. Dobbs <i>et al.</i>	
LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11H	PRL 107 221803	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11L	PR D84 092003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
WANG	11B	PR D84 071107	X.L. Wang <i>et al.</i>	(BELLE Collab.)
BONVICINI	10	PR D81 031104	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
LEES	10B	PRL 104 151802	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AUBERT	09AQ	PRL 103 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09Z	PRL 103 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)
BHARI	09	PR D79 011103	S.R. Bhari <i>et al.</i>	(CLEO Collab.)
AUBERT	08BP	PR D78 112002	B. Aubert <i>et al.</i>	(BABAR Collab.)
HE	08A	PRL 101 192001	Q. He <i>et al.</i>	(CLEO Collab.)
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
ASNER	07	PR D75 012009	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BESSON	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
BESSON	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
EDWARDS	99	PR D59 032003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
BARU	96	PRPL 267 71	S.E. Baru <i>et al.</i>	(NOVO)
KOBEL	92	ZPHY C53 193	M. Kobel <i>et al.</i>	(Crystal Ball Collab.)
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	89	ZPHY C42 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)
BUCHMUEL...	88	HE $e^+ e^-$ Physics 412	W. Buchmueller, S. Cooper	(HANN, DESY, MIT)
Editors: A. Ali and P. Soeding, World Scientific, Singapore				
JAKUBOWSKI	88	ZPHY C40 49	Z. Jakubowski <i>et al.</i>	(Crystal Ball Collab.) IGJPC
ALBRECHT	87	ZPHY C35 283	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
LURZ	87	ZPHY C36 383	B. Lurz <i>et al.</i>	(Crystal Ball Collab.)
BARU	86B	ZPHY C32 622 (erratum)	S.E. Baru <i>et al.</i>	(NOVO)
ALBRECHT	85	ZPHY C28 45	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	85E	PL 160B 331	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
GELPHMAN	85	PR D32 2893	D. Gelpman <i>et al.</i>	(Crystal Ball Collab.)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
Translated from YAF 41 733.				

NERNST	85	PRL 54 2195	R. Nernst <i>et al.</i>	(Crystal Ball Collab.)
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)
BARBER	84	PL 135B 498	D.P. Barber <i>et al.</i>	(DESY, ARGUS Collab.+)
BESSON	84	PR D30 1433	D. Besson <i>et al.</i>	(CLEO Collab.)
FONSECA	84	NP B242 31	V. Fonseca <i>et al.</i>	(CUSB Collab.)
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)
HAAS	84	PRL 52 799	J. Haas <i>et al.</i>	(CLEO Collab.)
HAAS	84B	PR D30 1996	J. Haas <i>et al.</i>	(CLEO Collab.)
KLOPFEN...	83	PRL 51 160	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)
ALBRECHT	82	PL 116B 383	H. Albrecht <i>et al.</i>	(DESY, DORT, HEIDH+)
NICZYPORUK	81B	PL 100B 95	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
NICZYPORUK	81C	PL 99B 169	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
BOCK	80	ZPHY C6 125	P. Bock <i>et al.</i>	(HEIDP, MPIM, DESY, HAMB)
