

$a_1(1640)$


$I^G(J^{PC}) = 1^-(1^{++})$

OMMITTED FROM SUMMARY TABLE

Seen in the amplitude analysis of the $3\pi^0$ system produced in $\bar{p}p \rightarrow 4\pi^0$. Possibly seen in the study of the hadronic structure in decay $\tau \rightarrow 3\pi\nu_\tau$ (ABREU 98G and ASNER 00). Needs confirmation.

$a_1(1640)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1647 ± 22 OUR AVERAGE		Error includes scale factor of 1.4.		See the ideogram below.
1630 ± 20	35280	1 BAKER	03 SPEC	$\bar{p}p \rightarrow \omega\pi^+\pi^-\pi^0$
$1714 \pm 9 \pm 36$		CHUNG	02 B852	$18.3\pi^- p \rightarrow \pi^+\pi^-\pi^- p$
$1640 \pm 12 \pm 30$		BAKER	99 SPEC	$1.94\bar{p}p \rightarrow 4\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1670 ± 90		BELLINI	85 SPEC	$40\pi^- A \rightarrow \pi^-\pi^+\pi^- A$



¹ Using the $a_1(1260)$ mass and width results of BOWLER 88.

$a_1(1640)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
254± 27 OUR AVERAGE				Error includes scale factor of 1.1.
225± 30	35280	² BAKER	03	SPEC $\bar{p}p \rightarrow \omega\pi^+\pi^-\pi^0$
308± 37±62		CHUNG	02	B852 18.3 $\pi^- p \rightarrow \pi^+\pi^-\pi^- p$
300± 22±40		BAKER	99	SPEC 1.94 $\bar{p}p \rightarrow 4\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
300±100		BELLINI	85	SPEC 40 $\pi^- A \rightarrow \pi^-\pi^+\pi^- A$

² Using the $a_1(1260)$ mass and width results of BOWLER 88.

$a_1(1640)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi\pi\pi$	seen
$\Gamma_2 f_2(1270)\pi$	seen
$\Gamma_3 \sigma\pi$	seen
$\Gamma_4 \rho\pi S-wave$	seen
$\Gamma_5 \rho\pi D-wave$	seen
$\Gamma_6 \omega\pi\pi$	seen
$\Gamma_7 f_1(1285)\pi$	seen
$\Gamma_8 a_1(1260)\eta$	not seen

$a_1(1640)$ BRANCHING RATIOS

$\Gamma(f_2(1270)\pi)/\Gamma(\sigma\pi)$ Γ_2/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.24±0.07	BAKER	99	SPEC 1.94 $\bar{p}p \rightarrow 4\pi^0$

$\Gamma(\rho\pi D-wave)/\Gamma_{total}$ Γ_5/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
seen	CHUNG	02	B852 18.3 $\pi^- p \rightarrow \pi^+\pi^-\pi^- p$
seen	AMELIN	95B	VES 36 $\pi^- A \rightarrow \pi^+\pi^-\pi^- A$

$\Gamma(\omega\pi\pi)/\Gamma_{total}$ Γ_6/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	35280	³ BAKER	03	SPEC $\bar{p}p \rightarrow \omega\pi^+\pi^-\pi^0$

$\Gamma(f_1(1285)\pi)/\Gamma_{total}$ Γ_7/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	KUHN	04	B852 18 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$
seen	LEE	94	MPS2 18 $\pi^- p \rightarrow K^+\bar{K}^0\pi^-\pi^- p$

$\Gamma(a_1(1260)\eta)/\Gamma_{\text{total}}$				Γ_8/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
not seen	KUHN	04	B852	$18 \pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$
³ Assuming the $\omega\rho$ mechanism for the $\omega\pi\pi$ state.				

a₁(1640) REFERENCES

KUHN	04	PL B595 109	J. Kuhn <i>et al.</i>	(BNL E852 Collab.)
BAKER	03	PL B563 140	C.A. Baker <i>et al.</i>	
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)
ASNER	00	PR D61 012002	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BAKER	99	PL B449 114	C.A. Baker <i>et al.</i>	
ABREU	98G	PL B426 411	P. Abreu <i>et al.</i>	(DELPHI Collab.)
AMELIN	95B	PL B356 595	D.V. Amelin <i>et al.</i>	(SERP, TBIL)
LEE	94	PL B323 227	J.H. Lee <i>et al.</i>	(BNL, IND, KYUN, MASD+)
BOWLER	88	PL B209 99	M.G. Bowler	(OXF)
BELLINI	85	SJNP 41 781	D. Bellini <i>et al.</i>	

Translated from YAF 41 1223.