

$$I(J^P) = 0(\frac{1}{2}^+) \text{ Status: } ***$$

In the quark model, a Λ_b^0 is an isospin-0 udb state. The lowest Λ_b^0 ought to have $J^P = 1/2^+$. None of I , J , or P have actually been measured.

Λ_b^0 MASS

$m_{\Lambda_b^0}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
5619.51 ± 0.23 OUR AVERAGE				
5619.30 ± 0.34		¹ AAIJ	14AA LHCb	pp at 7 TeV
5620.15 ± 0.31 ± 0.47		² AALTONEN	14B CDF	$p\bar{p}$ at 1.96 TeV
5619.7 ± 0.7 ± 1.1		² AAD	13U ATLS	pp at 7 TeV
5619.44 ± 0.13 ± 0.38		² AAIJ	13AV LHCb	pp at 7 TeV
5621 ± 4 ± 3		³ ABE	97B CDF	$p\bar{p}$ at 1.8 TeV
5668 ± 16 ± 8	4	⁴ ABREU	96N DLPH	$e^+e^- \rightarrow Z$
5614 ± 21 ± 4	4	⁴ BUSKULIC	96L ALEP	$e^+e^- \rightarrow Z$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
5619.19 ± 0.70 ± 0.30		² AAIJ	12E LHCb	Repl. by AAIJ 13AV
5619.7 ± 1.2 ± 1.2		⁵ ACOSTA	06 CDF	Repl. by AALTONEN 14B
not seen		⁶ ABE	93B CDF	Repl. by ABE 97B
5640 ± 50 ± 30	16	⁷ ALBAJAR	91E UA1	$p\bar{p}$ 630 GeV
5640 ⁺¹⁰⁰ / ₋₂₁₀	52	BARI	91 SFM	$\Lambda_b^0 \rightarrow p D^0 \pi^-$
5650 ⁺¹⁵⁰ / ₋₂₀₀	90	BARI	91 SFM	$\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^+ \pi^- \pi^-$

¹ Uses exclusively reconstructed final states $\Lambda_b^0 \rightarrow \Lambda_c^+ D_s^-$, $\Lambda_c^+ D^-$ and $\bar{B}^0 \rightarrow D^+ D_s^-$ decays. The uncertainty includes both statistical and systematic contributions.

² Uses $\Lambda_b^0 \rightarrow J/\psi \Lambda$ fully reconstructed decays.

³ ABE 97B observed 38 events with a background of 18 ± 1.6 events in the mass range 5.60–5.65 GeV/ c^2 , a significance of > 3.4 standard deviations.

⁴ Uses 4 fully reconstructed Λ_b events.

⁵ Uses exclusively reconstructed final states containing a $J/\psi \rightarrow \mu^+ \mu^-$ decays.

⁶ ABE 93B states that, based on the signal claimed by ALBAJAR 91E, CDF should have found $30 \pm 23 \Lambda_b^0 \rightarrow J/\psi(1S) \Lambda$ events. Instead, CDF found not more than 2 events.

⁷ ALBAJAR 91E claims 16 ± 5 events above a background of 9 ± 1 events, a significance of about 5 standard deviations.

$m_{\Lambda_b^0} - m_{B^0}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
339.2 ± 1.4 ± 0.1	¹ ACOSTA	06 CDF	$p\bar{p}$ at 1.96 TeV

¹ Uses exclusively reconstructed final states containing $J/\psi \rightarrow \mu^+ \mu^-$ decays.

$m_{\Lambda_b^0} - m_{B^+}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
339.72 ± 0.28 OUR AVERAGE			
339.72 ± 0.24 ± 0.18	¹ AAIJ	14AA LHCb	pp at 7 TeV
339.71 ± 0.71 ± 0.09	² AAIJ	12E LHCb	pp at 7 TeV
¹ Uses exclusively reconstructed final states $\Lambda_b^0 \rightarrow \Lambda_c^+ D_s^-$, $\Lambda_c^+ D^-$ and $\bar{B}^0 \rightarrow D^+ D_s^-$ decays.			
² Uses exclusively reconstructed final states containing $J/\psi \rightarrow \mu^+ \mu^-$ decays.			

Λ_b^0 MEAN LIFE

See b -baryon Admixture section for data on b -baryon mean life average over species of b -baryon particles.

“OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements and asymmetric lifetime errors.

VALUE (10^{-12} s)	EVTS	DOCUMENT ID	TECN	COMMENT
1.466 ± 0.010 OUR EVALUATION				
1.415 ± 0.027 ± 0.006	¹	AAIJ 14E	LHCb	pp at 7 TeV
1.479 ± 0.009 ± 0.010	²	AAIJ 14U	LHCb	pp at 7, 8 TeV
1.565 ± 0.035 ± 0.020	¹	AALTONEN 14B	CDF	$p\bar{p}$ at 1.96 TeV
1.449 ± 0.036 ± 0.017	¹	AAD 13U	ATLS	pp at 7 TeV
1.482 ± 0.018 ± 0.012	³	AAIJ 13BB	LHCb	pp at 7 TeV
1.503 ± 0.052 ± 0.031	¹	CHATRCHYAN 13AC	CMS	pp at 7 TeV
1.303 ± 0.075 ± 0.035	¹	ABAZOV 12U	D0	$p\bar{p}$ at 1.96 TeV
1.401 ± 0.046 ± 0.035	⁴	AALTONEN 10B	CDF	$p\bar{p}$ at 1.96 TeV
1.290 ^{+0.119 +0.087} _{-0.110 -0.091}	⁵	ABAZOV 07U	D0	$p\bar{p}$ at 1.96 TeV
1.11 ^{+0.19} _{-0.18} ± 0.05	⁶	ABREU 99W	DLPH	$e^+ e^- \rightarrow Z$
1.29 ^{+0.24} _{-0.22} ± 0.06	⁶	ACKERSTAFF 98G	OPAL	$e^+ e^- \rightarrow Z$
1.21 ± 0.11	⁶	BARATE 98D	ALEP	$e^+ e^- \rightarrow Z$
1.32 ± 0.15 ± 0.07	⁷	ABE 96M	CDF	$p\bar{p}$ at 1.8 TeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1.537 ± 0.045 ± 0.014	¹	AALTONEN 11	CDF	Repl. by AALTONEN 14B
1.218 ^{+0.130} _{-0.115} ± 0.042	¹	ABAZOV 07S	D0	Repl. by ABAZOV 12U
1.593 ^{+0.083} _{-0.078} ± 0.033	¹	ABULENCIA 07A	CDF	Repl. by AALTONEN 11
1.22 ^{+0.22} _{-0.18} ± 0.04	¹	ABAZOV 05C	D0	Repl. by ABAZOV 07S
1.19 ^{+0.21 +0.07} _{-0.18 -0.08}		ABREU 96D	DLPH	Repl. by ABREU 99W
1.14 ^{+0.22} _{-0.19} ± 0.07	69	AKERS 95K	OPAL	Repl. by ACKERSTAFF 98G
1.02 ^{+0.23} _{-0.18} ± 0.06	44	BUSKULIC 95L	ALEP	Repl. by BARATE 98D

- ¹ Measured mean life using fully reconstructed $\Lambda_b^0 \rightarrow J/\psi \Lambda$ decays.
- ² Used $\Lambda_b^0 \rightarrow J/\psi p K^-$ decays.
- ³ Measured the lifetime ratio of decays $\Lambda_b^0 \rightarrow J/\psi p K^-$ to $B^0 \rightarrow J/\psi \pi^+ K^-$ to be $0.976 \pm 0.012 \pm 0.006$ with $\tau_{B^0} = 1.519 \pm 0.007$ ps.
- ⁴ Measured mean life using fully reconstructed $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ decays.
- ⁵ Measured using semileptonic decays $\Lambda_b^0 \rightarrow \Lambda_c^+ \mu \nu X$ and $\Lambda_c^+ \rightarrow K_S^0 p$.
- ⁶ Measured using $\Lambda_c \ell^-$ and $\Lambda \ell^+ \ell^-$.
- ⁷ Excess $\Lambda_c \ell^-$, decay lengths.

$\tau_{\Lambda_b^0}/\tau_{B^0}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.940 ± 0.035 ± 0.006	¹ AAIJ	14E LHCB	pp at 7 TeV

- ¹ Measured using $\Lambda_b^0 \rightarrow J/\psi \Lambda$ decays.

$\tau_{\Lambda_b^0}/\tau_{B^0}$ MEAN LIFE RATIO

$\tau_{\Lambda_b^0}/\tau_{B^0}$ (direct measurements)

"OUR EVALUATION" has been obtained by the Heavy Flavor Averaging Group (HFAG) by including both B^0 and B^+ decays.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.964 ± 0.007 OUR EVALUATION


0.970 ± 0.008 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.

0.929 ± 0.018 ± 0.004	¹ AAIJ	14E LHCB	pp at 7 TeV
0.974 ± 0.006 ± 0.004	² AAIJ	14U LHCB	pp at 7, 8 TeV
0.960 ± 0.025 ± 0.016	³ AAD	13U ATLS	pp at 7 TeV
0.976 ± 0.012 ± 0.006	⁴ AAIJ	13BB LHCB	pp at 7 TeV
0.864 ± 0.052 ± 0.033	^{5,6} ABAZOV	12U D0	$p\bar{p}$ at 1.96 TeV
1.020 ± 0.030 ± 0.008	⁵ AALTONEN	11 CDF	$p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.811 ^{+0.096} _{-0.087} ± 0.034	^{5,6} ABAZOV	07S D0	Repl. by ABAZOV 12U
1.041 ± 0.057	⁷ ABULENCIA	07A CDF	Repl. by AALTONEN 11
0.87 ^{+0.17} _{-0.14} ± 0.03	⁷ ABAZOV	05C D0	Repl. by ABAZOV 07S

- ¹ Measured using $\Lambda_b^0 \rightarrow J/\psi \Lambda$ and $B^0 \rightarrow J/\psi K^{*0}$ decays.
- ² Used $\Lambda_b^0 \rightarrow J/\psi p K^-$ and $B^0 \rightarrow J/\psi K^{*0}$ decays.
- ³ Measured with $\Lambda_b^0 \rightarrow J/\psi (\mu^+ \mu^-) \Lambda^0 (p \pi^-)$ decays.
- ⁴ Measures $1/\tau_{\Lambda_b^0} - 1/\tau_{B^0}$ and uses $\tau_{B^0} = 1.519 \pm 0.007$ ps to extract lifetime ratio.
- ⁵ Uses fully reconstructed $\Lambda_b \rightarrow J/\psi \Lambda$ decays.
- ⁶ Uses $B^0 \rightarrow J/\psi K_S^0$ decays for denominator.
- ⁷ Measured mean life ratio using fully reconstructed decays.



Λ_b^0 DECAY MODES

The branching fractions $B(b\text{-baryon} \rightarrow \Lambda \ell^- \bar{\nu}_\ell \text{ anything})$ and $B(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{ anything})$ are not pure measurements because the underlying measured products of these with $B(b \rightarrow b\text{-baryon})$ were used to determine $B(b \rightarrow b\text{-baryon})$, as described in the note “Production and Decay of b -Flavored Hadrons.”

For inclusive branching fractions, e.g., $\Lambda_b \rightarrow \bar{\Lambda}_c \text{ anything}$, the values usually are multiplicities, not branching fractions. They can be greater than one.

Mode	Fraction (Γ_i / Γ)	Scale factor/ Confidence level
Γ_1 $J/\psi(1S)\Lambda \times B(b \rightarrow \Lambda_b^0)$	$(5.8 \pm 0.8) \times 10^{-5}$	
Γ_2 $pD^0\pi^-$	$(6.6 \pm 0.8) \times 10^{-4}$	
Γ_3 pD^0K^-	$(4.8 \pm 0.9) \times 10^{-5}$	
Γ_4 $pJ/\psi\pi^-$	seen	
Γ_5 $pJ/\psi K^-$	seen	
Γ_6 $p\bar{K}^0\pi^-$	$(1.3 \pm 0.4) \times 10^{-5}$	
Γ_7 pK^0K^-	$< 3.5 \times 10^{-6}$	CL=90%
Γ_8 $\Lambda_c^+\pi^-$	$(4.7 \pm 0.4) \times 10^{-3}$	S=1.4
Γ_9 $\Lambda_c^+K^-$	$(3.42 \pm 0.33) \times 10^{-4}$	S=1.4
Γ_{10} $\Lambda_c^+ a_1(1260)^-$	seen	

Γ_{11}	$\Lambda_c^+ D^-$		$(4.6 \pm 0.6) \times 10^{-4}$	
Γ_{12}	$\Lambda_c^+ D_s^-$		$(1.10 \pm 0.10) \%$	
Γ_{13}	$\Lambda_c^+ \pi^+ \pi^- \pi^-$		$(7.3 \pm 1.1) \times 10^{-3}$	S=1.1
Γ_{14}	$\Lambda_c(2595)^+ \pi^-, \Lambda_c(2595)^+ \rightarrow \Lambda_c^+ \pi^+ \pi^-$		$(3.2 \pm 1.4) \times 10^{-4}$	
Γ_{15}	$\Lambda_c(2625)^+ \pi^-, \Lambda_c(2625)^+ \rightarrow \Lambda_c^+ \pi^+ \pi^-$		$(3.1 \pm 1.2) \times 10^{-4}$	
Γ_{16}	$\Sigma_c(2455)^0 \pi^+ \pi^-, \Sigma_c^0 \rightarrow \Lambda_c^+ \pi^-$		$(5.4 \pm 2.1) \times 10^{-4}$	
Γ_{17}	$\Sigma_c(2455)^{++} \pi^- \pi^-, \Sigma_c^{++} \rightarrow \Lambda_c^+ \pi^+$		$(3.1 \pm 1.5) \times 10^{-4}$	
Γ_{18}	$\Lambda K^0 2\pi^+ 2\pi^-$			
Γ_{19}	$\Lambda_c^+ \ell^- \bar{\nu}_\ell$ anything	[a]	$(10.7 \pm 2.2) \%$	
Γ_{20}	$\Lambda_c^+ \ell^- \bar{\nu}_\ell$		$(6.2^{+1.4}_{-1.2}) \%$	
Γ_{21}	$\Lambda_c^+ \pi^+ \pi^- \ell^- \bar{\nu}_\ell$		$(5.6 \pm 3.1) \%$	
Γ_{22}	$\Lambda_c(2595)^+ \ell^- \bar{\nu}_\ell$		$(7.8^{+4.0}_{-3.5}) \times 10^{-3}$	
Γ_{23}	$\Lambda_c(2625)^+ \ell^- \bar{\nu}_\ell$		$(1.3^{+0.6}_{-0.5}) \%$	
Γ_{24}	$\Sigma_c(2455)^0 \pi^+ \ell^- \bar{\nu}_\ell$			
Γ_{25}	$\Sigma_c(2455)^{++} \pi^- \ell^- \bar{\nu}_\ell$			
Γ_{26}	ρh^-	[b]	$< 2.3 \times 10^{-5}$	CL=90%
Γ_{27}	$\rho \pi^-$		$(4.4 \pm 0.8) \times 10^{-6}$	
Γ_{28}	ρK^-		$(5.3 \pm 1.0) \times 10^{-6}$	
Γ_{29}	ρD_s^-		$< 4.8 \times 10^{-4}$	CL=90%
Γ_{30}	$\Lambda \mu^+ \mu^-$		$(1.08 \pm 0.28) \times 10^{-6}$	
Γ_{31}	$\Lambda \gamma$		$< 1.3 \times 10^{-3}$	CL=90%

[a] Not a pure measurement. See note at head of Λ_b^0 Decay Modes.

[b] Here h^- means π^- or K^- .

CONSTRAINED FIT INFORMATION

An overall fit to 10 branching ratios uses 12 measurements and one constraint to determine 7 parameters. The overall fit has a $\chi^2 = 11.6$ for 6 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x ₉	95				
x ₁₃	56	53			
x ₂₀	18	18	10		
x ₂₇	0	0	0	0	
x ₂₈	0	0	0	0	82
	x ₈	x ₉	x ₁₃	x ₂₀	x ₂₇

Λ_b^0 BRANCHING RATIOS

$\Gamma(J/\psi(1S)\Lambda \times B(b \rightarrow \Lambda_b^0))/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

5.8 ± 0.8 OUR AVERAGE

6.01 ± 0.60 ± 0.58 ± 0.28

4.7 ± 2.3 ± 0.2

¹ ABAZOV

110

D0

$p\bar{p}$ at 1.96 TeV

² ABE

97B

CDF

$p\bar{p}$ at 1.8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

180 ± 60 ± 90 16 ALBAJAR 91E UA1 $p\bar{p}$ at 630 GeV

¹ ABAZOV 110 uses $B(B^0 \rightarrow J/\psi K_S^0) \times B(b \rightarrow B^0) = (1.74 \pm 0.08) \times 10^{-4}$ to obtain the result. The $(\pm 0.08) \times 10^{-4}$ uncertainty of this product is listed as the last uncertainty of the measurement, $(\pm 0.28) \times 10^{-5}$.

² ABE 97B reports $[B(\Lambda_b^0 \rightarrow J/\psi \Lambda) \times B(b \rightarrow \Lambda_b^0)] / [B(B^0 \rightarrow J/\psi K_S^0) \times B(b \rightarrow B^0)] = 0.27 \pm 0.12 \pm 0.05$. We multiply by our best value $B(B^0 \rightarrow J/\psi K_S^0) \times B(b \rightarrow B^0) = (1.74 \pm 0.08) \times 10^{-4}$. Our first error is their experiment error and our second error is the systematic error from using our best value.

$\Gamma(\rho D^0 \pi^-)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen 52 BARI 91 SFM $D^0 \rightarrow K^- \pi^+$

seen BASILE 81 SFM $D^0 \rightarrow K^- \pi^+$

$\Gamma(\rho D^0 K^-)/\Gamma(\rho D^0 \pi^-)$ Γ_3/Γ_2

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

7.3 ± 0.8^{+0.5}_{-0.6}

AAIJ

14H

LHCB

pp at 7 TeV

$\Gamma(\rho J/\psi \pi^-)/\Gamma(\rho J/\psi K^-)$ Γ_4/Γ_5

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

8.24 ± 0.25 ± 0.42

AAIJ

14K

LHCB

pp at 7, 8 TeV

$\Gamma(\rho \bar{K}^0 \pi^-)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

1.26 ± 0.19 ± 0.36

¹ AAIJ

14Q

LHCB

pp at 7 TeV

¹ Used the normalizing mode branching fraction value of $B(B^0 \rightarrow K^0 \pi^+ \pi^-) = (4.96 \pm 0.20) \times 10^{-5}$.

$\Gamma(\rho K^0 K^-)/\Gamma_{\text{total}}$ Γ_7/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.5 \times 10^{-6}$	90	AAIJ	14Q	LHCB pp at 7 TeV

$\Gamma(\Lambda_c^+ \pi^-)/\Gamma_{\text{total}}$ Γ_8/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

4.7 \pm 0.4 OUR FIT Error includes scale factor of 1.4.

4.6 \pm 0.6 OUR AVERAGE Error includes scale factor of 1.8.

$4.30 \pm 0.03^{+0.36}_{-0.35}$		¹ AAIJ	14I	LHCB pp at 7 TeV
$5.97 \pm 0.28 \pm 0.81$		² AAIJ	14Q	LHCB pp at 7 TeV
$8.8 \pm 2.8 \pm 1.5$		³ ABULENCIA	07B	CDF $p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	3	ABREU	96N	DLPH $\Lambda_c^+ \rightarrow \rho K^- \pi^+$
seen	4	BUSKULIC	96L	ALEP $\Lambda_c^+ \rightarrow \rho K^- \pi^+$, $\rho \bar{K}^0, \Lambda \pi^+ \pi^+ \pi^-$

¹ AAIJ 14I reports $(4.30 \pm 0.03^{+0.12}_{-0.11} \pm 0.26 \pm 0.21) \times 10^{-3}$ from a measurement of $[\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)/\Gamma_{\text{total}}] \times [B(B^0 \rightarrow D^- \pi^+)]$ assuming $B(B^0 \rightarrow D^- \pi^+) = (2.68 \pm 0.13) \times 10^{-3}$. Uses information on f_{baryon}/f_d from measurement in semileptonic decays by the same authors.

² Obtained using the branching fraction of $\Lambda_c^+ \rightarrow \rho K^- \pi^+$ decay.

³ The result is obtained from $(f_{\text{baryon}}/f_d) (B(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)/B(\bar{B}^0 \rightarrow D^+ \pi^-)) = 0.82 \pm 0.08 \pm 0.11 \pm 0.22$, assuming $f_{\text{baryon}}/f_d = 0.25 \pm 0.04$ and $B(\bar{B}^0 \rightarrow D^+ \pi^-) = (2.68 \pm 0.13) \times 10^{-3}$.

$\Gamma(\rho D^0 \pi^-)/\Gamma(\Lambda_c^+ \pi^-)$ Γ_2/Γ_8

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

0.140 \pm 0.007 $^{+0.007}_{-0.008}$ ¹ AAIJ 14H LHCB pp at 7 TeV

¹ AAIJ 14H reports $[\Gamma(\Lambda_b^0 \rightarrow \rho D^0 \pi^-)/\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)] \times [B(D^0 \rightarrow K^- \pi^+)] / [B(\Lambda_c^+ \rightarrow \rho K^- \pi^+)] = (8.06 \pm 0.23 \pm 0.35) \times 10^{-2}$ which we multiply or divide by our best values $B(D^0 \rightarrow K^- \pi^+) = (3.93 \pm 0.04) \times 10^{-2}$, $B(\Lambda_c^+ \rightarrow \rho K^- \pi^+) = (6.84^{+0.32}_{-0.40}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(\Lambda_c^+ K^-)/\Gamma_{\text{total}}$ Γ_9/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

3.42 \pm 0.33 OUR FIT Error includes scale factor of 1.4.

3.55 \pm 0.44 \pm 0.50 ¹ AAIJ 14Q LHCB pp at 7 TeV

¹ Obtained using the branching fraction of $\Lambda_c^+ \rightarrow \rho K^- \pi^+$ decay.

$\Gamma(\Lambda_c^+ K^-)/\Gamma(\Lambda_c^+ \pi^-)$ Γ_9/Γ_8

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

7.32 \pm 0.22 OUR FIT

7.31 \pm 0.16 \pm 0.16 AAIJ 14H LHCB pp at 7 TeV

$\Gamma(\Lambda_c^+ a_1(1260)^-)/\Gamma_{\text{total}}$					Γ_{10}/Γ
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
seen	1	ABREU	96N DLPH	$\Lambda_c^+ \rightarrow p K^- \pi^+, a_1^- \rightarrow \rho^0 \pi^- \rightarrow \pi^+ \pi^- \pi^-$	

$\Gamma(\Lambda_c^+ D_s^-)/\Gamma_{\text{total}}$					Γ_{12}/Γ
<u>VALUE (units 10^{-2})</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
1.1 ± 0.1		¹ AAIJ	14AA LHCB	pp at 7 TeV	
¹ Uses $B(\bar{B}^0 \rightarrow D^+ D_s^-) = (7.2 \pm 0.8) \times 10^{-3}$ and their measured $B(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)/B(\bar{B}^0 \rightarrow D^+ \pi^-)$ values.					

$\Gamma(\Lambda_c^+ D^-)/\Gamma(\Lambda_c^+ D_s^-)$					Γ_{11}/Γ_{12}
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.042 ± 0.003 ± 0.003		AAIJ	14AA LHCB	pp at 7 TeV	

$\Gamma(\Lambda_c^+ \pi^+ \pi^- \pi^-)/\Gamma_{\text{total}}$					Γ_{13}/Γ
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
7.3 ± 1.1 OUR FIT		Error includes scale factor of 1.1.			
14.2^{+3.6}_{-3.0} ± 1.3		¹ AALTONEN	12A CDF	$p\bar{p}$ at 1.96 TeV	

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	90	BARI	91 SFM	$\Lambda_c^+ \rightarrow p K^- \pi^+$	
¹ AALTONEN 12A reports $[\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^+ \pi^- \pi^-)/\Gamma_{\text{total}}] / [B(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)] = 3.04 \pm 0.33^{+0.70}_{-0.55}$ which we multiply by our best value $B(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-) = (4.7 \pm 0.4) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

$\Gamma(\Lambda_c^+ \pi^+ \pi^- \pi^-)/\Gamma(\Lambda_c^+ \pi^-)$					Γ_{13}/Γ_8
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
1.56 ± 0.21 OUR FIT					
1.43 ± 0.16 ± 0.13		AAIJ	11E LHCB	pp at 7 TeV	

$\Gamma(\Lambda_c(2595)^+ \pi^-, \Lambda_c(2595)^+ \rightarrow \Lambda_c^+ \pi^+ \pi^-)/\Gamma(\Lambda_c^+ \pi^+ \pi^- \pi^-)$					Γ_{14}/Γ_{13}
<u>VALUE (units 10^{-2})</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
4.4 ± 1.7^{+0.6}_{-0.4}		AAIJ	11E LHCB	pp at 7 TeV	

$\Gamma(\Lambda_c(2625)^+ \pi^-, \Lambda_c(2625)^+ \rightarrow \Lambda_c^+ \pi^+ \pi^-)/\Gamma(\Lambda_c^+ \pi^+ \pi^- \pi^-)$					Γ_{15}/Γ_{13}
<u>VALUE (units 10^{-2})</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
4.3 ± 1.5 ± 0.4		AAIJ	11E LHCB	pp at 7 TeV	

$\Gamma(\Sigma_c(2455)^0 \pi^+ \pi^-, \Sigma_c^0 \rightarrow \Lambda_c^+ \pi^-)/\Gamma(\Lambda_c^+ \pi^+ \pi^- \pi^-)$					Γ_{16}/Γ_{13}
<u>VALUE (units 10^{-2})</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
7.4 ± 2.4 ± 1.2		AAIJ	11E LHCB	pp at 7 TeV	

$\Gamma(\Sigma_c(2455)^{++}\pi^-\pi^-, \Sigma_c^{++} \rightarrow \Lambda_c^+\pi^+)/\Gamma(\Lambda_c^+\pi^+\pi^-\pi^-)$ Γ_{17}/Γ_{13}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$4.2 \pm 1.8 \pm 0.7$	AAIJ	11E	LHCB pp at 7 TeV

$\Gamma(\Lambda K^0 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	4	¹ ARENTON	86	FMPS $\Lambda K_S^0 2\pi^+ 2\pi^-$
------	---	----------------------	----	------------------------------------

¹ See the footnote to the ARENTON 86 mass value.

$\Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}$ Γ_{19}/Γ

The values and averages in this section serve only to show what values result if one assumes our $B(b \rightarrow b\text{-baryon})$. They cannot be thought of as measurements since the underlying product branching fractions were also used to determine $B(b \rightarrow b\text{-baryon})$ as described in the note on "Production and Decay of b -Flavored Hadrons."

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.107 ± 0.022 OUR AVERAGE

$0.101 \pm 0.018 \pm 0.013$		¹ BARATE	98D	ALEP $e^+e^- \rightarrow Z$
-----------------------------	--	---------------------	-----	-----------------------------

$0.14^{+0.05}_{-0.04} \pm 0.02$	29	² ABREU	95S	DLPH $e^+e^- \rightarrow Z$
---------------------------------	----	--------------------	-----	-----------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.089 \pm 0.022 \pm 0.011$	55	³ BUSKULIC	95L	ALEP Repl. by BARATE 98D
-----------------------------	----	-----------------------	-----	--------------------------

$0.18 \pm 0.07 \pm 0.02$	21	⁴ BUSKULIC	92E	ALEP $\Lambda_c^+ \rightarrow pK^-\pi^+$
--------------------------	----	-----------------------	-----	--

¹ BARATE 98D reports $[\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.0086 \pm 0.0007 \pm 0.0014$ which we divide by our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (8.5 \pm 1.1) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Measured using $\Lambda_c \ell^-$ and $\Lambda \ell^+ \ell^-$.

² ABREU 95S reports $[\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.0118 \pm 0.0026^{+0.0031}_{-0.0021}$ which we divide by our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (8.5 \pm 1.1) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ BUSKULIC 95L reports $[\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.00755 \pm 0.0014 \pm 0.0012$ which we divide by our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (8.5 \pm 1.1) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ BUSKULIC 92E reports $[\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.015 \pm 0.0035 \pm 0.0045$ which we divide by our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (8.5 \pm 1.1) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Superseded by BUSKULIC 95L.

$\Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell)/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

$0.062^{+0.014}_{-0.012}$ OUR FIT

$0.050^{+0.011+0.016}_{-0.008-0.012}$	¹ ABDALLAH	04A	DLPH $e^+e^- \rightarrow Z^0$
---	-----------------------	-----	-------------------------------

¹ Derived from a combined likelihood and event rate fit to the distribution of the Isgur-Wise variable and using HQET. The slope of the form factor is measured to be $\rho^2 = 2.03 \pm 0.46^{+0.72}_{-1.00}$.

$\Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell) / \Gamma(\Lambda_c^+ \pi^-)$ Γ_{20} / Γ_8
VALUE DOCUMENT ID TECN COMMENT

13.2^{+3.1}_{-2.8} OUR FIT

16.6 \pm 3.0^{+2.8}_{-3.6} AALTONEN 09E CDF $\rho\bar{p}$ at 1.96 TeV

$\Gamma(\Lambda_c^+ \pi^+ \pi^- \ell^- \bar{\nu}_\ell) / \Gamma_{\text{total}}$ Γ_{21} / Γ
VALUE DOCUMENT ID TECN COMMENT

0.056^{+0.031}_{-0.030} ¹ ABDALLAH 04A DLPH $e^+ e^- \rightarrow Z^0$

¹ Derived from the fraction of $\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell) / (\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell) + \Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^+ \pi^- \ell^- \bar{\nu}_\ell)) = 0.47^{+0.10+0.07}_{-0.08-0.06}$.

$\Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell) / [\Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell) + \Gamma(\Lambda_c^+ \pi^+ \pi^- \ell^- \bar{\nu}_\ell)]$ $\Gamma_{20} / (\Gamma_{20} + \Gamma_{21})$
VALUE DOCUMENT ID TECN COMMENT

0.47^{+0.10+0.07}_{-0.08-0.06} ABDALLAH 04A DLPH $e^+ e^- \rightarrow Z^0$

$\Gamma(\Lambda_c(2595)^+ \ell^- \bar{\nu}_\ell) / \Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell)$ $\Gamma_{22} / \Gamma_{20}$
VALUE DOCUMENT ID TECN COMMENT

0.126 \pm 0.033^{+0.047}_{-0.038} AALTONEN 09E CDF $\rho\bar{p}$ at 1.96 TeV

$\Gamma(\Lambda_c(2625)^+ \ell^- \bar{\nu}_\ell) / \Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell)$ $\Gamma_{23} / \Gamma_{20}$
VALUE DOCUMENT ID TECN COMMENT

0.210 \pm 0.042^{+0.071}_{-0.050} AALTONEN 09E CDF $\rho\bar{p}$ at 1.96 TeV

$[\frac{1}{2}\Gamma(\Sigma_c(2455)^0 \pi^+ \ell^- \bar{\nu}_\ell) + \frac{1}{2}\Gamma(\Sigma_c(2455)^{++} \pi^- \ell^- \bar{\nu}_\ell)] / \Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell)$ $(\frac{1}{2}\Gamma_{24} + \frac{1}{2}\Gamma_{25}) / \Gamma_{20}$
VALUE DOCUMENT ID TECN COMMENT

0.054 \pm 0.022^{+0.021}_{-0.018} AALTONEN 09E CDF $\rho\bar{p}$ at 1.96 TeV

$\Gamma(\rho h^-) / \Gamma_{\text{total}}$ Γ_{26} / Γ
VALUE CL% DOCUMENT ID TECN COMMENT

<2.3 \times 10⁻⁵ 90 ¹ ACOSTA 05O CDF $\rho\bar{p}$ at 1.96 TeV

¹ Assumes $f_\Lambda / f_d = 0.25$, and equal momentum distribution for Λ_b and B mesons.

$\Gamma(\rho\pi^-) / \Gamma_{\text{total}}$ Γ_{27} / Γ
VALUE (units 10⁻⁶) CL% DOCUMENT ID TECN COMMENT

4.4 \pm 0.8 OUR FIT
3.9 \pm 0.9 \pm 0.5 ¹ AALTONEN 09C CDF $\rho\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<50 90 ² BUSKULIC 96V ALEP $e^+ e^- \rightarrow Z$

¹ AALTONEN 09C reports $[\Gamma(\Lambda_b^0 \rightarrow p\pi^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^+\pi^-)] \times [B(\bar{b} \rightarrow b\text{-baryon})] / [B(\bar{b} \rightarrow B^0)] = 0.042 \pm 0.007 \pm 0.006$ which we multiply or divide by our best values $B(B^0 \rightarrow K^+\pi^-) = (1.96 \pm 0.05) \times 10^{-5}$, $B(\bar{b} \rightarrow b\text{-baryon}) = (8.5 \pm 1.1) \times 10^{-2}$, $B(\bar{b} \rightarrow B^0) = (40.5 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

² BUSKULIC 96V assumes PDG 96 production fractions for B^0 , B^+ , B_s , b baryons.

$\Gamma(pK^-)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
5.3±1.0 OUR FIT				
6.2±1.1±0.8		¹ AALTONEN 09C	CDF	$p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<360	90	² ADAM 96D	DLPH	$e^+e^- \rightarrow Z$
< 50	90	³ BUSKULIC 96V	ALEP	$e^+e^- \rightarrow Z$

¹ AALTONEN 09C reports $[\Gamma(\Lambda_b^0 \rightarrow pK^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^+\pi^-)] \times [B(\bar{b} \rightarrow b\text{-baryon})] / [B(\bar{b} \rightarrow B^0)] = 0.066 \pm 0.009 \pm 0.008$ which we multiply or divide by our best values $B(B^0 \rightarrow K^+\pi^-) = (1.96 \pm 0.05) \times 10^{-5}$, $B(\bar{b} \rightarrow b\text{-baryon}) = (8.5 \pm 1.1) \times 10^{-2}$, $B(\bar{b} \rightarrow B^0) = (40.5 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

² ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

³ BUSKULIC 96V assumes PDG 96 production fractions for B^0 , B^+ , B_s , b baryons.

$\Gamma(p\pi^-)/\Gamma(pK^-)$ Γ_{27}/Γ_{28}

VALUE	DOCUMENT ID	TECN	COMMENT
0.84±0.09 OUR FIT			
0.86±0.08±0.05	AAIJ	12AR	LHCB pp at 7 TeV

$\Gamma(pD_s^-)/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<4.8 × 10⁻⁴	90	AAIJ	14Q	LHCB pp at 7 TeV

$\Gamma(\Lambda\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
10.8±2.8 OUR AVERAGE			
9.6±1.6±2.5	¹ AAIJ	13AJ	LHCB pp at 7 TeV
17.3±4.2±5.5	AALTONEN	11AI	CDF $p\bar{p}$ at 1.96 TeV

¹ Uses $B(\Lambda_b^0 \rightarrow J/\psi\Lambda) = (6.2 \pm 1.4) \times 10^{-4}$.

$\Gamma(\Lambda\gamma)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.3 × 10⁻³	90	ACOSTA	02G	CDF $p\bar{p}$ at 1.8 TeV

PARTIAL BRANCHING FRACTIONS IN $\Lambda_b \rightarrow \Lambda \mu^+ \mu^-$

$B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-)$ ($q^2 < 2.0 \text{ GeV}^2/c^4$)

<u>VALUE (units 10^{-7})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1 ±1 OUR AVERAGE			
0.56 ± 0.76 ± 0.80	¹ AAIJ	13AJ LHCB	pp at 7 TeV
0.15 ± 2.01 ± 0.05	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
¹ Uses $B(\Lambda_b^0 \rightarrow J/\psi \Lambda) = (6.2 \pm 1.4) \times 10^{-4}$.			

$B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-)$ ($2.0 < q^2 < 4.3 \text{ GeV}^2/c^4$)

<u>VALUE (units 10^{-7})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.8 ±0.6 OUR AVERAGE			
0.71 ± 0.60 ± 0.23	¹ AAIJ	13AJ LHCB	pp at 7 TeV
1.8 ± 1.7 ± 0.6	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
¹ Uses $B(\Lambda_b^0 \rightarrow J/\psi \Lambda) = (6.2 \pm 1.4) \times 10^{-4}$.			

$B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-)$ ($4.3 < q^2 < 8.68 \text{ GeV}^2/c^4$)

<u>VALUE (units 10^{-7})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.5 ±0.7 OUR AVERAGE			
0.66 ± 0.74 ± 0.18	¹ AAIJ	13AJ LHCB	pp at 7 TeV
-0.2 ± 1.6 ± 0.1	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
¹ Uses $B(\Lambda_b^0 \rightarrow J/\psi \Lambda) = (6.2 \pm 1.4) \times 10^{-4}$.			

$B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-)$ ($10.09 < q^2 < 12.86 \text{ GeV}^2/c^4$)

<u>VALUE (units 10^{-7})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.8 ±0.7 OUR AVERAGE			
1.55 ± 0.58 ± 0.55	¹ AAIJ	13AJ LHCB	pp at 7 TeV
3.0 ± 1.5 ± 1.0	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
¹ Uses $B(\Lambda_b^0 \rightarrow J/\psi \Lambda) = (6.2 \pm 1.4) \times 10^{-4}$.			

$B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-)$ ($14.18 < q^2 < 16.0 \text{ GeV}^2/c^4$)

<u>VALUE (units 10^{-7})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.3 ±0.5 OUR AVERAGE			
1.44 ± 0.44 ± 0.42	¹ AAIJ	13AJ LHCB	pp at 7 TeV
1.0 ± 0.7 ± 0.3	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
¹ Uses $B(\Lambda_b^0 \rightarrow J/\psi \Lambda) = (6.2 \pm 1.4) \times 10^{-4}$.			

$B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-)$ ($16.0 < q^2 \text{ GeV}^2/c^4$)

<u>VALUE (units 10^{-7})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.2 ±1.3 OUR AVERAGE			
4.73 ± 0.77 ± 1.25	^{1,2} AAIJ	13AJ LHCB	pp at 7 TeV
7.0 ± 1.9 ± 2.2	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
¹ Uses $B(\Lambda_b^0 \rightarrow J/\psi \Lambda) = (6.2 \pm 1.4) \times 10^{-4}$.			
² Requires $16.00 < q^2 < 20.30 \text{ GeV}^2/c^4$.			

$B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-)$ ($1.0 < q^2 < 6.0 \text{ GeV}^2/c^4$)

<u>VALUE (units 10^{-7})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.3 ± 2.1 ± 0.4	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV

$B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-)$ ($0.0 < q^2 < 4.3 \text{ GeV}^2/c^4$)

<u>VALUE (units 10^{-7})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.7 \pm 2.5 \pm 0.9$	AALTONEN 11A1	CDF	$p\bar{p}$ at 1.96 TeV

CP VIOLATION

A_{CP} is defined as

$$A_{CP} = \frac{B(\Lambda_b^0 \rightarrow f) - B(\bar{\Lambda}_b^0 \rightarrow \bar{f})}{B(\Lambda_b^0 \rightarrow f) + B(\bar{\Lambda}_b^0 \rightarrow \bar{f})},$$

the CP-violation asymmetry of exclusive Λ_b^0 and $\bar{\Lambda}_b^0$ decay.

$A_{CP}(\Lambda_b \rightarrow p\pi^-)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.06 ± 0.07 OUR AVERAGE			
$0.06 \pm 0.07 \pm 0.03$	AALTONEN 14P	CDF	$p\bar{p}$ at 1.96 TeV
$0.03 \pm 0.17 \pm 0.05$	AALTONEN 11N	CDF	$p\bar{p}$ at 1.96 TeV

$A_{CP}(\Lambda_b \rightarrow pK^-)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.00 ± 0.19 OUR AVERAGE	Error includes scale factor of 2.4.		
$-0.10 \pm 0.08 \pm 0.04$	AALTONEN 14P	CDF	$p\bar{p}$ at 1.96 TeV
$0.37 \pm 0.17 \pm 0.03$	AALTONEN 11N	CDF	$p\bar{p}$ at 1.96 TeV

$A_{CP}(\Lambda_b \rightarrow p\bar{K}^0\pi^-)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.22 \pm 0.13 \pm 0.03$	¹ AAIJ 14Q	LHCB	pp at 7 TeV

¹AAIJ 14Q definition of A_{CP} has the opposite sign to ours.

$\Delta A_{CP}(J/\psi p\pi^- / K^-) \equiv A_{CP}(J/\psi p\pi^-) - A_{CP}(J/\psi pK^-)$

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$5.7 \pm 2.4 \pm 1.2$	AAIJ 14K	LHCB	pp at 7, 8 TeV

Λ_b^0 DECAY PARAMETERS

See the note on "Baryon Decay Parameters" in the neutron Listings.

α decay parameter for $\Lambda_b \rightarrow J/\psi \Lambda$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.18 ± 0.13 OUR AVERAGE			
$0.30 \pm 0.16 \pm 0.06$	¹ AAD 14L	ATLS	pp at 7 TeV
$0.05 \pm 0.17 \pm 0.07$	² AAIJ 13AG	LHCB	pp at 7 TeV

¹An angular analysis of $\Lambda_b \rightarrow J/\psi \Lambda$ decay is performed and magnitudes of all helicity amplitudes are also reported.

²An angular analysis of $\Lambda_b \rightarrow J/\psi \Lambda$ decay is performed and a Λ_b transverse production polarization of $0.06 \pm 0.07 \pm 0.02$ is also reported.

REFERENCES

- | | | | | |
|------------|------|------------------------|-----------------------------|---------------------|
| AAD | 14L | PR D89 092009 | G. Aad <i>et al.</i> | (ATLAS Collab.) |
| AAIJ | 14AA | PRL 112 202001 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14E | JHEP 1404 114 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14H | PR D89 032001 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14I | JHEP 1408 143 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14K | JHEP 1407 103 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14Q | JHEP 1404 087 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14U | PL B734 122 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AALTONEN | 14B | PR D89 072014 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 14P | PRL 113 242001 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AAD | 13U | PR D87 032002 | G. Aad <i>et al.</i> | (ATLAS Collab.) |
| AAIJ | 13AG | PL B724 27 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13AJ | PL B725 25 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13AV | PRL 110 182001 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13BB | PRL 111 102003 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| CHATRCHYAN | 13AC | JHEP 1307 163 | S. Chatrchyan <i>et al.</i> | (CMS Collab.) |
| AAIJ | 12AR | JHEP 1210 037 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 12E | PL B708 241 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AALTONEN | 12A | PR D85 032003 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| ABAZOV | 12U | PR D85 112003 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| AAIJ | 11E | PR D84 092001 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| Also | | PR D85 039904 (errata) | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AALTONEN | 11 | PRL 106 121804 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 11AI | PRL 107 201802 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 11N | PRL 106 181802 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| ABAZOV | 11O | PR D84 031102 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| AALTONEN | 10B | PRL 104 102002 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 09C | PRL 103 031801 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 09E | PR D79 032001 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| ABAZOV | 07S | PRL 99 142001 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 07U | PRL 99 182001 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABULENCIA | 07A | PRL 98 122001 | A. Abulencia <i>et al.</i> | (FNAL CDF Collab.) |
| ABULENCIA | 07B | PRL 98 122002 | A. Abulencia <i>et al.</i> | (FNAL CDF Collab.) |
| ACOSTA | 06 | PRL 96 202001 | D. Acosta <i>et al.</i> | (CDF Collab.) |
| ABAZOV | 05C | PRL 94 102001 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ACOSTA | 05O | PR D72 051104 | D. Acosta <i>et al.</i> | (CDF Collab.) |
| ABDALLAH | 04A | PL B585 63 | J. Abdallah <i>et al.</i> | (DELPHI Collab.) |
| ACOSTA | 02G | PR D66 112002 | D. Acosta <i>et al.</i> | (CDF Collab.) |
| ABREU | 99W | EPJ C10 185 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ACKERSTAFF | 98G | PL B426 161 | K. Ackerstaff <i>et al.</i> | (OPAL Collab.) |
| BARATE | 98D | EPJ C2 197 | R. Barate <i>et al.</i> | (ALEPH Collab.) |
| ABE | 97B | PR D55 1142 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 96M | PRL 77 1439 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABREU | 96D | ZPHY C71 199 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ABREU | 96N | PL B374 351 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ADAM | 96D | ZPHY C72 207 | W. Adam <i>et al.</i> | (DELPHI Collab.) |
| BUSKULIC | 96L | PL B380 442 | D. Buskalic <i>et al.</i> | (ALEPH Collab.) |
| BUSKULIC | 96V | PL B384 471 | D. Buskalic <i>et al.</i> | (ALEPH Collab.) |
| PDG | 96 | PR D54 1 | R. M. Barnett <i>et al.</i> | (PDG Collab.) |
| ABREU | 95S | ZPHY C68 375 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| AKERS | 95K | PL B353 402 | R. Akers <i>et al.</i> | (OPAL Collab.) |
| BUSKULIC | 95L | PL B357 685 | D. Buskalic <i>et al.</i> | (ALEPH Collab.) |
| ABE | 93B | PR D47 R2639 | F. Abe <i>et al.</i> | (CDF Collab.) |
| BUSKULIC | 92E | PL B294 145 | D. Buskalic <i>et al.</i> | (ALEPH Collab.) |
| ALBAJAR | 91E | PL B273 540 | C. Albajar <i>et al.</i> | (UA1 Collab.) |
| BARI | 91 | NC 104A 1787 | G. Bari <i>et al.</i> | (CERN R422 Collab.) |
| ARENTON | 86 | NP B274 707 | M.W. Arenton <i>et al.</i> | (ARIZ, NDAM, VAND) |
| BASILE | 81 | LNC 31 97 | M. Basile <i>et al.</i> | (CERN R415 Collab.) |