

$\Upsilon(10860)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

$\Upsilon(10860)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10876 ± 11	OUR EVALUATION		Weighted-average of Belle and BaBar results, but tripling the scaling S -factors applied to the uncertainties to account for model-dependence, handling of radiative corrections, and interference effects.
• • •	We do not use the following data for averages, fits, limits, etc.	• • •	
10879 ± 3	^{1,2} CHEN	10	BELL $e^+e^- \rightarrow$ hadrons
10888.4 ⁺ ₋ $\frac{2.7}{2.6} \pm 1.2$	³ CHEN	10	BELL $e^+e^- \rightarrow \Upsilon(1S, 2S, 3S)\pi^+\pi^-$
10876 ± 2	¹ AUBERT	09E	BABR $e^+e^- \rightarrow$ hadrons
10869 ± 2	⁴ AUBERT	09E	BABR $e^+e^- \rightarrow$ hadrons
10868 ± 6 ± 5	⁵ BESSON	85	CLEO $e^+e^- \rightarrow$ hadrons
10845 ± 20	⁶ LOVELOCK	85	CUSB $e^+e^- \rightarrow$ hadrons

¹ In a model where a flat non-resonant $b\bar{b}$ -continuum is incoherently added to a second flat component interfering with two Breit-Wigner resonances. Systematic uncertainties not estimated.

² The parameters of the $\Upsilon(11020)$ are fixed to those in AUBERT 09E.

³ In a model where a flat nonresonant $\Upsilon(1S, 2S, 3S)\pi^+\pi^-$ continuum interferes with a single Breit-Wigner resonance.

⁴ In a model where a non-resonant $b\bar{b}$ -continuum represented by a threshold function at $\sqrt{s}=2m_B$ is incoherently added to a flat component interfering with two Breit-Wigner resonances. Not independent of other AUBERT 09E results. Systematic uncertainties not estimated.

⁵ Assuming four Gaussians with radiative tails and a single step in R .

⁶ In a coupled-channel model with three resonances and a smooth step in R .

$\Upsilon(10860)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
55 ± 28	OUR EVALUATION		Weighted-average of Belle and BaBar results, but tripling the scaling S -factors applied to the uncertainties to account for model-dependence, handling of radiative corrections, and interference effects.
• • •	We do not use the following data for averages, fits, limits, etc.	• • •	
46 ⁺ ₋ $\frac{9}{7}$	^{7,8} CHEN	10	BELL $e^+e^- \rightarrow$ hadrons
30.7 ⁺ ₋ $\frac{8.3}{7.0} \pm 3.1$	⁹ CHEN	10	BELL $e^+e^- \rightarrow \Upsilon(1S, 2S, 3S)\pi^+\pi^-$
43 ± 4	⁷ AUBERT	09E	BABR $e^+e^- \rightarrow$ hadrons
74 ± 4	¹⁰ AUBERT	09E	BABR $e^+e^- \rightarrow$ hadrons
112 ± 17 ± 23	¹¹ BESSON	85	CLEO $e^+e^- \rightarrow$ hadrons
110 ± 15	¹² LOVELOCK	85	CUSB $e^+e^- \rightarrow$ hadrons

- ⁷ In a model where a flat non-resonant $b\bar{b}$ -continuum is incoherently added to a second flat component interfering with two Breit-Wigner resonances. Systematic uncertainties not estimated.
- ⁸ The parameters of the $\Upsilon(11020)$ are fixed to those in AUBERT 09E.
- ⁹ In a model where a flat nonresonant $\Upsilon(1S, 2S, 3S)\pi^+\pi^-$ continuum interferes with a single Breit-Wigner resonance.
- ¹⁰ In a model where a non-resonant $b\bar{b}$ -continuum represented by a threshold function at $\sqrt{s}=2m_B$ is incoherently added to a flat component interfering with two Breit-Wigner resonances. Not independent of other AUBERT 09E results. Systematic uncertainties not estimated.
- ¹¹ Assuming four Gaussians with radiative tails and a single step in R .
- ¹² In a coupled-channel model with three resonances and a smooth step in R .

$\Upsilon(10860)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $B\bar{B}X$	(76.2 $^{+2.7}_{-4.0}$) %	
Γ_2 $B\bar{B}$	(5.5 ± 1.0) %	
Γ_3 $B\bar{B}^* + \text{c.c.}$	(13.7 ± 1.6) %	
Γ_4 $B^*\bar{B}^*$	(38.1 ± 3.4) %	
Γ_5 $B\bar{B}^{(*)}\pi$	< 19.7 %	90%
Γ_6 $B\bar{B}\pi$	(0.0 ± 1.2) %	
Γ_7 $B^*\bar{B}\pi + B\bar{B}^*\pi$	(7.3 ± 2.3) %	
Γ_8 $B^*\bar{B}^*\pi$	(1.0 ± 1.4) %	
Γ_9 $B\bar{B}\pi\pi$	< 8.9 %	90%
Γ_{10} $B_s^{(*)}\bar{B}_s^{(*)}$	(20.1 ± 3.1) %	
Γ_{11} $B_s\bar{B}_s$	(5 ± 5) $\times 10^{-3}$	
Γ_{12} $B_s\bar{B}_s^* + \text{c.c.}$	(1.35 ± 0.32) %	
Γ_{13} $B_s^*\bar{B}_s^*$	(17.6 ± 2.7) %	
Γ_{14} no open-bottom	(3.8 $^{+5.0}_{-0.5}$) %	
Γ_{15} e^+e^-	(5.6 ± 3.1) $\times 10^{-6}$	
Γ_{16} $K^*(892)^0\bar{K}^0$	< 1.0 $\times 10^{-5}$	90%
Γ_{17} $\Upsilon(1S)\pi^+\pi^-$	(5.3 ± 0.6) $\times 10^{-3}$	
Γ_{18} $\Upsilon(2S)\pi^+\pi^-$	(7.8 ± 1.3) $\times 10^{-3}$	
Γ_{19} $\Upsilon(3S)\pi^+\pi^-$	(4.8 $^{+1.9}_{-1.7}$) $\times 10^{-3}$	
Γ_{20} $\Upsilon(1S)K^+K^-$	(6.1 ± 1.8) $\times 10^{-4}$	
Γ_{21} $h_b(1P)\pi^+\pi^-$	(3.5 $^{+1.0}_{-1.3}$) $\times 10^{-3}$	
Γ_{22} $h_b(2P)\pi^+\pi^-$	(6.0 $^{+2.1}_{-1.8}$) $\times 10^{-3}$	
Γ_{23} $\chi_{b0}(1P)\pi^+\pi^-\pi^0$	< 6.3 $\times 10^{-3}$	90%
Γ_{24} $\chi_{b0}(1P)\omega$	< 3.9 $\times 10^{-3}$	90%
Γ_{25} $\chi_{b0}(1P)(\pi^+\pi^-\pi^0)_{\text{non-}\omega}$	< 4.8 $\times 10^{-3}$	90%
Γ_{26} $\chi_{b1}(1P)\pi^+\pi^-\pi^0$	(1.85 ± 0.33) $\times 10^{-3}$	

Γ_{27}	$\chi_{b1}(1P)\omega$	$(1.57 \pm 0.30) \times 10^{-3}$	
Γ_{28}	$\chi_{b1}(1P)(\pi^+\pi^-\pi^0)_{\text{non-}\omega}$	$(5.2 \pm 1.9) \times 10^{-4}$	
Γ_{29}	$\chi_{b2}(1P)\pi^+\pi^-\pi^0$	$(1.17 \pm 0.30) \times 10^{-3}$	
Γ_{30}	$\chi_{b2}(1P)\omega$	$(6.0 \pm 2.7) \times 10^{-4}$	
Γ_{31}	$\chi_{b2}(1P)(\pi^+\pi^-\pi^0)_{\text{non-}\omega}$	$(6 \pm 4) \times 10^{-4}$	
Γ_{32}	$\gamma X_b \rightarrow \gamma \Upsilon(1S)\omega$	$< 3.8 \times 10^{-5}$	90%

Inclusive Decays.

These decay modes are submodes of one or more of the decay modes above.

Γ_{33}	ϕ anything	$(13.8^{+2.4}_{-1.7})\%$
Γ_{34}	D^0 anything + c.c.	$(108 \pm 8)\%$
Γ_{35}	D_s anything + c.c.	$(46 \pm 6)\%$
Γ_{36}	J/ψ anything	$(2.06 \pm 0.21)\%$
Γ_{37}	B^0 anything + c.c.	$(77 \pm 8)\%$
Γ_{38}	B^+ anything + c.c.	$(72 \pm 6)\%$

$\Upsilon(10860)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$					Γ_{15}
<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
0.31 ± 0.07 OUR AVERAGE	Error includes scale factor of 1.3.				
$0.22 \pm 0.05 \pm 0.07$	BESSION	85	CLEO	$e^+e^- \rightarrow$ hadrons	
0.365 ± 0.070	LOVELOCK	85	CUSB	$e^+e^- \rightarrow$ hadrons	

$\Upsilon(10860)$ BRANCHING RATIOS

“OUR EVALUATION” is obtained based on averages of rescaled data listed below. The averages and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>.

$\Gamma(B\bar{B}X)/\Gamma_{\text{total}}$					Γ_1/Γ
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$0.762^{+0.027}_{-0.043}$ OUR EVALUATION					
0.71 ± 0.06 OUR AVERAGE					
$0.737 \pm 0.032 \pm 0.051$	1063	¹³ DRUTSKOY	10	BELL	$\Upsilon(5S) \rightarrow B^+X, B^0X$
$0.589 \pm 0.100 \pm 0.092$		¹⁴ HUANG	07	CLEO	$\Upsilon(5S) \rightarrow$ hadrons

$\Gamma(B\bar{B})/\Gamma_{\text{total}}$					Γ_2/Γ
<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$5.5^{+1.0}_{-0.9} \pm 0.4$		¹⁵ DRUTSKOY	10	BELL	$\Upsilon(5S) \rightarrow B^+X, B^0X$
••• We do not use the following data for averages, fits, limits, etc. •••					
<13.8	90	¹⁴ HUANG	07	CLEO	$\Upsilon(5S) \rightarrow$ hadrons

$\Gamma(B\bar{B})/\Gamma(B\bar{B}X)$					Γ_2/Γ_1
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<0.22	90	AQUINES	06	CLE3	$\Upsilon(5S) \rightarrow$ hadrons

$\Gamma(B\bar{B}^* + \text{c.c.})/\Gamma_{\text{total}}$					Γ_3/Γ
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.137±0.016 OUR AVERAGE					
0.137±0.013±0.011		15 DRUTSKOY	10	BELL	$\Upsilon(5S) \rightarrow B^+ X, B^0 X$
0.143±0.053±0.027		14 HUANG	07	CLEO	$\Upsilon(5S) \rightarrow$ hadrons

$\Gamma(B\bar{B}^* + \text{c.c.})/\Gamma(B\bar{B}X)$					Γ_3/Γ_1
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.24±0.09±0.03	10	AQUINES	06	CLE3	$\Upsilon(5S) \rightarrow$ hadrons

$\Gamma(B^* \bar{B}^*)/\Gamma_{\text{total}}$					Γ_4/Γ
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.381±0.034 OUR AVERAGE					
0.375 ^{+0.021} _{-0.019} ±0.030		15 DRUTSKOY	10	BELL	$\Upsilon(5S) \rightarrow B^+ X, B^0 X$
0.436±0.083±0.072		14 HUANG	07	CLEO	$\Upsilon(5S) \rightarrow$ hadrons

$\Gamma(B^* \bar{B}^*)/\Gamma(B\bar{B}X)$					Γ_4/Γ_1
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.74±0.15±0.08	31	AQUINES	06	CLE3	$\Upsilon(5S) \rightarrow$ hadrons

$\Gamma(B\bar{B}^*(*)\pi)/\Gamma_{\text{total}}$					Γ_5/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<0.197	90	14 HUANG	07	CLEO	$\Upsilon(5S) \rightarrow$ hadrons

$\Gamma(B\bar{B}^*(*)\pi)/\Gamma(B\bar{B}X)$					Γ_5/Γ_1
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<0.32	90	AQUINES	06	CLE3	$\Upsilon(5S) \rightarrow$ hadrons

$\Gamma(B\bar{B}\pi)/\Gamma_{\text{total}}$					Γ_6/Γ
<u>VALUE (units 10⁻²)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.0±1.2±0.3	0	15 DRUTSKOY	10	BELL	$\Upsilon(5S) \rightarrow B^{+,0}\pi^- X$

$[\Gamma(B^* \bar{B}\pi) + \Gamma(B\bar{B}^*\pi)]/\Gamma_{\text{total}}$					Γ_7/Γ
<u>VALUE (units 10⁻²)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
7.3^{+2.3}_{-2.1}±0.8	38	15 DRUTSKOY	10	BELL	$\Upsilon(5S) \rightarrow B^{+,0}\pi^- X$

$\Gamma(B^* \bar{B}^*\pi)/\Gamma_{\text{total}}$					Γ_8/Γ
<u>VALUE (units 10⁻²)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
1.0^{+1.4}_{-1.3}±0.4	5	15 DRUTSKOY	10	BELL	$\Upsilon(5S) \rightarrow B^{+,0}\pi^- X$

$\Gamma(B\bar{B}\pi\pi)/\Gamma_{\text{total}}$					Γ_9/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<0.089	90	14 HUANG	07	CLEO	$\Upsilon(5S) \rightarrow$ hadrons

$\Gamma(B\bar{B}\pi\pi)/\Gamma(B\bar{B}X)$					Γ_9/Γ_1
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<0.14	90	AQUINES	06	CLE3	$\Upsilon(5S) \rightarrow \text{hadrons}$

$\Gamma(B_s^{(*)}\bar{B}_s^{(*)})/\Gamma_{\text{total}}$					$\Gamma_{10}/\Gamma = (\Gamma_{11}+\Gamma_{12}+\Gamma_{13})/\Gamma$
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	

0.201^{+0.030}_{-0.031} OUR EVALUATION

0.189^{+0.027}_{-0.021} OUR AVERAGE

0.172±0.030 16 ESEN 13 BELL $\Upsilon(5S) \rightarrow D^0 X, D_s X$

0.21^{+0.06}_{-0.03} 17 HUANG 07 CLEO $\Upsilon(5S) \rightarrow D_s X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.180±0.013±0.032 18 DRUTSKOY 07 BELL $\Upsilon(5S) \rightarrow D^0 X, D_s X$

0.160±0.026±0.058 19 ARTUSO 05B CLEO $e^+e^- \rightarrow D_X X$

$\Gamma(B_s^{(*)}\bar{B}_s^{(*)})/\Gamma(B\bar{B}X)$					Γ_{10}/Γ_1
<u>VALUE</u>		<u>DOCUMENT ID</u>			

0.264^{+0.052}_{-0.045} OUR EVALUATION

$\Gamma(B_s^*\bar{B}_s^*)/\Gamma(B_s^{(*)}\bar{B}_s^{(*)})$					$\Gamma_{13}/\Gamma_{10} = \Gamma_{13}/(\Gamma_{11}+\Gamma_{12}+\Gamma_{13})$
<u>VALUE (units 10⁻²)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	

87.8±1.5 OUR AVERAGE

87.0±1.7 20,21 ESEN 13 BELL $B_s^0 \rightarrow D_s^- \pi^+$

90.5±3.2±0.1 227 21,22 LI 12 BELL $B_s^0 \rightarrow J/\psi \eta^{(\prime)}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

90.1^{+3.8}_{-4.0}±0.2 23 LOUVOT 09 BELL $10.86 e^+e^- \rightarrow B_s^{(*)}\bar{B}_s^{(*)}$

93⁺⁷₋₉ ±1 23 DRUTSKOY 07A BELL Superseded by LOUVOT 09

$\Gamma(B_s\bar{B}_s)/\Gamma(B_s^{(*)}\bar{B}_s^{(*)})$					$\Gamma_{11}/\Gamma_{10} = \Gamma_{11}/(\Gamma_{11}+\Gamma_{12}+\Gamma_{13})$
<u>VALUE (units 10⁻²)</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	

2.6^{+2.6}_{-2.5} LOUVOT 09 BELL $10.86 e^+e^- \rightarrow B_s^{(*)}\bar{B}_s^{(*)}$

$\Gamma(B_s\bar{B}_s)/\Gamma(B_s^*\bar{B}_s^*)$					Γ_{11}/Γ_{13}
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	

<0.16 90 BONVICINI 06 CLE3 e^+e^-

$\Gamma(B_s\bar{B}_s^* + \text{c.c.})/\Gamma(B_s^{(*)}\bar{B}_s^{(*)})$					$\Gamma_{12}/\Gamma_{10} = \Gamma_{12}/(\Gamma_{11}+\Gamma_{12}+\Gamma_{13})$
<u>VALUE (units 10⁻²)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	

6.7±1.2 OUR AVERAGE

7.3±1.4 20,21 ESEN 13 BELL $B_s^0 \rightarrow D_s^- \pi^+$

4.9±2.5±0.0 227 21,22 LI 12 BELL $B_s^0 \rightarrow J/\psi \eta^{(\prime)}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.3^{+3.3}_{-3.0}±0.1 LOUVOT 09 BELL $10.86 e^+e^- \rightarrow B_s^{(*)}\bar{B}_s^{(*)}$

$\Gamma(B_s \bar{B}_s^* + \text{c.c.})/\Gamma(B_s^* \bar{B}_s^*)$					Γ_{12}/Γ_{13}
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.16	90	BONVICINI	06	CLE3	$e^+ e^-$

$\Gamma(\text{no open-bottom})/\Gamma_{\text{total}}$		Γ_{14}/Γ
VALUE	DOCUMENT ID	

0.038^{+0.051}_{-0.005} OUR EVALUATION

$\Gamma(K^*(892)^0 \bar{K}^0)/\Gamma_{\text{total}}$					Γ_{16}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<1.0 × 10 ⁻⁵	90	SHEN	13A	BELL	$e^+ e^- \rightarrow K^*(892)^0 \bar{K}^0$

$\Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{17}/Γ
VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT	
5.3 ± 0.3 ± 0.5	325	24 CHEN	08	BELL	10.87 $e^+ e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$

$\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{18}/Γ
VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT	
7.8 ± 0.6 ± 1.1	186	24 CHEN	08	BELL	10.87 $e^+ e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$

$\Gamma(\Upsilon(3S)\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{19}/Γ
VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT	
4.8 ^{+1.8} _{-1.5} ± 0.7	10	24 CHEN	08	BELL	10.87 $e^+ e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$

$\Gamma(\Upsilon(1S)K^+K^-)/\Gamma_{\text{total}}$					Γ_{20}/Γ
VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT	
6.1 ^{+1.6} _{-1.4} ± 1.0	20	24 CHEN	08	BELL	10.87 $e^+ e^- \rightarrow \Upsilon(1S)K^+K^-$

$\Gamma(h_b(1P)\pi^+\pi^-)/\Gamma(\Upsilon(2S)\pi^+\pi^-)$					Γ_{21}/Γ_{18}
VALUE		DOCUMENT ID	TECN	COMMENT	
0.45 ± 0.08 ^{+0.07} _{-0.12}		ADACHI	12	BELL	10.86 $e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(h_b(2P)\pi^+\pi^-)/\Gamma(\Upsilon(2S)\pi^+\pi^-)$					Γ_{22}/Γ_{18}
VALUE		DOCUMENT ID	TECN	COMMENT	
0.77 ± 0.08 ^{+0.22} _{-0.17}		ADACHI	12	BELL	10.86 $e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\chi_{b0}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$					Γ_{23}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<6.3 × 10 ⁻³	90	25 HE	14	BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0 \gamma \Upsilon(1S)$

$\Gamma(\chi_{b0}(1P)\omega)/\Gamma_{\text{total}}$					Γ_{24}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<3.9 × 10 ⁻³	90	25 HE	14	BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0 \gamma \Upsilon(1S)$

$\Gamma(\chi_{b0}(1P)(\pi^+\pi^-\pi^0)_{\text{non-}\omega})/\Gamma_{\text{total}}$						Γ_{25}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<4.8 \times 10^{-3}$	90	25 HE	14	BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$	

$\Gamma(\chi_{b1}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$						Γ_{26}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT		
$1.85 \pm 0.23 \pm 0.23$	80	25 HE	14	BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$	

$\Gamma(\chi_{b1}(1P)\omega)/\Gamma_{\text{total}}$						Γ_{27}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT		
$1.57 \pm 0.22 \pm 0.21$	60	25 HE	14	BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$	

$\Gamma(\chi_{b1}(1P)(\pi^+\pi^-\pi^0)_{\text{non-}\omega})/\Gamma_{\text{total}}$						Γ_{28}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT		
$0.52 \pm 0.15 \pm 0.11$	24	25 HE	14	BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$	

$\Gamma(\chi_{b2}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$						Γ_{29}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT		
$1.17 \pm 0.27 \pm 0.14$	29	25 HE	14	BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$	

$\Gamma(\chi_{b2}(1P)\omega)/\Gamma_{\text{total}}$						Γ_{30}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT		
$0.60 \pm 0.23 \pm 0.15$	13	25 HE	14	BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$	

$\Gamma(\chi_{b2}(1P)\omega)/\Gamma(\chi_{b1}(1P)\omega)$						Γ_{30}/Γ_{27}
VALUE	DOCUMENT ID	TECN	COMMENT			

• • • We do not use the following data for averages, fits, limits, etc. • • •
 $0.38 \pm 0.16 \pm 0.09$ 26 HE 14 BELL $\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$

$\Gamma(\chi_{b2}(1P)(\pi^+\pi^-\pi^0)_{\text{non-}\omega})/\Gamma_{\text{total}}$						Γ_{31}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT		
$0.61 \pm 0.22 \pm 0.28$	16	25 HE	14	BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$	

$\Gamma(\chi_{b2}(1P)(\pi^+\pi^-\pi^0)_{\text{non-}\omega})/\Gamma(\chi_{b1}(1P)(\pi^+\pi^-\pi^0)_{\text{non-}\omega})$						Γ_{31}/Γ_{28}
VALUE	DOCUMENT ID	TECN	COMMENT			

• • • We do not use the following data for averages, fits, limits, etc. • • •
 $1.20 \pm 0.55 \pm 0.65$ 26 HE 14 BELL $\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$

$\Gamma(\gamma\chi_b \rightarrow \gamma\Upsilon(1S)\omega)/\Gamma_{\text{total}}$						Γ_{32}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<3.8 \times 10^{-5}$	90	27 HE	14	BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$	

$\Gamma(\phi \text{ anything})/\Gamma_{\text{total}}$						Γ_{33}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT			
$0.138 \pm 0.007^{+0.023}_{-0.015}$	HUANG	07	CLEO	$\Upsilon(5S) \rightarrow \phi X$		

$\Gamma(D^0 \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
1.076 ± 0.040 ± 0.068	DRUTSKOY 07	BELL	$\Upsilon(5S) \rightarrow D^0 X$

$\Gamma(D_s \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.46 ± 0.06 OUR AVERAGE			
0.472 ± 0.024 ± 0.072	¹⁸ DRUTSKOY 07	BELL	$\Upsilon(5S) \rightarrow D_s X$
0.44 ± 0.09 ± 0.04	²⁸ ARTUSO 05B	CLE3	$e^+ e^- \rightarrow D_s X$

$\Gamma(J/\psi \text{ anything})/\Gamma_{\text{total}}$ Γ_{36}/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
2.060 ± 0.160 ± 0.134	DRUTSKOY 07	BELL	$\Upsilon(5S) \rightarrow J/\psi X$

$\Gamma(B^0 \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.770^{+0.058}_{-0.056} ± 0.061	352	DRUTSKOY 10	BELL	$\Upsilon(5S) \rightarrow B^0 X$

$\Gamma(B^+ \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.721^{+0.039}_{-0.038} ± 0.050	711	DRUTSKOY 10	BELL	$\Upsilon(5S) \rightarrow B^+ X$

¹³ Not independent of DRUTSKOY 10 values for $\Upsilon(5S) \rightarrow B^{\pm,0}$ anything.

¹⁴ Using measurements or limits from AQUINES 06.

¹⁵ Assuming isospin conservation.

¹⁶ Supersedes DRUTSKOY 07.

¹⁷ Supersedes ARTUSO 05B. Combining inclusive ϕ , D_s , and B measurements. Using $B(D_s^+ \rightarrow \phi\pi^+) = 4.4 \pm 0.6\%$ from PDG 06.

¹⁸ Using $B(D_s^+ \rightarrow \phi\pi^+) = (4.4 \pm 0.6)\%$ from PDG 06.

¹⁹ Uses a model-dependent estimate $B(B_s \rightarrow D_s X) = (92 \pm 11)\%$.

²⁰ Supersedes LOUVOT 09.

²¹ With $N(B_s^{(*)}\bar{B}_s^{(*)}) = (7.11 \pm 1.30) \times 10^6$.

²² The ratios $N(B_s^*\bar{B}_s^*) / N(B_s^{(*)}\bar{B}_s^{(*)})$ and $N(B_s^*\bar{B}_s^0) / N(B_s^{(*)}\bar{B}_s^{(*)})$ are measured with a correlation coefficient of -0.72 .

²³ From a measurement of $\sigma(e^+e^- \rightarrow B_s^*\bar{B}_s^*) / \sigma(e^+e^- \rightarrow B_s^{(*)}\bar{B}_s^{(*)})$ at $\sqrt{s} = 10.86$ GeV.

²⁴ Assuming that the observed events are solely due to the $\Upsilon(5S)$ resonance.

²⁵ Assuming that all the $b\bar{b}$ events are from $\Upsilon(5S)$ resonance decays and using $\sigma(e^+e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016$ nb from ESEN 13. Correlated with other results from HE 14.

²⁶ Accounting for correlated systematics.

²⁷ Assuming that all the $b\bar{b}$ events are from $\Upsilon(5S)$ resonance decays and using $\sigma(e^+e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016$ nb from ESEN 13. Correlated with other results from HE 14. For a state X_b with mass between $10.55 \text{ GeV}/c^2$ and $10.65 \text{ GeV}/c^2$, the obtained 90% upper limit as a function of m_{X_b} varies from 2.6×10^{-5} to 3.8×10^{-5} .

²⁸ ARTUSO 05B reports $[\Gamma(\Upsilon(10860) \rightarrow D_s \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)] = 0.0198 \pm 0.0019 \pm 0.0038$ which we divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Upsilon(10860)$ REFERENCES

HE	14	PRL 113 142001	X.H. He <i>et al.</i>	(BELLE Collab.)
ESEN	13	PR D87 031101	S. Esen <i>et al.</i>	(BELLE Collab.)
SHEN	13A	PR D88 052019	C.P. Shen <i>et al.</i>	(BELLE Collab.)
ADACHI	12	PRL 108 032001	I. Adachi <i>et al.</i>	(BELLE Collab.)
LI	12	PRL 108 181808	J. Li <i>et al.</i>	(BELLE Collab.)
CHEN	10	PR D82 091106	K.-F. Chen <i>et al.</i>	(BELLE Collab.)
DRUTSKOY	10	PR D81 112003	A. Drutskoy <i>et al.</i>	(BELLE Collab.)
AUBERT	09E	PRL 102 012001	B. Aubert <i>et al.</i>	(BABAR Collab.)
LOUVOT	09	PRL 102 021801	R. Louvot <i>et al.</i>	(BELLE Collab.)
CHEN	08	PRL 100 112001	K.-F. Chen <i>et al.</i>	(BELLE Collab.)
DRUTSKOY	07	PRL 98 052001	A. Drutskoy <i>et al.</i>	(BELLE Collab.)
DRUTSKOY	07A	PR D76 012002	A. Drutskoy <i>et al.</i>	(BELLE Collab.)
HUANG	07	PR D75 012002	G.S. Huang <i>et al.</i>	(CLEO Collab.)
AQUINES	06	PRL 96 152001	O. Aquines <i>et al.</i>	(CLEO Collab.)
BONVICINI	06	PRL 96 022002	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)
ARTUSO	05B	PRL 95 261801	M. Artuso <i>et al.</i>	(CLEO Collab.)
BESSON	85	PRL 54 381	D. Besson <i>et al.</i>	(CLEO Collab.)
LOVELOCK	85	PRL 54 377	D.M.J. Lovelock <i>et al.</i>	(CUSB Collab.)
