

**$f_2(2150)$** 

$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

This entry was previously called  $T_0$ . **$f_2(2150)$  MASS** **$f_2(2150)$  MASS, COMBINED MODES (MeV)**

| <u>VALUE (MeV)</u>         | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>  |
|----------------------------|-------------|--------------------|-------------|---|
| <b>2157±12 OUR AVERAGE</b> |             |                    |             | Includes data from the 2 datablocks that follow this one. |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|        |     |                   |         |  |
|--------|-----|-------------------|---------|--|
| 2170±6 | 80k | <sup>1</sup> UMAN | 06 E835 | 5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$ |
|--------|-----|-------------------|---------|--|

<sup>1</sup> Statistical error only.

 **$\eta\eta$  MODE**

| <u>VALUE (MeV)</u>         | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>  |
|----------------------------|--------------------|-------------|---|
| <b>2157±12 OUR AVERAGE</b> |                    |             | The data in this block is included in the average printed for a previous datablock. |

**2157±12 OUR AVERAGE**

|         |            |     |   |
|---------|------------|-----|---|
| 2151±16 | BARBERIS   | 00E | 450 $p p \rightarrow p_f \eta \eta p_s$   |
| 2175±20 | PROKOSHKIN | 95D | GAM4 300 $\pi^- N \rightarrow \pi^- N 2\eta$ ,<br>450 $p p \rightarrow p p 2\eta$ |
| 2130±35 | SINGOVSKI  | 94  | GAM4 450 $p p \rightarrow p p 2\eta$  |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|         |                        |     |   |
|---------|------------------------|-----|---|
| 2140±30 | <sup>2</sup> ABELE     | 99B | CBAR  |
| 2104±20 | <sup>3</sup> ARMSTRONG | 93C | E760 $\bar{p}p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$ |

<sup>2</sup> Spin not determined.<sup>3</sup> No  $J^{PC}$  determination. **$\eta\pi\pi$  MODE**

| <u>VALUE (MeV)</u> | <u>DOCUMENT ID</u>   | <u>TECN</u> | <u>CHG</u> | <u>COMMENT</u>                          |
|--------------------|----------------------|-------------|------------|---|
| <b>2135±20±45</b>  | <sup>4</sup> ADOMEIT | 96          | CBAR 0     | 1.94 $\bar{p}p \rightarrow \eta 3\pi^0$ |

The data in this block is included in the average printed for a previous datablock.

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>4</sup> ANISOVICH 00E recommends to withdraw ADOMEIT 96 that assumed a single  $J^P = 2^+$  resonance.

 **$\bar{p}p \rightarrow \pi\pi$** 

| <u>VALUE (MeV)</u> | <u>DOCUMENT ID</u>  | <u>TECN</u> | <u>COMMENT</u>                               |
|--------------------|---------------------|-------------|--|
| <b>~ 2090</b>      | <sup>5</sup> OAKDEN | 94          | RVUE 0.36–1.55 $\bar{p}p \rightarrow \pi\pi$ |
| <b>~ 2120</b>      | <sup>6</sup> OAKDEN | 94          | RVUE 0.36–1.55 $\bar{p}p \rightarrow \pi\pi$ |
| <b>~ 2170</b>      | <sup>7</sup> MARTIN | 80B         | RVUE   |
| <b>~ 2150</b>      | <sup>7</sup> MARTIN | 80C         | RVUE   |
| <b>~ 2150</b>      | <sup>8</sup> DULUDE | 78B         | OSPK 1–2 $\bar{p}p \rightarrow \pi^0 \pi^0$  |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>5</sup> OAKDEN 94 makes an amplitude analysis of LEAR data on  $\bar{p}p \rightarrow \pi\pi$  using a method based on Barrelet zeros. This is solution A. The amplitude analysis of HASAN 94 includes earlier data as well, and assume that the data can be parametrized in terms of towers of nearly degenerate resonances on the leading Regge trajectory. See also KLOET 96 and MARTIN 97 who make related analyses.

<sup>6</sup> From solution B of amplitude analysis of data on  $\bar{p}p \rightarrow \pi\pi$ .

<sup>7</sup>  $I(J^P) = 0(2^+)$  from simultaneous analysis of  $p\bar{p} \rightarrow \pi^- \pi^+$  and  $\pi^0 \pi^0$ .

<sup>8</sup>  $I^G(J^P) = 0^+(2^+)$  from partial-wave amplitude analysis.

### S-CHANNEL $\bar{p}p$ , $\bar{N}N$ or $\bar{K}K$

| VALUE (MeV)   | DOCUMENT ID               | TECN | CHG    | COMMENT                                    |
|---|---------------------------|------|--------|--|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                           |      |        |  |
| $2139^{+8}_{-9}$  | <sup>9</sup> EVANGELIS... | 97   | SPEC   | 0.6-2.4 $\bar{p}p \rightarrow K_S^0 K_S^0$ |
| $\sim 2190$   | <sup>9</sup> CUTTS        | 78B  | CNTR   | 0.97-3 $\bar{p}p \rightarrow \bar{N}N$     |
| $2155 \pm 15$   | <sup>9,10</sup> COUPLAND  | 77   | CNTR 0 | 0.7-2.4 $\bar{p}p \rightarrow \bar{p}p$    |
| $2193 \pm 2$  | <sup>9,11</sup> ALSPECTOR | 73   | CNTR   | $\bar{p}p$ S channel                       |

<sup>9</sup> Isospins 0 and 1 not separated.

<sup>10</sup> From a fit to the total elastic cross section.

<sup>11</sup> Referred to as  $T$  or  $T$  region by ALSPECTOR 73.

### K $\bar{K}$ MODE

| VALUE (MeV)   | DOCUMENT ID     | TECN | COMMENT                                   |
|---|-----------------|------|---|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                 |      |   |
| $2200 \pm 13$   | VLADIMIRSK...06 | SPEC | 40 $\pi^- p \rightarrow K_S^0 K_S^0 n$    |
| $2150 \pm 20$   | ABLIKIM         | 04E  | BES2 $J/\psi \rightarrow \omega K^+ K^-$  |
| $2130 \pm 35$   | BARBERIS        | 99   | OMEG 450 $pp \rightarrow p_s p_f K^+ K^-$ |

### $f_2(2150)$ WIDTH

#### $f_2(2150)$ WIDTH, COMBINED MODES (MeV)


| VALUE (MeV)               | EVTS | DOCUMENT ID | TECN | COMMENT   |
|---------------------------|------|-------------|------|---|
| <b>152±30 OUR AVERAGE</b> |      |             |      | Includes data from the 2 datablocks that follow this one. Error includes scale factor of 1.4. See the ideogram below. |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|              |     |                    |    |      |  |
|--------------|-----|--------------------|----|------|--|
| $182 \pm 11$ | 80k | <sup>12</sup> UMAN | 06 | E835 | 5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$ |
|--------------|-----|--------------------|----|------|--|

<sup>12</sup> Statistical error only.

WEIGHTED AVERAGE  
152±30 (Error scaled by 1.4)



$f_2(2150)$  WIDTH, COMBINED MODES (MeV)

## $\eta\eta$ MODE

VALUE (MeV)                      DOCUMENT ID      TECN      COMMENT  
 The data in this block is included in the average printed for a previous datablock.


**152±30 OUR AVERAGE** Error includes scale factor of 1.4. See the ideogram below.

|        |   |     |      |  |
|--------|---|-----|------|--|
| 280±70 | BARBERIS  | 00E |      | 450 $p p \rightarrow p_f \eta \eta p_s$                                      |
| 150±35 | PROKOSHKIN  | 95D | GAM4 | 300 $\pi^- N \rightarrow \pi^- N 2\eta$ ,<br>450 $p p \rightarrow p p 2\eta$ |
| 130±30 | SINGOVSKI   | 94  | GAM4 | 450 $p p \rightarrow p p 2\eta$  |
| • • •  | We do not use the following data for averages, fits, limits, etc. • • • |     |      |  |
| 310±50 | <sup>13</sup> ABELE   | 99B | CBAR |  |
| 203±10 | <sup>14</sup> ARMSTRONG   | 93C | E760 | $\bar{p} p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$                  |

<sup>13</sup> Spin not determined.

<sup>14</sup> No  $J^{PC}$  determination.

WEIGHTED AVERAGE  
 152±30 (Error scaled by 1.4)



## $\eta\pi\pi$ MODE

VALUE (MeV)                      DOCUMENT ID      TECN      CHG      COMMENT  
 The data in this block is included in the average printed for a previous datablock.

• • • We do not use the following data for averages, fits, limits, etc. • • •

|           |                         |     |      |   |  |
|-----------|-------------------------|-----|------|---|--|
| 250±25±45 | <sup>15</sup> ADOMEIT   | 96  | CBAR | 0 | 1.94 $\bar{p} p \rightarrow \eta 3\pi^0$                                       |
|           | <sup>15</sup> ANISOVICH | 00E |      |   | recommends to withdraw ADOMEIT 96 that assumed a single $J^P = 2^+$ resonance. |

### $\bar{p}p \rightarrow \pi\pi$

| VALUE (MeV)   | DOCUMENT ID          | TECN | COMMENT                                      |
|---|----------------------|------|--|
| <b>250 OUR ESTIMATE</b>   |                      |      |  |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |                      |      |  |
| ~ 70  | <sup>16</sup> OAKDEN | 94   | RVUE 0.36–1.55 $\bar{p}p \rightarrow \pi\pi$ |
| ~ 250   | <sup>17</sup> MARTIN | 80B  | RVUE   |
| ~ 250   | <sup>17</sup> MARTIN | 80C  | RVUE   |
| ~ 250   | <sup>18</sup> DULUDE | 78B  | OSPK 1–2 $\bar{p}p \rightarrow \pi^0\pi^0$   |

<sup>16</sup>See however KLOET 96 who fit  $\pi^+\pi^-$  only and find waves only up to  $J = 3$  to be important but not significantly resonant.

<sup>17</sup> $I(J^P) = 0(2^+)$  from simultaneous analysis of  $p\bar{p} \rightarrow \pi^-\pi^+$  and  $\pi^0\pi^0$ .

<sup>18</sup> $I^G(J^P) = 0^+(2^+)$  from partial-wave amplitude analysis.

### S-CHANNEL $\bar{p}p, \bar{N}N$ or $\bar{K}K$

| VALUE (MeV)   | DOCUMENT ID                | TECN | CHG    | COMMENT                                    |
|---|----------------------------|------|--------|--|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |                            |      |        |  |
| $56^{+31}_{-16}$  | <sup>19</sup> EVANGELIS... | 97   | SPEC   | 0.6–2.4 $\bar{p}p \rightarrow K_S^0 K_S^0$ |
| $135 \pm 75$  | <sup>20,21</sup> COUPLAND  | 77   | CNTR 0 | 0.7–2.4 $\bar{p}p \rightarrow \bar{p}p$    |
| $98 \pm 8$  | <sup>21</sup> ALSPECTOR    | 73   | CNTR   | $\bar{p}p$ S channel                       |

<sup>19</sup>Isospin 0 and 2 not separated.

<sup>20</sup>From a fit to the total elastic cross section.

<sup>21</sup>Isospins 0 and 1 not separated.

### $K\bar{K}$ MODE

| VALUE (MeV)   | DOCUMENT ID     | TECN | COMMENT   |
|---|-----------------|------|---|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |                 |      |   |
| $91 \pm 62$   | VLADIMIRSK...06 | SPEC | 40 $\pi^- p \rightarrow K_S^0 K_S^0 n$          |
| $150 \pm 30$  | ABLIKIM         | 04E  | BES2 $J/\psi \rightarrow \omega K^+ K^-$        |
| $270 \pm 50$  | BARBERIS        | 99   | OMEG 450 $p\bar{p} \rightarrow p_S p_f K^+ K^-$ |

## $f_2(2150)$ DECAY MODES

| Mode                       | Fraction ( $\Gamma_i/\Gamma$ ) |
|----------------------------|--------------------------------|
| $\Gamma_1$ $\pi\pi$        |                                |
| $\Gamma_2$ $\eta\eta$      | seen                           |
| $\Gamma_3$ $K\bar{K}$      | seen                           |
| $\Gamma_4$ $f_2(1270)\eta$ | seen                           |
| $\Gamma_5$ $a_2(1320)\pi$  | seen                           |
| $\Gamma_6$ $p\bar{p}$      | seen                           |

## $f_2(2150)$ BRANCHING RATIOS

### $\Gamma(K\bar{K})/\Gamma(\eta\eta)$

### $\Gamma_3/\Gamma_2$

| VALUE              | CL% | DOCUMENT ID | TECN | COMMENT                                      |
|--------------------|-----|-------------|------|--|
| <b>1.28 ± 0.23</b> |     | BARBERIS    | 00E  | 450 $p\bar{p} \rightarrow p_f \eta \eta p_S$ |

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

|      |    |                              |      |  |
|------|----|------------------------------|------|--|
| <0.1 | 95 | <sup>22</sup> PROKOSHKIN 95D | GAM4 | 300 $\pi^- N \rightarrow \pi^- N 2\eta,$<br>450 $p\bar{p} \rightarrow p p 2\eta$ |
|------|----|------------------------------|------|--|

<sup>22</sup>Using data from ARMSTRONG 89D.

| $\Gamma(\pi\pi)/\Gamma(\eta\eta)$ |     |                              |      |   | $\Gamma_1/\Gamma_2$ |
|-----------------------------------|-----|------------------------------|------|---|---------------------|
| VALUE                             | CL% | DOCUMENT ID                  | TECN | COMMENT   |                     |
| <0.33                             | 95  | <sup>23</sup> PROKOSHKIN 95D | GAM4 | 300 $\pi^- N \rightarrow \pi^- N 2\eta$ ,<br>450 $pp \rightarrow p p 2\eta$ |                     |

<sup>23</sup> Derived from a  $\pi^0\pi^0/\eta\eta$  limit.

| $\Gamma(f_2(1270)\eta)/\Gamma(a_2(1320)\pi)$ |  |                          |      |   | $\Gamma_4/\Gamma_5$ |
|--|--|--------------------------|------|---|---------------------|
| VALUE  |  | DOCUMENT ID              | TECN | COMMENT                                 |                     |
| <b>0.79±0.11</b>                             |  | <sup>24</sup> ADOMEIT 96 | CBAR | 1.94 $\bar{p}p \rightarrow \eta 3\pi^0$ |                     |

<sup>24</sup> Using  $B(a_2(1320) \rightarrow \eta\pi) = 0.145$

| $\Gamma(p\bar{p})/\Gamma_{\text{total}}$ |      |              |      |  | $\Gamma_6/\Gamma$ |
|--|------|--------------|------|--|-------------------|
| VALUE                                    | EVTS | DOCUMENT ID  | TECN | COMMENT                                |                   |
| <b>seen</b>                              | 73   | ALEXANDER 10 | CLEO | $\psi(2S) \rightarrow \gamma p\bar{p}$ |                   |

### $f_2(2150)$ REFERENCES

|                  |                               |                                  |                          |
|------------------|-------------------------------|----------------------------------|--------------------------|
| ALEXANDER 10     | PR D82 092002                 | J.P. Alexander <i>et al.</i>     | (CLEO Collab.)           |
| UMAN 06          | PR D73 052009                 | I. Uman <i>et al.</i>            | (FNAL E835)              |
| VLADIMIRSK... 06 | PAN 69 493                    | V.V. Vladimirovsky <i>et al.</i> | (ITEP, Moscow)           |
|                  | Translated from YAF 69 515.   |                                  |                          |
| ABLIKIM 04E      | PL B603 138                   | M. Ablikim <i>et al.</i>         | (BES Collab.)            |
| ANISOVICH 00E    | PL B477 19                    | A.V. Anisovich <i>et al.</i>     |                          |
| BARBERIS 00E     | PL B479 59                    | D. Barberis <i>et al.</i>        | (WA 102 Collab.)         |
| ABELE 99B        | EPJ C8 67                     | A. Abele <i>et al.</i>           | (Crystal Barrel Collab.) |
| BARBERIS 99      | PL B453 305                   | D. Barberis <i>et al.</i>        | (Omega Expt.)            |
| EVANGELIS... 97  | PR D56 3803                   | C. Evangelista <i>et al.</i>     | (LEAR Collab.)           |
| MARTIN 97        | PR C56 1114                   | B.R. Martin, G.C. Oades          | (LOUC, AARH)             |
| ADOMEIT 96       | ZPHY C71 227                  | J. Adomeit <i>et al.</i>         | (Crystal Barrel Collab.) |
| KLOET 96         | PR D53 6120                   | W.M. Kloet, F. Myhrer            | (RUTG, NORD)             |
| PROKOSHKIN 95D   | SPD 40 495                    | Y.D. Prokoshkin                  | (SERP) IGJPC             |
|                  | Translated from DANS 344 469. |                                  |                          |
| HASAN 94         | PL B334 215                   | A. Hasan, D.V. Bugg              | (LOQM)                   |
| OAKDEN 94        | NP A574 731                   | M.N. Oakden, M.R. Pennington     | (DURH)                   |
| SINGOVSKI 94     | NC 107A 1911                  | A.V. Singovsky                   | (SERP)                   |
| ARMSTRONG 93C    | PL B307 394                   | T.A. Armstrong <i>et al.</i>     | (FNAL, FERR, GENO+)      |
| ARMSTRONG 89D    | PL B227 186                   | T.A. Armstrong, M. Benayoun      | (ATHU, BARI, BIRM+)      |
| MARTIN 80B       | NP B176 355                   | B.R. Martin, D. Morgan           | (LOUC, RHEL) JP          |
| MARTIN 80C       | NP B169 216                   | A.D. Martin, M.R. Pennington     | (DURH) JP                |
| CUTTS 78B        | PR D17 16                     | D. Cutts <i>et al.</i>           | (STON, WISC)             |
| DULUDE 78B       | PL 79B 335                    | R.S. Dulude <i>et al.</i>        | (BROW, MIT, BARI) JP     |
| COUPLAND 77      | PL 71B 460                    | M. Coupland <i>et al.</i>        | (LOQM, RHEL)             |
| ALSPECTOR 73     | PRL 30 511                    | J. Alspector <i>et al.</i>       | (RUTG, UPNJ)             |