

CHARMED BARYONS ($C = +1$)

$$\Lambda_c^+ = udc, \quad \Sigma_c^{++} = uuc, \quad \Sigma_c^+ = udc, \quad \Sigma_c^0 = ddc,$$

$$\Xi_c^+ = usc, \quad \Xi_c^0 = dsc, \quad \Omega_c^0 = ssc$$

Λ_c^+

$$I(J^P) = 0(\frac{1}{2}^+)$$

J is not well measured; $\frac{1}{2}$ is the quark-model prediction.

Mass $m = 2286.46 \pm 0.14$ MeV

Mean life $\tau = (200 \pm 6) \times 10^{-15}$ s ($S = 1.6$)

$c\tau = 59.9$ μ m

Decay asymmetry parameters

$\Lambda\pi^+$ $\alpha = -0.91 \pm 0.15$

$\Sigma^+\pi^0$ $\alpha = -0.45 \pm 0.32$

$\Lambda\ell^+\nu_\ell$ $\alpha = -0.86 \pm 0.04$

$(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha})$ in $\Lambda_c^+ \rightarrow \Lambda\pi^+, \bar{\Lambda}_c^- \rightarrow \bar{\Lambda}\pi^- = -0.07 \pm 0.31$

$(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha})$ in $\Lambda_c^+ \rightarrow \Lambda e^+\nu_e, \bar{\Lambda}_c^- \rightarrow \bar{\Lambda}e^-\bar{\nu}_e = 0.00 \pm 0.04$

Λ_c^+ DECAY MODES

Scale factor/
Confidence level p
(MeV/c)

Hadronic modes with a p : $S = -1$ final states

pK_S^0	(1.58 \pm 0.08) %	S=1.2	873
$pK^-\pi^+$	(6.35 \pm 0.33) %	S=1.4	823
$p\bar{K}^*(892)^0$	[a] (1.98 \pm 0.28) %		685
$\Delta(1232)^{++}K^-$	(1.09 \pm 0.25) %		710
$\Lambda(1520)\pi^+$	[a] (2.2 \pm 0.5) %		627
$pK^-\pi^+$ nonresonant	(3.5 \pm 0.4) %		823
$pK_S^0\pi^0$	(1.99 \pm 0.13) %	S=1.1	823
$p\bar{K}^0\eta$	(1.6 \pm 0.4) %		568
$pK_S^0\pi^+\pi^-$	(1.66 \pm 0.12) %	S=1.1	754
$pK^-\pi^+\pi^0$	(4.9 \pm 0.4) %	S=1.3	759
$pK^*(892)^-\pi^+$	[a] (1.5 \pm 0.5) %		580
$p(K^-\pi^+)_{\text{nonresonant}}\pi^0$	(4.6 \pm 0.9) %		759
$\Delta(1232)\bar{K}^*(892)$	seen		419
$pK^-2\pi^+\pi^-$	(1.4 \pm 1.0) $\times 10^{-3}$		671
$pK^-\pi^+2\pi^0$	(1.0 \pm 0.5) %		678

Hadronic modes with a p : $S = 0$ final states

$p\pi^+\pi^-$		$(4.4 \pm 2.3) \times 10^{-3}$	927
$pf_0(980)$	[a]	$(3.5 \pm 2.3) \times 10^{-3}$	614
$p2\pi^+2\pi^-$		$(2.3 \pm 1.5) \times 10^{-3}$	852
pK^+K^-		$(10 \pm 4) \times 10^{-4}$	616
$p\phi$	[a]	$(1.04 \pm 0.21) \times 10^{-3}$	590
$pK^+K^- \text{ non-}\phi$		$(4.4 \pm 1.8) \times 10^{-4}$	616

Hadronic modes with a hyperon: $S = -1$ final states

$\Lambda\pi^+$		$(1.30 \pm 0.07) \%$	S=1.2	864
$\Lambda\pi^+\pi^0$		$(7.1 \pm 0.4) \%$	S=1.2	844
$\Lambda\rho^+$		$< 6 \%$	CL=95%	636
$\Lambda\pi^-2\pi^+$		$(3.7 \pm 0.4) \%$	S=1.9	807
$\Sigma(1385)^+\pi^+\pi^-, \Sigma^{*+} \rightarrow$ $\Lambda\pi^+$		$(1.0 \pm 0.5) \%$		688
$\Sigma(1385)^-2\pi^+, \Sigma^{*-} \rightarrow$ $\Lambda\pi^-$		$(7.8 \pm 1.6) \times 10^{-3}$		688
$\Lambda\pi^+\rho^0$		$(1.5 \pm 0.6) \%$		524
$\Sigma(1385)^+\rho^0, \Sigma^{*+} \rightarrow \Lambda\pi^+$		$(5 \pm 4) \times 10^{-3}$		363
$\Lambda\pi^-2\pi^+ \text{ nonresonant}$		$< 1.1 \%$	CL=90%	807
$\Lambda\pi^-\pi^02\pi^+ \text{ total}$		$(2.3 \pm 0.8) \%$		757
$\Lambda\pi^+\eta$	[a]	$(2.3 \pm 0.5) \%$		691
$\Sigma(1385)^+\eta$	[a]	$(1.08 \pm 0.32) \%$		570
$\Lambda\pi^+\omega$	[a]	$(1.5 \pm 0.5) \%$		517
$\Lambda\pi^-\pi^02\pi^+, \text{ no } \eta \text{ or } \omega$		$< 8 \times 10^{-3}$	CL=90%	757
$\Lambda K^+\bar{K}^0$		$(5.7 \pm 1.1) \times 10^{-3}$	S=2.0	443
$\Xi(1690)^0 K^+, \Xi^{*0} \rightarrow \Lambda\bar{K}^0$		$(1.6 \pm 0.5) \times 10^{-3}$		286
$\Sigma^0\pi^+$		$(1.29 \pm 0.07) \%$	S=1.1	825
$\Sigma^+\pi^0$		$(1.24 \pm 0.10) \%$		827
$\Sigma^+\eta$		$(7.0 \pm 2.3) \times 10^{-3}$		713
$\Sigma^+\pi^+\pi^-$		$(4.57 \pm 0.29) \%$	S=1.2	804
$\Sigma^+\rho^0$		$< 1.7 \%$	CL=95%	575
$\Sigma^-2\pi^+$		$(2.1 \pm 0.4) \%$		799
$\Sigma^0\pi^+\pi^0$		$(2.3 \pm 0.9) \%$		803
$\Sigma^0\pi^-2\pi^+$		$(1.13 \pm 0.29) \%$		763
$\Sigma^+\pi^+\pi^-\pi^0$		—		767
$\Sigma^+\omega$	[a]	$(1.74 \pm 0.21) \%$		569
$\Sigma^+K^+K^-$		$(3.6 \pm 0.4) \times 10^{-3}$		349
$\Sigma^+\phi$	[a]	$(4.0 \pm 0.6) \times 10^{-3}$	S=1.1	295
$\Xi(1690)^0 K^+, \Xi^{*0} \rightarrow$ Σ^+K^-		$(1.03 \pm 0.26) \times 10^{-3}$		286
$\Sigma^+K^+K^- \text{ nonresonant}$		$< 8 \times 10^{-4}$	CL=90%	349

$\Xi^0 K^+$	$(5.0 \pm 1.2) \times 10^{-3}$		653
$\Xi^- K^+ \pi^+$	$(6.2 \pm 0.6) \times 10^{-3}$	S=1.1	565
$\Xi(1530)^0 K^+$	[a] $(3.3 \pm 0.9) \times 10^{-3}$		473

Hadronic modes with a hyperon: $S = 0$ final states

ΛK^+	$(6.1 \pm 1.2) \times 10^{-4}$		781
$\Lambda K^+ \pi^+ \pi^-$	$< 5 \times 10^{-4}$	CL=90%	637
$\Sigma^0 K^+$	$(5.2 \pm 0.8) \times 10^{-4}$		735
$\Sigma^0 K^+ \pi^+ \pi^-$	$< 2.6 \times 10^{-4}$	CL=90%	574
$\Sigma^+ K^+ \pi^-$	$(2.1 \pm 0.6) \times 10^{-3}$		670
$\Sigma^+ K^*(892)^0$	[a] $(3.6 \pm 1.0) \times 10^{-3}$		470
$\Sigma^- K^+ \pi^+$	$< 1.2 \times 10^{-3}$	CL=90%	664

Doubly Cabibbo-suppressed modes

$p K^+ \pi^-$	$< 2.9 \times 10^{-4}$	CL=90%	823
---------------	------------------------	--------	-----

Semileptonic modes

$\Lambda e^+ \nu_e$	$(3.6 \pm 0.4) \%$		871
---------------------	----------------------	--	-----

Inclusive modes

e^+ anything	$(4.5 \pm 1.7) \%$		—
$p e^+$ anything	$(1.8 \pm 0.9) \%$		—
p anything	$(50 \pm 16) \%$		—
p anything (no Λ)	$(12 \pm 19) \%$		—
n anything	$(50 \pm 16) \%$		—
n anything (no Λ)	$(29 \pm 17) \%$		—
Λ anything	$(35 \pm 11) \%$	S=1.4	—
Σ^\pm anything	[b] $(10 \pm 5) \%$		—
3prongs	$(24 \pm 8) \%$		—

$\Delta C = 1$ weak neutral current ($C1$) modes, or Lepton Family number (LF), or Lepton number (L), or Baryon number (B) violating modes

$p e^+ e^-$	$C1$	$< 5.5 \times 10^{-6}$	CL=90%	951
$p \mu^+ \mu^-$	$C1$	$< 4.4 \times 10^{-5}$	CL=90%	937
$p e^+ \mu^-$	LF	$< 9.9 \times 10^{-6}$	CL=90%	947
$p e^- \mu^+$	LF	$< 1.9 \times 10^{-5}$	CL=90%	947
$\bar{p} 2e^+$	L, B	$< 2.7 \times 10^{-6}$	CL=90%	951
$\bar{p} 2\mu^+$	L, B	$< 9.4 \times 10^{-6}$	CL=90%	937
$\bar{p} e^+ \mu^+$	L, B	$< 1.6 \times 10^{-5}$	CL=90%	947
$\Sigma^- \mu^+ \mu^+$	L	$< 7.0 \times 10^{-4}$	CL=90%	812

$\Lambda_c(2595)^+$

$$I(J^P) = 0(\frac{1}{2}^-)$$

The spin-parity follows from the fact that $\Sigma_c(2455)\pi$ decays, with little available phase space, are dominant. This assumes that $J^P = 1/2^+$ for the $\Sigma_c(2455)$.

$$\text{Mass } m = 2592.25 \pm 0.28 \text{ MeV}$$

$$m - m_{\Lambda_c^+} = 305.79 \pm 0.24 \text{ MeV}$$

$$\text{Full width } \Gamma = 2.6 \pm 0.6 \text{ MeV}$$

$\Lambda_c^+ \pi \pi$ and its submode $\Sigma_c(2455)\pi$ — the latter just barely — are the only strong decays allowed to an excited Λ_c^+ having this mass; and the submode seems to dominate.

$\Lambda_c(2595)^+$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_c^+ \pi^+ \pi^-$	[c] —	117
$\Sigma_c(2455)^{++} \pi^-$	$24 \pm 7 \%$	†
$\Sigma_c(2455)^0 \pi^+$	$24 \pm 7 \%$	†
$\Lambda_c^+ \pi^+ \pi^-$ 3-body	$18 \pm 10 \%$	117
$\Lambda_c^+ \pi^0$	[d] not seen	258
$\Lambda_c^+ \gamma$	not seen	288

 $\Lambda_c(2625)^+$

$$I(J^P) = 0(\frac{3}{2}^-)$$

J^P has not been measured; $\frac{3}{2}^-$ is the quark-model prediction.

$$\text{Mass } m = 2628.11 \pm 0.19 \text{ MeV} \quad (S = 1.1)$$

$$m - m_{\Lambda_c^+} = 341.65 \pm 0.13 \text{ MeV} \quad (S = 1.1)$$

$$\text{Full width } \Gamma < 0.97 \text{ MeV, CL} = 90\%$$

$\Lambda_c^+ \pi \pi$ and its submode $\Sigma(2455)\pi$ are the only strong decays allowed to an excited Λ_c^+ having this mass.

$\Lambda_c(2625)^+$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$\Lambda_c^+ \pi^+ \pi^-$	[c] $\approx 67\%$		184
$\Sigma_c(2455)^{++} \pi^-$	< 5	90%	102
$\Sigma_c(2455)^0 \pi^+$	< 5	90%	102
$\Lambda_c^+ \pi^+ \pi^-$ 3-body	large		184
$\Lambda_c^+ \pi^0$	[d] not seen		293
$\Lambda_c^+ \gamma$	not seen		319

$\Lambda_c(2880)^+$

$$I(J^P) = 0(\frac{5}{2}^+)$$

There is some good evidence that indeed $J^P = 5/2^+$

Mass $m = 2881.53 \pm 0.35$ MeV

$m - m_{\Lambda_c^+} = 595.1 \pm 0.4$ MeV

Full width $\Gamma = 5.8 \pm 1.1$ MeV

$\Lambda_c(2880)^+$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_c^+ \pi^+ \pi^-$	seen	471
$\Sigma_c(2455)^{0,++} \pi^\pm$	seen	376
$\Sigma_c(2520)^{0,++} \pi^\pm$	seen	317
pD^0	seen	316

 $\Lambda_c(2940)^+$

$$I(J^P) = 0(?^?)$$

Mass $m = 2939.3^{+1.4}_{-1.5}$ MeV

Full width $\Gamma = 17^{+8}_{-6}$ MeV

$\Lambda_c(2940)^+$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
pD^0	seen	420
$\Sigma_c(2455)^{0,++} \pi^\pm$	seen	—

 $\Sigma_c(2455)$

$$I(J^P) = 1(\frac{1}{2}^+)$$

$\Sigma_c(2455)^{++}$ mass $m = 2453.97 \pm 0.14$ MeV

$\Sigma_c(2455)^+$ mass $m = 2452.9 \pm 0.4$ MeV

$\Sigma_c(2455)^0$ mass $m = 2453.75 \pm 0.14$ MeV

$m_{\Sigma_c^{++}} - m_{\Lambda_c^+} = 167.510 \pm 0.017$ MeV

$m_{\Sigma_c^+} - m_{\Lambda_c^+} = 166.4 \pm 0.4$ MeV

$m_{\Sigma_c^0} - m_{\Lambda_c^+} = 167.290 \pm 0.017$ MeV

$m_{\Sigma_c^{++}} - m_{\Sigma_c^0} = 0.220 \pm 0.013$ MeV

$m_{\Sigma_c^+} - m_{\Sigma_c^0} = -0.9 \pm 0.4$ MeV

$\Sigma_c(2455)^{++}$ full width $\Gamma = 1.89^{+0.09}_{-0.18}$ MeV ($S = 1.1$)

$\Sigma_c(2455)^+$ full width $\Gamma < 4.6$ MeV, CL = 90%

$\Sigma_c(2455)^0$ full width $\Gamma = 1.83^{+0.11}_{-0.19}$ MeV ($S = 1.2$)

 $\Lambda_c^+ \pi$ is the only strong decay allowed to a Σ_c having this mass.

$\Sigma_c(2455)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_c^+ \pi$	$\approx 100\%$	94

$\Sigma_c(2520)$

$$I(J^P) = 1(\frac{3}{2}^+)$$

J^P has not been measured; $\frac{3}{2}^+$ is the quark-model prediction.

$$\Sigma_c(2520)^{++} \text{ mass } m = 2518.41^{+0.21}_{-0.19} \text{ MeV } (S = 1.1)$$

$$\Sigma_c(2520)^+ \text{ mass } m = 2517.5 \pm 2.3 \text{ MeV}$$

$$\Sigma_c(2520)^0 \text{ mass } m = 2518.48 \pm 0.20 \text{ MeV } (S = 1.1)$$

$$m_{\Sigma_c(2520)^{++}} - m_{\Lambda_c^+} = 231.95^{+0.17}_{-0.12} \text{ MeV } (S = 1.3)$$

$$m_{\Sigma_c(2520)^+} - m_{\Lambda_c^+} = 231.0 \pm 2.3 \text{ MeV}$$

$$m_{\Sigma_c(2520)^0} - m_{\Lambda_c^+} = 232.02^{+0.15}_{-0.14} \text{ MeV } (S = 1.3)$$

$$m_{\Sigma_c(2520)^{++}} - m_{\Sigma_c(2520)^0} = 0.01 \pm 0.15 \text{ MeV}$$

$$\Sigma_c(2520)^{++} \text{ full width } \Gamma = 14.78^{+0.30}_{-0.40} \text{ MeV}$$

$$\Sigma_c(2520)^+ \text{ full width } \Gamma < 17 \text{ MeV, CL} = 90\%$$

$$\Sigma_c(2520)^0 \text{ full width } \Gamma = 15.3^{+0.4}_{-0.5} \text{ MeV}$$

$\Lambda_c^+ \pi$ is the only strong decay allowed to a Σ_c having this mass.

$\Sigma_c(2520)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_c^+ \pi$	$\approx 100\%$	179

$\Sigma_c(2800)$

$$I(J^P) = 1(?^?)$$

$$\Sigma_c(2800)^{++} \text{ mass } m = 2801^{+4}_{-6} \text{ MeV}$$

$$\Sigma_c(2800)^+ \text{ mass } m = 2792^{+14}_{-5} \text{ MeV}$$

$$\Sigma_c(2800)^0 \text{ mass } m = 2806^{+5}_{-7} \text{ MeV } (S = 1.3)$$

$$m_{\Sigma_c(2800)^{++}} - m_{\Lambda_c^+} = 514^{+4}_{-6} \text{ MeV}$$

$$m_{\Sigma_c(2800)^+} - m_{\Lambda_c^+} = 505^{+14}_{-5} \text{ MeV}$$


$$m_{\Sigma_c(2800)^0} - m_{\Lambda_c^+} = 519^{+5}_{-7} \text{ MeV } (S = 1.3)$$

$$\Sigma_c(2800)^{++} \text{ full width } \Gamma = 75^{+22}_{-17} \text{ MeV}$$

$$\Sigma_c(2800)^+ \text{ full width } \Gamma = 62^{+60}_{-40} \text{ MeV}$$

$$\Sigma_c(2800)^0 \text{ full width } \Gamma = 72^{+22}_{-15} \text{ MeV}$$

$\Sigma_c(2800)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_c^+ \pi$	seen	443



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

$$\text{Mass } m = 2467.93_{-0.40}^{+0.28} \text{ MeV}$$

$$\text{Mean life } \tau = (442 \pm 26) \times 10^{-15} \text{ s} \quad (S = 1.3)$$

$$c\tau = 132 \mu\text{m}$$

Ξ_c^+ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	ρ (MeV/c)
-----------------------	--------------------------------	------------------	-------------------

**No absolute branching fractions have been measured.
The following are branching *ratios* relative to $\Xi^- 2\pi^+$.**

Cabibbo-favored ($S = -2$) decays — relative to $\Xi^- 2\pi^+$

$p 2K_S^0$	0.087 ± 0.021		767
$\Lambda \bar{K}^0 \pi^+$	—		852
$\Sigma(1385)^+ \bar{K}^0$	[a] 1.0 ± 0.5		746
$\Lambda K^- 2\pi^+$	0.323 ± 0.033		787
$\Lambda \bar{K}^*(892)^0 \pi^+$	[a] < 0.16	90%	608
$\Sigma(1385)^+ K^- \pi^+$	[a] < 0.23	90%	678
$\Sigma^+ K^- \pi^+$	0.94 ± 0.10		811
$\Sigma^+ \bar{K}^*(892)^0$	[a] 0.81 ± 0.15		658
$\Sigma^0 K^- 2\pi^+$	0.27 ± 0.12		735
$\Xi^0 \pi^+$	0.55 ± 0.16		877
$\Xi^- 2\pi^+$	DEFINED AS 1		851
$\Xi(1530)^0 \pi^+$	[a] < 0.10	90%	750
$\Xi^0 \pi^+ \pi^0$	2.3 ± 0.7		856
$\Xi^0 \pi^- 2\pi^+$	1.7 ± 0.5		818
$\Xi^0 e^+ \nu_e$	$2.3 \begin{smallmatrix} +0.7 \\ -0.8 \end{smallmatrix}$		884
$\Omega^- K^+ \pi^+$	0.07 ± 0.04		399

Cabibbo-suppressed decays — relative to $\Xi^- 2\pi^+$

$p K^- \pi^+$	0.21 ± 0.04		944
$p \bar{K}^*(892)^0$	[a] 0.116 ± 0.030		828
$\Sigma^+ \pi^+ \pi^-$	0.48 ± 0.20		922
$\Sigma^- 2\pi^+$	0.18 ± 0.09		918
$\Sigma^+ K^+ K^-$	0.15 ± 0.06		580
$\Sigma^+ \phi$	[a] < 0.11	90%	549
$\Xi(1690)^0 K^+, \Xi^0 \rightarrow \Sigma^+ K^-$	< 0.05	90%	501



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

$$\text{Mass } m = 2470.85^{+0.28}_{-0.40} \text{ MeV}$$

$$m_{\Xi_c^0} - m_{\Xi_c^+} = 2.93 \pm 0.24 \text{ MeV}$$

$$\text{Mean life } \tau = (112^{+13}_{-10}) \times 10^{-15} \text{ s}$$

$$c\tau = 33.6 \mu\text{m}$$

Decay asymmetry parameters

$$\Xi^- \pi^+ \quad \alpha = -0.6 \pm 0.4$$

No absolute branching fractions have been measured. Several measurements of ratios of fractions may be found in the Listings that follow.

Ξ_c^0 DECAY MODES

Fraction (Γ_i/Γ)

p (MeV/c)


**No absolute branching fractions have been measured.
The following are branching *ratios* relative to $\Xi^- \pi^+$.**

Cabibbo-favored ($S = -2$) decays — relative to $\Xi^- \pi^+$

$pK^- K^- \pi^+$	0.34 ± 0.04	676
$pK^- \bar{K}^*(892)^0$	0.21 ± 0.05	413
$pK^- K^- \pi^+$ (no \bar{K}^{*0})	0.21 ± 0.04	676
ΛK_S^0	0.210 ± 0.028	906
$\Lambda K^- \pi^+$	1.07 ± 0.14	856
$\Lambda \bar{K}^0 \pi^+ \pi^-$	seen	787
$\Lambda K^- \pi^+ \pi^+ \pi^-$	seen	703
$\Xi^- \pi^+$	DEFINED AS 1	875
$\Xi^- \pi^+ \pi^+ \pi^-$	3.3 ± 1.4	816
$\Omega^- K^+$	0.297 ± 0.024	522
$\Xi^- e^+ \nu_e$	3.1 ± 1.1	882
$\Xi^- \ell^+$ anything	1.0 ± 0.5	—

Cabibbo-suppressed decays — relative to $\Xi^- \pi^+$

$\Xi^- K^+$	0.028 ± 0.006	790
$\Lambda K^+ K^-$ (no ϕ)	0.029 ± 0.007	648
$\Lambda \phi$	0.034 ± 0.007	621



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$


J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

$$\text{Mass } m = 2575.7 \pm 3.0 \text{ MeV}$$

$$m_{\Xi_c^{'+}} - m_{\Xi_c^+} = 107.8 \pm 3.0 \text{ MeV}$$

The $\Xi_c^{'+} - \Xi_c^+$ mass difference is too small for any strong decay to occur.

$\Xi_c^{'+}$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c^+ \gamma$	seen	106



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$


J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

Mass $m = 2577.9 \pm 2.9$ MeV

$$m_{\Xi_c^{'0}} - m_{\Xi_c^0} = 107.0 \pm 2.9 \text{ MeV}$$

The $\Xi_c^{'0} - \Xi_c^0$ mass difference is too small for any strong decay to occur.

$\Xi_c^{'0}$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c^0 \gamma$	seen	105



$$I(J^P) = \frac{1}{2}(\frac{3}{2}^+)$$

J^P has not been measured; $\frac{3}{2}^+$ is the quark-model prediction.

$\Xi_c(2645)^+$ mass $m = 2645.9 \pm 0.5$ MeV ($S = 1.1$)

$\Xi_c(2645)^0$ mass $m = 2645.9 \pm 0.5$ MeV

$$m_{\Xi_c(2645)^+} - m_{\Xi_c^0} = 175.0 \pm 0.6 \text{ MeV} \quad (S = 1.1)$$

$$m_{\Xi_c(2645)^0} - m_{\Xi_c^+} = 178.0 \pm 0.6 \text{ MeV}$$

$$m_{\Xi_c(2645)^+} - m_{\Xi_c(2645)^0} = 0.0 \pm 0.5 \text{ MeV}$$

$\Xi_c(2645)^+$ full width $\Gamma = 2.6 \pm 0.4$ MeV

$\Xi_c(2645)^0$ full width $\Gamma < 5.5$ MeV, CL = 90%

$\Xi_c \pi$ is the only strong decay allowed to a Ξ_c resonance having this mass.

$\Xi_c(2645)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c^0 \pi^+$	seen	102
$\Xi_c^+ \pi^-$	seen	107

$\Xi_c(2790)$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^-)$$

J^P has not been measured; $\frac{1}{2}^-$ is the quark-model prediction.

$$\Xi_c(2790)^+ \text{ mass} = 2789.1 \pm 3.2 \text{ MeV}$$

$$\Xi_c(2790)^0 \text{ mass} = 2791.9 \pm 3.3 \text{ MeV}$$

$$m_{\Xi_c(2790)^+} - m_{\Xi_c^0} = 318.2 \pm 3.2 \text{ MeV}$$

$$m_{\Xi_c(2790)^0} - m_{\Xi_c^+} = 324.0 \pm 3.3 \text{ MeV}$$

$$\Xi_c(2790)^+ \text{ width} < 15 \text{ MeV, CL} = 90\%$$

$$\Xi_c(2790)^0 \text{ width} < 12 \text{ MeV, CL} = 90\%$$

 $\Xi_c(2790)$ DECAY MODES

	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c^+ \pi$	seen	159

 $\Xi_c(2815)$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$$

J^P has not been measured; $\frac{3}{2}^-$ is the quark-model prediction.

$$\Xi_c(2815)^+ \text{ mass } m = 2816.6 \pm 0.9 \text{ MeV}$$

$$\Xi_c(2815)^0 \text{ mass } m = 2819.6 \pm 1.2 \text{ MeV}$$

$$m_{\Xi_c(2815)^+} - m_{\Xi_c^+} = 348.7 \pm 0.9 \text{ MeV}$$

$$m_{\Xi_c(2815)^0} - m_{\Xi_c^0} = 348.8 \pm 1.2 \text{ MeV}$$

$$m_{\Xi_c(2815)^+} - m_{\Xi_c(2815)^0} = -3.0 \pm 1.3 \text{ MeV}$$

$$\Xi_c(2815)^+ \text{ full width } \Gamma < 3.5 \text{ MeV, CL} = 90\%$$

$$\Xi_c(2815)^0 \text{ full width } \Gamma < 6.5 \text{ MeV, CL} = 90\%$$

The $\Xi_c \pi \pi$ modes are consistent with being entirely via $\Xi_c(2645) \pi$.

 $\Xi_c(2815)$ DECAY MODES

	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c^+ \pi^+ \pi^-$	seen	196
$\Xi_c^0 \pi^+ \pi^-$	seen	191

 $\Xi_c(2970)$ **was $\Xi_c(2980)$**

$$I(J^P) = \frac{1}{2}(?)^?$$

$$\Xi_c(2970)^+ \text{ } m = 2970.7 \pm 2.2 \text{ MeV } (S = 1.5)$$

$$\Xi_c(2970)^0 \text{ } m = 2968.0 \pm 2.6 \text{ MeV } (S = 1.2)$$

$$\Xi_c(2970)^+ \text{ width } \Gamma = 17.9 \pm 3.5 \text{ MeV}$$

$$\Xi_c(2970)^0 \text{ width } \Gamma = 20 \pm 7 \text{ MeV } (S = 1.3)$$

$\Xi_c(2970)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_c^+ \bar{K} \pi$	seen	231
$\Sigma_c(2455) \bar{K}$	seen	134
$\Lambda_c^+ \bar{K}$	not seen	414
$\Xi_c 2\pi$	seen	385
$\Xi_c(2645) \pi$	seen	277

 $\Xi_c(3055)$

$$I(J^P) = ?(??)$$

Mass $m = 3055.1 \pm 1.7$ MeV (S = 1.5)Full width $\Gamma = 11 \pm 4$ MeV **$\Xi_c(3080)$**

$$I(J^P) = \frac{1}{2}(??)$$

 $\Xi_c(3080)^+ m = 3076.94 \pm 0.28$ MeV $\Xi_c(3080)^0 m = 3079.9 \pm 1.4$ MeV (S = 1.3) $\Xi_c(3080)^+$ width $\Gamma = 4.3 \pm 1.5$ MeV (S = 1.3) $\Xi_c(3080)^0$ width $\Gamma = 5.6 \pm 2.2$ MeV

$\Xi_c(3080)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_c^+ \bar{K} \pi$	seen	415
$\Sigma_c(2455) \bar{K}$	seen	342
$\Sigma_c(2455) \bar{K} + \Sigma_c(2520) \bar{K}$	seen	—
$\Lambda_c^+ \bar{K}$	not seen	536
$\Lambda_c^+ \bar{K} \pi^+ \pi^-$	not seen	143

 Ω_c^0

$$I(J^P) = 0(\frac{1}{2}^+)$$

 J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.Mass $m = 2695.2 \pm 1.7$ MeV (S = 1.3)Mean life $\tau = (69 \pm 12) \times 10^{-15}$ s

$$c\tau = 21 \mu\text{m}$$

No absolute branching fractions have been measured.

Ω_c^0 DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Sigma^+ K^- K^- \pi^+$	seen	689
$\Xi^0 K^- \pi^+$	seen	901
$\Xi^- K^- \pi^+ \pi^+$	seen	830

$\Omega^- e^+ \nu_e$	seen	829
$\Omega^- \pi^+$	seen	821
$\Omega^- \pi^+ \pi^0$	seen	797
$\Omega^- \pi^- \pi^+ \pi^+$	seen	753

$\Omega_c(2770)^0$

$$I(J^P) = 0(\frac{3}{2}^+)$$

J^P has not been measured; $\frac{3}{2}^+$ is the quark-model prediction.

$$\text{Mass } m = 2765.9 \pm 2.0 \text{ MeV} \quad (S = 1.2)$$

$$m_{\Omega_c(2770)^0} - m_{\Omega_c^0} = 70.7^{+0.8}_{-0.9} \text{ MeV}$$

The $\Omega_c(2770)^0 - \Omega_c^0$ mass difference is too small for any strong decay to occur.

$\Omega_c(2770)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Omega_c^0 \gamma$	presumably 100%	70

NOTES

- [a] This branching fraction includes all the decay modes of the final-state resonance.
- [b] The value is for the sum of the charge states or particle/antiparticle states indicated.
- [c] See AALTONEN 11H, Fig. 8, for the calculated ratio of $\Lambda_c^+ \pi^0 \pi^0$ and $\Lambda_c^+ \pi^+ \pi^-$ partial widths as a function of the $\Lambda_c(2595)^+ - \Lambda_c^+$ mass difference. At our value of the mass difference, the ratio is about 4.
- [d] A test that the isospin is indeed 0, so that the particle is indeed a Λ_c^+ .