

95. Spectroscopy of Mesons Containing Two Heavy Quarks

Updated October 2017 by S. Eidelman (Budker Inst. and Novosibirsk State Univ.), C. Hanhart (Forschungszentrum Jülich), B.K. Heltsley (Cornell Univ.), J.J. Hernandez-Rey (Univ. Valencia–CSIC), R.E. Mitchell (Indiana Univ.), S. Navas (Univ. Granada), and C. Patrignani (Univ. Bologna, INFN).

A golden age for heavy quarkonium physics dawned at the turn of this century, initiated by the confluence of exciting advances in quantum chromodynamics (QCD) and an explosion of related experimental activity. The subsequent broad spectrum of breakthroughs, surprises, and continuing puzzles had not been anticipated. Since that time CLEO-c, BESIII and the B-factories, recently joined by ATLAS, CMS and LHCb, have continued to make groundbreaking observations. For an extensive presentation of the status of heavy quarkonium physics, the reader is referred to several reviews [1–13]. This note focuses on experimental developments in heavy quarkonium spectroscopy with very few theoretical comments. Some other comments on possible theoretical interpretations of the states not predicted by the quark model are presented in the minireview on non- $q\bar{q}$ states. Note that in this review we follow the new naming scheme for hadrons (see the review “Naming scheme for hadrons” in the current edition).

This minireview covers the newly discovered states, where “newly” is interpreted to include the period since 2002. In earlier versions of this write-up the particles were sorted according to an assumed *conventional* or *unconventional* nature with respect to the quark model. However, since this classification is not always unambiguous, we here follow Ref. [13] and sort the states into three groups, namely states below (*cf.* Table 95.1), near (*cf.* Table 95.2) and above (*cf.* Table 95.3) the lowest open flavor thresholds.

Table 95.1 lists properties of newly observed heavy quarkonium states located below the lowest open flavor thresholds. Those are expected to be (at least prominently) conventional quarkonia. The $h_c(1P)$ is the 1P_1 state of charmonium, singlet partner of the long-known χ_{cJ} triplet 3P_J . The $\eta_c(2S)$ is the first excited state of the pseudoscalar ground state $\eta_c(1S)$, lying just below the mass of its vector counterpart, $\psi(2S)$.

Although $\eta_c(2S)$ measurements began to converge towards a mass and a width some time ago, refinements are still in progress. In particular, Belle [21] has revisited its analysis of $B \rightarrow K\eta_c(2S)$, $\eta_c(2S) \rightarrow KK\pi$ decays with more data and methods that account for interference between the above decay chain, an equivalent one with the $\eta_c(1S)$ instead, and one with no intermediate resonance. The net effect of this interference is far from trivial; it shifts the apparent mass by $\sim +10$ MeV and blows up the apparent width by a factor of six. The updated $\eta_c(2S)$ mass and width are in better accordance with other measurements than the previous treatment [20], which did not include interference. Complementing this measurement in B -decay, BaBar [22] updated their previous [23] $\eta_c(2S)$ mass and width measurements in two-photon production, where interference effects, judging from studies of $\eta_c(1S)$, appear to be small. In combination, precision on the $\eta_c(2S)$ mass has improved dramatically.

2 95. Spectroscopy of mesons containing two heavy quarks

Belle reported an observation of the $\psi_2(1D)$ decaying to $\gamma\chi_{c1}$ with J^{PC} presumed to be 2^{--} [28]. This state is listed in Table 95.1 as $\psi_2(3823)$. Its existence was confirmed with high significance by BESIII [29]. While the negative C-parity is indeed established by its observed decay channel, the assignment of $J = 2$ was done by matching to the closest quark model state. This assignment therefore requires experimental confirmation.

A new $c\bar{b}$ state was discovered by the ATLAS Collaboration [33]. Its properties are consistent with expectations for the first excited state of the B_c^\pm meson, the $B_c^\pm(2S)$.

A new structure, dubbed $X(5568)$, was reported in the $B_s^0\pi^\pm$ final state by the $D0$ collaboration [34]. It was not, however, confirmed either by LHCb [35] or CMS [36]. In Ref. [37] it was suggested that the $D0$ signal might have emerged from a combination of cross-channel contamination and the particular choice of event selection criteria. To clarify the situation, more studies appear to be necessary.




Figure 95.1: From Belle [44], the mass recoiling against $\pi^+\pi^-$ pairs, M_{miss} , in e^+e^- collision data taken near the peak of the $\Upsilon(10860)$ (points with error bars). The smooth combinatorial and $K_S^0 \rightarrow \pi^+\pi^-$ background contributions have been subtracted. The fit to the various labeled signal contributions is overlaid (curve). Adapted from [44] with kind permission, copyright (2011) The American Physical Society.

The ground state of bottomonium, $\eta_b(1S)$, was confirmed with a second observation of more than 5σ significance at Belle. In addition, the same experiment collected strong evidence for the $\eta_b(2S)$ [41], but it still needs experimental confirmation at the 5σ level.

95. Spectroscopy of mesons containing two heavy quarks 3

Using dipion transitions from the $\Upsilon(10860)$ (Fig. 95.1), Belle simultaneously discovered the $h_b(1P)$, the bottomonium counterpart of the $h_c(1P)$, and the next excited state, the $h_b(2P)$ [44]. The same analysis also showed the $\Upsilon(1D)$, the lowest-lying D -wave triplet of the $b\bar{b}$ system. The search for the $h_b(1P)$ was directly inspired by a CLEO result [136], which found a surprisingly copious production of $e^+e^- \rightarrow \pi^+\pi^-h_c(1P)$ and an indication that $Y(4260) \rightarrow \pi^+\pi^-h_c(1P)$ occurs at a comparable rate with the signature mode, $Y(4260) \rightarrow \pi^+\pi^-J/\psi$. The presence of $\Upsilon(nS)$ peaks in Fig. 95.1 at rates two orders of magnitude larger than expected, along with separate studies with exclusive decays $\Upsilon(nS) \rightarrow \mu^+\mu^-$, allow precise calibration of the $\pi^+\pi^-$ recoil mass spectrum and very accurate measurements of $h_b(1P)$ and $h_b(2P)$ masses. Both corresponding hyperfine splittings are consistent with zero within an uncertainty of about 1.5 MeV (lowered to ± 1.1 MeV for $h_b(1P)$ in Ref. [43]) .




Figure 95.2: From ATLAS [48] pp collision data (points with error bars) taken at $\sqrt{s} = 7$ TeV, the effective mass of $\chi_{bJ}(1P, 2P, 3P) \rightarrow \gamma\Upsilon(1S, 2S)$ candidates in which $\Upsilon(1S, 2S) \rightarrow \mu^+\mu^-$ and the photon is reconstructed as an e^+e^- conversion in the tracking system. Fits (smooth curves) show significant signals for each triplet (J -merged) on top of a smooth background. From [48] with kind permission, copyright (2012) The American Physical Society.

4 95. Spectroscopy of mesons containing two heavy quarks

We no longer mention a hypothetical $Y_b(10888)$ state since a new analysis of the $\Upsilon(10860)$ energy range does not show evidence for an additional state with mass shifted from the $\Upsilon(10860)$ [129]. After the mass of the $\eta_b(1S)$ was shifted upwards by about 10 MeV based on the new Belle measurements [41,42], all of the bottomonium states mentioned above fit into their respective spectroscopies roughly where expected. An independent experimental confirmation of the shifted masses came from the Belle observation of the $\Upsilon(4S) \rightarrow \eta h_b(1P)$ [42].

The $\chi_{bJ}(nP)$ states have been observed at the LHC by ATLAS [48] and confirmed by D0 [49] for $n = 1, 2, 3$, although in each case the three J states are not distinguished from one another. Events are sought which have both a photon and an $\Upsilon(1S, 2S) \rightarrow \mu^+ \mu^-$ candidate which together form a mass in the χ_b region. All three J -merged peaks are observed with a significance in excess of 6σ for both unconverted and converted photons. The mass plot for converted photons, which provide better mass resolution, is shown in Fig. 95.2. This marks the first observation of the $\chi_{bJ}(3P)$ triplet, quite near the expected mass. A precise confirmation of this result came from LHCb [50].




Figure 95.3: $J/\psi\pi$ invariant mass distributions from BES-III [70] e^+e^- collision data taken near the peak of the $Y(4260)$. Adapted from [70] with kind permission, copyright (2013) The American Physical Society.

95. Spectroscopy of mesons containing two heavy quarks 5

There is a large number of newly discovered states both near and above the lowest open-flavor thresholds. They are displayed in Table 95.2 and Table 95.3, respectively. With the exception of the tensor state located at 3930 MeV, now called $\chi_{c2}(3930)$, which has properties consistent with those expected for the $\chi_{c2}(2P)$, none of these states can easily be assigned a place in the quark model spectrum of charmonia or bottomonia. At the same time, these states have no universally accepted unconventional interpretation either. The $\chi_{c1}(3872)$, also known as $X(3872)$, is widely studied and seen in many transitions — *c.f.* Table 95.2. Yet its interpretation demands additional experimental attention: after the quantum numbers were fixed at LHCb [68,69], the next experimental challenge will be a measurement of its lineshape.

Another state (referred to here as the $X(3915)$), was discovered at 3915 MeV [85] and from a subsequent measurement its quantum numbers were determined to be $J^{PC} = 0^{++}$ [87]. This suggests it may be the $\chi_{c0}(2P)$ quark model state, but this interpretation is not generally accepted [132,133]. In addition, it was pointed out in Ref. [134] that if the assumption of helicity-2 dominance is abandoned and instead one allows for a sizable helicity-0 component, a $J^{PC} = 2^{++}$ assignment is possible. This could imply that the state at 3930 MeV (referred to here as the $\chi_{c2}(3930)$) is actually identical to the one at 3915 MeV—but to explain the large helicity-0 component a sizable portion of non- $q\bar{q}$ is necessary [134]. Because of this analysis, the name of the state was changed from $\chi_{c0}(3915)$ back to $X(3915)$. An alternative candidate for the $\chi_{c0}(2P)$ was reported in Ref. [135] with properties more consistent with expectation: its mass is close to the potential model expectations, it decays to $D\bar{D}$, and the preferred quantum numbers are $J^{PC} = 0^{++}$ (this hypothesis is favored over the 2^{++} one with a 2.5σ significance).

The $\psi(4260)$, also known as $Y(4260)$, and the $\psi(4360)$, also known as $Y(4360)$, are vector states decaying to $\pi^+\pi^-J/\psi$ and $\pi^+\pi^-\psi(2S)$, respectively, yet, unlike most conventional vector charmonia, they do not correspond to enhancements in the e^+e^- hadronic cross section. Furthermore, BESIII observed $\chi_{c1}(3872)$, also known as $X(3872)$, in $e^+e^- \rightarrow \gamma\chi_{c1}(3872)$ [116]—an observation claimed to allow for additional insight into the structure of both $\psi(4260)$ as well as $\chi_{c1}(3872)$ (*c.f.* the minireview on non- $q\bar{q}$ states).

Recently BESIII produced a high-accuracy data set for $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ [115], not only demonstrating that the mass of the $\psi(4260)$ is significantly lower than previously believed, but also that the lineshape is highly non-trivial. The latter observation was interpreted by the authors as the presence of two states. However, this lineshape is also consistent with other possible interpretations, such as one assuming a molecular structure for the $\psi(4260)$ [117]. Note that the data of Ref. [115] does not show any indication of the $Y(4008)$ reported by Belle — the data in this region can either be fit with a non-resonant background component or a much wider resonance at lower mass. Also see the analysis of the $Y(4008)$ region in Ref. [118], where a wide resonance is also extracted.

Another interesting question is whether a heavier $\pi^+\pi^-\psi(2S)$ state, the $\psi(4660)$, discovered by Belle [119,120] and confirmed by BaBar [114], is identical to the $\Lambda_c^+\Lambda_c^-$ state observed by Belle with a nearby mass and width [126].

6 95. Spectroscopy of mesons containing two heavy quarks




Figure 95.4: From Belle [80] e^+e^- collision data taken near the peak of the $\Upsilon(10860)$ for events with a $\pi^+\pi^-$ -missing mass consistent with an $\Upsilon(2S) \rightarrow \mu^+\mu^-$, (a) the maximum of the two possible single π^\pm -missing-mass-squared combinations vs. the $\pi^+\pi^-$ -mass-squared; and (b) projection of the maximum of the two possible single π^\pm -missing-mass combinations (*points with error bars*) overlaid with a fit (*curve*). Events to the left of the vertical line in (a) are excluded from amplitude analysis. The hatched histogram in (b) corresponds to the combinatorial background. The two horizontal stripes in (a) and two peaks in (b) correspond to the two Z_b states. Adapted from [80] with kind permission, copyright (2011) The American Physical Society.




Figure 95.5: From Belle [43] e^+e^- collision data taken near the peak of the $\Upsilon(10860)$, the $h_b(1P)$ event yield vs. the mass recoiling against the $\pi^+\pi^-\gamma$ (corrected for misreconstructed $\pi^+\pi^-$), where the $h_b(1P)$ yield is obtained by fitting the mass recoiling against the $\pi^+\pi^-$ (*points with error bars*). The fit results (*solid histograms*) for signal plus background and background alone are superimposed.

95. Spectroscopy of mesons containing two heavy quarks 7

Table 95.1: New states below the open-flavor thresholds in the $c\bar{c}$, $b\bar{c}$, and $b\bar{b}$ regions, ordered by mass. Masses m and widths Γ represent the PDG16 weighted averages. Ellipses (...) in the Process column indicate inclusively selected event topologies; *i.e.*, additional particles not required by the Experiments to be present. A question mark (?) indicates an unmeasured value. For each Experiment a citation is given, as well as the statistical significance ($\#\sigma$), or “(np)” for “not provided”. The Year column gives the date of the first measurement cited. The Status column indicates that the state has been observed by at most one (NC!-needs confirmation) or at least two independent experiments with significance of $>5\sigma$ (OK).

State	m (MeV)	Γ (MeV)	J^{PC}	Process (mode)	Experiment ($\#\sigma$)	Year Stat.
$h_c(1P)$	3525.38 ± 0.11	0.7 ± 0.35	1^{+-}	$\psi(2S) \rightarrow \pi^0 (\gamma\eta_c(1S))$ $\psi(2S) \rightarrow \pi^0 (\gamma\dots)$	CLEO [14–16] (13.2) CLEO [14–16] (10), BES [17] (19)	2004 OK
$\eta_c(2S)$	3639.2 ± 1.2	$11.3^{+3.2}_{-2.9}$	0^{-+}	$p\bar{p} \rightarrow (\gamma\eta_c) \rightarrow (\gamma\gamma\gamma)$ $\psi(2S) \rightarrow \pi^0 (\gamma\eta_c(1S))$ $B \rightarrow K (K_S^0 K^- \pi^+)$ $e^+e^- \rightarrow e^+e^- (K_S^0 K^- \pi^+)$	Belle [18] (3.1) BESIII [19] (np) Belle [20,21] (6.0) BaBar [22,23] (7.8), CLEO [24] (6.5), Belle [25] (6)	2002 OK
$\psi_2(3823)$	3822.5 ± 1.2	< 16	$?^{? -}$	$B \rightarrow K (\gamma \chi_{c1})$ $e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$	Belle [28] (3.8) BESIII [29] (6.2)	2013 NC!
B_c^+	6277 ± 6	?	0^-	$p\bar{p} \rightarrow (\pi^+ J/\psi)\dots$	CDF [30,31] (8.0), D0 [32] (5.2)	2007 OK 2007 OK
$B_c^+(2S)$	6842 ± 6	?	0^-	$pp \rightarrow (B_c^+ \pi^+ \pi^-)\dots$	ATLAS [33] (5.2)	2014 NC!
$\eta_b(1S)$	9399.2 ± 1.9	$9.8^{+4.4}_{-3.6}$	0^{-+}	$\Upsilon(3S) \rightarrow \gamma(\dots)$ $\Upsilon(2S) \rightarrow \gamma(\dots)$ $h_b(1P, 2P) \rightarrow \gamma(\dots)$ $\Upsilon(4S) \rightarrow \eta h_b(1P)$ $\Upsilon(10860) \rightarrow \pi^+\pi^-\gamma(\dots)$	BaBar [38] (10), CLEO [39] (4.0) BaBar [40] (3.0) Belle [41] (14) Belle [42] (9) Belle [43] (14)	2008 OK 2008 OK
$h_b(1P)$	9899.3 ± 0.7	?	1^{+-}	$\Upsilon(10860) \rightarrow \pi^+\pi^-(\dots)$ $\Upsilon(3S) \rightarrow \pi^0(\dots)$ $\Upsilon(4S) \rightarrow \eta h_b(1P)$	Belle [44,43] (5.5) BaBar [45] (3.0) Belle [42] (11)	2011 NC!
$\eta_b(2S)$	$9999.0^{+4.5}_{-4.0}$	< 24	0^{-+}	$h_b(2P) \rightarrow \gamma(\dots)$	Belle [41] (4.2)	2012 NC!
$\Upsilon(1^3D_2)$	10163.7 ± 1.4	?	2^{--}	$\Upsilon(3S) \rightarrow \gamma\gamma (\gamma\gamma\Upsilon(1S))$ $\Upsilon(3S) \rightarrow \gamma\gamma (\pi^+\pi^-\Upsilon(1S))$ $\Upsilon(10860) \rightarrow \pi^+\pi^-(\dots)$	CLEO [46] (10.2) BaBar [47] (5.8) Belle [44] (2.4)	2004 OK
$h_b(2P)$	$10259.8^{+1.5}_{-1.2}$?	1^{+-}	$\Upsilon(10860) \rightarrow \pi^+\pi^-(\dots)$	Belle [44,43] (11.2)	2011 NC!
$\chi_{bJ}(3P)$	10512.1 ± 2.3	?	$?^{?+}$	$pp \rightarrow (\gamma\mu^+\mu^-)\dots$	ATLAS [48] (>6), D0 [49] (3.6) LHCb [50] (6.9)	2011 OK 2011 OK

8 95. Spectroscopy of mesons containing two heavy quarks

Table 95.2: As in Table 95.1, but for new states near the first open-flavor thresholds in the $c\bar{c}$ and $b\bar{b}$ regions, ordered by mass. For $X(3872)$, the values given are based only upon decays to $\pi^+\pi^-J/\psi$. Updated from [12] with kind permission, copyright (2011), Springer, and [13] with kind permission from the authors.

State	m (MeV)	Γ (MeV)	J^{PC}	Process (mode)	Experiment ($\#\sigma$)	Year	Stat.
$X(3872)$	3871.68 ± 0.17	< 1.2	1^{++}	$B \rightarrow K(\pi^+\pi^-J/\psi)$	Belle [51,52] (10.3), BaBar [53] (8.6)	2003	OK
				$p\bar{p} \rightarrow (\pi^+\pi^-J/\psi) + \dots$	CDF [54–56] (np), D0 [57] (5.2)		
				$B \rightarrow K(\omega J/\psi)$	Belle [58] (4.3), BaBar [59] (4.9)		
				$B \rightarrow K(D^{*0}\bar{D}^0)$	Belle [60,61] (6.4), BaBar [62] (4.9)		
				$B \rightarrow K(\gamma J/\psi)$	Belle [63] (4.0), BaBar [64,65] (3.6), LHCb [66] (>10)		
				$B \rightarrow K(\gamma\psi(2S))$	BaBar [65] (3.5), Belle [63] (0.4), LHCb [66] (4.4)		
				$pp \rightarrow (\pi^+\pi^-J/\psi) + \dots$	LHCb [67,68,69] (np)		
$Z_c(3900)$	3891.2 ± 3.3	40 ± 8	1^{+-}	$Y(4260) \rightarrow \pi^-(\pi^+J/\psi)$	BESIII [70] (> 8), Belle [71] (5.2)	2013	OK
				$Y(4260) \rightarrow \pi^0(\pi^0J/\psi)$	CLEO [72] (>5) BESIII [73] (10.4) CLEO [72] (3.5)		
				$Y(4260) \rightarrow \pi^-(D\bar{D}^*)^+$	BESIII [74] (18)		
				$Y(4260) \rightarrow \pi^0(D\bar{D}^*)^0$	BESIII [75] (> 10)		
				$Y(4260) \rightarrow \pi^-(D^*\bar{D}^*)^+$	BESIII [76] (8.9)	2013	NC!
$X(4020)^\pm$	4022.9 ± 2.8	7.9 ± 3.7	1^{+-}	$Y(4260, 4360) \rightarrow \pi^-(\pi^+h_c)$	BESIII [76] (8.9)		
				$Y(4260, 4360) \rightarrow \pi^0(\pi^0h_c)$	BESIII [77] (> 5)		
				$Y(4260) \rightarrow \pi^-(D^*\bar{D}^*)^+$	BESIII [78] (10)		
				$Y(4260) \rightarrow \pi^0(D^*\bar{D}^*)^0$	BESIII [79] (5.9)		
$Z_b(10610)$	10607.2 ± 2.0	18.4 ± 2.4	1^{+-}	$\Upsilon(10860) \rightarrow \pi^-(\pi^+\Upsilon(1S, 2S, 3S))$	Belle [80] (> 10) Belle [81]	2011	NC!
				$\Upsilon(10860) \rightarrow \pi^-(\pi^+h_b(1P, 2P))$	Belle [80] (16)		
				$\Upsilon(10860) \rightarrow \pi^0(\pi^0\Upsilon(1S, 2S, 3S))$	Belle [82] (6.5)		
				$\Upsilon(10860) \rightarrow \pi^-(B\bar{B}^*)^+$	Belle [83] (> 8)		
				$\Upsilon(10860) \rightarrow \pi^-(\pi^+\Upsilon(1S, 2S, 3S))$	Belle [80] (> 10)	2011	OK
$Z_b(10650)$	10652.2 ± 1.5	11.5 ± 2.2	1^{+-}	$\Upsilon(10860) \rightarrow \pi^-(\pi^+h_b(1P, 2P))$	Belle [80] (16)		
				$\Upsilon(10860) \rightarrow \pi^-(B^*\bar{B}^*)^+$	Belle [83] (6.8)		

95. Spectroscopy of mesons containing two heavy quarks 9

Table 95.3: As in Table 95.1, but for new states above the first open-flavor thresholds in the $c\bar{c}$ and $b\bar{b}$ regions, ordered by mass. $X(3945)$ and $Y(3940)$ have been subsumed under $X(3940)$ due to compatible properties. The $\chi_{c0}(3915)$ is now changed back to $X(3915)$ as explained in the main text. In some cases experiment still allows two J^{PC} values, in which case both appear. See also the review [13].

State	m (MeV)	Γ (MeV)	J^{PC}	Process (mode)	Experiment ($\#\sigma$)	Year	Status
$X(3915)$	3917.4 ± 2.7	28_{-9}^{+10}	$0/2^{++}$	$B \rightarrow K(\omega J/\psi)$	Belle [85] (8.1), BaBar [59] (np)	2004	OK
				$e^+e^- \rightarrow e^+e^-\omega J/\psi$	Belle [86] (7.7), BaBar [87] (19)		OK
$\chi_{c2}(3930)$	3927.2 ± 2.6	24 ± 6	2^{++}	$e^+e^- \rightarrow e^+e^-(D\bar{D})$	Belle [88] (5.3), BaBar [89]	2005	OK
$X(3940)$	3942_{-8}^{+9}	37_{-17}^{+27}	$?^{?+}$	$e^+e^- \rightarrow J/\psi(D\bar{D}^*)$ $e^+e^- \rightarrow J/\psi(\dots)$	Belle [90] (6.0) Belle [27] (5.0)	2007	NC!
$Y(4008)$	4008_{-49}^{+121}	226 ± 97	1^{--}	$e^+e^- \rightarrow \gamma(\pi^+\pi^- J/\psi)$	Belle [91] (7.4)	2007	NC!
$X(4050)^\pm$	4051_{-43}^{+24}	82_{-55}^{+51}	$?$	$B \rightarrow K(\pi^+\chi_{c1}(1P))$	Belle [92] (5.0), BaBar [93] (1.1)	2008	NC!
$\chi_{c1}(4140)$	4145.8 ± 2.6	18 ± 8	$?^{?+}$	$B^+ \rightarrow K^+(\phi J/\psi)$	CDF [96,97] (5.0) D0 [98] (3.1), CMS [99](>5) Belle [100] (1.9), LHCb [101] (1.4), BaBar [102]	2009	NC!
				$e^+e^- \rightarrow e^+e^-(\phi J/\psi)$	Belle [103] (3.2)	2009	NC!
$X(4160)$	4156_{-25}^{+29}	139_{-65}^{+113}	$?^{?+}$	$e^+e^- \rightarrow J/\psi(D\bar{D}^*)$	Belle [90] (5.5)	2007	NC!
$Z_c(4200)$	4196_{-32}^{+35}	370_{-149}^{+99}	1^+	$\bar{B}^0 \rightarrow K^-(J/\psi\pi^+)$	Belle [106] (6.2)	2014	NC!
$X(4250)^\pm$	4248_{-45}^{+185}	177_{-72}^{+321}	$?$	$B \rightarrow K(\pi^+\chi_{c1}(1P))$	Belle [92] (5.0), BaBar [93] (2.0)	2008	NC!
$\psi(4260)$	4263_{-9}^{+8}	95 ± 14	1^{--}	$e^+e^- \rightarrow \gamma(\pi^+\pi^- J/\psi)$	BaBar [107,108] (8.0) CLEO [109] (5.4), Belle [91] (15)	2005	OK
				$e^+e^- \rightarrow (\pi^+\pi^- J/\psi)$	CLEO [110] (11)		
				$e^+e^- \rightarrow (\pi^0\pi^0 J/\psi)$	CLEO [110] (5.1)		
				$e^+e^- \rightarrow (f_0(980)J/\psi)$	BaBar [111] (np), Belle [71] (np)		
				$e^+e^- \rightarrow (\pi^- Z_c(3900)^+)$	BESIII [70] (8), Belle [71] (5.2)		
				$e^+e^- \rightarrow (\gamma X(3872))$	BESIII [112] (5.3)		
$\chi_{c1}(4274)$	4293 ± 20	35 ± 16	$?^{?+}$	$B^+ \rightarrow K^+(\phi J/\psi)$	CDF [97] (3.1), LHCb [101] (1.0), CMS [99] (>3), D0 [98] (np)	2011	NC!

10 95. Spectroscopy of mesons containing two heavy quarks

State	m (MeV)	Γ (MeV)	J^{PC}	Process (mode)	Experiment ($\#\sigma$)	Year Status
$X(4350)$	$4350.6^{+4.6}_{-5.1}$	$13.3^{+18.4}_{-10.0}$	$0/2^{++}$	$e^+e^- \rightarrow e^+e^- (\phi J/\psi)$	Belle [103] (3.2)	2009 NC!
$\psi(4360)$	4361 ± 13	74 ± 18	1^{--}	$e^+e^- \rightarrow \gamma(\pi^+\pi^-\psi(2S))$	BaBar [113,114] (np), Belle [119,120] (8.0)	2007 OK
$Z_c(4430)$	4458 ± 15	166^{+37}_{-32}	1^+	$\bar{B}^0 \rightarrow K^-(\pi^+\psi(2S))$	Belle [121,122,123] (6.4), BaBar [124] (2.4), LHCb [125] (13.9)	2007 OK
				$\bar{B}^0 \rightarrow (J/\psi\pi^+)K^-$	Belle [106] (4.0)	
$X(4630)$	4634^{+9}_{-11}	92^{+41}_{-32}	1^{--}	$e^+e^- \rightarrow \gamma(\Lambda_c^+\Lambda_c^-)$	Belle [126] (8.2)	2007 NC!
$\psi(4660)$	4664 ± 12	48 ± 15	1^{--}	$e^+e^- \rightarrow \gamma(\pi^+\pi^-\psi(2S))$	Belle [119,120] (5.8), BaBar [114] (np)	2007 NC!
$\Upsilon(10860)$	10876 ± 11	55 ± 28	1^{--}	$e^+e^- \rightarrow (B_{(s)}^{(*)}\bar{B}_{(s)}^{(*)})(\pi)$	PDG [127] (> 10)	1985 OK
				$e^+e^- \rightarrow (\pi\pi\Upsilon(1S, 2S, 3S))$	Belle [128,80,82,129] (> 10)	
				$e^+e^- \rightarrow (f_0(980)\Upsilon(1S))$	Belle [80,82] (> 5)	
				$e^+e^- \rightarrow (\pi Z_b(10610, 10650))$	Belle [80,82] (> 10)	
				$e^+e^- \rightarrow (\eta\Upsilon(1S, 2S))$	Belle [42] (10)	
				$e^+e^- \rightarrow (\pi^+\pi^-\Upsilon(1D))$	Belle [130] (9)	
				$e^+e^- \rightarrow (\pi^+\pi^-h_b(1P, 2P))$	Belle [131] (9)	
$\Upsilon(11020)$	$10987.5^{+11.1}_{-3.3}$	$61.0^{+9.2}_{-27.7}$	1^{--}	$e^+e^- \rightarrow (B_{(s)}^{(*)}\bar{B}_{(s)}^{(*)})(\pi)$	PDG [127] (> 10)	1985 OK
				$e^+e^- \rightarrow (\pi\pi\Upsilon(1S, 2S, 3S))$	Belle [129] (> 10)	
				$e^+e^- \rightarrow (\pi^+\pi^-h_b(1P, 2P))$	Belle [131] (9)	

Based on a full amplitude analysis of $B^0 \rightarrow K^+\pi^-\psi(2S)$ decays, Belle determined the spin-parity of the $Z_c(4430)$ to be $J^P = 1^+$ [123]. From their study of $B^0 \rightarrow K^+\pi^-J/\psi$ decays, Belle also found evidence for the decay mode $Z_c(4430) \rightarrow \pi J/\psi$ [106], which has an order of magnitude lower branching fraction than the discovery mode $Z_c(4430) \rightarrow \pi\psi(2S)$. In the same analysis, Belle also reported evidence for one more charged state, dubbed $Z_c(4200)$, decaying to $\pi J/\psi$. The existence of the $Z_c(4430)$ in $\pi\psi(2S)$ as well as its quantum number assignments were confirmed at LHCb [125] with much higher statistics. Improved values for the mass and width of the $Z_c(4430)$ from LHCb are consistent with earlier measurements; the experiment even reports a resonant behavior of the $Z_c(4430)$ amplitude. The $Z_c(4430)$ was not confirmed (or excluded) by BaBar [124].

Belle also reported an observation of two charged states decaying to $\pi\chi_{c1}$ in an analysis of $B^0 \rightarrow K^+\pi^-\chi_{c1}$ decays [92]. These were originally called the $Z_1(4050)^\pm$ and the $Z_2(4250)^\pm$, but are referred to in Table 95.3 as $X(4050)^\pm$ and $X(4250)^\pm$. These were also not confirmed by BaBar [93]. Belle observes signals with 5.0σ significance for both the $Z_1(4050)^\pm$ and $Z_2(4250)^\pm$, whereas BABAR reports 1.1σ and 2.0σ effects, respectively, setting upper limits on product branching fractions that are not inconsistent with Belle's measured rates. The situation remains unresolved.

In addition to the Z_c states discussed above, in 2013 a state named $Z_c(3900)$ was unearthed in the charmonium region at BESIII [70] and Belle [71]- –the corresponding

spectrum from BESIII is shown in Fig. 95.3. Ref. [72] confirmed this finding and also provided evidence for a neutral partner. A nearby signal was also seen in the $D\bar{D}^*$ channel [74] whose quantum numbers were fixed to 1^{+-} . BESIII reported its neutral partner in both $J/\psi\pi^0$ [73] and $D\bar{D}^*$ [75] decay modes. The masses extracted from these experiments in different decay modes have differences reaching up to 2σ . However, since the extraction of the mass and width parameters did not allow for an interference with the background and used Breit-Wigner line shapes, which is not justified near thresholds, there might be some additional systematic uncertainty in the mass values. Therefore in the RPP listings as well as Table 95.2, both structures appear under the name $Z_c(3900)$. BESIII also reported an observation of another charged state, the $X(4020)^\pm$ (originally called $Z_c(4020)^\pm$), in two decay modes — $h_c\pi^\pm$ [76] and $(D^*\bar{D}^*)^\pm$ [78]. The neutral partners have also been observed by BESIII in the $h_c\pi^0$ [77] and $(D^*\bar{D}^*)^0$ [79] final states. The Z_c states show some remarkable similarities to the Z_b states (discussed below), e.g. they decay dominantly to $D^{(*)}\bar{D}^*$ channels. However, current analyses suggest that the mass of the $Z_c(3900)$ might be somewhat above the $D\bar{D}^*$ threshold. If confirmed, this feature would clearly challenge a possible $D\bar{D}^*$ -molecular interpretation. Finally, 3.5σ evidence for one more charged charmoniumlike state at 4055 MeV decaying into $\psi(2S)\pi^\pm$ was reported by Belle in their analysis of the process $e^+e^- \rightarrow \psi(2S)\pi^+\pi^-$ [120]. This state was confirmed by BESIII, although there appears to be complications in the Dalitz plot requiring further investigation [95].

The $Y(4140)$ observed in 2008 by CDF [96,97] was confirmed at D0 and CMS [98,99]. However, a second structure, the $Y(4274)$, could not be established unambiguously. Neither of the two states were seen in B decays at Belle [100], LHCb [101] and BaBar [102] or in $\gamma\gamma$ collisions at Belle [103]. The real breakthrough happened recently when LHCb performed a full amplitude analysis of $B^+ \rightarrow J/\psi\phi K^+$ with $J/\psi \rightarrow \mu^+\mu^-$, $\phi \rightarrow K^+K^-$ decays and showed that the data cannot be described in a model that contains only excited kaon states decaying into ϕK^+ [105,104]. They observe two 1^{++} states with masses close to those originally reported by CDF (the $\chi_{c1}(4140)$ and $\chi_{c1}(4274)$), but the width of the one at 4140 MeV is much larger. In addition, they find two significant 0^{++} structures at 4500 and 4700 MeV (the $\chi_{c0}(4500)$ and $\chi_{c0}(4700)$).

New results on the η_b , h_b , and Z_b mostly come from Belle [41–44], [80–83], [128–131], all from analyses of 121.4 fb^{-1} of e^+e^- collision data collected near the peak of the $\Upsilon(10860)$ resonance as well as from an additional 25 fb^{-1} of data collected during the scans of the c.m. energy range 10.63–11.05 GeV. They all appear in the decay chains: $\Upsilon(10860) \rightarrow \pi^- Z_b^+$, $Z_b^+ \rightarrow \pi^+(b\bar{b})$, and, when the $b\bar{b}$ forms an $h_b(1P)$, frequently decaying as $h_b(1P) \rightarrow \gamma\eta_b$.

Belle soon noticed that, for events in the peaks of Fig. 95.1, there seemed to be two intermediate charged states. For example, Fig. 95.4 shows a Dalitz plot for events restricted to the $\Upsilon(2S)$ region of $\pi^+\pi^-$ recoil mass, with $\Upsilon(2S) \rightarrow \mu^+\mu^-$. The two bands observed in the maximum of the two $M[\pi^\pm\Upsilon(2S)]^2$ values also appear for $\Upsilon(1S)$, $\Upsilon(3S)$, $h_b(1P)$, and $h_b(2P)$ samples. Belle fits all subsamples to resonant plus non-resonant amplitudes, allowing for interference (notably, between $\pi^- Z_b^+$ and $\pi^+ Z_b^-$), and finds consistent pairs of Z_b masses for all bottomonium transitions, and comparable strengths of the two states. A recent angular analysis assigned $J^P = 1^+$ for both Z_b states [81],

12 95. Spectroscopy of mesons containing two heavy quarks

which must also have negative G -parity. Transitions through Z_b to the $h_b(nP)$ saturate the observed $\pi^+\pi^-h_b(nP)$ cross sections. While the two masses of the Z_b states as extracted from Breit-Wigner fits for the various channels are just a few MeV above the $B^*\bar{B}$ and $B^*\bar{B}^*$ thresholds, respectively, more refined analyses find pole locations right below the corresponding thresholds either on the physical [137] or the unphysical sheet [138]. Regardless of their proximity to the corresponding thresholds, both states predominantly decay into these open-flavor channels [83,84] with branching fractions that exceed 80% and 70%, respectively, at 90% CL. This feature provides strong evidence for their molecular nature.

References:

1. A. Esposito *et al.*, Int. J. Mod. Phys. A **30**,1530002(2015) arXiv:1411.5997 [hep-ph].
2. S.L. Olsen, Front. Phys. (Beijing) **10**, no. 2, 121 (2015) arXiv:1411.7738 [hep-ex].
3. H.X. Chen, W. Chen, X. Liu, and S.L. Zhu, Phys. Rept. **639**, 1 (2016) arXiv:1601.02092 [hep-ph].
4. R.F. Lebed, R.E. Mitchell, and E.S. Swanson, Prog. Part. Nucl. Phys. **93**, 143 (2017) arXiv:1610.04528 [hep-ph].
5. A. Ali, J.S. Lange, and S. Stone, Prog. Part. Nucl. Phys. **97**, 123 (2017) arXiv:1706.00610 [hep-ph].
6. F. K. Guo *et al.*, arXiv:1705.00141 [hep-ph].
7. S. L. Olsen, T. Skwarnicki and D. Zieminska, arXiv:1708.04012 [hep-ph].
8. E. Eichten *et al.*, Rev. Mod. Phys. **80**, 1161 (2008) arXiv:hep-ph/0701208.
9. S. Godfrey and S.L. Olsen, Ann. Rev. Nucl. Part. Sci. **58** 51 (2008), arXiv:0801.3867 [hep-ph].
10. T. Barnes and S.L. Olsen, Int. J. Mod. Phys. A **24S1**(2009).
11. G.V. Pakhlova, P.N. Pakhlov, and S.I. Eidelman, Phys. Usp. **53**, 219 (2010), [Usp. Fiz. Nauk **180**, 225 (2010)].
12. N. Brambilla *et al.*, Eur. Phys. J. **C71**, 1534 (2011) arXiv:1010.5827 [hep-ph].
13. N. Brambilla *et al.*, Eur. Phys. J. C **74**, 2981 (2014) arXiv:1404.3723 [hep-ph].
14. P. Rubin *et al.* (CLEO Collab.), Phys. Rev. **D72**, 092004 (2005) arXiv:hep-ex/0508037.
15. J.L. Rosner *et al.* (CLEO Collab.), Phys. Rev. Lett. **95**, 102003 (2005) arXiv:hep-ex/0505073.
16. S. Dobbs *et al.* (CLEO Collab.), Phys. Rev. Lett. **101**, 182003 (2008) arXiv:0805.4599 [hep-ex].
17. M. Ablikim *et al.* (BESIII Collab.), Phys. Rev. Lett. **104**, 132003 (2010) arXiv:1002.0501 [hep-ex].
18. M. Andreotti *et al.* (E835 Collab.), Phys. Rev. **D72**, 032001 (2005).
19. M. Ablikim *et al.* (BESIII Collab.), Phys. Rev. D **86**, 092009 (2012) arXiv:1209.4963 [hep-ex].
20. S.K. Choi *et al.* (Belle Collab.), Phys. Rev. Lett. **89**, 102001 (2002) [Erratum-ibid. **89**, 129901 (2002)], arXiv:hep-ex/0206002.

21. A. Vinokurova *et al.* (Belle Collab.), Phys. Lett. **B706**, 139 (2011) arXiv:1105.0978 [hep-ex].
22. P. del Amo Sanchez *et al.* (BaBar Collab.), Phys. Rev. **D84**, 021004 (2011) arXiv:1103.3971 [hep-ex].
23. B. Aubert *et al.* (BaBar Collab.), Phys. Rev. Lett. **92**, 142002 (2004) arXiv:hep-ex/0311038.
24. D.M. Asner *et al.* (CLEO Collab.), Phys. Rev. Lett. **92**, 142001 (2004) arXiv:hep-ex/0312058.
25. H. Nakazawa (Belle Collab.), Nucl. Phys. (Proc. Supp.) **184**, 220 (2008).
26. B. Aubert *et al.* (BaBar Collab.), Phys. Rev. **D72**, 031101 (2005) arXiv:hep-ex/0506062.
27. K. Abe *et al.* (Belle Collab.), Phys. Rev. Lett. **98**, 082001 (2007) arXiv:hep-ex/0507019.
28. V. Bhardwaj *et al.* (Belle Collaboration), Phys. Rev. Lett. **111**, 032001 (2013), arXiv:1304.3975.
29. M. Ablikim *et al.*, Phys. Rev. Lett. **115**, 011803 (2015), arXiv:1503.08203 [hep-ex].
30. F. Abe *et al.* [CDF Collab.], Phys. Rev. Lett. **81**, 2432 (1998) arXiv:hep-ex/9805034.
31. T. Aaltonen *et al.* (CDF Collab.), Phys. Rev. Lett. **100**, 182002 (2008) arXiv:0712.1506 [hep-ex].
32. V.M. Abazov *et al.* (D0 Collab.), Phys. Rev. Lett. **101**, 012001 (2008) arXiv:0802.4258 [hep-ex].
33. G. Aad *et al.* (ATLAS Collab.), Phys. Rev. Lett. **113**, 212004 (2014), arXiv:1407.1032 [hep-ex].
34. V. M. Abazov *et al.* [D0 Collaboration], Phys. Rev. Lett. **117**, no. 2, 022003 (2016) arXiv:1602.07588 [hep-ex].
35. R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. Lett. **117**, no. 15, 152003 (2016) Addendum: [Phys. Rev. Lett. **118**, no. 10, 109904 (2017)] arXiv:1608.00435 [hep-ex].
36. CMS Collaboration [CMS Collaboration], CMS-PAS-BPH-16-002.
37. Z. Yang, Q. Wang and U.-G. Meißner, Phys. Lett. B **767**, 470 (2017) arXiv:1609.08807 [hep-ph].
38. B. Aubert *et al.* (BaBar Collab.), Phys. Rev. Lett. **101**, 071801 (2008) [Erratum-ibid. **102**, 029901 (2009)], arXiv:0807.1086 [hep-ex].
39. G. Bonvicini *et al.* (CLEO Collab.), Phys. Rev. **D81**, 031104 (2010) arXiv:0909.5474 [hep-ex].
40. B. Aubert *et al.* (BaBar Collab.), Phys. Rev. Lett. **103**, 161801 (2009) arXiv:0903.1124 [hep-ex].
41. R. Mizuk *et al.* (Belle Collaboration), Phys. Rev. Lett. **109**, 232002 (2012), arXiv:1205.6351.
42. U. Tamponi *et al.* (Belle Collab.), Phys. Rev. Lett. **115**, 142001 (2015), arXiv:1506.08914 [hep-ex].
43. I. Adachi *et al.* (Belle Collab.), arXiv:1110.3934 [hep-ex].

14 95. Spectroscopy of mesons containing two heavy quarks

44. I. Adachi *et al.* (Belle Collab.), Phys. Rev. Lett. **108**, 032001 (2012) arXiv:1103.3419 [hep-ex].
45. J.P. Lees *et al.* (BaBar Collab.), Phys. Rev. **D84**, 091101 (2011) arXiv:1102.4565 [hep-ex].
46. G. Bonvicini *et al.* (CLEO Collab.), Phys. Rev. **D70**, 032001 (2004) arXiv:hep-ex/0404021.
47. P. del Amo Sanchez *et al.* (BaBar Collab.), Phys. Rev. **D82**, 111102 (2010) arXiv:1004.0175 [hep-ex].
48. G. Aad *et al.* (ATLAS Collab.), Phys. Rev. Lett. **108**, 152001 (2012) arXiv:1112.5154 [hep-ex].
49. V. M. Abazov (D0 Collab.), Phys.Rev. D **86**, 031103 (2012) arXiv:1203.6034.
50. R. Aaij *et al.* (LHCb Collab.), JHEP **1410**, 088 (2014), arXiv:1409.1408 [hep-ex].
51. S.K. Choi *et al.* (Belle Collab.), Phys. Rev. Lett. **91**, 262001 (2003) arXiv:hep-ex/0309032.
52. S.-K. Choi *et al.* (Belle Collab.), Phys. Rev. **D84**, 052004R (2011) arXiv:1107.0163 [hep-ex].
53. B. Aubert *et al.* (BaBar Collab.), Phys. Rev. **D77**, 111101 (2008) arXiv:0803.2838 [hep-ex].
54. D.E. Acosta *et al.* (CDF Collab.), Phys. Rev. Lett. **93**, 072001 (2004) arXiv:hep-ex/0312021.
55. A. Abulencia *et al.* (CDF Collab.), Phys. Rev. Lett. **98**, 132002 (2007) arXiv:hep-ex/0612053.
56. T. Aaltonen *et al.* (CDF Collab.), Phys. Rev. Lett. **103**, 152001 (2009) arXiv:0906.5218 [hep-ex].
57. V.M. Abazov *et al.* (D0 Collab.), Phys. Rev. Lett. **93**, 162002 (2004) arXiv:hep-ex/0405004.
58. K. Abe *et al.* (Belle Collab.), arXiv:hep-ex/0505037.
59. P. del Amo Sanchez *et al.* (BaBar Collab.), Phys. Rev. **D82**, 011101R (2010) arXiv:1005.5190 [hep-ex].
60. G. Gokhroo *et al.* (Belle Collab.), Phys. Rev. Lett. **97**, 162002 (2006) arXiv:hep-ex/0606055.
61. T. Aushev *et al.* (Belle Collab.), Phys. Rev. **D81**, 031103R (2010), arXiv:0810.0358 [hep-ex].
62. B. Aubert *et al.* (BaBar Collab.), Phys. Rev. **D77**, 011102 (2008) arXiv:0708.1565 [hep-ex].
63. V. Bhardwaj *et al.* (Belle Collab.), Phys. Rev. Lett. **107**, 091803 (2011) arXiv:1105.0177 [hep-ex].
64. B. Aubert *et al.* (BaBar Collab.), Phys. Rev. **D74**, 071101 (2006) arXiv:hep-ex/0607050.
65. B. Aubert *et al.* (BaBar Collab.), Phys. Rev. Lett. **102**, 132001 (2009) arXiv:0809.0042 [hep-ex].
66. R. Aaij *et al.* (LHCb Collab.), Nucl. Phys. B **886**, 665 (2014) arXiv:1404.0275 [hep-ex].

95. Spectroscopy of mesons containing two heavy quarks 15

67. R. Aaij *et al.* (LHCb Collab.), Eur. Phys. J. **C72**, 1972 (2012) arXiv:1112.5310 [hep-ex].
68. R. Aaij *et al.* (LHCb Collab.), Phys. Rev. Lett. **110**, 222001 (2013) arXiv:1302.6269 [hep-ex].
69. R. Aaij *et al.* (LHCb Collab.), Phys. Rev. D **92**, 011102 (2015) arXiv:1504.06339 [hep-ex].
70. M. Ablikim *et al.* (BESIII Collab.), Phys. Rev. Lett. **110**, 252001 (2013) arXiv:1303.5949.
71. Z. Q. Liu *et al.* (Belle Collab.), Phys. Rev. Lett. **110**, 252002 (2013) arXiv:1304.0121.
72. T. Xiao, S. Dobbs, A. Tomaradze and K. K. Seth, Phys. Lett. **B727**, 366 (2013) arXiv:1304.3036.
73. M. Ablikim *et al.* (BESIII Collab.), Phys. Rev. Lett. **115**, 112003 (2015) arXiv:1506.06018 [hep-ex].
74. M. Ablikim *et al.* (BESIII Collab.), Phys. Rev. Lett. **112**, 022001 (2014) arXiv:1310.1163.
75. M. Ablikim *et al.* (BESIII Collab.), Phys. Rev. Lett. **115**, 222002 (2015) arXiv:1509.05620 [hep-ex].
76. M. Ablikim *et al.* (BESIII Collab.), Phys. Rev. Lett. **111**, 242001 (2013) arXiv:1309.1896.
77. M. Ablikim *et al.* (BESIII Collab.), Phys. Rev. Lett. **113**, 212002 (2014) arXiv:1409.6577 [hep-ex].
78. M. Ablikim *et al.* (BESIII Collab.), Phys. Rev. Lett. **112**, 132001 (2014), arXiv:1308.2760.
79. M. Ablikim *et al.* (BESIII Collab.), Phys. Rev. Lett. **115**, 182002 (2015) arXiv:1507.02404 [hep-ex].
80. A. Bondar *et al.* (Belle Collab.), Phys. Rev. Lett. **108**, 122001 (2012) arXiv:1110.2251 [hep-ex].
81. A. Garmash *et al.* (Belle Collab.), Phys. Rev. D **91**, 072003 (2015), arXiv:1403.0992.
82. P. Krokovny *et al.* (Belle Collab.), Phys. Rev. D **88**, 052016 (2013), arXiv:1308.2646.
83. I. Adachi *et al.* (Belle Collab.) (2012), arXiv:1209.6450.
84. A. Garmash *et al.* [Belle Collaboration], Phys. Rev. Lett. **116**, no. 21, 212001 (2016) arXiv:1512.07419 [hep-ex].
85. S.-K. Choi *et al.* (Belle Collab.), Phys. Rev. Lett. **94**, 182002 (2005) arXiv:hep-ex/0408126.
86. S. Uehara *et al.* (Belle Collab.), Phys. Rev. Lett. **104**, 092001 (2010) arXiv:0912.4451 [hep-ex].
87. J. P. Lees *et al.* (BaBar Collab.), Phys. Rev. D **86**, 072002 (2012) arXiv:1207.2651 [hep-ex].
88. S. Uehara *et al.* (Belle Collab.), Phys. Rev. Lett. **96**, 082003 (2006) arXiv:hep-ex/0512035.

16 95. Spectroscopy of mesons containing two heavy quarks

89. B. Aubert *et al.* (BaBar Collab.), Phys. Rev. **D81**, 092003 (2010) arXiv:1002.0281 [hep-ex].
90. P. Pakhlov *et al.* (Belle Collab.), Phys. Rev. Lett. **100**, 202001 (2008) arXiv:0708.3812 [hep-ex].
91. C.Z. Yuan *et al.* (Belle Collab.), Phys. Rev. Lett. **99**, 182004 (2007) arXiv:0707.2541 [hep-ex].
92. R. Mizuk *et al.* (Belle Collab.), Phys. Rev. **D78**, 072004 (2008) arXiv:0806.4098 [hep-ex].
93. J.P. Lees *et al.* (BaBar Collab.), Phys. Rev. **D85**, 052003 (2011) arXiv:1111.5919 [hep-ex].
94. X.L. Wang *et al.* (Belle Collab.), Phys. Rev. D 91,112007(2015) arXiv:1410.7641.
95. M. Ablikim *et al.* (BESIII Collab.), Phys. Rev. D 96,032004(2017) arXiv:1703.08787.
96. T. Aaltonen *et al.* (CDF Collab.), Phys. Rev. Lett. **102**, 242002 (2009) arXiv:0903.2229 [hep-ex].
97. T. Aaltonen *et al.* (CDF Collab.), Mod. Phys. Lett. **A32**, 1750139 (2017) arXiv:1101.6058 [hep-ex].
98. V. Abazov *et al.* (D0 Collab.), Phys.Rev. D **89**, 012004 (2014), arXiv:1309.6580.
99. S. Chatrchyan *et al.* (CMS Collab.), Phys. Lett. B **734**, 261 (2014) arXiv:1309.6920.
100. J. Brodzicka (Belle Collab.), Conf.Proc. C0908171, 299 (2009).
101. R. Aaij *et al.* (LHCb Collab.), Phys. Rev. D **85**,091103 (2012), arXiv:1202.5087.
102. J.P. Lees *et al.* (BaBar Collab.), Phys. Rev. D **91**,012003 (2015), arXiv:1407.7244.
103. C.P. Shen *et al.* (Belle Collab.), Phys. Rev. Lett. **104**, 112004 (2010) arXiv:0912.2383 [hep-ex].
104. R. Aaij *et al.* (LHCb Collab.), Phys. Rev. Lett. **118**, 022003 (2017) arXiv:1606.07895 [hep-ex].
105. R. Aaij *et al.* (LHCb Collab.), Phys. Rev. D 95,012002(2017) arXiv:1606.07898 [hep-ex].
106. K. Chilikin *et al.* (Belle Collab.), Phys. Rev. D **90**, 112009 (2014), arXiv:1408.6457 [hep-ex].
107. B. Aubert *et al.* (BaBar Collab.), Phys. Rev. Lett. **95**, 142001 (2005) arXiv:hep-ex/0506081.
108. B. Aubert *et al.* (BaBar Collab.), arXiv:0808.1543v2 [hep-ex].
109. Q. He *et al.* (CLEO Collab.), Phys. Rev. **D74**, 091104 (2006) arXiv:hep-ex/0611021.
110. T.E. Coan *et al.* (CLEO Collab.), Phys. Rev. Lett. **96**, 162003 (2006) arXiv:hep-ex/0602034.
111. J. Lees *et al.* (BaBar Collab.), Phys.Rev. D **86**, 051102 (2012), arXiv:1204.2158.
112. M. Ablikim *et al.* (BESIII Collab.), Phys.Rev.Lett. **112**, 092001 (2014), arXiv:1310.4101.
113. B. Aubert *et al.* (BaBar Collab.), Phys. Rev. Lett. **98**, 212001 (2007) arXiv:hep-ex/0610057.

95. Spectroscopy of mesons containing two heavy quarks 17

114. J.P. Lees *et al.* (BaBar Collab.), Phys. Rev. D 89,111103 (2014) arXiv:1211.6271 [hep-ex].
115. M. Ablikim *et al.* [BESIII Collaboration], Phys. Rev. Lett. **118**, no. 9, 092001 (2017) arXiv:1611.01317 [hep-ex].
116. M. Ablikim *et al.* [BESIII Collaboration], Phys. Rev. Lett. **112**, no. 9, 092001 (2014) arXiv:1310.4101 [hep-ex].
117. M. Cleven *et al.*, Phys. Rev. D **90**, no. 7, 074039 (2014) arXiv:1310.2190 [hep-ph].
118. X. Y. Gao, C. P. Shen and C. Z. Yuan, Phys. Rev. D **95**, no. 9, 092007 (2017) arXiv:1703.10351 [hep-ex].
119. X.L. Wang *et al.* (Belle Collab.), Phys. Rev. Lett. **99**, 142002 (2007) arXiv:0707.3699 [hep-ex].
120. X.L. Wang *et al.* (Belle Collab.), Phys. Rev. D 91,112007(2015) arXiv:1410.7641 [hep-ex].
121. S.K. Choi *et al.* (Belle Collab.), Phys. Rev. Lett. **100**, 142001 (2008) arXiv:0708.1790 [hep-ex].
122. R. Mizuk *et al.* (Belle Collab.), Phys. Rev. **D80**, 031104 (2009) arXiv:0905.2869 [hep-ex].
123. K. Chilikin *et al.* (Belle Collab.), Phys.Rev. D **88**,074026 (2013), arXiv:1306.4894.
124. B. Aubert *et al.* (BaBar Collab.), Phys. Rev. D **79**, 112001 (2009), arXiv:0811.0564.
125. R. Aaij *et al.* (LHCb Collab.), Phys. Rev. Lett. **112** no.22, 222002 (2014) arXiv:1404.1903 [hep-ex].
126. G. Pakhlova *et al.* (Belle Collab.), Phys. Rev. Lett. **101**, 172001 (2008) arXiv:0807.4458 [hep-ex].
127. K. Olive *et al.* (Particle Data Group), Chin. Phys. C **38**, 090001 (2014).
128. K.F. Chen *et al.* (Belle Collab.), Phys. Rev. Lett. **100**, 112001 (2008) arXiv:0710.2577 [hep-ex].
129. D. Santel *et al.* (Belle Collab.), Phys. Rev. D 93, 011101 (2016) arXiv:1501.01137 [hep-ex].
130. P. Krokovny (Belle Collab.), talk given at Les Rencontres de Physique de la Vallee d'Aoste, La Thuile, Aosta Valley, Italy, 2012.
131. A. Abdesselam *et al.* (Belle Collab.), arXiv:1508.06562.
132. F. -K. Guo and U.-G. Meißner, Phys. Rev. D 86, 091501 (2012) arXiv:1208.1134.
133. S.L. Olsen, Phys. Rev. D 91, 057501 (2015) arXiv:1410.6534.
134. Z. Y. Zhou, Z. Xiao and H. Q. Zhou, Phys. Rev. Lett. **115** (2015) no.2, 022001 arXiv:1501.00879 [hep-ph].
135. K. Chilikin *et al.* [Belle Collaboration], Phys. Rev. D **95**, 112003 (2017) arXiv:1704.01872 [hep-ex].
136. T.K. Pedlar *et al.* (CLEO Collab.), Phys. Rev. Lett. **107**, 041803 (2011) arXiv:1104.2025 [hep-ex].
137. M. Cleven *et al.*, Eur. Phys. J. A **47**, 120 (2011) arXiv:1107.0254 [hep-ph].

18 *95. Spectroscopy of mesons containing two heavy quarks*

138. F.-K. Guo et al., Phys. Rev. D **93**, no. 7, 074031 (2016) arXiv:1602.00940 [hep-ph].