

$f_0(1370)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

See also the mini-reviews on scalar mesons under $f_0(500)$ (see the index for the page number) and on non- $q\bar{q}$ candidates in PDG 06, Journal of Physics **G33** 1 (2006).

 $f_0(1370)$ T-MATRIX POLE POSITION

Note that $\Gamma \approx 2 \operatorname{Im}(\sqrt{s_{\text{pole}}})$.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(1200–1500)–i(150–250) OUR ESTIMATE			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$(1290 \pm 50) - i(170^{+20}_{-40})$	¹ ANISOVICH	09	RVUE $0.0 \bar{p}p, \pi N$
$(1373 \pm 15) - i(137 \pm 10)$	² BARGIOTTI	03	OBLX $\bar{p}p$
$(1302 \pm 17) - i(166 \pm 18)$	³ BARBERIS	00C	450 $pp \rightarrow p_f 4\pi p_S$
$(1312 \pm 25 \pm 10) - i(109 \pm 22 \pm 15)$	BARBERIS	99D	OMEG 450 $pp \rightarrow K^+ K^-, \pi^+ \pi^-$
$(1406 \pm 19) - i(80 \pm 6)$	⁴ KAMINSKI	99	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$
$(1300 \pm 20) - i(120 \pm 20)$	ANISOVICH	98B	RVUE Compilation
$(1290 \pm 15) - i(145 \pm 15)$	BARBERIS	97B	OMEG 450 $pp \rightarrow pp2(\pi^+ \pi^-)$
$(1548 \pm 40) - i(560 \pm 40)$	BERTIN	97C	OBLX $0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
$(1380 \pm 40) - i(180 \pm 25)$	ABELE	96B	CBAR $0.0 \bar{p}p \rightarrow \pi^0 K_L^0 K_L^0$
$(1300 \pm 15) - i(115 \pm 8)$	BUGG	96	RVUE
$(1330 \pm 50) - i(150 \pm 40)$	⁵ AMSLER	95B	CBAR $\bar{p}p \rightarrow 3\pi^0$
$(1360 \pm 35) - i(150-300)$	⁵ AMSLER	95C	CBAR $\bar{p}p \rightarrow \pi^0 \eta\eta$
$(1390 \pm 30) - i(190 \pm 40)$	⁶ AMSLER	95D	CBAR $\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta\eta, \pi^0 \pi^0 \eta$
1346 – i 249	^{7,8} JANSSEN	95	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
1214 – i 168	^{8,9} TORNQVIST	95	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
1364 – i 139	AMSLER	94D	CBAR $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$
$(1365^{+20}_{-55}) - i(134 \pm 35)$	ANISOVICH	94	CBAR $\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta\eta$
$(1340 \pm 40) - i(127^{+30}_{-20})$	¹⁰ BUGG	94	RVUE $\bar{p}p \rightarrow 3\pi^0, \eta\eta\pi^0, \eta\pi^0 \pi^0$
$(1430 \pm 5) - i(73 \pm 13)$	¹¹ KAMINSKI	94	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
1420 – i 220	¹² AU	87	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$

¹ Another pole is found at $(1510 \pm 130) - i(800^{+100}_{-150})$ MeV.

² Coupled channel analysis of $\pi^+ \pi^- \pi^0, K^+ K^- \pi^0$, and $K^\pm K_S^0 \pi^\mp$.

³ Average between $\pi^+ \pi^- 2\pi^0$ and $2(\pi^+ \pi^-)$.

⁴ T-matrix pole on sheet – – –.

⁵ Supersedes ANISOVICH 94.

⁶ Coupled-channel analysis of $\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta\eta$, and $\pi^0 \pi^0 \eta$ on sheet IV. Demonstrates explicitly that $f_0(500)$ and $f_0(1370)$ are two different poles.

⁷ Analysis of data from FALVARD 88.

⁸ The pole is on Sheet III. Demonstrates explicitly that $f_0(500)$ and $f_0(1370)$ are two different poles.

⁹ Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

¹⁰ Reanalysis of ANISOVICH 94 data.

¹¹ T-matrix pole on sheet III.

¹² Analysis of data from OCHS 73, GRAYER 74, BECKER 79, and CASON 83.

$f_0(1370)$ BREIT-WIGNER MASS OR K-MATRIX POLE PARAMETER

VALUE (MeV) DOCUMENT ID
1200 to 1500 OUR ESTIMATE

$\pi\pi$ MODE

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1400±40		¹ AUBERT	09L BABR	$B^\pm \rightarrow \pi^\pm \pi^\pm \pi^\mp$
1470 ⁺ ₋ $\frac{6+72}{7-255}$		² UEHARA	08A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
1259±55	2.6k	BONVICINI	07 CLEO	$D^+ \rightarrow \pi^- \pi^+ \pi^+$
1309± 1± 15		³ BUGG	07A RVUE	$0.0 p\bar{p} \rightarrow 3\pi^0$
1449±13	4.3k	⁴ GARMASH	06 BELL	$B^+ \rightarrow K^+ \pi^+ \pi^-$
1350±50		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-$
1265±30 ⁺ ₋ $\frac{20}{35}$		ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
1434±18± 9	848	AITALA	01A E791	$D_s^+ \rightarrow \pi^- \pi^+ \pi^+$
1308±10		BARBERIS	99B OMEG	$450 pp \rightarrow p_s p_f \pi^+ \pi^-$
1315±50		BELLAZZINI	99 GAM4	$450 pp \rightarrow pp \pi^0 \pi^0$
1315±30		ALDE	98 GAM4	$100 \pi^- p \rightarrow \pi^0 \pi^0 n$
1280±55		BERTIN	98 OBLX	$0.05-0.405 \bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$
1186		^{5,6} TORNQVIST	95 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
1472±12		ARMSTRONG	91 OMEG	$300 pp \rightarrow pp\pi\pi, ppK\bar{K}$
1275±20		BREAKSTONE	90 SFM	$62 pp \rightarrow pp\pi^+ \pi^-$
1420±20		AKESSON	86 SPEC	$63 pp \rightarrow pp\pi^+ \pi^-$
1256		FROGGATT	77 RVUE	$\pi^+ \pi^-$ channel

¹ Breit-Wigner mass.

² Breit-Wigner mass. May also be the $f_0(1500)$.

³ Reanalysis of ABELE 96C data.

⁴ Also observed by GARMASH 07 in $B^0 \rightarrow K_S^0 \pi^+ \pi^-$ decays. Supersedes GARMASH 05.

⁵ Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

⁶ Also observed by ASNER 00 in $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$ decays

$K\bar{K}$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$1360 \pm 31 \pm 28$	430	^{1,2} DOBBS	15	$J/\psi \rightarrow \gamma K^+ K^-$
$1350 \pm 48 \pm 15$	168	^{1,2} DOBBS	15	$\psi(2S) \rightarrow \gamma K^+ K^-$
1440 ± 6		VLADIMIRSK...06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1391 ± 10		TIKHOMIROV 03	SPEC	$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
1440 ± 50		BOLONKIN 88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1463 ± 9		ETKIN 82B	MPS	$23 \pi^- p \rightarrow n 2 K_S^0$
1425 ± 15		WICKLUND 80	SPEC	$6 \pi N \rightarrow K^+ K^- N$
~ 1300		POLYCHRO... 79	STRC	$7 \pi^- p \rightarrow n 2 K_S^0$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.² From a fit to a Breit-Wigner line shape with fixed $\Gamma = 346$ MeV. **4π MODE $2(\pi\pi)_S + \rho\rho$**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1395 ± 40		ABELE 01	CBAR	$0.0 \bar{p} d \rightarrow \pi^- 4\pi^0 p$
1374 ± 38		AMSLER 94	CBAR	$0.0 \bar{p} p \rightarrow \pi^+ \pi^- 3\pi^0$
1345 ± 12		ADAMO 93	OBLX	$\bar{n} p \rightarrow 3\pi^+ 2\pi^-$
1386 ± 30		GASPERO 93	DBC	$0.0 \bar{p} n \rightarrow 2\pi^+ 3\pi^-$
~ 1410	5751	¹ BETTINI 66	DBC	$0.0 \bar{p} n \rightarrow 2\pi^+ 3\pi^-$

¹ $\rho\rho$ dominant. **$\eta\eta$ MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$1262^{+51+82}_{-78-103}$	¹ UEHARA 10A	BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \eta\eta$
1430	AMSLER 92	CBAR	$0.0 \bar{p} p \rightarrow \pi^0 \eta\eta$
1220 ± 40	ALDE 86D	GAM4	$100 \pi^- p \rightarrow n 2\eta$

¹ Breit-Wigner mass. May also be the $f_0(1500)$.**COUPLED CHANNEL MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1306 ± 20	¹ ANISOVICH 03	RVUE	
¹ K-matrix pole from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K\bar{K}n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta\eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p} n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.			

 $f_0(1370)$ BREIT-WIGNER WIDTH

VALUE (MeV)	DOCUMENT ID
200 to 500 OUR ESTIMATE	

$\pi\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
300 ± 80		¹ AUBERT	09L BABR	$B^\pm \rightarrow \pi^\pm \pi^\pm \pi^\mp$
$90^+_{-1} \quad 2^+_{-1} \quad 50_{-22}$		² UEHARA	08A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
298 ± 21	2.6k	BONVICINI	07 CLEO	$D^+ \rightarrow \pi^- \pi^+ \pi^+$
126 ± 25	4286	³ GARMASH	06 BELL	$B^+ \rightarrow K^+ \pi^+ \pi^-$
265 ± 40		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-$
$350 \pm 100^+_{-60} \quad 105_{-60}$		ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
$173 \pm 32 \pm 6$	848	AITALA	01A E791	$D^+_{S^*} \rightarrow \pi^- \pi^+ \pi^+$
222 ± 20		BARBERIS	99B OMEG	$450 p p \rightarrow p_S p_f \pi^+ \pi^-$
255 ± 60		BELLAZZINI	99 GAM4	$450 p p \rightarrow p p \pi^0 \pi^0$
190 ± 50		ALDE	98 GAM4	$100 \pi^- p \rightarrow \pi^0 \pi^0 n$
323 ± 13		BERTIN	98 OBLX	$0.05-0.405 \bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$
350		^{4,5} TORNQVIST	95 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
195 ± 33		ARMSTRONG	91 OMEG	$300 p p \rightarrow p p \pi\pi, p p K\bar{K}$
285 ± 60		BREAKSTONE	90 SFM	$62 p p \rightarrow p p \pi^+ \pi^-$
460 ± 50		AKESSON	86 SPEC	$63 p p \rightarrow p p \pi^+ \pi^-$
~ 400		⁶ FROGGATT	77 RVUE	$\pi^+ \pi^-$ channel

¹ The systematic errors are not reported.

² Breit-Wigner width. May also be the $f_0(1500)$.

³ Also observed by GARMASH 07 in $B^0 \rightarrow K^0_S \pi^+ \pi^-$ decays. Supersedes GARMASH 05.

⁴ Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CA-SON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

⁵ Also observed by ASNER 00 in $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$ decays

⁶ Width defined as distance between 45 and 135° phase shift.

$K\bar{K}$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
121 ± 15	VLADIMIRSK...06	SPEC	$40 \pi^- p \rightarrow K^0_S K^0_S n$
55 ± 26	TIKHOMIROV 03	SPEC	$40.0 \pi^- C \rightarrow K^0_S K^0_S K^0_L X$
250 ± 80	BOLONKIN 88	SPEC	$40 \pi^- p \rightarrow K^0_S K^0_S n$
$118^+_{-16} \quad 138_{-16}$	ETKIN 82B	MPS	$23 \pi^- p \rightarrow n 2 K^0_S$
160 ± 30	WICKLUND 80	SPEC	$6 \pi N \rightarrow K^+ K^- N$
~ 150	POLYCHRO... 79	STRC	$7 \pi^- p \rightarrow n 2 K^0_S$

4π MODE $2(\pi\pi)_S + \rho\rho$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
275 ± 55		ABELE	01 CBAR	$0.0 \bar{p} d \rightarrow \pi^- 4\pi^0 p$
375 ± 61		AMSLER	94 CBAR	$0.0 \bar{p} p \rightarrow \pi^+ \pi^- 3\pi^0$
398 ± 26		ADAMO	93 OBLX	$\bar{n} p \rightarrow 3\pi^+ 2\pi^-$
310 ± 50		GASPERO	93 DBC	$0.0 \bar{p} n \rightarrow 2\pi^+ 3\pi^-$
~ 90	5751	¹ BETTINI	66 DBC	$0.0 \bar{p} n \rightarrow 2\pi^+ 3\pi^-$

¹ $\rho\rho$ dominant.

$\eta\eta$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$484^{+246+246}_{-170-263}$	¹ UEHARA	10A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \eta\eta$
250	AMSLER	92 CBAR	$0.0 \bar{p}p \rightarrow \pi^0 \eta\eta$
320 ± 40	ALDE	86D GAM4	$100 \pi^- p \rightarrow n 2\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Breit-Wigner width. May also be the $f_0(1500)$.

COUPLED CHANNEL MODE

VALUE (MeV)	DOCUMENT ID	TECN
147^{+30}_{-50}	¹ ANISOVICH	03 RVUE

¹ K-matrix pole from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K \bar{K} n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta\eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p}n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.

• • • We do not use the following data for averages, fits, limits, etc. • • •

$f_0(1370)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi$	seen
Γ_2 4π	seen
Γ_3 $4\pi^0$	seen
Γ_4 $2\pi^+ 2\pi^-$	seen
Γ_5 $\pi^+ \pi^- 2\pi^0$	seen
Γ_6 $\rho\rho$	dominant
Γ_7 $2(\pi\pi)_S\text{-wave}$	seen
Γ_8 $\pi(1300)\pi$	seen
Γ_9 $a_1(1260)\pi$	seen
Γ_{10} $\eta\eta$	seen
Γ_{11} $K\bar{K}$	seen
Γ_{12} $K\bar{K}n\pi$	not seen
Γ_{13} 6π	not seen
Γ_{14} $\omega\omega$	not seen
Γ_{15} $\gamma\gamma$	seen
Γ_{16} $e^+ e^-$	not seen

$f_0(1370)$ PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$ Γ_{15}
 See $\gamma\gamma$ widths under $f_0(500)$ and MORGAN 90.

$\Gamma(e^+ e^-)$ Γ_{16}

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<20	90	VOROBYEV	88 ND	$e^+ e^- \rightarrow \pi^0 \pi^0$

$f_0(1370) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{10}\Gamma_{15}/\Gamma$

VALUE (eV) DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

$121^{+133+169}_{-53-106}$ ¹ UEHARA 10A BELL 10.6 $e^+e^- \rightarrow e^+e^-\eta\eta$

¹ Including interference with the $f'_2(1525)$ (parameters fixed to the values from the 2008 edition of this review, PDG 08) and $f_2(1270)$. May also be the $f_0(1500)$.

$f_0(1370)$ BRANCHING RATIOS

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE CL% DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.10 95 OCHS 13 RVUE
 0.26 ± 0.09 BUGG 96 RVUE
 <0.15 ¹ AMSLER 94 CBAR $\bar{p}p \rightarrow \pi^+\pi^-3\pi^0$
 <0.06 GASPERO 93 DBC 0.0 $\bar{p}n \rightarrow \text{hadrons}$

¹ Using AMSLER 95B ($3\pi^0$).

$\Gamma(4\pi)/\Gamma_{\text{total}}$ $\Gamma_2/\Gamma = (\Gamma_3+\Gamma_4+\Gamma_5)/\Gamma$

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

>0.72 GASPERO 93 DBC 0.0 $\bar{p}n \rightarrow \text{hadrons}$

$\Gamma(4\pi^0)/\Gamma(4\pi)$ Γ_3/Γ_2

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen ABELE 96 CBAR 0.0 $\bar{p}p \rightarrow 5\pi^0$
 0.068 ± 0.005 ¹ GASPERO 93 DBC 0.0 $\bar{p}n \rightarrow \text{hadrons}$

¹ Model-dependent evaluation.

$\Gamma(2\pi^+2\pi^-)/\Gamma(4\pi)$ $\Gamma_4/\Gamma_2 = \Gamma_4/(\Gamma_3+\Gamma_4+\Gamma_5)$

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.420 ± 0.014 ¹ GASPERO 93 DBC 0.0 $\bar{p}n \rightarrow 2\pi^+3\pi^-$

¹ Model-dependent evaluation.

$\Gamma(\pi^+\pi^-2\pi^0)/\Gamma(4\pi)$ $\Gamma_5/\Gamma_2 = \Gamma_5/(\Gamma_3+\Gamma_4+\Gamma_5)$

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.512 ± 0.019 ¹ GASPERO 93 DBC 0.0 $\bar{p}n \rightarrow \text{hadrons}$

¹ Model-dependent evaluation.

$\Gamma(\rho\rho)/\Gamma(4\pi)$ Γ_6/Γ_2

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.26 ± 0.07 ABELE 01B CBAR 0.0 $\bar{p}d \rightarrow 5\pi p$

$\Gamma(2(\pi\pi)_{S\text{-wave}})/\Gamma(\pi\pi)$ Γ_7/Γ_1

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.6 ± 2.6 ¹ ABELE 01 CBAR 0.0 $\bar{p}d \rightarrow \pi^- 4\pi^0 p$

¹ From the combined data of ABELE 96 and ABELE 96C.

$\Gamma(2(\pi\pi)_{S\text{-wave}})/\Gamma(4\pi)$ Γ_7/Γ_2

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.51 ± 0.09 ABELE 01B CBAR 0.0 $\bar{p}d \rightarrow 5\pi p$

$\Gamma(\rho\rho)/\Gamma(2(\pi\pi)_{S\text{-wave}})$ Γ_6/Γ_7

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

large	BARBERIS	00C		450 $pp \rightarrow p_f 4\pi p_s$
1.6 ± 0.2	AMSLER	94	CBAR	$\bar{p}p \rightarrow \pi^+ \pi^- 3\pi^0$
~ 0.65	GASPERO	93	DBC	0.0 $\bar{p}n \rightarrow \text{hadrons}$

$\Gamma(\pi(1300)\pi)/\Gamma(4\pi)$ Γ_8/Γ_2

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.17 ± 0.06 ABELE 01B CBAR 0.0 $\bar{p}d \rightarrow 5\pi p$

$\Gamma(a_1(1260)\pi)/\Gamma(4\pi)$ Γ_9/Γ_2

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.06 ± 0.02 ABELE 01B CBAR 0.0 $\bar{p}d \rightarrow 5\pi p$

$\Gamma(\eta\eta)/\Gamma(4\pi)$ $\Gamma_{10}/\Gamma_2 = \Gamma_{10}/(\Gamma_3 + \Gamma_4 + \Gamma_5)$

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

(28 ± 11) × 10 ⁻³	¹ ANISOVICH	02D	SPEC	Combined fit
(4.7 ± 2.0) × 10 ⁻³	BARBERIS	00E		450 $pp \rightarrow p_f \eta \eta p_s$

¹ From a combined K-matrix analysis of Crystal Barrel (0. $p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta \eta, \pi^0 \pi^0 \eta$), GAMS ($\pi p \rightarrow \pi^0 \pi^0 n, \eta \eta n, \eta \eta' n$), and BNL ($\pi p \rightarrow K \bar{K} n$) data.

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE DOCUMENT ID TECN

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.35 ± 0.13 BUGG 96 RVUE

$\Gamma(K\bar{K})/\Gamma(\pi\pi)$

Γ_{11}/Γ_1

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.08±0.08	ABLIKIM	05	BES2	$J/\psi \rightarrow \phi\pi^+\pi^-, \phi K^+K^-$
0.91±0.20	¹ BARGIOTTI	03	OBLX	$\bar{p}p$
0.12±0.06	² ANISOVICH	02D	SPEC	Combined fit
0.46±0.15±0.11	BARBERIS	99D	OMEG	450 $pp \rightarrow K^+K^-, \pi^+\pi^-$

¹ Coupled channel analysis of $\pi^+\pi^-\pi^0, K^+K^-\pi^0,$ and $K^\pm K_S^0 \pi^\mp$.

² From a combined K-matrix analysis of Crystal Barrel ($0. \ p\bar{p} \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$), GAMS ($\pi p \rightarrow \pi^0\pi^0 n, \eta\eta n, \eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K} n$) data.

$\Gamma(K\bar{K}n\pi)/\Gamma_{total}$

Γ_{12}/Γ

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.03	GASPERO	93	DBC	0.0 $\bar{p}n \rightarrow$ hadrons
-------	---------	----	-----	------------------------------------

$\Gamma(6\pi)/\Gamma_{total}$

Γ_{13}/Γ

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.22	GASPERO	93	DBC	0.0 $\bar{p}n \rightarrow$ hadrons
-------	---------	----	-----	------------------------------------

$\Gamma(\omega\omega)/\Gamma_{total}$

Γ_{14}/Γ

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.13	GASPERO	93	DBC	0.0 $\bar{p}n \rightarrow$ hadrons
-------	---------	----	-----	------------------------------------

$f_0(1370)$ REFERENCES

DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
OCHS	13	JP G40 043001	W. Ochs	
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	
AUBERT	09L	PR D79 072006	B. Aubert <i>et al.</i>	(BABAR Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
UEHARA	08A	PR D78 052004	S. Uehara <i>et al.</i>	(BELLE Collab.)
BONVICINI	07	PR D76 012001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
BUGG	07A	JP G34 151	D.V. Bugg <i>et al.</i>	
GARMASH	07	PR D75 012006	A. Garmash <i>et al.</i>	(BELLE Collab.)
GARMASH	06	PRL 96 251803	A. Garmash <i>et al.</i>	(BELLE Collab.)
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)
VLADIMIRSK...	06	PAN 69 493	V.V. Vladimirov <i>et al.</i>	(ITEP, Moscow)
		Translated from YAF 69 515.		
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)
GARMASH	05	PR D71 092003	A. Garmash <i>et al.</i>	(BELLE Collab.)
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>	
BARGIOTTI	03	EPJ C26 371	M. Bargiotti <i>et al.</i>	(OBELIX Collab.)
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>	
		Translated from YAF 66 860.		
ANISOVICH	02D	PAN 65 1545	V.V. Anisovich <i>et al.</i>	
		Translated from YAF 65 1583.		

ABELE	01	EPJ C19 667	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE	01B	EPJ C21 261	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
AITALA	01A	PRL 86 765	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
ASNER	00	PR D61 012002	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	99B	PL B453 316	D. Barberis <i>et al.</i>	(Omega Expt.)
BARBERIS	99D	PL B462 462	D. Barberis <i>et al.</i>	(Omega Expt.)
BELLAZZINI	99	PL B467 296	R. Bellazzini <i>et al.</i>	
KAMINSKI	99	EPJ C9 141	R. Kaminski, L. Lesniak, B. Loiseau	(CRAC, PARIN)
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)
Also		PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)
		Translated from YAF 62 446.		
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>	
		Translated from UFN 168 481.		
BERTIN	98	PR D57 55	A. Bertin <i>et al.</i>	(OBELIX Collab.)
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)
ABELE	96	PL B380 453	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE	96B	PL B385 425	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE	96C	NP A609 562	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
BUGG	96	NP B471 59	D.V. Bugg, A.V. Sarantsev, B.S. Zou	(LOQM, PNPI)
AMSLER	95B	PL B342 433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	95C	PL B353 571	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	95D	PL B355 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
JANSEN	95	PR D52 2690	G. Janssen <i>et al.</i>	(STON, ADLD, JULI)
TORNQVIST	95	ZPHY C68 647	N.A. Tornqvist	(HELS)
AMSLER	94	PL B322 431	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.) JPC
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ANISOVICH	94	PL B323 233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.) JPC
BUGG	94	PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)
KAMINSKI	94	PR D50 3145	R. Kaminski, L. Lesniak, J.P. Maillet	(CRAC+)
ADAMO	93	NP A558 13C	A. Adamo <i>et al.</i>	(OBELIX Collab.) JPC
GASPERO	93	NP A562 407	M. Gaspero	(ROMA1) JPC
AMSLER	92	PL B291 347	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ARMSTRONG	91	ZPHY C51 351	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)
ARMSTRONG	91B	ZPHY C52 389	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)
BREAKSTONE	90	ZPHY C48 569	A.M. Breakstone <i>et al.</i>	(ISU, BGNA, CERN+)
MORGAN	90	ZPHY C48 623	D. Morgan, M.R. Pennington	(RAL, DURH)
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
BOLONKIN	88	NP B309 426	B.V. Bolonkin <i>et al.</i>	(ITEP, SERP)
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)
		Translated from YAF 48 436.		
AU	87	PR D35 1633	K.L. Au, D. Morgan, M.R. Pennington	(DURH, RAL)
AKESSON	86	NP B264 154	T. Akesson <i>et al.</i>	(Axial Field Spec. Collab.)
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)
CASON	83	PR D28 1586	N.M. Cason <i>et al.</i>	(NDAM, ANL)
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)
WICKLUND	80	PRL 45 1469	A.B. Wicklund <i>et al.</i>	(ANL)
BECKER	79	NP B151 46	H. Becker <i>et al.</i>	(MPIM, CERN, ZEEM, CRAC)
POLYCHRO...	79	PR D19 1317	V.A. Polychronakos <i>et al.</i>	(NDAM, ANL)
FROGGATT	77	NP B129 89	C.D. Froggatt, J.L. Petersen	(GLAS, NORD)
ROSSELET	77	PR D15 574	L. Rosselet <i>et al.</i>	(GEVA, SACL)
GRAYER	74	NP B75 189	G. Grayer <i>et al.</i>	(CERN, MPIM)
HYAMS	73	NP B64 134	B.D. Hyams <i>et al.</i>	(CERN, MPIM)
OCHS	73	Thesis	W. Ochs	(MPIM, MUNI)
BEIER	72B	PRL 29 511	E.W. Beier <i>et al.</i>	(PENN)
BETTINI	66	NC 42A 695	A. Bettini <i>et al.</i>	(PADO, PISA)