

New Heavy Bosons (W' , Z' , leptoquarks, etc.), Searches for

We list here various limits on charged and neutral heavy vector bosons (other than W 's and Z 's), heavy scalar bosons (other than Higgs bosons), vector or scalar leptoquarks, and axigluons. The latest unpublished results are described in " W' Searches" and " Z' Searches" reviews. For recent searches on scalar bosons which could be identified as Higgs bosons, see the listings in the Higgs boson section.

CONTENTS:

Mass Limits for W' (Heavy Charged Vector Boson Other Than W) in Hadron Collider Experiments

W_R (Right-Handed W Boson) Mass Limits

Limit on W_L - W_R Mixing Angle ζ

Mass Limits for Z' (Heavy Neutral Vector Boson Other Than Z)

- Limits for Z'_{SM}
- Limits for Z'_{LR}
- Limits for Z'_X
- Limits for Z'_ψ
- Limits for Z'_η
- Limits for other Z'
- Searches for Z' with Lepton-Flavor-Violating decays

Indirect Constraints on Kaluza-Klein Gauge Bosons

Mass Limits for Leptoquarks from Pair Production

Mass Limits for Leptoquarks from Single Production

Indirect Limits for Leptoquarks

Mass Limits for Diquarks

Mass Limits for g_A (axigluon) and Other Color-Octet Gauge Bosons

Mass Limits for Color-Octet Scalar Bosons

X^0 (Heavy Boson) Searches in Z Decays

Mass Limits for a Heavy Neutral Boson Coupling to e^+e^-

Search for X^0 Resonance in e^+e^- Collisions

Search for X^0 Resonance in $e p$ Collisions

Search for X^0 Resonance in Two-Photon Process

Search for X^0 Resonance in $e^+e^- \rightarrow X^0\gamma$

Search for X^0 Resonance in $Z \rightarrow f\bar{f}X^0$

Search for X^0 Resonance in WX^0 final state

Search for X^0 Resonance in Quarkonium Decays

See the related review(s):

[\$W'\$ -Boson Searches](#)

MASS LIMITS for W' (Heavy Charged Vector Boson Other Than W) in Hadron Collider Experiments

Couplings of W' to quarks and leptons are taken to be identical with those of W . The following limits are obtained from $p\bar{p}$ or $pp \rightarrow W'X$ with W' decaying to the mode

indicated in the comments. New decay channels (e.g., $W' \rightarrow WZ$) are assumed to be suppressed. The most recent preliminary results can be found in the “ W' -boson searches” review above.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
none 1200–3300	95	1 AABOUD 18F	ATLS	$W' \rightarrow WZ$
none 1000–3600	95	2 SIRUNYAN 18	CMS	$W' \rightarrow tb$
>3600	95	3 AABOUD 17AK	ATLS	$W' \rightarrow q\bar{q}$
none 1100–2500	95	4 AABOUD 17AO	ATLS	$W' \rightarrow hW$
>2220	95	5 AABOUD 17B	ATLS	$W' \rightarrow hW$
>2300	95	6 KHACHATRY...17J	CMS	$W' \rightarrow N_{\tau}\tau \rightarrow \tau\tau jj$
none 600–2700	95	7 KHACHATRY...17W	CMS	$W' \rightarrow q\bar{q}$
>4100	95	8 KHACHATRY...17Z	CMS	$W' \rightarrow e\nu, \mu\nu$
>2200	95	9 SIRUNYAN 17A	CMS	$W' \rightarrow WZ$
>2300	95	10 SIRUNYAN 17AK	CMS	$W' \rightarrow WZ, hW$
>2900	95	11 SIRUNYAN 17H	CMS	$W' \rightarrow \tau N$
>2600	95	12 SIRUNYAN 17I	CMS	$W' \rightarrow tb$
>2450	95	13 SIRUNYAN 17R	CMS	$W' \rightarrow hW$
none 2780–3150	95	13 SIRUNYAN 17R	CMS	$W' \rightarrow hW$
>2600	95	14 AABOUD 16AE	ATLS	$W' \rightarrow WZ$
>4070	95	15 AABOUD 16V	ATLS	$W' \rightarrow e\nu, \mu\nu$
>1810	95	16 AAD 16R	ATLS	$W' \rightarrow WZ$
>2600	95	17 AAD 16S	ATLS	$W' \rightarrow q\bar{q}$
>2150	95	18 KHACHATRY...16AO	CMS	$W' \rightarrow tb$
none 1000–1600	95	19 KHACHATRY...16AP	CMS	$W' \rightarrow hW$
none 800–1500	95	20 KHACHATRY...16BD	CMS	$W' \rightarrow hW \rightarrow b\bar{b}l\nu$
none 1500–2600	95	21 KHACHATRY...16K	CMS	$W' \rightarrow q\bar{q}$
none 500–1600	95	22 KHACHATRY...16L	CMS	$W' \rightarrow q\bar{q}$
none 300–2700	95	23 KHACHATRY...16O	CMS	$W' \rightarrow \tau\nu$
none 400–1590	95	24 AAD 15AU	ATLS	$W' \rightarrow WZ$
none 1500–1760	95	25 AAD 15AV	ATLS	$W' \rightarrow tb$
none 300–1490	95	26 AAD 15AZ	ATLS	$W' \rightarrow WZ$
none 1300–1500	95	27 AAD 15CP	ATLS	$W' \rightarrow WZ$
none 500–1920	95	28 AAD 15R	ATLS	$W' \rightarrow tb$
none 800–2450	95	29 AAD 15V	ATLS	$W' \rightarrow q\bar{q}$
>1470	95	30 KHACHATRY...15C	CMS	$W' \rightarrow WZ$
>3710	95	31 KHACHATRY...15T	CMS	$W' \rightarrow e\nu, \mu\nu$
none 1000–3010	95	32 KHACHATRY...14O	CMS	$W' \rightarrow Nl \rightarrow lljj$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
		33 KHACHATRY...17U	CMS	$W' \rightarrow hW$
		34 AAD 15BB	ATLS	$W' \rightarrow Wh$
none 300–880	95	35 AALTONEN 15C	CDF	$W' \rightarrow tb$
none 1200–1900 and 2000–2200	95	36 KHACHATRY...15V	CMS	$W' \rightarrow q\bar{q}$
>3240	95	AAD 14AI	ATLS	$W' \rightarrow e\nu, \mu\nu$
		37 AAD 14AT	ATLS	$W' \rightarrow W\gamma$
none 200–1520	95	38 AAD 14S	ATLS	$W' \rightarrow WZ$
none 1000–1700	95	39 KHACHATRY...14	CMS	$W' \rightarrow WZ$
		40 KHACHATRY...14A	CMS	$W' \rightarrow WZ$
none 500–950	95	41 AAD 13AO	ATLS	$W' \rightarrow WZ$
none 1100–1680	95	AAD 13D	ATLS	$W' \rightarrow q\bar{q}$

none 1000–1920	95	42	CHATRCHYAN 13A	CMS	$W' \rightarrow q\bar{q}$
		42	CHATRCHYAN 13AJ	CMS	$W' \rightarrow WZ$
>2900	95	43	CHATRCHYAN 13AQ	CMS	$W' \rightarrow e\nu, \mu\nu$
none 800–1510	95	44	CHATRCHYAN 13E	CMS	$W' \rightarrow tb$
none 700–940	95	45	CHATRCHYAN 13U	CMS	$W' \rightarrow WZ$
none 700–1130	95	46	AAD	12AV ATLS	$W' \rightarrow tb$
none 200–760	95	47	AAD	12BB ATLS	$W' \rightarrow WZ$
		48	AAD	12CK ATLS	$W' \rightarrow \bar{\tau}q$
>2550	95	49	AAD	12CR ATLS	$W' \rightarrow e\nu, \mu\nu$
		50	AAD	12M ATLS	$W' \rightarrow N\ell \rightarrow lljj$
		51	AALTONEN	12N CDF	$W' \rightarrow \bar{\tau}q$
none 200–1143	95	47	CHATRCHYAN 12AF	CMS	$W' \rightarrow WZ$
		52	CHATRCHYAN 12AR	CMS	$W' \rightarrow \bar{\tau}q$
		53	CHATRCHYAN 12BG	CMS	$W' \rightarrow N\ell \rightarrow lljj$
>1120	95		AALTONEN	11C CDF	$W' \rightarrow e\nu$
none 180–690	95	54	ABAZOV	11H D0	$W' \rightarrow WZ$
none 600–863	95	55	ABAZOV	11L D0	$W' \rightarrow tb$
none 285–516	95	56	AALTONEN	10N CDF	$W' \rightarrow WZ$
none 280–840	95	57	AALTONEN	09AC CDF	$W' \rightarrow q\bar{q}$
>1000	95		ABAZOV	08C D0	$W' \rightarrow e\nu$
none 300–800	95		ABAZOV	04C D0	$W' \rightarrow q\bar{q}$
none 225–536	95	58	ACOSTA	03B CDF	$W' \rightarrow tb$
none 200–480	95	59	AFFOLDER	02C CDF	$W' \rightarrow WZ$
> 786	95	60	AFFOLDER	01I CDF	$W' \rightarrow e\nu, \mu\nu$
none 300–420	95	61	ABE	97G CDF	$W' \rightarrow q\bar{q}$
> 720	95	62	ABACHI	96C D0	$W' \rightarrow e\nu$
> 610	95	63	ABACHI	95E D0	$W' \rightarrow e\nu, \tau\nu$
none 260–600	95	64	RIZZO	93 RVUE	$W' \rightarrow q\bar{q}$

¹ AABOUD 18F search for resonances decaying to WZ in pp collisions at $\sqrt{s} = 13$ TeV.

The quoted limit is for heavy-vector-triplet W' with $g_V = 3$. The limit becomes $M_{W'} > 3000$ GeV for $g_V = 1$. If we assume $M_{Z'} = M_{W'}$, the limit increases $M_{W'} > 3500$ GeV and $M_{W'} > 3100$ GeV for $g_V = 3$ and $g_V = 1$, respectively. See their Fig.5 for limits on $\sigma \cdot B$.

² SIRUNYAN 18 limit is for right-handed W' using pp collisions at $\sqrt{s} = 13$ TeV. $W' \rightarrow \ell\nu_R$ decay is assumed to be forbidden. The limit becomes $M_{W'} > 3.4$ TeV if $M_{\nu_R} \ll$

$M_{W'}$. See their Fig. 5 for exclusion limits on W' models having both left- and right-handed couplings.

³ AABOUD 17AK search for a new resonance decaying to dijets in pp collisions at $\sqrt{s} = 13$ TeV. The limit above is for a W' boson having axial-vector SM couplings and decaying to quarks with 75% branching fraction.

⁴ AABOUD 17AO search for resonances decaying to hW in pp collisions at $\sqrt{s} = 13$ TeV. The limit quoted above is for a W' in the heavy-vector-triplet model with $g_V = 3$. See their Fig.4 for limits on $\sigma \cdot B$.

⁵ AABOUD 17B search for resonances decaying to hW ($h \rightarrow b\bar{b}, c\bar{c}$; $W \rightarrow \ell\nu$) in pp collisions at $\sqrt{s} = 13$ TeV. The quoted limit is for heavy-vector-triplet W' with $g_V = 3$. The limit becomes $M_{W'} > 1750$ GeV for $g_V = 1$. If we assume $M_{W'} = M_{Z'}$, the limit increases $M_{W'} > 2310$ GeV and $M_{W'} > 1730$ GeV for $g_V = 3$ and $g_V = 1$, respectively. See their Fig.3 for limits on $\sigma \cdot B$.

- ⁶ KHACHATRYAN 17J search for right-handed W_R in pp collisions at $\sqrt{s} = 13$ TeV. W_R is assumed to decay into τ and hypothetical heavy neutrino N_τ , with N_τ decaying into τjj . The quoted limit is for $M_{N_\tau} = M_{W_R}/2$. The limit becomes $M_{W_R} > 2350$ GeV (1630 GeV) for $M_{W_R}/M_{N_\tau} = 0.8$ (0.2). See their Fig. 4 for excluded regions in the $M_{W_R} - M_{N_\tau}$ plane.
- ⁷ KHACHATRYAN 17W search for resonances decaying to dijets in pp collisions at $\sqrt{s} = 13$ TeV.
- ⁸ KHACHATRYAN 17Z limit is for W' with SM-like coupling using pp collisions at $\sqrt{s} = 13$ TeV. The bosonic decays of W' and the interference with SM W process are neglected.
- ⁹ SIRUNYAN 17A search for resonances decaying to WZ with $WZ \rightarrow \ell\nu q\bar{q}, q\bar{q}q\bar{q}$ in pp collisions at $\sqrt{s} = 13$ TeV. The quoted limit is for heavy-vector-triplet W' with $g_V = 3$. The limit becomes $M_{W'} > 2000$ GeV for $g_V = 1$. If we assume $M_{Z'} = M_{W'}$, the limit increases $M_{W'} > 2400$ GeV and $M_{W'} > 2300$ GeV for $g_V = 3$ and $g_V = 1$, respectively. See their Fig.6 for limits on $\sigma \cdot B$.
- ¹⁰ SIRUNYAN 17AK search for resonances decaying to WZ or hW in pp collisions at $\sqrt{s} = 8$ and 13 TeV. The quoted limit is for heavy-vector-triplet W' with $g_V = 3$. The limit becomes $M_{W'} > 2300$ GeV for $g_V = 1$. If we assume $M_{W'} = M_{Z'}$, the limit increases $M_{W'} > 2400$ GeV for both $g_V = 3$ and $g_V = 1$. See their Fig.1 and 2 for limits on $\sigma \cdot B$.
- ¹¹ SIRUNYAN 17H search for right-handed W' in pp collisions at $\sqrt{s} = 13$ TeV. W' is assumed to decay into τ and a heavy neutrino N , with N decaying to $\tau q\bar{q}$. The limit above assumes $M_N = M_{W'}/2$.
- ¹² SIRUNYAN 17I limit is for a right-handed W' using pp collisions at $\sqrt{s} = 13$ TeV. The limit becomes $M_{W'} > 2400$ GeV for $M_{\nu_R} \ll M_{W'}$.
- ¹³ SIRUNYAN 17R search for resonances decaying to hW in pp collisions at $\sqrt{s} = 13$ TeV. The quoted limit is for heavy-vector-triplet W' with $g_V = 3$. Mass regions $M_{W'} < 2370$ GeV and $2870 < M_{W'} < 2970$ GeV are excluded for $g_V = 1$. If we assume $M_{Z'} = M_{W'}$, the excluded mass regions are $1000 < M_{W'} < 2500$ GeV and $2760 < M_{W'} < 3300$ GeV for $g_V = 3$; $1000 < M_{W'} < 2430$ GeV and $2810 < M_{W'} < 3130$ GeV for $g_V = 1$. See their Fig.5 for limits on $\sigma \cdot B$.
- ¹⁴ AABOUD 16AE search for resonances decaying to VV ($V = W$ or Z) in pp collisions at $\sqrt{s} = 13$ TeV. Results from $\nu\nu qq, \nu\ell qq, \ell\ell qq$ and $qqqq$ final states are combined. The quoted limit is for a heavy-vector-triplet W' with $g_V = 3$ and $M_{W'} = M_{Z'}$.
- ¹⁵ AABOUD 16V limit is for W' with SM-like coupling using pp collisions at $\sqrt{s} = 13$ TeV. The bosonic decays of W' and the interference with SM W process are neglected.
- ¹⁶ AAD 16R search for $W' \rightarrow WZ$ in pp collisions at $\sqrt{s} = 8$ TeV. $\ell\nu\ell'\ell', \ell\ell q\bar{q}, \ell\nu q\bar{q}$, and all hadronic channels are combined. The quoted limit assumes $g_{W'WZ}/g_{WWZ} = (M_W/M_{W'})^2$.
- ¹⁷ AAD 16S search for a new resonance decaying to dijets in pp collisions at $\sqrt{s} = 13$ TeV. The limit quoted above is for a W' having SM-like couplings to quarks.
- ¹⁸ KHACHATRYAN 16AO limit is for a SM-like right-handed W' using pp collisions at $\sqrt{s} = 8$ TeV. The quoted limit combines $t \rightarrow qq\bar{b}$ and $t \rightarrow \ell\nu b$ events.
- ¹⁹ KHACHATRYAN 16AP search for a resonance decaying to hW in pp collisions at $\sqrt{s} = 8$ TeV. Both h and W are assumed to decay to fat jets. The quoted limit is for heavy-vector-triplet W' with $g_V = 3$.
- ²⁰ KHACHATRYAN 16BD search for resonance decaying to hW in pp collisions at $\sqrt{s} = 8$ TeV. The quoted limit is for heavy-vector-triplet (HVT) W' with $g_V = 3$. The HVT model $m_{W'} = m_{Z'} > 1.8$ TeV is also obtained by combining $W'/Z' \rightarrow Wh/Zh \rightarrow \ell\nu bb, qq\tau\tau, qqbb$, and $qqqqqq$ channels.

- 21 KHACHATRYAN 16K search for resonances decaying to dijets in pp collisions at $\sqrt{s} = 13$ TeV.
- 22 KHACHATRYAN 16L search for resonances decaying to dijets in pp collisions at $\sqrt{s} = 8$ TeV with the data scouting technique, increasing the sensitivity to the low mass resonances.
- 23 KHACHATRYAN 16O limit is for W' having universal couplings. Interferences with the SM amplitudes are assumed to be absent.
- 24 AAD 15AU search for W' decaying into the WZ final state with $W \rightarrow q\bar{q}'$, $Z \rightarrow \ell^+\ell^-$ using pp collisions at $\sqrt{s} = 8$ TeV. The quoted limit assumes $g_{W'WZ}/g_{WWZ} = (M_W/M_{W'})^2$.
- 25 AAD 15AV limit is for a SM like right-handed W' using pp collisions at $\sqrt{s} = 8$ TeV. $W' \rightarrow \ell\nu$ decay is assumed to be forbidden.
- 26 AAD 15AZ search for W' decaying into the WZ final state with $W \rightarrow \ell\nu$, $Z \rightarrow q\bar{q}$ using pp collisions at $\sqrt{s} = 8$ TeV. The quoted limit assumes $g_{W'WZ}/g_{WWZ} = (M_W/M_{W'})^2$.
- 27 AAD 15CP search for W' decaying into the WZ final state with $W \rightarrow q\bar{q}$, $Z \rightarrow q\bar{q}$ using pp collisions at $\sqrt{s} = 8$ TeV. The quoted limit assumes $g_{W'WZ}/g_{WWZ} = (M_W/M_{W'})^2$.
- 28 AAD 15R limit is for a SM like right-handed W' using pp collisions at $\sqrt{s} = 8$ TeV. $W' \rightarrow \ell\nu$ decay is assumed to be forbidden.
- 29 AAD 15V search for new resonance decaying to dijets in pp collisions at $\sqrt{s} = 8$ TeV.
- 30 KHACHATRYAN 15C search for W' decaying via WZ to fully leptonic final states using pp collisions at $\sqrt{s}=8$ TeV. The quoted limit assumes $g_{W'WZ}/g_{WWZ} = M_W M_Z/M_{W'}^2$.
- 31 KHACHATRYAN 15T limit is for W' with SM-like coupling which interferes the SM W boson constructively using pp collisions at $\sqrt{s} = 8$ TeV. For W' without interference, the limit becomes > 3280 GeV.
- 32 KHACHATRYAN 14O search for right-handed W_R in pp collisions at $\sqrt{s} = 8$ TeV. W_R is assumed to decay into ℓ and hypothetical heavy neutrino N , with N decaying into ℓjj . The quoted limit is for $M_{\nu_{eR}} = M_{\nu_{\mu R}} = M_{W_R}/2$. See their Fig. 3 and Fig. 5 for excluded regions in the $M_{W_R} - M_\nu$ plane.
- 33 KHACHATRYAN 17U search for resonances decaying to hW ($h \rightarrow b\bar{b}$; $W \rightarrow \ell\nu$) in pp collisions at $\sqrt{s} = 13$ TeV. The limit on the heavy-vector-triplet model is $M_{Z'} = M_{W'} > 2$ TeV for $g_V = 3$, in which constraints from the $Z' \rightarrow hZ$ ($h \rightarrow b\bar{b}$; $Z \rightarrow \ell^+\ell^-, \nu\bar{\nu}$) are combined. See their Fig.3 and Fig.4 for limits on $\sigma \cdot B$.
- 34 AAD 15BB search for W' decaying into Wh with $W \rightarrow \ell\nu$, $h \rightarrow b\bar{b}$. See their Fig. 4 for the exclusion limits in the heavy vector triplet benchmark model parameter space.
- 35 AALTONEN 15C limit is for a SM-like right-handed W' assuming $W' \rightarrow \ell\nu$ decays are forbidden, using $p\bar{p}$ collisions at $\sqrt{s}=1.96$ TeV. See their Fig. 3 for limit on $g_{W'}/g_W$.
- 36 KHACHATRYAN 15V search new resonance decaying to dijets in pp collisions at $\sqrt{s} = 8$ TeV.
- 37 AAD 14AT search for a narrow charged vector boson decaying to $W\gamma$. See their Fig. 3a for the exclusion limit in $m_{W'} - \sigma B$ plane.
- 38 AAD 14S search for W' decaying into the WZ final state with $W \rightarrow \ell\nu$, $Z \rightarrow \ell\ell$ using pp collisions at $\sqrt{s}=8$ TeV. The quoted limit assumes $g_{W'WZ}/g_{WWZ} = (M_W/M_{W'})^2$.
- 39 KHACHATRYAN 14 search for W' decaying into WZ final state with $W \rightarrow q\bar{q}$, $Z \rightarrow q\bar{q}$ using pp collisions at $\sqrt{s}=8$ TeV. The quoted limit assumes $g_{W'WZ}/g_{WWZ} = (M_W/M_{W'})^2$.

- 40 KHACHATRYAN 14A search for W' decaying into the WZ final state with $W \rightarrow \ell\nu$, $Z \rightarrow q\bar{q}$, or $W \rightarrow q\bar{q}$, $Z \rightarrow \ell\ell$. pp collisions data at $\sqrt{s}=8$ TeV are used for the search. See their Fig. 13 for the exclusion limit on the number of events in the mass–width plane.
- 41 AAD 13AO search for W' decaying into the WZ final state with $W \rightarrow \ell\nu$, $Z \rightarrow 2j$ using pp collisions at $\sqrt{s}=7$ TeV. The quoted limit assumes $g_{W'WZ}/g_{WWZ} = (M_W/M_{W'})^2$.
- 42 CHATRCHYAN 13AJ search for resonances decaying to WZ pair, using the hadronic decay modes of W and Z , in pp collisions at $\sqrt{s}=7$ TeV. See their Fig. 7 for the limit on the cross section.
- 43 CHATRCHYAN 13AQ limit is for W' with SM-like coupling which interferes with the SM W boson using pp collisions at $\sqrt{s}=7$ TeV.
- 44 CHATRCHYAN 13E limit is for W' with SM-like coupling which interfere with the SM W boson using pp collisions at $\sqrt{s}=7$ TeV. For W' with right-handed coupling, the bound becomes >1850 GeV (>1910 GeV) if W' decays to both leptons and quarks (only to quarks). If both left- and right-handed couplings are present, the limit becomes >1640 GeV.
- 45 CHATRCHYAN 13U search for W' decaying to the WZ final state, with W decaying into jets, in pp collisions at $\sqrt{s}=7$ TeV. The quoted limit assumes $g_{W'WZ}/g_{WWZ} = (M_W/M_{W'})^2$.
- 46 The AAD 12AV quoted limit is for a SM-like right-handed W' using pp collisions at $\sqrt{s}=7$ TeV. $W' \rightarrow \ell\nu$ decay is assumed to be forbidden.
- 47 AAD 12BB use pp collisions data at $\sqrt{s}=7$ TeV. The quoted limit assumes $g_{W'WZ}/g_{WWZ} = (M_W/M_{W'})^2$.
- 48 AAD 12CK search for $pp \rightarrow tW'$, $W' \rightarrow \bar{t}q$ events in pp collisions. See their Fig. 5 for the limit on $\sigma \cdot B$.
- 49 AAD 12CR use pp collisions at $\sqrt{s}=7$ TeV.
- 50 AAD 12M search for right-handed W_R in pp collisions at $\sqrt{s} = 7$ TeV. W_R is assumed to decay into ℓ and hypothetical heavy neutrino N , with N decaying into ℓjj . See their Fig. 4 for the limit in the $m_N - m_{W'}$ plane.
- 51 AALTONEN 12N search for $p\bar{p} \rightarrow tW'$, $W' \rightarrow \bar{t}d$ events in $p\bar{p}$ collisions. See their Fig. 3 for the limit on $\sigma \cdot B$.
- 52 CHATRCHYAN 12AR search for $pp \rightarrow tW'$, $W' \rightarrow \bar{t}d$ events in pp collisions. See their Fig. 2 for the limit on $\sigma \cdot B$.
- 53 CHATRCHYAN 12BG search for right-handed W_R in pp collisions $\sqrt{s} = 7$ TeV. W_R is assumed to decay into ℓ and hypothetical heavy neutrino N , with N decaying into ℓjj . See their Fig. 3 for the limit in the $m_N - m_{W'}$ plane.
- 54 ABAZOV 11H use data from $p\bar{p}$ collisions at $\sqrt{s}=1.96$ TeV. The quoted limit is obtained assuming $W'WZ$ coupling strength is the same as the ordinary WWZ coupling strength in the Standard Model.
- 55 ABAZOV 11L limit is for W' with SM-like coupling which interferes with the SM W boson, using $p\bar{p}$ collisions at $\sqrt{s}=1.96$ TeV. For W' with right-handed coupling, the bound becomes >885 GeV (>890 GeV) if W' decays to both leptons and quarks (only to quarks). If both left- and right-handed couplings present, the limit becomes >916 GeV.
- 56 AALTONEN 10N use $p\bar{p}$ collision data at $\sqrt{s}=1.96$ TeV. The quoted limit assumes $g_{W'WZ}/g_{WWZ} = (M_W/M_{W'})^2$. See their Fig. 4 for limits in mass-coupling plane.
- 57 AALTONEN 09AC search for new particle decaying to dijets using $p\bar{p}$ collisions at $\sqrt{s}=1.96$ TeV.
- 58 The ACOSTA 03B quoted limit is for $M_{W'} \gg M_{\nu_R}$, using $p\bar{p}$ collisions at $\sqrt{s}=1.8$ TeV. For $M_{W'} < M_{\nu_R}$, $M_{W'}$ between 225 and 566 GeV is excluded.

- 59 The quoted limit is obtained assuming $W' W Z$ coupling strength is the same as the ordinary $W W Z$ coupling strength in the Standard Model, using $p\bar{p}$ collisions at $\sqrt{s}=1.8$ TeV. See their Fig. 2 for the limits on the production cross sections as a function of the W' width.
- 60 AFFOLDER 01I combine a new bound on $W' \rightarrow e\nu$ of 754 GeV, using $p\bar{p}$ collisions at $\sqrt{s}=1.8$ TeV, with the bound of ABE 00 on $W' \rightarrow \mu\nu$ to obtain quoted bound.
- 61 ABE 97G search for new particle decaying to dijets using $p\bar{p}$ collisions at $\sqrt{s}=1.8$ TeV.
- 62 For bounds on W_R with nonzero right-handed mass, see Fig. 5 from ABACHI 96C.
- 63 ABACHI 95E assume that the decay $W' \rightarrow W Z$ is suppressed and that the neutrino from W' decay is stable and has a mass significantly less $m_{W'}$.
- 64 RIZZO 93 analyses CDF limit on possible two-jet resonances. The limit is sensitive to the inclusion of the assumed K factor.

W_R (Right-Handed W Boson) MASS LIMITS

Assuming a light right-handed neutrino, except for BEALL 82, LANGACKER 89B, and COLANGELO 91. $g_R = g_L$ assumed. [Limits in the section MASS LIMITS for W' below are also valid for W_R if $m_{\nu_R} \ll m_{W_R}$.] Some limits assume manifest left-right symmetry, *i.e.*, the equality of left- and right Cabibbo-Kobayashi-Maskawa matrices. For a comprehensive review, see LANGACKER 89B. Limits on the W_L - W_R mixing angle ζ are found in the next section. Values in brackets are from cosmological and astrophysical considerations and assume a light right-handed neutrino.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
> 592	90	1 BUENO	11	TWST μ decay
> 715	90	2 CZAKON	99	RVUE Electroweak
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
> 235	90	3 PRIEELS	14	PIE3 μ decay
> 245	90	4 WAUTERS	10	CNTR ^{60}Co β decay
>2500		5 ZHANG	08	THEO $m_{K_L^0} - m_{K_S^0}$
> 180	90	6 MELCONIAN	07	CNTR ^{37}K β^+ decay
> 290.7	90	7 SCHUMANN	07	CNTR Polarized neutron decay
[> 3300]	95	8 CYBURT	05	COSM Nucleosynthesis; light ν_R
> 310	90	9 THOMAS	01	CNTR β^+ decay
> 137	95	10 ACKERSTAFF	99D	OPAL τ decay
>1400	68	11 BARENBOIM	98	RVUE Electroweak, Z - Z' mixing
> 549	68	12 BARENBOIM	97	RVUE μ decay
> 220	95	13 STAHL	97	RVUE τ decay
> 220	90	14 ALLET	96	CNTR β^+ decay
> 281	90	15 KUZNETSOV	95	CNTR Polarized neutron decay
> 282	90	16 KUZNETSOV	94B	CNTR Polarized neutron decay
> 439	90	17 BHATTACH...	93	RVUE Z - Z' mixing
> 250	90	18 SEVERIJNS	93	CNTR β^+ decay
		19 IMAZATO	92	CNTR K^+ decay
> 475	90	20 POLAK	92B	RVUE μ decay
> 240	90	21 AQUINO	91	RVUE Neutron decay
> 496	90	21 AQUINO	91	RVUE Neutron and muon decay
> 700		22 COLANGELO	91	THEO $m_{K_L^0} - m_{K_S^0}$
> 477	90	23 POLAK	91	RVUE μ decay
[none 540–23000]		24 BARBIERI	89B	ASTR SN 1987A; light ν_R

> 300	90	25	LANGACKER	89B	RVUE	General
> 160	90	26	BALKE	88	CNTR	$\mu \rightarrow e\nu\bar{\nu}$
> 406	90	27	JODIDIO	86	ELEC	Any ζ
> 482	90	27	JODIDIO	86	ELEC	$\zeta = 0$
> 800			MOHAPATRA	86	RVUE	$SU(2)_L \times SU(2)_R \times U(1)$
> 400	95	28	STOKER	85	ELEC	Any ζ
> 475	95	28	STOKER	85	ELEC	$\zeta < 0.041$
		29	BERGSMA	83	CHRM	$\nu_\mu e \rightarrow \mu\nu_e$
> 380	90	30	CARR	83	ELEC	μ^+ decay
>1600		31	BEALL	82	THEO	$m_{K_L^0} - m_{K_S^0}$

¹ The quoted limit is for manifest left-right symmetric model.

² CZAKON 99 perform a simultaneous fit to charged and neutral sectors.

³ PRIEELS 14 limit is from $\mu^+ \rightarrow e^+ \nu \bar{\nu}$ decay parameter ξ'' , which is determined by the positron polarization measurement.

⁴ WAUTERS 10 limit is from a measurement of the asymmetry parameter of polarized ^{60}Co β decays. The listed limit assumes no mixing.

⁵ ZHANG 08 limit uses a lattice QCD calculation of the relevant hadronic matrix elements, while BEALL 82 limit used the vacuum saturation approximation.

⁶ MELCONIAN 07 measure the neutrino angular asymmetry in β^+ -decays of polarized ^{37}K , stored in a magneto-optical trap. Result is consistent with SM prediction and does not constrain the $W_L - W_R$ mixing angle appreciably.

⁷ SCHUMANN 07 limit is from measurements of the asymmetry $\langle \vec{p}_\nu \cdot \sigma_n \rangle$ in the β decay of polarized neutrons. Zero mixing is assumed.

⁸ CYBURT 05 limit follows by requiring that three light ν_R 's decouple when $T_{dec} > 140$ MeV. For different T_{dec} , the bound becomes $m_{W_R} > 3.3 \text{ TeV} (T_{dec} / 140 \text{ MeV})^{3/4}$.

⁹ THOMAS 01 limit is from measurement of β^+ polarization in decay of polarized ^{12}N . The listed limit assumes no mixing.

¹⁰ ACKERSTAFF 99D limit is from τ decay parameters. Limit increase to 145 GeV for zero mixing.

¹¹ BARENBOIM 98 assumes minimal left-right model with Higgs of $SU(2)_R$ in $SU(2)_L$ doublet. For Higgs in $SU(2)_L$ triplet, $m_{W_R} > 1100$ GeV. Bound calculated from effect of corresponding Z_{LR} on electroweak data through $Z-Z_{LR}$ mixing.

¹² The quoted limit is from μ decay parameters. BARENBOIM 97 also evaluate limit from $K_L - K_S$ mass difference.

¹³ STAHL 97 limit is from fit to τ -decay parameters.

¹⁴ ALLET 96 measured polarization-asymmetry correlation in $^{12}\text{N}\beta^+$ decay. The listed limit assumes zero L - R mixing.

¹⁵ KUZNETSOV 95 limit is from measurements of the asymmetry $\langle \vec{p}_\nu \cdot \sigma_n \rangle$ in the β decay of polarized neutrons. Zero mixing assumed. See also KUZNETSOV 94B.

¹⁶ KUZNETSOV 94B limit is from measurements of the asymmetry $\langle \vec{p}_\nu \cdot \sigma_n \rangle$ in the β decay of polarized neutrons. Zero mixing assumed.

¹⁷ BHATTACHARYYA 93 uses Z - Z' mixing limit from LEP '90 data, assuming a specific Higgs sector of $SU(2)_L \times SU(2)_R \times U(1)$ gauge model. The limit is for $m_t = 200$ GeV and slightly improves for smaller m_t .

¹⁸ SEVERIJNS 93 measured polarization-asymmetry correlation in $^{107}\text{In}\beta^+$ decay. The listed limit assumes zero L - R mixing. Value quoted here is from SEVERIJNS 94 erratum.

¹⁹ IMAZATO 92 measure positron asymmetry in $K^+ \rightarrow \mu^+ \nu_\mu$ decay and obtain $\xi_{P_\mu} > 0.990$ (90% CL). If W_R couples to $u\bar{s}$ with full weak strength ($V_{us}^R = 1$), the result corresponds to $m_{W_R} > 653$ GeV. See their Fig. 4 for m_{W_R} limits for general $|V_{us}^R|^2 = 1 - |V_{ud}^R|^2$.

- ²⁰ POLAK 92B limit is from fit to muon decay parameters and is essentially determined by JODIDIO 86 data assuming $\zeta=0$. Supersedes POLAK 91.
- ²¹ AQUINO 91 limits obtained from neutron lifetime and asymmetries together with unitarity of the CKM matrix. Manifest left-right symmetry assumed. Stronger of the two limits also includes muon decay results.
- ²² COLANGELO 91 limit uses hadronic matrix elements evaluated by QCD sum rule and is less restrictive than BEALL 82 limit which uses vacuum saturation approximation. Manifest left-right symmetry assumed.
- ²³ POLAK 91 limit is from fit to muon decay parameters and is essentially determined by JODIDIO 86 data assuming $\zeta=0$. Superseded by POLAK 92B.
- ²⁴ BARBIERI 89B limit holds for $m_{\nu_R} \leq 10$ MeV.
- ²⁵ LANGACKER 89B limit is for any ν_R mass (either Dirac or Majorana) and for a general class of right-handed quark mixing matrices.
- ²⁶ BALKE 88 limit is for $m_{\nu_{eR}} = 0$ and $m_{\nu_{\mu R}} \leq 50$ MeV. Limits come from precise measurements of the muon decay asymmetry as a function of the positron energy.
- ²⁷ JODIDIO 86 is the same TRIUMF experiment as STOKER 85 (and CARR 83); however, it uses a different technique. The results given here are combined results of the two techniques. The technique here involves precise measurement of the end-point e^+ spectrum in the decay of the highly polarized μ^+ .
- ²⁸ STOKER 85 is same TRIUMF experiment as CARR 83. Here they measure the decay e^+ spectrum asymmetry above 46 MeV/c using a muon-spin-rotation technique. Assumed a light right-handed neutrino. Quoted limits are from combining with CARR 83.
- ²⁹ BERGSMA 83 set limit $m_{W_2}/m_{W_1} > 1.9$ at CL = 90%.
- ³⁰ CARR 83 is TRIUMF experiment with a highly polarized μ^+ beam. Looked for deviation from $V-A$ at the high momentum end of the decay e^+ energy spectrum. Limit from previous world-average muon polarization parameter is $m_{W_R} > 240$ GeV. Assumes a light right-handed neutrino.
- ³¹ BEALL 82 limit is obtained assuming that W_R contribution to $K_L^0-K_S^0$ mass difference is smaller than the standard one, neglecting the top quark contributions. Manifest left-right symmetry assumed.

Limit on W_L - W_R Mixing Angle ζ

Lighter mass eigenstate $W_1 = W_L \cos \zeta - W_R \sin \zeta$. Light ν_R assumed unless noted.

Values in brackets are from cosmological and astrophysical considerations.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
-0.020 to 0.017	90	BUENO 11	TWST	$\mu \rightarrow e \nu \bar{\nu}$
< 0.022	90	MACDONALD 08	TWST	$\mu \rightarrow e \nu \bar{\nu}$
< 0.12	95	1 ACKERSTAFF 99D	OPAL	τ decay
< 0.013	90	2 CZAKON 99	RVUE	Electroweak
< 0.0333		3 BARENBOIM 97	RVUE	μ decay
< 0.04	90	4 MISHRA 92	CCFR	νN scattering
-0.0006 to 0.0028	90	5 AQUINO 91	RVUE	
[none 0.00001-0.02]		6 BARBIERI 89B	ASTR	SN 1987A
< 0.040	90	7 JODIDIO 86	ELEC	μ decay
-0.056 to 0.040	90	7 JODIDIO 86	ELEC	μ decay

- ¹ ACKERSTAFF 99D limit is from τ decay parameters.
- ² CZAKON 99 perform a simultaneous fit to charged and neutral sectors.
- ³ The quoted limit is from μ decay parameters. BARENBOIM 97 also evaluate limit from K_L-K_S mass difference.
- ⁴ MISHRA 92 limit is from the absence of extra large- x , large- y $\bar{\nu}_\mu N \rightarrow \bar{\nu}_\mu X$ events at Tevatron, assuming left-handed ν and right-handed $\bar{\nu}$ in the neutrino beam. The result gives $\zeta^2(1-2m_{W_1}^2/m_{W_2}^2) < 0.0015$. The limit is independent of ν_R mass.
- ⁵ AQUINO 91 limits obtained from neutron lifetime and asymmetries together with unitarity of the CKM matrix. Manifest left-right asymmetry is assumed.
- ⁶ BARBIERI 89B limit holds for $m_{\nu_R} \leq 10$ MeV.
- ⁷ First JODIDIO 86 result assumes $m_{W_R} = \infty$, second is for unconstrained m_{W_R} .

See the related review(s):

[Z'-Boson Searches](#)

MASS LIMITS for Z' (Heavy Neutral Vector Boson Other Than Z)

Limits for Z'_{SM}

Z'_{SM} is assumed to have couplings with quarks and leptons which are identical to those of Z, and decays only to known fermions. The most recent preliminary results can be found in the "Z'-boson searches" review above.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>2420	95	1 AABOUD	18G ATLS	$p\bar{p}; Z'_{SM} \rightarrow \tau^+ \tau^-$
>4500	95	2 AABOUD	17AT ATLS	$p\bar{p}; Z'_{SM} \rightarrow e^+ e^-, \mu^+ \mu^-$
>2100	95	3 KHACHATRY...17H	CMS	$p\bar{p}; Z'_{SM} \rightarrow \tau^+ \tau^-$
>3370	95	4 KHACHATRY...17T	CMS	$p\bar{p}; Z'_{SM} \rightarrow e^+ e^-, \mu^+ \mu^-$
none 600–2100, 2300–2600	95	5 KHACHATRY...17W	CMS	$p\bar{p}; Z'_{SM} \rightarrow q\bar{q}$
>3360	95	6 AABOUD	16U ATLS	$p\bar{p}; Z'_{SM} \rightarrow e^+ e^-, \mu^+ \mu^-$
>2900	95	7 KHACHATRY...15AE	CMS	$p\bar{p}; Z'_{SM} \rightarrow e^+ e^-, \mu^+ \mu^-$
none 1200–1700	95	8 KHACHATRY...15V	CMS	$p\bar{p}; Z'_{SM} \rightarrow q\bar{q}$
>2900	95	9 AAD	14V ATLS	$p\bar{p}; Z'_{SM} \rightarrow e^+ e^-, \mu^+ \mu^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
>1900	95	10 AABOUD	16AA ATLS	$p\bar{p}; Z'_{SM} \rightarrow \tau^+ \tau^-$
>2020	95	11 AAD	15AMATLS	$p\bar{p}; Z'_{SM} \rightarrow \tau^+ \tau^-$
>1400	95	12 AAD	13S ATLS	$p\bar{p}; Z'_{SM} \rightarrow \tau^+ \tau^-$
>1470	95	13 CHATRCHYAN 13A	CMS	$p\bar{p}; Z'_{SM} \rightarrow q\bar{q}$
>2590	95	14 CHATRCHYAN 13AF	CMS	$p\bar{p}; Z'_{SM} \rightarrow e^+ e^-, \mu^+ \mu^-$
>2220	95	15 AAD	12CC ATLS	$p\bar{p}; Z'_{SM} \rightarrow e^+ e^-, \mu^+ \mu^-$
>1400	95	16 CHATRCHYAN 12O	CMS	$p\bar{p}; Z'_{SM} \rightarrow \tau^+ \tau^-$
>1071	95	17 AALTONEN	11I CDF	$p\bar{p}; Z'_{SM} \rightarrow \mu^+ \mu^-$
>1023	95	18 ABAZOV	11A D0	$p\bar{p}; Z'_{SM} \rightarrow e^+ e^-$
none 247–544	95	19 AALTONEN	10N CDF	$Z' \rightarrow W W$
none 320–740	95	20 AALTONEN	09AC CDF	$Z' \rightarrow q\bar{q}$
> 963	95	18 AALTONEN	09T CDF	$p\bar{p}; Z'_{SM} \rightarrow e^+ e^-$

>1403	95	21 ERLER	09 RVUE	Electroweak
>1305	95	22 ABDALLAH	06C DLPH	e^+e^-
> 399	95	23 ACOSTA	05R CDF	$p\bar{p}: Z'_{SM} \rightarrow \tau^+\tau^-$
none 400–640	95	ABAZOV	04C D0	$p\bar{p}: Z'_{SM} \rightarrow q\bar{q}$
>1018	95	24 ABBIENDI	04G OPAL	e^+e^-
> 670	95	25 ABAZOV	01B D0	$p\bar{p}, Z'_{SM} \rightarrow e^+e^-$
>1500	95	26 CHEUNG	01B RVUE	Electroweak
> 710	95	27 ABREU	00S DLPH	e^+e^-
> 898	95	28 BARATE	00I ALEP	e^+e^-
> 809	95	29 ERLER	99 RVUE	Electroweak
> 690	95	30 ABE	97S CDF	$p\bar{p}; Z'_{SM} \rightarrow e^+e^-, \mu^+\mu^-$
> 398	95	31 VILAIN	94B CHM2	$\nu_\mu e \rightarrow \nu_\mu e$ and $\bar{\nu}_\mu e \rightarrow \bar{\nu}_\mu e$
> 237	90	32 ALITTI	93 UA2	$p\bar{p}; Z'_{SM} \rightarrow q\bar{q}$
none 260–600	95	33 RIZZO	93 RVUE	$p\bar{p}; Z'_{SM} \rightarrow q\bar{q}$
> 426	90	34 ABE	90F VNS	e^+e^-

- ¹ AABOUD 18G search for resonances decaying to $\tau^+\tau^-$ in pp collisions at $\sqrt{s} = 13$ TeV.
- ² AABOUD 17AT search for resonances decaying to $\ell^+\ell^-$ in pp collisions at $\sqrt{s} = 13$ TeV.
- ³ KHACHATRYAN 17H search for resonances decaying to $\tau^+\tau^-$ in pp collisions at $\sqrt{s} = 13$ TeV.
- ⁴ KHACHATRYAN 17T search for resonances decaying to $e^+e^-, \mu^+\mu^-$ in pp collisions at $\sqrt{s} = 8, 13$ TeV.
- ⁵ KHACHATRYAN 17W search for resonances decaying to dijets in pp collisions at $\sqrt{s} = 13$ TeV.
- ⁶ AABOUD 16U search for resonances decaying to $\ell^+\ell^-$ in pp collisions at $\sqrt{s} = 13$ TeV.
- ⁷ KHACHATRYAN 15AE search for resonances decaying to $e^+e^-, \mu^+\mu^-$ in pp collisions at $\sqrt{s} = 8$ TeV.
- ⁸ KHACHATRYAN 15V search for resonances decaying to dijets in pp collisions at $\sqrt{s} = 8$ TeV.
- ⁹ AAD 14V search for resonances decaying to $e^+e^-, \mu^+\mu^-$ in pp collisions at $\sqrt{s} = 8$ TeV.
- ¹⁰ AABOUD 16AA search for resonances decaying to $\tau^+\tau^-$ in pp collisions at $\sqrt{s} = 13$ TeV.
- ¹¹ AAD 15AM search for resonances decaying to $\tau^+\tau^-$ in pp collisions at $\sqrt{s} = 8$ TeV.
- ¹² AAD 13S search for resonances decaying to $\tau^+\tau^-$ in pp collisions at $\sqrt{s} = 7$ TeV.
- ¹³ CHATRCHYAN 13A use pp collisions at $\sqrt{s}=7$ TeV.
- ¹⁴ CHATRCHYAN 13AF search for resonances decaying to $e^+e^-, \mu^+\mu^-$ in pp collisions at $\sqrt{s} = 7$ TeV and 8 TeV.
- ¹⁵ AAD 12CC search for resonances decaying to $e^+e^-, \mu^+\mu^-$ in pp collisions at $\sqrt{s} = 7$ TeV.
- ¹⁶ CHATRCHYAN 120 search for resonances decaying to $\tau^+\tau^-$ in pp collisions at $\sqrt{s} = 7$ TeV.
- ¹⁷ AALTONEN 11I search for resonances decaying to $\mu^+\mu^-$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV.
- ¹⁸ ABAZOV 11A, AALTONEN 09T, AALTONEN 07H, and ABULENCIA 06L search for resonances decaying to e^+e^- in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV.
- ¹⁹ The quoted limit assumes $g_{WWZ'}/g_{WWZ} = (M_W/M_{Z'})^2$. See their Fig. 4 for limits in mass-coupling plane.
- ²⁰ AALTONEN 09AC search for new particle decaying to dijets.
- ²¹ ERLER 09 give 95% CL limit on the Z - Z' mixing $-0.0026 < \theta < 0.0006$.
- ²² ABDALLAH 06C use data $\sqrt{s} = 130$ – 207 GeV.

- ²³ ACOSTA 05R search for resonances decaying to tau lepton pairs in $\bar{p}p$ collisions at $\sqrt{s} = 1.96$ TeV.
- ²⁴ ABBIENDI 04G give 95% CL limit on Z - Z' mixing $-0.00422 < \theta < 0.00091$. $\sqrt{s} = 91$ to 207 GeV.
- ²⁵ ABAZOV 01B search for resonances in $p\bar{p} \rightarrow e^+e^-$ at $\sqrt{s}=1.8$ TeV. They find $\sigma \cdot B(Z' \rightarrow ee) < 0.06$ pb for $M_{Z'} > 500$ GeV.
- ²⁶ CHEUNG 01B limit is derived from bounds on contact interactions in a global electroweak analysis.
- ²⁷ ABREU 00S uses LEP data at $\sqrt{s}=90$ to 189 GeV.
- ²⁸ BARATE 00I search for deviations in cross section and asymmetries in $e^+e^- \rightarrow$ fermions at $\sqrt{s}=90$ to 183 GeV. Assume $\theta=0$. Bounds in the mass-mixing plane are shown in their Figure 18.
- ²⁹ ERLER 99 give 90%CL limit on the Z - Z' mixing $-0.0041 < \theta < 0.0003$. $\rho_0=1$ is assumed.
- ³⁰ ABE 97S find $\sigma(Z') \times B(e^+e^-, \mu^+\mu^-) < 40$ fb for $m_{Z'} > 600$ GeV at $\sqrt{s}=1.8$ TeV.
- ³¹ VILAIN 94B assume $m_t = 150$ GeV.
- ³² ALITTI 93 search for resonances in the two-jet invariant mass. The limit assumes $B(Z' \rightarrow q\bar{q})=0.7$. See their Fig. 5 for limits in the $m_{Z'}-B(q\bar{q})$ plane.
- ³³ RIZZO 93 analyses CDF limit on possible two-jet resonances.
- ³⁴ ABE 90F use data for $R, R_{\ell\ell},$ and $A_{\ell\ell}$. They fix $m_W = 80.49 \pm 0.43 \pm 0.24$ GeV and $m_Z = 91.13 \pm 0.03$ GeV.

Limits for Z_{LR}

Z_{LR} is the extra neutral boson in left-right symmetric models. $g_L = g_R$ is assumed unless noted. Values in parentheses assume stronger constraint on the Higgs sector, usually motivated by specific left-right symmetric models (see the Note on the W'). Values in brackets are from cosmological and astrophysical considerations and assume a light right-handed neutrino. Direct search bounds assume decays to Standard Model fermions only, unless noted.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>1162	95	¹ DEL-AGUILA	10 RVUE	Electroweak
> 630	95	² ABE	97S CDF	$p\bar{p}; Z'_{LR} \rightarrow e^+e^-, \mu^+\mu^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
> 998	95	³ ERLER	09 RVUE	Electroweak
> 600	95	SCHAEL	07A ALEP	e^+e^-
> 455	95	⁴ ABDALLAH	06C DLPH	e^+e^-
> 518	95	⁵ ABBIENDI	04G OPAL	e^+e^-
> 860	95	⁶ CHEUNG	01B RVUE	Electroweak
> 380	95	⁷ ABREU	00S DLPH	e^+e^-
> 436	95	⁸ BARATE	00I ALEP	Repl. by SCHAEL 07A
> 550	95	⁹ CHAY	00 RVUE	Electroweak
		¹⁰ ERLER	00 RVUE	Cs
		¹¹ CASALBUONI	99 RVUE	Cs
(> 1205)	90	¹² CZAKON	99 RVUE	Electroweak
> 564	95	¹³ ERLER	99 RVUE	Electroweak
(> 1673)	95	¹⁴ ERLER	99 RVUE	Electroweak
(> 1700)	68	¹⁵ BARENBOIM	98 RVUE	Electroweak
> 244	95	¹⁶ CONRAD	98 RVUE	$\nu_\mu N$ scattering

> 253	95	17 VILAIN	94B CHM2	$\nu_\mu e \rightarrow \nu_\mu e$ and $\bar{\nu}_\mu e \rightarrow \bar{\nu}_\mu e$
none 200–600	95	18 RIZZO	93 RVUE	$p\bar{p}; Z_{LR} \rightarrow q\bar{q}$
[> 2000]		WALKER	91 COSM	Nucleosynthesis; light ν_R
none 200–500		19 GRIFOLS	90 ASTR	SN 1987A; light ν_R
none 350–2400		20 BARBIERI	89B ASTR	SN 1987A; light ν_R

- ¹ DEL-AGUILA 10 give 95% CL limit on the Z - Z' mixing $-0.0012 < \theta < 0.0004$.
- ² ABE 97S find $\sigma(Z') \times B(e^+e^-, \mu^+\mu^-) < 40$ fb for $m_{Z'} > 600$ GeV at $\sqrt{s} = 1.8$ TeV.
- ³ ERLER 09 give 95% CL limit on the Z - Z' mixing $-0.0013 < \theta < 0.0006$.
- ⁴ ABDALLAH 06C give 95% CL limit $|\theta| < 0.0028$. See their Fig. 14 for limit contours in the mass-mixing plane.
- ⁵ ABBIENDI 04G give 95% CL limit on Z - Z' mixing $-0.00098 < \theta < 0.00190$. See their Fig. 20 for the limit contour in the mass-mixing plane. $\sqrt{s} = 91$ to 207 GeV.
- ⁶ CHEUNG 01B limit is derived from bounds on contact interactions in a global electroweak analysis.
- ⁷ ABREU 00S give 95% CL limit on Z - Z' mixing $|\theta| < 0.0018$. See their Fig. 6 for the limit contour in the mass-mixing plane. $\sqrt{s} = 90$ to 189 GeV.
- ⁸ BARATE 00I search for deviations in cross section and asymmetries in $e^+e^- \rightarrow$ fermions at $\sqrt{s} = 90$ to 183 GeV. Assume $\theta = 0$. Bounds in the mass-mixing plane are shown in their Figure 18.
- ⁹ CHAY 00 also find $-0.0003 < \theta < 0.0019$. For g_R free, $m_{Z'} > 430$ GeV.
- ¹⁰ ERLER 00 discuss the possibility that a discrepancy between the observed and predicted values of $Q_W(\text{Cs})$ is due to the exchange of Z' . The data are better described in a certain class of the Z' models including Z_{LR} and Z_χ .
- ¹¹ CASALBUONI 99 discuss the discrepancy between the observed and predicted values of $Q_W(\text{Cs})$. It is shown that the data are better described in a class of models including the Z_{LR} model.
- ¹² CZAKON 99 perform a simultaneous fit to charged and neutral sectors. Assumes manifest left-right symmetric model. Finds $|\theta| < 0.0042$.
- ¹³ ERLER 99 give 90% CL limit on the Z - Z' mixing $-0.0009 < \theta < 0.0017$.
- ¹⁴ ERLER 99 assumes 2 Higgs doublets, transforming as 10 of $SO(10)$, embedded in E_6 .
- ¹⁵ BARENBOIM 98 also gives 68% CL limits on the Z - Z' mixing $-0.0005 < \theta < 0.0033$. Assumes Higgs sector of minimal left-right model.
- ¹⁶ CONRAD 98 limit is from measurements at CCFR, assuming no Z - Z' mixing.
- ¹⁷ VILAIN 94B assume $m_t = 150$ GeV and $\theta = 0$. See Fig. 2 for limit contours in the mass-mixing plane.
- ¹⁸ RIZZO 93 analyses CDF limit on possible two-jet resonances.
- ¹⁹ GRIFOLS 90 limit holds for $m_{\nu_R} \lesssim 1$ MeV. A specific Higgs sector is assumed. See also GRIFOLS 90D, RIZZO 91.
- ²⁰ BARBIERI 89B limit holds for $m_{\nu_R} \leq 10$ MeV. Bounds depend on assumed supernova core temperature.

Limits for Z_χ

Z_χ is the extra neutral boson in $SO(10) \rightarrow SU(5) \times U(1)_\chi$. $g_\chi = e/\cos\theta_W$ is assumed unless otherwise stated. We list limits with the assumption $\rho = 1$ but with no further constraints on the Higgs sector. Values in parentheses assume stronger constraint on the Higgs sector motivated by superstring models. Values in brackets are from cosmological and astrophysical considerations and assume a light right-handed neutrino.

<u>VALUE (GeV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
>4100	95	¹ AABOUD	17AT ATLS	$p\bar{p}; Z'_\chi \rightarrow e^+e^-, \mu^+\mu^-$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

>3050	95	² AABOUD	16U ATLS	$pp; Z'_{\chi} \rightarrow e^+ e^-, \mu^+ \mu^-$
>2620	95	³ AAD	14V ATLS	$pp, Z'_{\chi} \rightarrow e^+ e^-, \mu^+ \mu^-$
>1970	95	⁴ AAD	12CC ATLS	$pp, Z'_{\chi} \rightarrow e^+ e^-, \mu^+ \mu^-$
> 930	95	⁵ AALTONEN	11I CDF	$p\bar{p}; Z'_{\chi} \rightarrow \mu^+ \mu^-$
> 903	95	⁶ ABAZOV	11A D0	$p\bar{p}, Z'_{\chi} \rightarrow e^+ e^-$
>1022	95	⁷ DEL-AGUILA	10 RVUE	Electroweak
> 862	95	⁶ AALTONEN	09T CDF	$p\bar{p}, Z'_{\chi} \rightarrow e^+ e^-$
> 892	95	⁸ AALTONEN	09V CDF	Repl. by AALTONEN 11I
>1141	95	⁹ ERLER	09 RVUE	Electroweak
> 822	95	⁶ AALTONEN	07H CDF	Repl. by AALTONEN 09T
> 680	95	SCHAEL	07A ALEP	$e^+ e^-$
> 545	95	¹⁰ ABDALLAH	06C DLPH	$e^+ e^-$
> 740		⁶ ABULENCIA	06L CDF	Repl. by AALTONEN 07H
> 690	95	¹¹ ABULENCIA	05A CDF	$p\bar{p}; Z'_{\chi} \rightarrow e^+ e^-, \mu^+ \mu^-$
> 781	95	¹² ABBIENDI	04G OPAL	$e^+ e^-$
>2100		¹³ BARGER	03B COSM	Nucleosynthesis; light ν_R
> 680	95	¹⁴ CHEUNG	01B RVUE	Electroweak
> 440	95	¹⁵ ABREU	00S DLPH	$e^+ e^-$
> 533	95	¹⁶ BARATE	00I ALEP	Repl. by SCHAEL 07A
> 554	95	¹⁷ CHO	00 RVUE	Electroweak
		¹⁸ ERLER	00 RVUE	Cs
		¹⁹ ROSNER	00 RVUE	Cs
> 545	95	²⁰ ERLER	99 RVUE	Electroweak
(> 1368)	95	²¹ ERLER	99 RVUE	Electroweak
> 215	95	²² CONRAD	98 RVUE	$\nu_{\mu} N$ scattering
> 595	95	²³ ABE	97S CDF	$p\bar{p}; Z'_{\chi} \rightarrow e^+ e^-, \mu^+ \mu^-$
> 190	95	²⁴ ARIMA	97 VNS	Bhabha scattering
> 262	95	²⁵ VILAIN	94B CHM2	$\nu_{\mu} e \rightarrow \nu_{\mu} e; \bar{\nu}_{\mu} e \rightarrow \bar{\nu}_{\mu} e$
[>1470]		²⁶ FARAGGI	91 COSM	Nucleosynthesis; light ν_R
> 231	90	²⁷ ABE	90F VNS	$e^+ e^-$
[> 1140]		²⁸ GONZALEZ...	90D COSM	Nucleosynthesis; light ν_R
[> 2100]		²⁹ GRIFOLS	90 ASTR	SN 1987A; light ν_R

¹ AABOUD 17AT search for resonances decaying to $\ell^+ \ell^-$ in pp collisions at $\sqrt{s} = 13$ TeV.

² AABOUD 16U search for resonances decaying to $\ell^+ \ell^-$ in pp collisions at $\sqrt{s} = 13$ TeV.

³ AAD 14V search for resonances decaying to $e^+ e^-, \mu^+ \mu^-$ in pp collisions at $\sqrt{s} = 8$ TeV.

⁴ AAD 12CC search for resonances decaying to $e^+ e^-, \mu^+ \mu^-$ in pp collisions at $\sqrt{s} = 7$ TeV.

⁵ AALTONEN 11I search for resonances decaying to $\mu^+ \mu^-$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV.

⁶ ABAZOV 11A, AALTONEN 09T, AALTONEN 07H, and ABULENCIA 06L search for resonances decaying to $e^+ e^-$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV.

⁷ DEL-AGUILA 10 give 95% CL limit on the Z - Z' mixing $-0.0011 < \theta < 0.0007$.

⁸ AALTONEN 09V search for resonances decaying to $\mu^+ \mu^-$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV.

⁹ ERLER 09 give 95% CL limit on the Z - Z' mixing $-0.0016 < \theta < 0.0006$.

- ¹⁰ ABDALLAH 06C give 95% CL limit $|\theta| < 0.0031$. See their Fig. 14 for limit contours in the mass-mixing plane.
- ¹¹ ABULENCIA 05A search for resonances decaying to electron or muon pairs in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV.
- ¹² ABBIENDI 04G give 95% CL limit on Z - Z' mixing $-0.00099 < \theta < 0.00194$. See their Fig. 20 for the limit contour in the mass-mixing plane. $\sqrt{s} = 91$ to 207 GeV.
- ¹³ BARGER 03B limit is from the nucleosynthesis bound on the effective number of light neutrino $\delta N_\nu < 1$. The quark-hadron transition temperature $T_c=150$ MeV is assumed. The limit with $T_c=400$ MeV is >4300 GeV.
- ¹⁴ CHEUNG 01B limit is derived from bounds on contact interactions in a global electroweak analysis.
- ¹⁵ ABREU 00S give 95% CL limit on Z - Z' mixing $|\theta| < 0.0017$. See their Fig. 6 for the limit contour in the mass-mixing plane. $\sqrt{s}=90$ to 189 GeV.
- ¹⁶ BARATE 00I search for deviations in cross section and asymmetries in $e^+e^- \rightarrow$ fermions at $\sqrt{s}=90$ to 183 GeV. Assume $\theta=0$. Bounds in the mass-mixing plane are shown in their Figure 18.
- ¹⁷ CHO 00 use various electroweak data to constrain Z' models assuming $m_H=100$ GeV. See Fig. 3 for limits in the mass-mixing plane.
- ¹⁸ ERLER 00 discuss the possibility that a discrepancy between the observed and predicted values of $Q_W(\text{Cs})$ is due to the exchange of Z' . The data are better described in a certain class of the Z' models including Z_{LR} and Z_χ .
- ¹⁹ ROSNER 00 discusses the possibility that a discrepancy between the observed and predicted values of $Q_W(\text{Cs})$ is due to the exchange of Z' . The data are better described in a certain class of the Z' models including Z_χ .
- ²⁰ ERLER 99 give 90% CL limit on the Z - Z' mixing $-0.0020 < \theta < 0.0015$.
- ²¹ ERLER 99 assumes 2 Higgs doublets, transforming as 10 of $SO(10)$, embedded in E_6 .
- ²² CONRAD 98 limit is from measurements at CCFR, assuming no Z - Z' mixing.
- ²³ ABE 97S find $\sigma(Z') \times B(e^+e^-, \mu^+\mu^-) < 40$ fb for $m_{Z'} > 600$ GeV at $\sqrt{s}=1.8$ TeV.
- ²⁴ Z - Z' mixing is assumed to be zero. $\sqrt{s}=57.77$ GeV.
- ²⁵ VILAIN 94B assume $m_t = 150$ GeV and $\theta=0$. See Fig. 2 for limit contours in the mass-mixing plane.
- ²⁶ FARAGGI 91 limit assumes the nucleosynthesis bound on the effective number of neutrinos $\Delta N_\nu < 0.5$ and is valid for $m_{\nu_R} < 1$ MeV.
- ²⁷ ABE 90F use data for R , $R_{\ell\ell}$, and $A_{\ell\ell}$. ABE 90F fix $m_W = 80.49 \pm 0.43 \pm 0.24$ GeV and $m_Z = 91.13 \pm 0.03$ GeV.
- ²⁸ Assumes the nucleosynthesis bound on the effective number of light neutrinos ($\delta N_\nu < 1$) and that ν_R is light ($\lesssim 1$ MeV).
- ²⁹ GRIFOLS 90 limit holds for $m_{\nu_R} \lesssim 1$ MeV. See also GRIFOLS 90D, RIZZO 91.

Limits for Z_ψ

Z_ψ is the extra neutral boson in $E_6 \rightarrow SO(10) \times U(1)_\psi$. $g_\psi = e/\cos\theta_W$ is assumed unless otherwise stated. We list limits with the assumption $\rho = 1$ but with no further constraints on the Higgs sector. Values in brackets are from cosmological and astrophysical considerations and assume a light right-handed neutrino.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>3800	95	¹ AABOUD 17AT ATLS		$p\bar{p}; Z'_\psi \rightarrow e^+e^-, \mu^+\mu^-$
>2820	95	² KHACHATRY...17T CMS		$p\bar{p}; Z'_\psi \rightarrow e^+e^-, \mu^+\mu^-$
>1100	95	³ CHATRCHYAN 120 CMS		$p\bar{p}; Z'_\psi \rightarrow \tau^+\tau^-$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

>2740	95	⁴ AABOUD	16U ATLS	$pp; Z'_\psi \rightarrow e^+ e^-, \mu^+ \mu^-$
>2570	95	⁵ KHACHATRYAN...15AE	CMS	$pp; Z'_\psi \rightarrow e^+ e^-, \mu^+ \mu^-$
>2510	95	⁶ AAD	14V ATLS	$pp; Z'_\psi \rightarrow e^+ e^-, \mu^+ \mu^-$
>2260	95	⁷ CHATRCHYAN	13AF CMS	$pp; Z'_\psi \rightarrow e^+ e^-, \mu^+ \mu^-$
>1790	95	⁸ AAD	12CC ATLS	$pp; Z'_\psi \rightarrow e^+ e^-, \mu^+ \mu^-$
>2000	95	⁹ CHATRCHYAN	12M CMS	Repl. by CHATRCHYAN 13AF
> 917	95	¹⁰ AALTONEN	11I CDF	$p\bar{p}; Z'_\psi \rightarrow \mu^+ \mu^-$
> 891	95	¹¹ ABAZOV	11A D0	$p\bar{p}; Z'_\psi \rightarrow e^+ e^-$
> 476	95	¹² DEL-AGUILA	10 RVUE	Electroweak
> 851	95	¹¹ AALTONEN	09T CDF	$p\bar{p}; Z'_\psi \rightarrow e^+ e^-$
> 878	95	¹³ AALTONEN	09V CDF	Repl. by AALTONEN 11I
> 147	95	¹⁴ ERLER	09 RVUE	Electroweak
> 822	95	¹¹ AALTONEN	07H CDF	Repl. by AALTONEN 09T
> 410	95	SCHAEL	07A ALEP	$e^+ e^-$
> 475	95	¹⁵ ABDALLAH	06C DLPH	$e^+ e^-$
> 725		¹¹ ABULENCIA	06L CDF	Repl. by AALTONEN 07H
> 675	95	¹⁶ ABULENCIA	05A CDF	Repl. by AALTONEN 11I and AALTONEN 09T
> 366	95	¹⁷ ABBIENDI	04G OPAL	$e^+ e^-$
> 600		¹⁸ BARGER	03B COSM	Nucleosynthesis; light ν_R
> 350	95	¹⁹ ABREU	00S DLPH	$e^+ e^-$
> 294	95	²⁰ BARATE	00I ALEP	Repl. by SCHAEL 07A
> 137	95	²¹ CHO	00 RVUE	Electroweak
> 146	95	²² ERLER	99 RVUE	Electroweak
> 54	95	²³ CONRAD	98 RVUE	$\nu_\mu N$ scattering
> 590	95	²⁴ ABE	97S CDF	$p\bar{p}; Z'_\psi \rightarrow e^+ e^-, \mu^+ \mu^-$
> 135	95	²⁵ VILAIN	94B CHM2	$\nu_\mu e \rightarrow \nu_\mu e; \bar{\nu}_\mu e \rightarrow \bar{\nu}_\mu e$
> 105	90	²⁶ ABE	90F VNS	$e^+ e^-$
[> 160]		²⁷ GONZALEZ...	90D COSM	Nucleosynthesis; light ν_R
[> 2000]		²⁸ GRIFOLS	90D ASTR	SN 1987A; light ν_R

¹ AABOUD 17AT search for resonances decaying to $\ell^+ \ell^-$ in pp collisions at $\sqrt{s} = 13$ TeV.

² KHACHATRYAN 17T search for resonances decaying to $e^+ e^-, \mu^+ \mu^-$ in pp collisions at $\sqrt{s} = 8, 13$ TeV.

³ CHATRCHYAN 120 search for resonances decaying to $\tau^+ \tau^-$ in pp collisions at $\sqrt{s} = 7$ TeV.

⁴ AABOUD 16U search for resonances decaying to $\ell^+ \ell^-$ in pp collisions at $\sqrt{s} = 13$ TeV.

⁵ KHACHATRYAN 15AE search for resonances decaying to $e^+ e^-, \mu^+ \mu^-$ in pp collisions at $\sqrt{s} = 8$ TeV.

⁶ AAD 14V search for resonances decaying to $e^+ e^-, \mu^+ \mu^-$ in pp collisions at $\sqrt{s} = 8$ TeV.

⁷ CHATRCHYAN 13AF search for resonances decaying to $e^+ e^-, \mu^+ \mu^-$ in pp collisions at $\sqrt{s} = 7$ TeV and 8 TeV.

⁸ AAD 12CC search for resonances decaying to $e^+ e^-, \mu^+ \mu^-$ in pp collisions at $\sqrt{s} = 7$ TeV.

⁹ CHATRCHYAN 12M search for resonances decaying to $e^+ e^-$ or $\mu^+ \mu^-$ in pp collisions at $\sqrt{s} = 7$ TeV.

- 10 AALTONEN 11I search for resonances decaying to $\mu^+ \mu^-$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV.
- 11 ABAZOV 11A, AALTONEN 09T, AALTONEN 07H, and ABULENCIA 06L search for resonances decaying to $e^+ e^-$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV.
- 12 DEL-AGUILA 10 give 95% CL limit on the Z - Z' mixing $-0.0019 < \theta < 0.0007$.
- 13 AALTONEN 09V search for resonances decaying to $\mu^+ \mu^-$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV.
- 14 ERLER 09 give 95% CL limit on the Z - Z' mixing $-0.0018 < \theta < 0.0009$.
- 15 ABDALLAH 06C give 95% CL limit $|\theta| < 0.0027$. See their Fig. 14 for limit contours in the mass-mixing plane.
- 16 ABULENCIA 05A search for resonances decaying to electron or muon pairs in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV.
- 17 ABBIENDI 04G give 95% CL limit on Z - Z' mixing $-0.00129 < \theta < 0.00258$. See their Fig. 20 for the limit contour in the mass-mixing plane. $\sqrt{s} = 91$ to 207 GeV.
- 18 BARGER 03B limit is from the nucleosynthesis bound on the effective number of light neutrino $\delta N_\nu < 1$. The quark-hadron transition temperature $T_c=150$ MeV is assumed. The limit with $T_c=400$ MeV is >1100 GeV.
- 19 ABREU 00S give 95% CL limit on Z - Z' mixing $|\theta| < 0.0018$. See their Fig. 6 for the limit contour in the mass-mixing plane. $\sqrt{s}=90$ to 189 GeV.
- 20 BARATE 00I search for deviations in cross section and asymmetries in $e^+ e^- \rightarrow$ fermions at $\sqrt{s}=90$ to 183 GeV. Assume $\theta=0$. Bounds in the mass-mixing plane are shown in their Figure 18.
- 21 CHO 00 use various electroweak data to constrain Z' models assuming $m_H=100$ GeV. See Fig. 3 for limits in the mass-mixing plane.
- 22 ERLER 99 give 90% CL limit on the Z - Z' mixing $-0.0013 < \theta < 0.0024$.
- 23 CONRAD 98 limit is from measurements at CCFR, assuming no Z - Z' mixing.
- 24 ABE 97S find $\sigma(Z') \times B(e^+ e^-, \mu^+ \mu^-) < 40$ fb for $m_{Z'} > 600$ GeV at $\sqrt{s}=1.8$ TeV.
- 25 VILAIN 94B assume $m_t = 150$ GeV and $\theta=0$. See Fig. 2 for limit contours in the mass-mixing plane.
- 26 ABE 90F use data for R , $R_{\ell\ell}$, and $A_{\ell\ell}$. ABE 90F fix $m_W = 80.49 \pm 0.43 \pm 0.24$ GeV and $m_Z = 91.13 \pm 0.03$ GeV.
- 27 Assumes the nucleosynthesis bound on the effective number of light neutrinos ($\delta N_\nu < 1$) and that ν_R is light ($\lesssim 1$ MeV).
- 28 GRIFOLS 90D limit holds for $m_{\nu_R} \lesssim 1$ MeV. See also RIZZO 91.

Limits for Z_η

Z_η is the extra neutral boson in E_6 models, corresponding to $Q_\eta = \sqrt{3/8} Q_\chi - \sqrt{5/8} Q_\psi$. $g_\eta = e/\cos\theta_W$ is assumed unless otherwise stated. We list limits with the assumption $\rho = 1$ but with no further constraints on the Higgs sector. Values in parentheses assume stronger constraint on the Higgs sector motivated by superstring models. Values in brackets are from cosmological and astrophysical considerations and assume a light right-handed neutrino.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>3900	95	1 AABOUD	17AT ATLS	$p\bar{p}; Z'_\eta \rightarrow e^+ e^-, \mu^+ \mu^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
>2810	95	2 AABOUD	16U ATLS	$p\bar{p}; Z'_\eta \rightarrow e^+ e^-, \mu^+ \mu^-$
>1870	95	3 AAD	12CC ATLS	$p\bar{p}; Z'_\eta \rightarrow e^+ e^-, \mu^+ \mu^-$
> 938	95	4 AALTONEN	11I CDF	$p\bar{p}; Z'_\eta \rightarrow \mu^+ \mu^-$
> 923	95	5 ABAZOV	11A D0	$p\bar{p}; Z'_\eta \rightarrow e^+ e^-$

> 488	95	⁶ DEL-AGUILA	10	RVUE	Electroweak
> 877	95	⁵ AALTONEN	09T	CDF	$p\bar{p}, Z'_\eta \rightarrow e^+ e^-$
> 904	95	⁷ AALTONEN	09V	CDF	Repl. by AALTONEN 11i
> 427	95	⁸ ERLER	09	RVUE	Electroweak
> 891	95	⁵ AALTONEN	07H	CDF	Repl. by AALTONEN 09T
> 350	95	SCHAEL	07A	ALEP	$e^+ e^-$
> 360	95	⁹ ABDALLAH	06C	DLPH	$e^+ e^-$
> 745		⁵ ABULENCIA	06L	CDF	Repl. by AALTONEN 07H
> 720	95	¹⁰ ABULENCIA	05A	CDF	Repl. by AALTONEN 11i and AALTONEN 09T
> 515	95	¹¹ ABBIENDI	04G	OPAL	$e^+ e^-$
>1600		¹² BARGER	03B	COSM	Nucleosynthesis; light ν_R
> 310	95	¹³ ABREU	00S	DLPH	$e^+ e^-$
> 329	95	¹⁴ BARATE	00i	ALEP	Repl. by SCHAEL 07A
> 619	95	¹⁵ CHO	00	RVUE	Electroweak
> 365	95	¹⁶ ERLER	99	RVUE	Electroweak
> 87	95	¹⁷ CONRAD	98	RVUE	$\nu_\mu N$ scattering
> 620	95	¹⁸ ABE	97S	CDF	$p\bar{p}; Z'_\eta \rightarrow e^+ e^-, \mu^+ \mu^-$
> 100	95	¹⁹ VILAIN	94B	CHM2	$\nu_\mu e \rightarrow \nu_\mu e; \bar{\nu}_\mu e \rightarrow \bar{\nu}_\mu e$
> 125	90	²⁰ ABE	90F	VNS	$e^+ e^-$
[> 820]		²¹ GONZALEZ...	90D	COSM	Nucleosynthesis; light ν_R
[> 3300]		²² GRIFOLS	90	ASTR	SN 1987A; light ν_R
[> 1040]		²¹ LOPEZ	90	COSM	Nucleosynthesis; light ν_R

¹ AABOUD 17AT search for resonances decaying to $\ell^+ \ell^-$ in pp collisions at $\sqrt{s} = 13$ TeV.

² AABOUD 16U search for resonances decaying to $\ell^+ \ell^-$ in pp collisions at $\sqrt{s} = 13$ TeV.

³ AAD 12CC search for resonances decaying to $e^+ e^-, \mu^+ \mu^-$ in pp collisions at $\sqrt{s} = 7$ TeV.

⁴ AALTONEN 11i search for resonances decaying to $\mu^+ \mu^-$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV.

⁵ ABAZOV 11A, AALTONEN 09T, AALTONEN 07H, and ABULENCIA 06L search for resonances decaying to $e^+ e^-$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV.

⁶ DEL-AGUILA 10 give 95% CL limit on the Z - Z' mixing $-0.0023 < \theta < 0.0027$.

⁷ AALTONEN 09V search for resonances decaying to $\mu^+ \mu^-$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV.

⁸ ERLER 09 give 95% CL limit on the Z - Z' mixing $-0.0047 < \theta < 0.0021$.

⁹ ABDALLAH 06C give 95% CL limit $|\theta| < 0.0092$. See their Fig. 14 for limit contours in the mass-mixing plane.

¹⁰ ABULENCIA 05A search for resonances decaying to electron or muon pairs in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV.

¹¹ ABBIENDI 04G give 95% CL limit on Z - Z' mixing $-0.00447 < \theta < 0.00331$. See their Fig. 20 for the limit contour in the mass-mixing plane. $\sqrt{s} = 91$ to 207 GeV.

¹² BARGER 03B limit is from the nucleosynthesis bound on the effective number of light neutrino $\delta N_\nu < 1$. The quark-hadron transition temperature $T_c = 150$ MeV is assumed. The limit with $T_c = 400$ MeV is > 3300 GeV.

¹³ ABREU 00S give 95% CL limit on Z - Z' mixing $|\theta| < 0.0024$. See their Fig. 6 for the limit contour in the mass-mixing plane. $\sqrt{s} = 90$ to 189 GeV.

¹⁴ BARATE 00i search for deviations in cross section and asymmetries in $e^+ e^- \rightarrow$ fermions at $\sqrt{s} = 90$ to 183 GeV. Assume $\theta = 0$. Bounds in the mass-mixing plane are shown in their Figure 18.

¹⁵ CHO 00 use various electroweak data to constrain Z' models assuming $m_H = 100$ GeV. See Fig. 3 for limits in the mass-mixing plane.

- ¹⁶ ERLER 99 give 90% CL limit on the Z - Z' mixing $-0.0062 < \theta < 0.0011$.
¹⁷ CONRAD 98 limit is from measurements at CCFR, assuming no Z - Z' mixing.
¹⁸ ABE 97S find $\sigma(Z') \times B(e^+ e^-, \mu^+ \mu^-) < 40$ fb for $m_{Z'} > 600$ GeV at $\sqrt{s} = 1.8$ TeV.
¹⁹ VILAIN 94B assume $m_t = 150$ GeV and $\theta = 0$. See Fig. 2 for limit contours in the mass-mixing plane.
²⁰ ABE 90F use data for R , $R_{\ell\ell}$, and $A_{\ell\ell}$. ABE 90F fix $m_W = 80.49 \pm 0.43 \pm 0.24$ GeV and $m_Z = 91.13 \pm 0.03$ GeV.
²¹ These authors claim that the nucleosynthesis bound on the effective number of light neutrinos ($\delta N_\nu < 1$) constrains Z' masses if ν_R is light ($\lesssim 1$ MeV).
²² GRIFOLS 90 limit holds for $m_{\nu_R} \lesssim 1$ MeV. See also GRIFOLS 90D, RIZZO 91.

Limits for other Z'

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>1300	95	1 AABOUD	18B ATLS	$Z' \rightarrow W W$
none 1200–2800	95	2 AABOUD	18F ATLS	$Z' \rightarrow W W$
>2900	95	3 AABOUD	17AK ATLS	$Z' \rightarrow q\bar{q}$
none 1100–2600	95	4 AABOUD	17AO ATLS	$Z' \rightarrow h Z$
>2300	95	5 SIRUNYAN	17AK CMS	$Z' \rightarrow W W, h Z$
>2500	95	6 SIRUNYAN	17Q CMS	$Z' \rightarrow t\bar{t}$
>1190	95	7 SIRUNYAN	17R CMS	$Z' \rightarrow h Z$
none 1210–2260	95	7 SIRUNYAN	17R CMS	$Z' \rightarrow h Z$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
		8 SIRUNYAN	18G CMS	$Z' \rightarrow q\bar{q}$
>1580	95	9 AABOUD	17B ATLS	$Z' \rightarrow h Z$
		10 KHACHATRY...17AX	CMS	$Z' \rightarrow \ell\ell\ell\ell$
		11 KHACHATRY...17U	CMS	$Z' \rightarrow h Z$
>1700	95	12 SIRUNYAN	17A CMS	$Z' \rightarrow W W$
		13 SIRUNYAN	17AP CMS	$Z' \rightarrow h A$
		14 SIRUNYAN	17T CMS	$Z' \rightarrow q\bar{q}$
		15 SIRUNYAN	17V CMS	$Z' \rightarrow T t$
none 1100–1500	95	16 AABOUD	16 ATLS	$Z' \rightarrow b\bar{b}$
		17 AAD	16L ATLS	$Z' \rightarrow a\gamma, a \rightarrow \gamma\gamma$
none 1500–2600	95	18 AAD	16S ATLS	$Z' \rightarrow q\bar{q}$
none 1000–1100, none 1300–1500	95	19 KHACHATRY...16AP	CMS	$Z' \rightarrow h Z$
>2400	95	20 KHACHATRY...16E	CMS	$Z' \rightarrow t\bar{t}$
		21 AAD	15AO ATLS	$Z' \rightarrow t\bar{t}$
		22 AAD	15AT ATLS	monotop
		23 AAD	15CD ATLS	$h \rightarrow Z Z', Z' Z'; Z' \rightarrow \ell^+ \ell^-$
		24 KHACHATRY...15F	CMS	monotop
		25 KHACHATRY...15O	CMS	$Z' \rightarrow h Z$
		26 AAD	14AT ATLS	$Z' \rightarrow Z\gamma$
		27 KHACHATRY...14A	CMS	$Z' \rightarrow V V$
		28 MARTINEZ	14 RVUE	Electroweak
none 500–1740	95	29 AAD	13AQ ATLS	$Z' \rightarrow t\bar{t}$
>1320 or 1000–1280	95	30 AAD	13G ATLS	$Z' \rightarrow t\bar{t}$
> 915	95	30 AALTONEN	13A CDF	$Z' \rightarrow t\bar{t}$

>1300	95	31	CHATRCHYAN 13AP CMS	$Z' \rightarrow t\bar{t}$
>2100	95	30	CHATRCHYAN 13BMCMS	$Z' \rightarrow t\bar{t}$
		32	AAD 12BV ATLS	$Z' \rightarrow t\bar{t}$
		33	AAD 12K ATLS	$Z' \rightarrow t\bar{t}$
		34	AALTONEN 12AR CDF	Chromophilic
		35	AALTONEN 12N CDF	$Z' \rightarrow \bar{t}u$
> 835	95	36	ABAZOV 12R D0	$Z' \rightarrow t\bar{t}$
		37	CHATRCHYAN 12AI CMS	$Z' \rightarrow t\bar{u}$
		38	CHATRCHYAN 12AQ CMS	$Z' \rightarrow t\bar{t}$
>1490	95	30	CHATRCHYAN 12BL CMS	$Z' \rightarrow t\bar{t}$
		39	AALTONEN 11AD CDF	$Z' \rightarrow t\bar{t}$
		40	AALTONEN 11AE CDF	$Z' \rightarrow t\bar{t}$
		41	CHATRCHYAN 11O CMS	$pp \rightarrow tt$
		42	AALTONEN 08D CDF	$Z' \rightarrow t\bar{t}$
		42	AALTONEN 08Y CDF	$Z' \rightarrow t\bar{t}$
		42	ABAZOV 08AA D0	$Z' \rightarrow t\bar{t}$
		43	ABAZOV 04A D0	Repl. by ABAZOV 08AA
		44	BARGER 03B COSM	Nucleosynthesis; light ν_R
		45	CHO 00 RVUE	E_6 -motivated
		46	CHO 98 RVUE	E_6 -motivated
		47	ABE 97G CDF	$Z' \rightarrow \bar{q}q$

¹ AABOUD 18B search for resonances decaying to WW in pp collisions at $\sqrt{s} = 13$ TeV. The quoted limit is for heavy-vector-triplet Z' with $g_V = 1$. See their Fig.11 for limits on $\sigma \cdot B$.

² AABOUD 18F search for resonances decaying to WW in pp collisions at $\sqrt{s} = 13$ TeV. The quoted limit is for heavy-vector-triplet Z' with $g_V = 3$. The limit becomes $M_{Z'} > 2200$ GeV for $g_V = 1$. If we assume $M_{Z'} = M_{W'}$, the limit increases $M_{Z'} > 3500$ GeV and $M_{Z'} > 3100$ GeV for $g_V = 3$ and $g_V = 1$, respectively. See their Fig.5 for limits on $\sigma \cdot B$.

³ AABOUD 17AK search for a new resonance decaying to dijets in pp collisions at $\sqrt{s} = 13$ TeV. The limit quoted above is for a leptophobic Z' boson having axial-vector coupling strength with quarks $g_q = 0.2$. The limit is 2100 GeV if $g_q = 0.1$.

⁴ AABOUD 17AO search for resonances decaying to hZ in pp collisions at $\sqrt{s} = 13$ TeV. The limit quoted above is for a Z' in the heavy-vector-triplet model with $g_V = 3$. See their Fig.4 for limits on $\sigma \cdot B$.

⁵ SIRUNYAN 17AK search for resonances decaying to WW or hZ in pp collisions at $\sqrt{s} = 8$ and 13 TeV. The quoted limit is for heavy-vector-triplet Z' with $g_V = 3$. The limit becomes $M_{Z'} > 2200$ GeV for $g_V = 1$. If we assume $M_{Z'} = M_{W'}$, the limit increases $M_{Z'} > 2400$ GeV for both $g_V = 3$ and $g_V = 1$. See their Fig.1 and 2 for limits on $\sigma \cdot B$.

⁶ SIRUNYAN 17Q search for a resonance decaying to $t\bar{t}$ in pp collisions at $\sqrt{s} = 13$ TeV. The limit quoted above is for a resonance with relative width $\Gamma_{Z'} / M_{Z'} = 0.01$. Limits for wider resonances are available. See their Fig.6 for limits on $\sigma \cdot B$.

⁷ SIRUNYAN 17R search for resonances decaying to hZ in pp collisions at $\sqrt{s} = 13$ TeV. The quoted limit is for heavy-vector-triplet Z' with $g_V = 3$. Mass regions $M_{Z'} < 1150$ GeV and 1250 GeV $< M_{Z'} < 1670$ GeV are excluded for $g_V = 1$. If we assume $M_{Z'} = M_{W'}$, the excluded mass regions are $1000 < M_{Z'} < 2500$ GeV and $2760 < M_{Z'} < 3300$ GeV for $g_V = 3$; $1000 < M_{Z'} < 2430$ GeV and $2810 < M_{Z'} < 3130$ GeV for $g_V = 1$. See their Fig.5 for limits on $\sigma \cdot B$.

- ⁸ SIRUNYAN 18G search for a new resonance decaying to dijets in pp collisions at $\sqrt{s} = 13$ TeV in the mass range 50–300 GeV. See their Fig.7 for limits in the mass-coupling plane.
- ⁹ AABOUD 17B search for resonances decaying to hZ ($h \rightarrow b\bar{b}, c\bar{c}; Z \rightarrow \ell^+\ell^-, \nu\bar{\nu}$) in pp collisions at $\sqrt{s} = 13$ TeV. The quoted limit is for heavy-vector-triplet Z' with $g_V = 3$. The limit becomes $M_{Z'} > 1490$ GeV for $g_V = 1$. If we assume $M_{Z'} = M_{W'}$, the limit increases $M_{Z'} > 2310$ GeV and $M_{Z'} > 1730$ GeV for $g_V = 3$ and $g_V = 1$, respectively. See their Fig.3 for limits on $\sigma \cdot B$.
- ¹⁰ KHACHATRYAN 17AX search for lepto-phobic resonances decaying to four leptons in pp collisions at $\sqrt{s} = 8$ TeV.
- ¹¹ KHACHATRYAN 17U search for resonances decaying to hZ ($h \rightarrow b\bar{b}; Z \rightarrow \ell^+\ell^-, \nu\bar{\nu}$) in pp collisions at $\sqrt{s} = 13$ TeV. The limit on the heavy-vector-triplet model is $M_{Z'} = M_{W'} > 2$ TeV for $g_V = 3$, in which constraints from the $W' \rightarrow hW$ ($h \rightarrow b\bar{b}; W \rightarrow \ell\nu$) are combined. See their Fig.3 and Fig.4 for limits on $\sigma \cdot B$.
- ¹² SIRUNYAN 17A search for resonances decaying to WW with $WW \rightarrow \ell\nu q\bar{q}, q\bar{q}q\bar{q}$ in pp collisions at $\sqrt{s} = 13$ TeV. The quoted limit is for heavy-vector-triplet Z' with $g_V = 3$. The limit becomes $M_{Z'} > 1600$ GeV for $g_V = 1$. If we assume $M_{Z'} = M_{W'}$, the limit increases $M_{Z'} > 2400$ GeV and $M_{Z'} > 2300$ GeV for $g_V = 3$ and $g_V = 1$, respectively. See their Fig.6 for limits on $\sigma \cdot B$.
- ¹³ SIRUNYAN 17AP search for resonances decaying into a SM-like Higgs scalar h and a light pseudo scalar A . A is assumed to decay invisibly. See their Fig.9 for limits on $\sigma \cdot B$.
- ¹⁴ SIRUNYAN 17T search for a new resonance decaying to dijets in pp collisions at $\sqrt{s} = 13$ TeV in the mass range 100–300 GeV. See their Fig.3 for limits in the mass-coupling plane.
- ¹⁵ SIRUNYAN 17V search for a new resonance decaying to a top quark and a heavy vector-like top partner T in pp collisions at $\sqrt{s} = 13$ TeV. See their table 5 for limits on the Z' production cross section for various values of $M_{Z'}$ and M_T in the range of $M_{Z'} = 1500$ –2500 GeV and $M_T = 700$ –1500 GeV.
- ¹⁶ AABOUD 16 search for a narrow resonance decaying into $b\bar{b}$ in pp collisions at $\sqrt{s} = 13$ TeV. The limit quoted above is for a leptophobic Z' with SM-like couplings to quarks. See their Fig.6 for limits on $\sigma \cdot B$.
- ¹⁷ AAD 16L search for $Z' \rightarrow a\gamma$, $a \rightarrow \gamma\gamma$ in pp collisions at $\sqrt{s} = 8$ TeV. See their Table 6 for limits on $\sigma \cdot B$.
- ¹⁸ AAD 16S search for a new resonance decaying to dijets in pp collisions at $\sqrt{s} = 13$ TeV. The limit quoted above is for a leptophobic Z' having coupling strength with quark $g_q = 0.3$ and is taken from their Figure 3.
- ¹⁹ KHACHATRYAN 16AP search for a resonance decaying to hZ in pp collisions at $\sqrt{s} = 8$ TeV. Both h and Z are assumed to decay to fat jets. The quoted limit is for heavy-vector-triplet Z' with $g_V = 3$.
- ²⁰ KHACHATRYAN 16E search for a leptophobic top-color Z' decaying to $t\bar{t}$ using pp collisions at $\sqrt{s} = 8$ TeV. The quoted limit assumes that $\Gamma_{Z'}/m_{Z'} = 0.012$. Also $m_{Z'} < 2.9$ TeV is excluded for wider topcolor Z' with $\Gamma_{Z'}/m_{Z'} = 0.1$.
- ²¹ AAD 15AO search for narrow resonance decaying to $t\bar{t}$ using pp collisions at $\sqrt{s} = 8$ TeV. See Fig. 11 for limit on σB .
- ²² AAD 15AT search for monotop production plus large missing E_T events in pp collisions at $\sqrt{s} = 8$ TeV and give constraints on a Z' model having $Z' u\bar{t}$ coupling. Z' is assumed to decay invisibly. See their Fig. 6 for limits on $\sigma \cdot B$.
- ²³ AAD 15CD search for decays of Higgs bosons to 4 ℓ states via Z' bosons, $h \rightarrow ZZ' \rightarrow 4\ell$ or $h \rightarrow Z'Z' \rightarrow 4\ell$. See Fig. 5 for the limit on the signal strength of the $h \rightarrow ZZ' \rightarrow 4\ell$ process and Fig. 16 for the limit on $h \rightarrow Z'Z' \rightarrow 4\ell$.

- 24 KHACHATRYAN 15F search for monotop production plus large missing E_T events in pp collisions at $\sqrt{s} = 8$ TeV and give constraints on a Z' model having $Z' u \bar{t}$ coupling. Z' is assumed to decay invisibly. See Fig. 3 for limits on σB .
- 25 KHACHATRYAN 15O search for narrow Z' resonance decaying to $Z h$ in pp collisions at $\sqrt{s} = 8$ TeV. See their Fig. 6 for limit on σB .
- 26 AAD 14AT search for a narrow neutral vector boson decaying to $Z \gamma$. See their Fig. 3b for the exclusion limit in $m_{Z'} - \sigma B$ plane.
- 27 KHACHATRYAN 14A search for new resonance in the $W W (\ell \nu q \bar{q})$ and the $Z Z (\ell \ell q \bar{q})$ channels using pp collisions at $\sqrt{s}=8$ TeV. See their Fig.13 for the exclusion limit on the number of events in the mass-width plane.
- 28 MARTINEZ 14 use various electroweak data to constrain the Z' boson in the 3-3-1 models.
- 29 AAD 13AQ search for a leptophobic top-color Z' decaying to $t \bar{t}$. The quoted limit assumes that $\Gamma_{Z'}/m_{Z'} = 0.012$.
- 30 CHATRCHYAN 13BM search for top-color Z' decaying to $t \bar{t}$ using pp collisions at $\sqrt{s}=8$ TeV. The quoted limit is for $\Gamma_{Z'}/m_{Z'} = 0.012$.
- 31 CHATRCHYAN 13AP search for top-color leptophobic Z' decaying to $t \bar{t}$ using pp collisions at $\sqrt{s}=7$ TeV. The quoted limit is for $\Gamma_{Z'}/m_{Z'} = 0.012$.
- 32 AAD 12BV search for narrow resonance decaying to $t \bar{t}$ using pp collisions at $\sqrt{s}=7$ TeV. See their Fig. 7 for limit on $\sigma \cdot B$.
- 33 AAD 12K search for narrow resonance decaying to $t \bar{t}$ using pp collisions at $\sqrt{s}=7$ TeV. See their Fig. 5 for limit on $\sigma \cdot B$.
- 34 AALTONEN 12AR search for chromophilic Z' in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV. See their Fig. 5 for limit on $\sigma \cdot B$.
- 35 AALTONEN 12N search for $p\bar{p} \rightarrow t Z', Z' \rightarrow \bar{t} u$ events in $p\bar{p}$ collisions. See their Fig. 3 for the limit on $\sigma \cdot B$.
- 36 ABAZOV 12R search for top-color Z' boson decaying exclusively to $t \bar{t}$. The quoted limit is for $\Gamma_{Z'}/m_{Z'} = 0.012$.
- 37 CHATRCHYAN 12AI search for $pp \rightarrow tt$ events and give constraints on a Z' model having $Z' \bar{u} t$ coupling. See their Fig. 4 for the limit in mass-coupling plane.
- 38 Search for resonance decaying to $t \bar{t}$. See their Fig. 6 for limit on $\sigma \cdot B$.
- 39 Search for narrow resonance decaying to $t \bar{t}$. See their Fig. 4 for limit on $\sigma \cdot B$.
- 40 Search for narrow resonance decaying to $t \bar{t}$. See their Fig. 3 for limit on $\sigma \cdot B$.
- 41 CHATRCHYAN 11O search for same-sign top production in pp collisions induced by a hypothetical FCNC Z' at $\sqrt{s} = 7$ TeV. See their Fig. 3 for limit in mass-coupling plane.
- 42 Search for narrow resonance decaying to $t \bar{t}$. See their Fig. 3 for limit on $\sigma \cdot B$.
- 43 Search for narrow resonance decaying to $t \bar{t}$. See their Fig. 2 for limit on $\sigma \cdot B$.
- 44 BARGER 03B use the nucleosynthesis bound on the effective number of light neutrino δN_ν . See their Figs. 4–5 for limits in general E_6 motivated models.
- 45 CHO 00 use various electroweak data to constrain Z' models assuming $m_H=100$ GeV. See Fig. 2 for limits in general E_6 -motivated models.
- 46 CHO 98 study constraints on four-Fermi contact interactions obtained from low-energy electroweak experiments, assuming no Z - Z' mixing.
- 47 Search for Z' decaying to dijets at $\sqrt{s}=1.8$ TeV. For Z' with electromagnetic strength coupling, no bound is obtained.

Searches for Z' with Lepton-Flavor-Violating decays

The following limits are obtained from $p\bar{p}$ or $pp \rightarrow Z' X$ with Z' decaying to the mode indicated in the comments.

VALUE	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
	¹ AABOUD 16P	ATLS	$Z' \rightarrow e\mu, e\tau, \mu\tau$
	² KHACHATRYAN 16BE	CMS	$Z' \rightarrow e\mu$
	³ AAD 150	ATLS	$Z' \rightarrow e\mu, e\tau, \mu\tau$
	⁴ AAD 11H	ATLS	$Z' \rightarrow e\mu$
	⁵ AAD 11Z	ATLS	$Z' \rightarrow e\mu$
	⁶ ABULENCIA 06M	CDF	$Z' \rightarrow e\mu$

¹ AABOUD 16P search for new particle with lepton flavor violating decay in pp collisions at $\sqrt{s} = 13$ TeV. See their Figs.2, 3, and 4 for limits on $\sigma \cdot B$.

² KHACHATRYAN 16BE search for new particle Z' with lepton flavor violating decay in pp collisions at $\sqrt{s} = 8$ TeV in the range of $200 \text{ GeV} < M_{Z'} < 2000 \text{ GeV}$. See their Fig.4 for limits on $\sigma \cdot B$ and their Table 5 for bounds on various masses.

³ AAD 150 search for new particle Z' with lepton flavor violating decay in pp collisions at $\sqrt{s} = 8$ TeV in the range of $500 \text{ GeV} < M_{Z'} < 3000 \text{ GeV}$. See their Fig. 2 for limits on σB .

⁴ AAD 11H search for new particle Z' with lepton flavor violating decay in pp collisions at $\sqrt{s} = 7$ TeV in the range of $700 \text{ GeV} < M_{Z'} < 1000 \text{ GeV}$. See their Fig. 3 for limits on $\sigma \cdot B$.

⁵ AAD 11Z search for new particle Z' with lepton flavor violating decay in pp collisions at $\sqrt{s} = 7$ TeV in the range $700 \text{ GeV} < M_{Z'} < 2000 \text{ GeV}$. See their Fig. 3 for limits on $\sigma \cdot B$.

⁶ ABULENCIA 06M search for new particle Z' with lepton flavor violating decay in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV in the range of $100 \text{ GeV} < M_{Z'} < 800 \text{ GeV}$. See their Fig. 4 for limits in the mass-coupling plane.

Indirect Constraints on Kaluza-Klein Gauge Bosons

Bounds on a Kaluza-Klein excitation of the Z boson or photon in $d=1$ extra dimension. These bounds can also be interpreted as a lower bound on $1/R$, the size of the extra dimension. Unless otherwise stated, bounds assume all fermions live on a single brane and all gauge fields occupy the $4+d$ -dimensional bulk. See also the section on "Extra Dimensions" in the "Searches" Listings in this Review.

VALUE (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
> 4.7		¹ MUECK 02	RVUE	Electroweak
> 3.3	95	² CORNET 00	RVUE	$e\nu q q'$
>5000		³ DELGADO 00	RVUE	ϵ_K
> 2.6	95	⁴ DELGADO 00	RVUE	Electroweak
> 3.3	95	⁵ RIZZO 00	RVUE	Electroweak
> 2.9	95	⁶ MARCIANO 99	RVUE	Electroweak
> 2.5	95	⁷ MASIP 99	RVUE	Electroweak
> 1.6	90	⁸ NATH 99	RVUE	Electroweak
> 3.4	95	⁹ STRUMIA 99	RVUE	Electroweak

- ¹ MUECK 02 limit is 2σ and is from global electroweak fit ignoring correlations among observables. Higgs is assumed to be confined on the brane and its mass is fixed. For scenarios of bulk Higgs, of brane-SU(2)_L, bulk-U(1)_Y, and of bulk-SU(2)_L, brane-U(1)_Y, the corresponding limits are > 4.6 TeV, > 4.3 TeV and > 3.0 TeV, respectively.
- ² Bound is derived from limits on $e\nu qq'$ contact interaction, using data from HERA and the Tevatron.
- ³ Bound holds only if first two generations of quarks lives on separate branes. If quark mixing is not complex, then bound lowers to 400 TeV from Δm_K .
- ⁴ See Figs. 1 and 2 of DELGADO 00 for several model variations. Special boundary conditions can be found which permit KK states down to 950 GeV and that agree with the measurement of $Q_W(\text{Cs})$. Quoted bound assumes all Higgs bosons confined to brane; placing one Higgs doublet in the bulk lowers bound to 2.3 TeV.
- ⁵ Bound is derived from global electroweak analysis assuming the Higgs field is trapped on the matter brane. If the Higgs propagates in the bulk, the bound increases to 3.8 TeV.
- ⁶ Bound is derived from global electroweak analysis but considering only presence of the KK W bosons.
- ⁷ Global electroweak analysis used to obtain bound independent of position of Higgs on brane or in bulk.
- ⁸ Bounds from effect of KK states on G_F , α , M_W , and M_Z . Hard cutoff at string scale determined using gauge coupling unification. Limits for $d=2,3,4$ rise to 3.5, 5.7, and 7.8 TeV.
- ⁹ Bound obtained for Higgs confined to the matter brane with $m_H=500$ GeV. For Higgs in the bulk, the bound increases to 3.5 TeV.

See the related review(s):

[Leptoquarks](#)

MASS LIMITS for Leptoquarks from Pair Production

These limits rely only on the color or electroweak charge of the leptoquark.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
> 740	95	1 KHACHATRY...17J	CMS	Third generation
> 850	95	2 SIRUNYAN 17H	CMS	Third generation
>1050	95	3 AAD 16G	ATLS	First generation
>1000	95	4 AAD 16G	ATLS	Second generation
> 625	95	5 AAD 16G	ATLS	Third generation
none 200–640	95	6 AAD 16G	ATLS	Third generation
>1010	95	7 KHACHATRY...16AF	CMS	First generation
>1080	95	8 KHACHATRY...16AF	CMS	Second generation
> 685	95	9 KHACHATRY...15AJ	CMS	Third generation
> 740	95	10 KHACHATRY...14T	CMS	Third generation

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

> 534	95	11 AAD 13AE	ATLS	Third generation
> 525	95	12 CHATRCHYAN 13M	CMS	Third generation
> 660	95	13 AAD 12H	ATLS	First generation
> 685	95	14 AAD 120	ATLS	Second generation
> 830	95	15 CHATRCHYAN 12AG	CMS	First generation
> 840	95	16 CHATRCHYAN 12AG	CMS	Second generation
> 450	95	17 CHATRCHYAN 12BO	CMS	Third generation
> 376	95	18 AAD 11D	ATLS	Superseded by AAD 12H
> 422	95	19 AAD 11D	ATLS	Superseded by AAD 120
> 326	95	20 ABAZOV 11V	D0	First generation
> 339	95	21 CHATRCHYAN 11N	CMS	Superseded by CHATRCHYAN 12AG

> 384	95	22	KHACHATRY...11D	CMS	Superseded by CHA-TRCHYAN 12AG
> 394	95	23	KHACHATRY...11E	CMS	Superseded by CHA-TRCHYAN 12AG
> 247	95	24	ABAZOV	10L D0	Third generation
> 316	95	25	ABAZOV	09 D0	Second generation
> 299	95	26	ABAZOV	09AF D0	Superseded by ABAZOV 11V
		27	AALTONEN	08P CDF	Third generation
> 153	95	28	AALTONEN	08Z CDF	Third generation
> 205	95	29	ABAZOV	08AD D0	All generations
> 210	95	28	ABAZOV	08AN D0	Third generation
> 229	95	30	ABAZOV	07J D0	Superseded by ABAZOV 10L
> 251	95	31	ABAZOV	06A D0	Superseded by ABAZOV 09
> 136	95	32	ABAZOV	06L D0	Superseded by ABAZOV 08AD
> 226	95	33	ABULENCIA	06T CDF	Second generation
> 256	95	34	ABAZOV	05H D0	First generation
> 117	95	29	ACOSTA	05I CDF	First generation
> 236	95	35	ACOSTA	05P CDF	First generation
> 99	95	36	ABBIENDI	03R OPAL	First generation
> 100	95	36	ABBIENDI	03R OPAL	Second generation
> 98	95	36	ABBIENDI	03R OPAL	Third generation
> 98	95	37	ABAZOV	02 D0	All generations
> 225	95	38	ABAZOV	01D D0	First generation
> 85.8	95	39	ABBIENDI	00M OPAL	Superseded by ABBIENDI 03R
> 85.5	95	39	ABBIENDI	00M OPAL	Superseded by ABBIENDI 03R
> 82.7	95	39	ABBIENDI	00M OPAL	Superseded by ABBIENDI 03R
> 200	95	40	ABBOTT	00C D0	Second generation
> 123	95	41	AFFOLDER	00K CDF	Second generation
> 148	95	42	AFFOLDER	00K CDF	Third generation
> 160	95	43	ABBOTT	99J D0	Second generation
> 225	95	44	ABBOTT	98E D0	First generation
> 94	95	45	ABBOTT	98J D0	Third generation
> 202	95	46	ABE	98S CDF	Second generation
> 242	95	47	GROSS-PILCH.98		First generation
> 99	95	48	ABE	97F CDF	Third generation
> 213	95	49	ABE	97X CDF	First generation
> 45.5	95	50,51	ABREU	93J DLPH	First + second generation
> 44.4	95	52	ADRIANI	93M L3	First generation
> 44.5	95	52	ADRIANI	93M L3	Second generation
> 45	95	52	DECAMP	92 ALEP	Third generation
none 8.9–22.6	95	53	KIM	90 AMY	First generation
none 10.2–23.2	95	53	KIM	90 AMY	Second generation
none 5–20.8	95	54	BARTEL	87B JADE	
none 7–20.5	95	55	BEHREND	86B CELL	

¹ KHACHATRYAN 17J search for scalar leptoquarks decaying to τb using pp collisions at $\sqrt{s} = 13$ TeV. The limit above assumes $B(\tau b) = 1$.

² SIRUNYAN 17H search for scalar leptoquarks using $\tau\tau bb$ events in pp collisions at $\sqrt{s} = 8$ TeV. The limit above assumes $B(\tau b) = 1$.

³ AAD 16G search for scalar leptoquarks using $eejj$ events in collisions at $\sqrt{s} = 8$ TeV. The limit above assumes $B(eq) = 1$.

⁴ AAD 16G search for scalar leptoquarks using $\mu\mu jj$ events in collisions at $\sqrt{s} = 8$ TeV. The limit above assumes $B(\mu q) = 1$.

- ⁵ AAD 16G search for scalar leptoquarks decaying to $b\nu$. The limit above assumes $B(b\nu) = 1$.
- ⁶ AAD 16G search for scalar leptoquarks decaying to $t\nu$. The limit above assumes $B(t\nu) = 1$.
- ⁷ KHACHATRYAN 16AF search for scalar leptoquarks using $eejj$ and $e\nu jj$ events in pp collisions at $\sqrt{s} = 8$ TeV. The limit above assumes $B(eq) = 1$. For $B(eq) = 0.5$, the limit becomes 850 GeV.
- ⁸ KHACHATRYAN 16AF search for scalar leptoquarks using $\mu\mu jj$ and $\mu\nu jj$ events in pp collisions at $\sqrt{s} = 8$ TeV. The limit above assumes $B(\mu q) = 1$. For $B(\mu q) = 0.5$, the limit becomes 760 GeV.
- ⁹ KHACHATRYAN 15AJ search for scalar leptoquarks using $\tau\tau tt$ events in pp collisions at $\sqrt{s} = 8$ TeV. The limit above assumes $B(\tau t) = 1$.
- ¹⁰ KHACHATRYAN 14T search for scalar leptoquarks decaying to τb using pp collisions at $\sqrt{s} = 8$ TeV. The limit above assumes $B(\tau b) = 1$. See their Fig. 5 for the exclusion limit as function of $B(\tau b)$.
- ¹¹ AAD 13AE search for scalar leptoquarks using $\tau\tau bb$ events in pp collisions at $E_{\text{cm}} = 7$ TeV. The limit above assumes $B(\tau b) = 1$.
- ¹² CHATRCHYAN 13M search for scalar and vector leptoquarks decaying to τb in pp collisions at $E_{\text{cm}} = 7$ TeV. The limit above is for scalar leptoquarks with $B(\tau b) = 1$.
- ¹³ AAD 12H search for scalar leptoquarks using $eejj$ and $e\nu jj$ events in pp collisions at $E_{\text{cm}} = 7$ TeV. The limit above assumes $B(eq) = 1$. For $B(eq) = 0.5$, the limit becomes 607 GeV.
- ¹⁴ AAD 12O search for scalar leptoquarks using $\mu\mu jj$ and $\mu\nu jj$ events in pp collisions at $E_{\text{cm}} = 7$ TeV. The limit above assumes $B(\mu q) = 1$. For $B(\mu q) = 0.5$, the limit becomes 594 GeV.
- ¹⁵ CHATRCHYAN 12AG search for scalar leptoquarks using $eejj$ and $e\nu jj$ events in pp collisions at $E_{\text{cm}} = 7$ TeV. The limit above assumes $B(eq) = 1$. For $B(eq) = 0.5$, the limit becomes 640 GeV.
- ¹⁶ CHATRCHYAN 12AG search for scalar leptoquarks using $\mu\mu jj$ and $\mu\nu jj$ events in pp collisions at $E_{\text{cm}} = 7$ TeV. The limit above assumes $B(\mu q) = 1$. For $B(\mu q) = 0.5$, the limit becomes 650 GeV.
- ¹⁷ CHATRCHYAN 12BO search for scalar leptoquarks decaying to νb in pp collisions at $\sqrt{s} = 7$ TeV. The limit above assumes $B(\nu b) = 1$.
- ¹⁸ AAD 11D search for scalar leptoquarks using $eejj$ and $e\nu jj$ events in pp collisions at $E_{\text{cm}} = 7$ TeV. The limit above assumes $B(eq) = 1$. For $B(eq) = 0.5$, the limit becomes 319 GeV.
- ¹⁹ AAD 11D search for scalar leptoquarks using $\mu\mu jj$ and $\mu\nu jj$ events in pp collisions at $E_{\text{cm}} = 7$ TeV. The limit above assumes $B(\mu q) = 1$. For $B(\mu q) = 0.5$, the limit becomes 362 GeV.
- ²⁰ ABAZOV 11V search for scalar leptoquarks using $e\nu jj$ events in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.96$ TeV. The limit above assumes $B(eq) = 0.5$.
- ²¹ CHATRCHYAN 11N search for scalar leptoquarks using $e\nu jj$ events in pp collisions at $E_{\text{cm}} = 7$ TeV. The limit above assumes $B(eq) = 0.5$.
- ²² KHACHATRYAN 11D search for scalar leptoquarks using $eejj$ events in pp collisions at $E_{\text{cm}} = 7$ TeV. The limit above assumes $B(eq) = 1$.
- ²³ KHACHATRYAN 11E search for scalar leptoquarks using $\mu\mu jj$ events in pp collisions at $E_{\text{cm}} = 7$ TeV. The limit above assumes $B(\mu q) = 1$.
- ²⁴ ABAZOV 10L search for pair productions of scalar leptoquark state decaying to νb in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.96$ TeV. The limit above assumes $B(\nu b) = 1$.
- ²⁵ ABAZOV 09 search for scalar leptoquarks using $\mu\mu jj$ and $\mu\nu jj$ events in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.96$ TeV. The limit above assumes $B(\mu q) = 1$. For $B(\mu q) = 0.5$, the limit becomes 270 GeV.
- ²⁶ ABAZOV 09AF search for scalar leptoquarks using $eejj$ and $e\nu jj$ events in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.96$ TeV. The limit above assumes $B(eq) = 1$. For $B(eq) = 0.5$ the bound becomes 284 GeV.

- 27 AALTONEN 08P search for vector leptoquarks using $\tau^+ \tau^- b \bar{b}$ events in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.96$ TeV. Assuming Yang-Mills (minimal) couplings, the mass limit is >317 GeV (251 GeV) at 95% CL for $B(\tau b) = 1$.
- 28 Search for pair production of scalar leptoquark state decaying to τb in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.96$ TeV. The limit above assumes $B(\tau b) = 1$.
- 29 Search for scalar leptoquarks using $\nu\nu jj$ events in $\bar{p}p$ collisions at $E_{\text{cm}} = 1.96$ TeV. The limit above assumes $B(\nu q) = 1$.
- 30 ABAZOV 07J search for pair productions of scalar leptoquark state decaying to νb in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.96$ TeV. The limit above assumes $B(\nu b) = 1$.
- 31 ABAZOV 06A search for scalar leptoquarks using $\mu\mu jj$ events in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.8$ TeV and 1.96 TeV. The limit above assumes $B(\mu q) = 1$. For $B(\mu q) = 0.5$, the limit becomes 204 GeV.
- 32 ABAZOV 06L search for scalar leptoquarks using $\nu\nu jj$ events in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.8$ TeV and at 1.96 TeV. The limit above assumes $B(\nu q) = 1$.
- 33 ABULENCIA 06T search for scalar leptoquarks using $\mu\mu jj$, $\mu\nu jj$, and $\nu\nu jj$ events in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.96$ TeV. The quoted limit assumes $B(\mu q) = 1$. For $B(\mu q) = 0.5$ or 0.1, the bound becomes 208 GeV or 143 GeV, respectively. See their Fig. 4 for the exclusion limit as a function of $B(\mu q)$.
- 34 ABAZOV 05H search for scalar leptoquarks using $eejj$ and $e\nu jj$ events in $\bar{p}p$ collisions at $E_{\text{cm}} = 1.8$ TeV and 1.96 TeV. The limit above assumes $B(eq) = 1$. For $B(eq) = 0.5$ the bound becomes 234 GeV.
- 35 ACOSTA 05P search for scalar leptoquarks using $eejj$, $e\nu jj$ events in $\bar{p}p$ collisions at $E_{\text{cm}} = 1.96$ TeV. The limit above assumes $B(eq) = 1$. For $B(eq) = 0.5$ and 0.1, the bound becomes 205 GeV and 145 GeV, respectively.
- 36 ABBIENDI 03R search for scalar/vector leptoquarks in $e^+ e^-$ collisions at $\sqrt{s} = 189\text{--}209$ GeV. The quoted limits are for charge $-4/3$ isospin 0 scalar-leptoquark with $B(\ell q) = 1$. See their table 12 for other cases.
- 37 ABAZOV 02 search for scalar leptoquarks using $\nu\nu jj$ events in $\bar{p}p$ collisions at $E_{\text{cm}} = 1.8$ TeV. The bound holds for all leptoquark generations. Vector leptoquarks are likewise constrained to lie above 200 GeV.
- 38 ABAZOV 01D search for scalar leptoquarks using $e\nu jj$, $eejj$, and $\nu\nu jj$ events in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.8$ TeV. The limit above assumes $B(eq) = 1$. For $B(eq) = 0.5$ and 0, the bound becomes 204 and 79 GeV, respectively. Bounds for vector leptoquarks are also given. Supersedes ABBOTT 98E.
- 39 ABBIENDI 00M search for scalar/vector leptoquarks in $e^+ e^-$ collisions at $\sqrt{s} = 183$ GeV. The quoted limits are for charge $-4/3$ isospin 0 scalar-leptoquarks with $B(\ell q) = 1$. See their Table 8 and Figs. 6–9 for other cases.
- 40 ABBOTT 00C search for scalar leptoquarks using $\mu\mu jj$, $\mu\nu jj$, and $\nu\nu jj$ events in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.8$ TeV. The limit above assumes $B(\mu q) = 1$. For $B(\mu q) = 0.5$ and 0, the bound becomes 180 and 79 GeV respectively. Bounds for vector leptoquarks are also given.
- 41 AFFOLDER 00K search for scalar leptoquark using $\nu\nu cc$ events in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.8$ TeV. The quoted limit assumes $B(\nu c) = 1$. Bounds for vector leptoquarks are also given.
- 42 AFFOLDER 00K search for scalar leptoquark using $\nu\nu bb$ events in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.8$ TeV. The quoted limit assumes $B(\nu b) = 1$. Bounds for vector leptoquarks are also given.
- 43 ABBOTT 99J search for leptoquarks using $\mu\nu jj$ events in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.8$ TeV. The quoted limit is for a scalar leptoquark with $B(\mu q) = B(\nu q) = 0.5$. Limits on vector leptoquarks range from 240 to 290 GeV.
- 44 ABBOTT 98E search for scalar leptoquarks using $e\nu jj$, $eejj$, and $\nu\nu jj$ events in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.8$ TeV. The limit above assumes $B(eq) = 1$. For $B(eq) = 0.5$ and 0, the bound becomes 204 and 79 GeV, respectively.
- 45 ABBOTT 98J search for charge $-1/3$ third generation scalar and vector leptoquarks in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.8$ TeV. The quoted limit is for scalar leptoquark with $B(\nu b) = 1$.
- 46 ABE 98S search for scalar leptoquarks using $\mu\mu jj$ events in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.8$ TeV. The limit is for $B(\mu q) = 1$. For $B(\mu q) = B(\nu q) = 0.5$, the limit is > 160 GeV.

- 47 GROSS-PILCHER 98 is the combined limit of the CDF and DØ Collaborations as determined by a joint CDF/DØ working group and reported in this FNAL Technical Memo. Original data published in ABE 97X and ABBOTT 98E.
- 48 ABE 97F search for third generation scalar and vector leptoquarks in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.8$ TeV. The quoted limit is for scalar leptoquark with $B(\tau b) = 1$.
- 49 ABE 97X search for scalar leptoquarks using $e e j j$ events in $p\bar{p}$ collisions at $E_{\text{cm}}=1.8$ TeV. The limit is for $B(e q)=1$.
- 50 Limit is for charge $-1/3$ isospin-0 leptoquark with $B(\ell q) = 2/3$.
- 51 First and second generation leptoquarks are assumed to be degenerate. The limit is slightly lower for each generation.
- 52 Limits are for charge $-1/3$, isospin-0 scalar leptoquarks decaying to $\ell^- q$ or νq with any branching ratio. See paper for limits for other charge-isospin assignments of leptoquarks.
- 53 KIM 90 assume pair production of charge $2/3$ scalar-leptoquark via photon exchange. The decay of the first (second) generation leptoquark is assumed to be any mixture of $d e^+$ and $u \bar{\nu}$ ($s \mu^+$ and $c \bar{\nu}$). See paper for limits for specific branching ratios.
- 54 BARTEL 87B limit is valid when a pair of charge $2/3$ spinless leptoquarks X is produced with point coupling, and when they decay under the constraint $B(X \rightarrow c \bar{\nu}_\mu) + B(X \rightarrow s \mu^+) = 1$.
- 55 BEHREND 86B assumed that a charge $2/3$ spinless leptoquark, χ , decays either into $s \mu^+$ or $c \bar{\nu}$: $B(\chi \rightarrow s \mu^+) + B(\chi \rightarrow c \bar{\nu}) = 1$.

MASS LIMITS for Leptoquarks from Single Production

These limits depend on the q - ℓ -leptoquark coupling g_{LQ} . It is often assumed that $g_{LQ}^2/4\pi=1/137$. Limits shown are for a scalar, weak isoscalar, charge $-1/3$ leptoquark.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>1755	95	¹ KHACHATRY...16AG	CMS	First generation
> 660	95	² KHACHATRY...16AG	CMS	Second generation
> 304	95	³ ABRAMOWICZ12A	ZEUS	First generation
> 73	95	⁴ ABREU 93J	DLPH	Second generation
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
		⁵ DEY 16	ICCB	$\nu q \rightarrow LQ \rightarrow \nu q$
		⁶ AARON 11A	H1	Lepton-flavor violation
> 300	95	⁷ AARON 11B	H1	First generation
		⁸ ABAZOV 07E	D0	Second generation
> 295	95	⁹ AKTAS 05B	H1	First generation
		¹⁰ CHEKANOV 05A	ZEUS	Lepton-flavor violation
> 298	95	¹¹ CHEKANOV 03B	ZEUS	First generation
> 197	95	¹² ABBIENDI 02B	OPAL	First generation
		¹³ CHEKANOV 02	ZEUS	Repl. by CHEKANOV 05A
> 290	95	¹⁴ ADLOFF 01C	H1	First generation
> 204	95	¹⁵ BREITWEG 01	ZEUS	First generation
		¹⁶ BREITWEG 00E	ZEUS	First generation
> 161	95	¹⁷ ABREU 99G	DLPH	First generation
> 200	95	¹⁸ ADLOFF 99	H1	First generation
		¹⁹ DERRICK 97	ZEUS	Lepton-flavor violation
> 168	95	²⁰ DERRICK 93	ZEUS	First generation

- ¹ KHACHATRYAN 16AG search for single production of charge $\pm 1/3$ scalar leptoquarks using $e e j$ events in $p p$ collisions at $\sqrt{s} = 8$ TeV. The limit above assumes $B(e q) = 1$ and the leptoquark coupling strength $\lambda = 1$.
 - ² KHACHATRYAN 16AG search for single production of charge $\pm 1/3$ scalar leptoquarks using $\mu \mu j$ events in $p p$ collisions at $\sqrt{s} = 8$ TeV. The limit above assumes $B(\mu q) = 1$ and the leptoquark coupling strength $\lambda = 1$.
 - ³ ABRAMOWICZ 12A limit is for a scalar, weak isoscalar, charge $-1/3$ leptoquark coupled with e_R . See their Figs. 12–17 and Table 4 for states with different quantum numbers.
 - ⁴ Limit from single production in Z decay. The limit is for a leptoquark coupling of electromagnetic strength and assumes $B(\ell q) = 2/3$. The limit is 77 GeV if first and second leptoquarks are degenerate.
 - ⁵ DEY 16 use the 2010-2012 IceCube PeV energy data set to constrain the leptoquark production cross section through the $\nu q \rightarrow LQ \rightarrow \nu q$ process. See their Figure 4 for the exclusion limit in the mass-coupling plane.
 - ⁶ AARON 11A search for various leptoquarks with lepton-flavor violating couplings. See their Figs. 2–3 and Tables 1–4 for detailed limits.
 - ⁷ The quoted limit is for a scalar, weak isoscalar, charge $-1/3$ leptoquark coupled with e_R . See their Figs. 3–5 for limits on states with different quantum numbers.
 - ⁸ ABAZOV 07E search for leptoquark single production through $q g$ fusion process in $p \bar{p}$ collisions. See their Fig. 4 for exclusion plot in mass-coupling plane.
 - ⁹ AKTAS 05B limit is for a scalar, weak isoscalar, charge $-1/3$ leptoquark coupled with e_R . See their Fig. 3 for limits on states with different quantum numbers.
 - ¹⁰ CHEKANOV 05 search for various leptoquarks with lepton-flavor violating couplings. See their Figs. 6–10 and Tables 1–8 for detailed limits.
 - ¹¹ CHEKANOV 03B limit is for a scalar, weak isoscalar, charge $-1/3$ leptoquark coupled with e_R . See their Figs. 11–12 and Table 5 for limits on states with different quantum numbers.
 - ¹² For limits on states with different quantum numbers and the limits in the mass-coupling plane, see their Fig. 4 and Fig. 5.
 - ¹³ CHEKANOV 02 search for various leptoquarks with lepton-flavor violating couplings. See their Figs. 6–7 and Tables 5–6 for detailed limits.
 - ¹⁴ For limits on states with different quantum numbers and the limits in the mass-coupling plane, see their Fig. 3.
 - ¹⁵ See their Fig. 14 for limits in the mass-coupling plane.
 - ¹⁶ BREITWEG 00E search for $F=0$ leptoquarks in $e^+ p$ collisions. For limits in mass-coupling plane, see their Fig. 11.
 - ¹⁷ ABREU 99G limit obtained from process $e \gamma \rightarrow LQ + q$. For limits on vector and scalar states with different quantum numbers and the limits in the coupling-mass plane, see their Fig. 4 and Table 2.
 - ¹⁸ For limits on states with different quantum numbers and the limits in the mass-coupling plane, see their Fig. 13 and Fig. 14. ADLOFF 99 also search for leptoquarks with lepton-flavor violating couplings. ADLOFF 99 supersedes AID 96B.
 - ¹⁹ DERRICK 97 search for various leptoquarks with lepton-flavor violating couplings. See their Figs. 5–8 and Table 1 for detailed limits.
 - ²⁰ DERRICK 93 search for single leptoquark production in $e p$ collisions with the decay $e q$ and νq . The limit is for leptoquark coupling of electromagnetic strength and assumes $B(e q) = B(\nu q) = 1/2$. The limit for $B(e q) = 1$ is 176 GeV. For limits on states with different quantum numbers, see their Table 3.
-

Indirect Limits for Leptoquarks

VALUE (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
		¹ BARRANCO	16	RVUE D decays
		² KUMAR	16	RVUE neutral K mixing, rare K decays
		³ BESSAA	15	RVUE $q\bar{q} \rightarrow e^+e^-$
> 14	95	⁴ SAHOO	15A	RVUE $B_{s,d} \rightarrow \mu^+\mu^-$
		⁵ SAKAKI	13	RVUE $B \rightarrow D^{(*)}\tau\bar{\nu}, B \rightarrow X_S\nu\bar{\nu}$
		⁶ KOSNIK	12	RVUE $b \rightarrow s\ell^+\ell^-$
> 2.5	95	⁷ AARON	11C	H1 First generation
		⁸ DORSNER	11	RVUE scalar, weak singlet, charge 4/3
		⁹ AKTAS	07A	H1 Lepton-flavor violation
> 0.49	95	¹⁰ SCHAEEL	07A	ALEP $e^+e^- \rightarrow q\bar{q}$
		¹¹ SMIRNOV	07	RVUE $K \rightarrow e\mu, B \rightarrow e\tau$
		¹² CHEKANOV	05A	ZEUS Lepton-flavor violation
> 1.7	96	¹³ ADLOFF	03	H1 First generation
> 46	90	¹⁴ CHANG	03	BELL Pati-Salam type
		¹⁵ CHEKANOV	02	ZEUS Repl. by CHEKANOV 05A
> 1.7	95	¹⁶ CHEUNG	01B	RVUE First generation
> 0.39	95	¹⁷ ACCIARRI	00P	L3 $e^+e^- \rightarrow qq$
> 1.5	95	¹⁸ ADLOFF	00	H1 First generation
> 0.2	95	¹⁹ BARATE	00I	ALEP Repl. by SCHAEEL 07A
		²⁰ BARGER	00	RVUE Cs
		²¹ GABRIELLI	00	RVUE Lepton flavor violation
> 0.74	95	²² ZARNECKI	00	RVUE S_1 leptoquark
		²³ ABBIENDI	99	OPAL
> 19.3	95	²⁴ ABE	98V	CDF $B_s \rightarrow e^\pm\mu^\mp, \text{Pati-Salam type}$
		²⁵ ACCIARRI	98J	L3 $e^+e^- \rightarrow q\bar{q}$
		²⁶ ACKERSTAFF	98V	OPAL $e^+e^- \rightarrow q\bar{q}, e^+e^- \rightarrow b\bar{b}$
> 0.76	95	²⁷ DEANDREA	97	RVUE \tilde{R}_2 leptoquark
		²⁸ DERRICK	97	ZEUS Lepton-flavor violation
		²⁹ GROSSMAN	97	RVUE $B \rightarrow \tau^+\tau^-(X)$
		³⁰ JADACH	97	RVUE $e^+e^- \rightarrow q\bar{q}$
>1200		³¹ KUZNETSOV	95B	RVUE Pati-Salam type
		³² MIZUKOSHI	95	RVUE Third generation scalar leptoquark
> 0.3	95	³³ BHATTACH...	94	RVUE Spin-0 leptoquark coupled to $\bar{e}_R t_L$
		³⁴ DAVIDSON	94	RVUE
> 18		³⁵ KUZNETSOV	94	RVUE Pati-Salam type
> 0.43	95	³⁶ LEURER	94	RVUE First generation spin-1 leptoquark
> 0.44	95	³⁶ LEURER	94B	RVUE First generation spin-0 leptoquark
		³⁷ MAHANTA	94	RVUE P and T violation
> 1		³⁸ SHANKER	82	RVUE Nonchiral spin-0 leptoquark
> 125		³⁸ SHANKER	82	RVUE Nonchiral spin-1 leptoquark

¹ BARRANCO 16 give bounds on leptoquark induced four-fermion interactions from $D \rightarrow K\ell\nu$ and $D_s \rightarrow \ell\nu$.

² KUMAR 16 gives bound on SU(2) singlet scalar leptoquark with charge $-1/3$ from $K^0 - \bar{K}^0$ mixing, $K \rightarrow \pi\nu\bar{\nu}$, $K_L^0 \rightarrow \mu^+\mu^-$, and $K_L^0 \rightarrow \mu^\pm e^\mp$ decays.

³ BESSAA 15 obtain limit on leptoquark induced four-fermion interactions from the ATLAS and CMS limit on the $\bar{q}q\bar{e}e$ contact interactions.

- ⁴ SAHOO 15A obtain limit on leptoquark induced four-fermion interactions from $B_{s,d} \rightarrow \mu^+ \mu^-$ for $\lambda \simeq O(1)$.
- ⁵ SAKAKI 13 explain the $B \rightarrow D^{(*)} \tau \bar{\nu}$ anomaly using Wilson coefficients of leptoquark-induced four-fermion operators.
- ⁶ KOSNIK 12 obtains limits on leptoquark induced four-fermion interactions from $b \rightarrow s \ell^+ \ell^-$ decays.
- ⁷ AARON 11C limit is for weak isotriplet spin-0 leptoquark at strong coupling $\lambda = \sqrt{4\pi}$. For the limits of leptoquarks with different quantum numbers, see their Table 3. Limits are derived from bounds of $e q$ contact interactions.
- ⁸ DORSNER 11 give bounds on scalar, weak singlet, charge 4/3 leptoquark from K, B, τ decays, meson mixings, $LFV, g-2$ and $Z \rightarrow b \bar{b}$.
- ⁹ AKTAS 07A search for lepton-flavor violation in $e p$ collision. See their Tables 4–7 for limits on lepton-flavor violating four-fermion interactions induced by various leptoquarks.
- ¹⁰ SCHAEEL 07A limit is for the weak-isoscalar spin-0 left-handed leptoquark with the coupling of electromagnetic strength. For the limits of leptoquarks with different quantum numbers, see their Table 35.
- ¹¹ SMIRNOV 07 obtains mass limits for the vector and scalar chiral leptoquark states from $K \rightarrow e \mu, B \rightarrow e \tau$ decays.
- ¹² CHEKANOV 05 search for various leptoquarks with lepton-flavor violating couplings. See their Figs.6–10 and Tables 1–8 for detailed limits.
- ¹³ ADLOFF 03 limit is for the weak isotriplet spin-0 leptoquark at strong coupling $\lambda = \sqrt{4\pi}$. For the limits of leptoquarks with different quantum numbers, see their Table 3. Limits are derived from bounds on $e^\pm q$ contact interactions.
- ¹⁴ The bound is derived from $B(B^0 \rightarrow e^\pm \mu^\mp) < 1.7 \times 10^{-7}$.
- ¹⁵ CHEKANOV 02 search for lepton-flavor violation in $e p$ collisions. See their Tables 1–4 for limits on lepton-flavor violating and four-fermion interactions induced by various leptoquarks.
- ¹⁶ CHEUNG 01B quoted limit is for a scalar, weak isoscalar, charge $-1/3$ leptoquark with a coupling of electromagnetic strength. The limit is derived from bounds on contact interactions in a global electroweak analysis. For the limits of leptoquarks with different quantum numbers, see Table 5.
- ¹⁷ ACCIARRI 00P limit is for the weak isoscalar spin-0 leptoquark with the coupling of electromagnetic strength. For the limits of leptoquarks with different quantum numbers, see their Table 4.
- ¹⁸ ADLOFF 00 limit is for the weak isotriplet spin-0 leptoquark at strong coupling, $\lambda = \sqrt{4\pi}$. For the limits of leptoquarks with different quantum numbers, see their Table 2. ADLOFF 00 limits are from the Q^2 spectrum measurement of $e^+ p \rightarrow e^+ X$.
- ¹⁹ BARATE 00i search for deviations in cross section and jet-charge asymmetry in $e^+ e^- \rightarrow \bar{q} q$ due to t -channel exchange of a leptoquark at $\sqrt{s} = 130$ to 183 GeV. Limits for other scalar and vector leptoquarks are also given in their Table 22.
- ²⁰ BARGER 00 explain the deviation of atomic parity violation in cesium atoms from prediction is explained by scalar leptoquark exchange.
- ²¹ GABRIELLI 00 calculate various process with lepton flavor violation in leptoquark models.
- ²² ZARNECKI 00 limit is derived from data of HERA, LEP, and Tevatron and from various low-energy data including atomic parity violation. Leptoquark coupling with electromagnetic strength is assumed.
- ²³ ABBIENDI 99 limits are from $e^+ e^- \rightarrow q \bar{q}$ cross section at 130–136, 161–172, 183 GeV. See their Fig. 8 and Fig. 9 for limits in mass-coupling plane.
- ²⁴ ABE 98v quoted limit is from $B(B_s \rightarrow e^\pm \mu^\mp) < 8.2 \times 10^{-6}$. ABE 98v also obtain a similar limit on $M_{LQ} > 20.4$ TeV from $B(B_d \rightarrow e^\pm \mu^\mp) < 4.5 \times 10^{-6}$. Both bounds assume the non-canonical association of the b quark with electrons or muons under SU(4).
- ²⁵ ACCIARRI 98J limit is from $e^+ e^- \rightarrow q \bar{q}$ cross section at $\sqrt{s} = 130$ –172 GeV which can be affected by the t - and u -channel exchanges of leptoquarks. See their Fig. 4 and Fig. 5 for limits in the mass-coupling plane.

- 26 ACKERSTAFF 98V limits are from $e^+e^- \rightarrow q\bar{q}$ and $e^+e^- \rightarrow b\bar{b}$ cross sections at $\sqrt{s} = 130\text{--}172$ GeV, which can be affected by the t - and u -channel exchanges of leptoquarks. See their Fig. 21 and Fig. 22 for limits of leptoquarks in mass-coupling plane.
- 27 DEANDREA 97 limit is for \tilde{R}_2 leptoquark obtained from atomic parity violation (APV). The coupling of leptoquark is assumed to be electromagnetic strength. See Table 2 for limits of the four-fermion interactions induced by various scalar leptoquark exchange. DEANDREA 97 combines APV limit and limits from Tevatron and HERA. See Fig. 1–4 for combined limits of leptoquark in mass-coupling plane.
- 28 DERRICK 97 search for lepton-flavor violation in $e p$ collision. See their Tables 2–5 for limits on lepton-flavor violating four-fermion interactions induced by various leptoquarks.
- 29 GROSSMAN 97 estimate the upper bounds on the branching fraction $B \rightarrow \tau^+ \tau^- (X)$ from the absence of the B decay with large missing energy. These bounds can be used to constrain leptoquark induced four-fermion interactions.
- 30 JADACH 97 limit is from $e^+e^- \rightarrow q\bar{q}$ cross section at $\sqrt{s}=172.3$ GeV which can be affected by the t - and u -channel exchanges of leptoquarks. See their Fig. 1 for limits on vector leptoquarks in mass-coupling plane.
- 31 KUZNETSOV 95B use π, K, B, τ decays and μe conversion and give a list of bounds on the leptoquark mass and the fermion mixing matrix in the Pati-Salam model. The quoted limit is from $K_L \rightarrow \mu e$ decay assuming zero mixing.
- 32 MIZUKOSHI 95 calculate the one-loop radiative correction to the Z -physics parameters in various scalar leptoquark models. See their Fig. 4 for the exclusion plot of third generation leptoquark models in mass-coupling plane.
- 33 BHATTACHARYYA 94 limit is from one-loop radiative correction to the leptonic decay width of the Z . $m_H=250$ GeV, $\alpha_s(m_Z)=0.12$, $m_t=180$ GeV, and the electroweak strength of leptoquark coupling are assumed. For leptoquark coupled to $\bar{e}_L t_R, \bar{\mu} t$, and $\bar{\tau} t$, see Fig. 2 in BHATTACHARYYA 94B erratum and Fig. 3.
- 34 DAVIDSON 94 gives an extensive list of the bounds on leptoquark-induced four-fermion interactions from π, K, D, B, μ, τ decays and meson mixings, *etc.* See Table 15 of DAVIDSON 94 for detail.
- 35 KUZNETSOV 94 gives mixing independent bound of the Pati-Salam leptoquark from the cosmological limit on $\pi^0 \rightarrow \bar{\nu}\nu$.
- 36 LEURER 94, LEURER 94B limits are obtained from atomic parity violation and apply to any chiral leptoquark which couples to the first generation with electromagnetic strength. For a nonchiral leptoquark, universality in $\pi_{\ell 2}$ decay provides a much more stringent bound.
- 37 MAHANTA 94 gives bounds of P - and T -violating scalar-leptoquark couplings from atomic and molecular experiments.
- 38 From $(\pi \rightarrow e\nu)/(\pi \rightarrow \mu\nu)$ ratio. SHANKER 82 assumes the leptoquark induced four-fermion coupling $4g^2/M^2 (\bar{\nu}_{eL} u_R) (\bar{d}_L e_R)$ with $g=0.004$ for spin-0 leptoquark and $g^2/M^2 (\bar{\nu}_{eL} \gamma_\mu u_L) (\bar{d}_R \gamma^\mu e_R)$ with $g \simeq 0.6$ for spin-1 leptoquark.

MASS LIMITS for Diquarks

<u>VALUE (GeV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
>6000 (CL = 95%) OUR LIMIT				
none 600–6900	95	1 KHACHATRY...17W	CMS	E_6 diquark
none 1500–6000	95	2 KHACHATRY...16K	CMS	E_6 diquark
none 500–1600	95	3 KHACHATRY...16L	CMS	E_6 diquark
none 1200–4700	95	4 KHACHATRY...15V	CMS	E_6 diquark

• • • We do not use the following data for averages, fits, limits, etc. • • •

>3750	95	⁵ CHATRCHYAN 13A	CMS	E_6 diquark
none 1000–4280	95	⁶ CHATRCHYAN 13AS	CMS	Superseded by KHACHA- TRYAN 15V
>3520	95	⁷ CHATRCHYAN 11Y	CMS	Superseded by CHA- TRCHYAN 13A
none 970–1080, 1450–1600	95	⁸ KHACHATRY...10	CMS	Superseded by CHA- TRCHYAN 13A
none 290–630	95	⁹ AALTONEN 09AC	CDF	E_6 diquark
none 290–420	95	¹⁰ ABE 97G	CDF	E_6 diquark
none 15–31.7	95	¹¹ ABREU 940	DLPH	SUSY E_6 diquark

¹ KHACHATRYAN 17W search for resonances decaying to dijets in pp collisions at $\sqrt{s} = 13$ TeV.

² KHACHATRYAN 16K search for resonances decaying to dijets in pp collisions at $\sqrt{s} = 13$ TeV.

³ KHACHATRYAN 16L search for resonances decaying to dijets in pp collisions at $\sqrt{s} = 8$ TeV with the data scouting technique, increasing the sensitivity to the low mass resonances.

⁴ KHACHATRYAN 15V search for resonances decaying to dijets in pp collisions at $\sqrt{s} = 8$ TeV.

⁵ CHATRCHYAN 13A search for new resonance decaying to dijets in pp collisions at $\sqrt{s} = 7$ TeV.

⁶ CHATRCHYAN 13AS search for new resonance decaying to dijets in pp collisions at $\sqrt{s} = 8$ TeV.

⁷ CHATRCHYAN 11Y search for new resonance decaying to dijets in pp collisions at $\sqrt{s} = 7$ TeV.

⁸ KHACHATRYAN 10 search for new resonance decaying to dijets in pp collisions at $\sqrt{s} = 7$ TeV.

⁹ AALTONEN 09AC search for new narrow resonance decaying to dijets.

¹⁰ ABE 97G search for new particle decaying to dijets.

¹¹ ABREU 940 limit is from $e^+e^- \rightarrow \bar{c}c s$. Range extends up to 43 GeV if diquarks are degenerate in mass.

MASS LIMITS for g_A (axigluon) and Other Color-Octet Gauge Bosons

Axigluons are massive color-octet gauge bosons in chiral color models and have axial-vector coupling to quarks with the same coupling strength as gluons.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>5500 (CL = 95%) OUR LIMIT				
none 600–5500	95	¹ KHACHATRY...17W	CMS	$pp \rightarrow g_A X, g_A \rightarrow 2j$
none 1500–5100	95	² KHACHATRY...16K	CMS	$pp \rightarrow g_A X, g_A \rightarrow 2j$
none 500–1600	95	³ KHACHATRY...16L	CMS	$pp \rightarrow g_A X, g_A \rightarrow 2j$
none 1300–3600	95	⁴ KHACHATRY...15V	CMS	$pp \rightarrow g_A X, g_A \rightarrow 2j$

• • • We do not use the following data for averages, fits, limits, etc. • • •

		⁵ KHACHATRY...17Y	CMS	$pp \rightarrow g_A g_A \rightarrow 8j$
		⁶ AAD 16W	ATLS	$pp \rightarrow g_A X, g_A \rightarrow b\bar{b}b\bar{b}$
>2800	95	⁷ KHACHATRY...16E	CMS	$pp \rightarrow g_{KK} X, g_{KK} \rightarrow t\bar{t}$
		⁸ KHACHATRY...15AV	CMS	$pp \rightarrow \Theta^0 \Theta^0 \rightarrow b\bar{b} Z g$
		⁹ AALTONEN 13R	CDF	$p\bar{p} \rightarrow g_A X, g_A \rightarrow \sigma\sigma, \sigma \rightarrow 2j$
>3360	95	¹⁰ CHATRCHYAN 13A	CMS	$pp \rightarrow g_A X, g_A \rightarrow 2j$
none 1000–3270	95	¹¹ CHATRCHYAN 13AS	CMS	Superseded by KHACHA- TRYAN 15V

none 250–740	95	12	CHATRCHYAN 13AU	CMS	$pp \rightarrow 2g_A X, g_A \rightarrow 2j$
> 775	95	13	ABAZOV 12R	D0	$p\bar{p} \rightarrow g_A X, g_A \rightarrow t\bar{t}$
>2470	95	14	CHATRCHYAN 11Y	CMS	Superseded by CHA- TRCHYAN 13A
		15	AALTONEN 10L	CDF	$p\bar{p} \rightarrow g_A X, g_A \rightarrow t\bar{t}$
none 1470–1520	95	16	KHACHATRYAN 10	CMS	Superseded by CHA- TRCHYAN 13A
none 260–1250	95	17	AALTONEN 09AC	CDF	$p\bar{p} \rightarrow g_A X, g_A \rightarrow 2j$
> 910	95	18	CHOUDHURY 07	RVUE	$p\bar{p} \rightarrow t\bar{t} X$
> 365	95	19	DONCHESKI 98	RVUE	$\Gamma(Z \rightarrow \text{hadron})$
none 200–980	95	20	ABE 97G	CDF	$p\bar{p} \rightarrow g_A X, g_A \rightarrow 2j$
none 200–870	95	21	ABE 95N	CDF	$p\bar{p} \rightarrow g_A X, g_A \rightarrow q\bar{q}$
none 240–640	95	22	ABE 93G	CDF	$p\bar{p} \rightarrow g_A X, g_A \rightarrow 2j$
> 50	95	23	CUYPERS 91	RVUE	$\sigma(e^+ e^- \rightarrow \text{hadrons})$
none 120–210	95	24	ABE 90H	CDF	$p\bar{p} \rightarrow g_A X, g_A \rightarrow 2j$
> 29		25	ROBINETT 89	THEO	Partial-wave unitarity
none 150–310	95	26	ALBAJAR 88B	UA1	$p\bar{p} \rightarrow g_A X, g_A \rightarrow 2j$
> 20			BERGSTROM 88	RVUE	$p\bar{p} \rightarrow \gamma X$ via $g_A g$
> 9		27	CUYPERS 88	RVUE	γ decay
> 25		28	DONCHESKI 88B	RVUE	γ decay

¹ KHACHATRYAN 17W search for resonances decaying to dijets in pp collisions at $\sqrt{s} = 13$ TeV.

² KHACHATRYAN 16K search for resonances decaying to dijets in pp collisions at $\sqrt{s} = 13$ TeV.

³ KHACHATRYAN 16L search for resonances decaying to dijets in pp collisions at $\sqrt{s} = 8$ TeV with the data scouting technique, increasing the sensitivity to the low mass resonances.

⁴ KHACHATRYAN 15V search for resonances decaying to dijets in pp collisions at $\sqrt{s} = 8$ TeV.

⁵ KHACHATRYAN 17Y search for pair production of color-octet gauge boson g_A each decaying to $4j$ in pp collisions at $\sqrt{s} = 8$ TeV.

⁶ AAD 16W search for a new resonance decaying to a pair of b and B_H in pp collisions at $\sqrt{s} = 8$ TeV. The vector-like quark B_H is assumed to decay to bH . See their Fig. 3 and Fig. 4 for limits on $\sigma \cdot B$.

⁷ KHACHATRYAN 16E search for KK gluon decaying to $t\bar{t}$ in pp collisions at $\sqrt{s} = 8$ TeV.

⁸ KHACHATRYAN 15AV search for pair productions of neutral color-octet weak-triplet scalar particles (Θ^0), decaying to $b\bar{b}$, Zg or γg , in pp collisions at $\sqrt{s} = 8$ TeV. The Θ^0 particle is often predicted in coloron (G' , color-octet gauge boson) models and appear in the pp collisions through $G' \rightarrow \Theta^0 \Theta^0$ decays. Assuming $B(\Theta^0 \rightarrow b\bar{b}) = 0.5$, they give limits $m_{\Theta^0} > 623$ GeV (426 GeV) for $m_{G'} = 2.3 m_{\Theta^0}$ ($m_{G'} = 5 m_{\Theta^0}$).

⁹ AALTONEN 13R search for new resonance decaying to $\sigma\sigma$, with hypothetical strongly interacting σ particle subsequently decaying to 2 jets, in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV, using data corresponding to an integrated luminosity of 6.6 fb^{-1} . For $50 \text{ GeV} < m_\sigma < m_{g_A}/2$, axigluons in mass range 150–400 GeV are excluded.

¹⁰ CHATRCHYAN 13A search for new resonance decaying to dijets in pp collisions at $\sqrt{s} = 7$ TeV.

¹¹ CHATRCHYAN 13AS search for new resonance decaying to dijets in pp collisions at $\sqrt{s} = 8$ TeV.

¹² CHATRCHYAN 13AU search for the pair produced color-octet vector bosons decaying to $q\bar{q}$ pairs in pp collisions. The quoted limit is for $B(g_A \rightarrow q\bar{q}) = 1$.

¹³ ABAZOV 12R search for massive color octet vector particle decaying to $t\bar{t}$. The quoted limit assumes g_A couplings with light quarks are suppressed by 0.2.

- 14 CHATRCHYAN 11Y search for new resonance decaying to dijets in pp collisions at $\sqrt{s} = 7$ TeV.
- 15 AALTONEN 10L search for massive color octet non-chiral vector particle decaying into $t\bar{t}$ pair with mass in the range $400 \text{ GeV} < M < 800 \text{ GeV}$. See their Fig. 6 for limit in the mass-coupling plane.
- 16 KHACHATRYAN 10 search for new resonance decaying to dijets in pp collisions at $\sqrt{s} = 7$ TeV.
- 17 AALTONEN 09AC search for new narrow resonance decaying to dijets.
- 18 CHOUDHURY 07 limit is from the $t\bar{t}$ production cross section measured at CDF.
- 19 DONCHESKI 98 compare α_s derived from low-energy data and that from $\Gamma(Z \rightarrow \text{hadrons})/\Gamma(Z \rightarrow \text{leptons})$.
- 20 ABE 97G search for new particle decaying to dijets.
- 21 ABE 95N assume axigluons decaying to quarks in the Standard Model only.
- 22 ABE 93G assume $\Gamma(g_A) = N\alpha_s m_{g_A}/6$ with $N = 10$.
- 23 CUYPERS 91 compare α_s measured in Υ decay and that from R at PEP/PETRA energies.
- 24 ABE 90H assumes $\Gamma(g_A) = N\alpha_s m_{g_A}/6$ with $N = 5$ ($\Gamma(g_A) = 0.09m_{g_A}$). For $N = 10$, the excluded region is reduced to 120–150 GeV.
- 25 ROBINETT 89 result demands partial-wave unitarity of $J = 0$ $t\bar{t} \rightarrow t\bar{t}$ scattering amplitude and derives a limit $m_{g_A} > 0.5 m_t$. Assumes $m_t > 56$ GeV.
- 26 ALBAJAR 88B result is from the nonobservation of a peak in two-jet invariant mass distribution. $\Gamma(g_A) < 0.4 m_{g_A}$ assumed. See also BAGGER 88.
- 27 CUYPERS 88 requires $\Gamma(\Upsilon \rightarrow g g_A) < \Gamma(\Upsilon \rightarrow g g g)$. A similar result is obtained by DONCHESKI 88.
- 28 DONCHESKI 88B requires $\Gamma(\Upsilon \rightarrow g q\bar{q})/\Gamma(\Upsilon \rightarrow g g g) < 0.25$, where the former decay proceeds via axigluon exchange. A more conservative estimate of < 0.5 leads to $m_{g_A} > 21$ GeV.

MASS LIMITS for Color-Octet Scalar Bosons

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
none 150–287	95	¹ KHACHATRY...15AV	CMS	$pp \rightarrow \Theta^0 \Theta^0 \rightarrow b\bar{b}Zg$
		² AAD	13K ATLS	$pp \rightarrow S_8 S_8 X, S_8 \rightarrow 2 \text{ jets}$
¹ KHACHATRYAN 15AV search for pair productions of neutral color-octet weak-triplet scalar particles (Θ^0), decaying to $b\bar{b}$, Zg or γg , in pp collisions at $\sqrt{s} = 8$ TeV. The Θ^0 particle is often predicted in coloron (G' , color-octet gauge boson) models and appear in the pp collisions through $G' \rightarrow \Theta^0 \Theta^0$ decays. Assuming $B(\Theta^0 \rightarrow b\bar{b}) = 0.5$, they give limits $m_{\Theta^0} > 623$ GeV (426 GeV) for $m_{G'} = 2.3 m_{\Theta^0}$ ($m_{G'} = 5 m_{\Theta^0}$).				
² AAD 13K search for pair production of color-octet scalar particles in pp collisions at $\sqrt{s} = 7$ TeV. Cross section limits are interpreted as mass limits on scalar partners of a Dirac gluino.				

X^0 (Heavy Boson) Searches in Z Decays

Searches for radiative transition of Z to a lighter spin-0 state X^0 decaying to hadrons, a lepton pair, a photon pair, or invisible particles as shown in the comments. The limits are for the product of branching ratios.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
		1 BARATE	98U ALEP	$X^0 \rightarrow \ell\bar{\ell}, q\bar{q}, gg, \gamma\gamma, \nu\bar{\nu}$
		2 ACCIARRI	97Q L3	$X^0 \rightarrow$ invisible particle(s)
		3 ACTON	93E OPAL	$X^0 \rightarrow \gamma\gamma$
		4 ABREU	92D DLPH	$X^0 \rightarrow$ hadrons
		5 ADRIANI	92F L3	$X^0 \rightarrow$ hadrons
		6 ACTON	91 OPAL	$X^0 \rightarrow$ anything
$<1.1 \times 10^{-4}$	95	7 ACTON	91B OPAL	$X^0 \rightarrow e^+e^-$
$<9 \times 10^{-5}$	95	7 ACTON	91B OPAL	$X^0 \rightarrow \mu^+\mu^-$
$<1.1 \times 10^{-4}$	95	7 ACTON	91B OPAL	$X^0 \rightarrow \tau^+\tau^-$
$<2.8 \times 10^{-4}$	95	8 ADEVA	91D L3	$X^0 \rightarrow e^+e^-$
$<2.3 \times 10^{-4}$	95	8 ADEVA	91D L3	$X^0 \rightarrow \mu^+\mu^-$
$<4.7 \times 10^{-4}$	95	9 ADEVA	91D L3	$X^0 \rightarrow$ hadrons
$<8 \times 10^{-4}$	95	10 AKRAWY	90J OPAL	$X^0 \rightarrow$ hadrons

¹ BARATE 98U obtain limits on $B(Z \rightarrow \gamma X^0)B(X^0 \rightarrow \ell\bar{\ell}, q\bar{q}, gg, \gamma\gamma, \nu\bar{\nu})$. See their Fig. 17.

² See Fig. 4 of ACCIARRI 97Q for the upper limit on $B(Z \rightarrow \gamma X^0; E_\gamma > E_{\min})$ as a function of E_{\min} .

³ ACTON 93E give $\sigma(e^+e^- \rightarrow X^0\gamma) \cdot B(X^0 \rightarrow \gamma\gamma) < 0.4$ pb (95%CL) for $m_{X^0} = 60 \pm 2.5$ GeV. If the process occurs via s -channel γ exchange, the limit translates to $\Gamma(X^0) \cdot B(X^0 \rightarrow \gamma\gamma)^2 < 20$ MeV for $m_{X^0} = 60 \pm 1$ GeV.

⁴ ABREU 92D give $\sigma_Z \cdot B(Z \rightarrow \gamma X^0) \cdot B(X^0 \rightarrow \text{hadrons}) < (3-10)$ pb for $m_{X^0} = 10-78$ GeV. A very similar limit is obtained for spin-1 X^0 .

⁵ ADRIANI 92F search for isolated γ in hadronic Z decays. The limit $\sigma_Z \cdot B(Z \rightarrow \gamma X^0) \cdot B(X^0 \rightarrow \text{hadrons}) < (2-10)$ pb (95%CL) is given for $m_{X^0} = 25-85$ GeV.

⁶ ACTON 91 searches for $Z \rightarrow Z^* X^0$, $Z^* \rightarrow e^+e^-, \mu^+\mu^-,$ or $\nu\bar{\nu}$. Excludes any new scalar X^0 with $m_{X^0} < 9.5$ GeV/ c if it has the same coupling to $Z Z^*$ as the MSM Higgs boson.

⁷ ACTON 91B limits are for $m_{X^0} = 60-85$ GeV.

⁸ ADEVA 91D limits are for $m_{X^0} = 30-89$ GeV.

⁹ ADEVA 91D limits are for $m_{X^0} = 30-86$ GeV.

¹⁰ AKRAWY 90J give $\Gamma(Z \rightarrow \gamma X^0) \cdot B(X^0 \rightarrow \text{hadrons}) < 1.9$ MeV (95%CL) for $m_{X^0} = 32-80$ GeV. We divide by $\Gamma(Z) = 2.5$ GeV to get product of branching ratios. For nonresonant transitions, the limit is $B(Z \rightarrow \gamma q\bar{q}) < 8.2$ MeV assuming three-body phase space distribution.

MASS LIMITS for a Heavy Neutral Boson Coupling to e^+e^-

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
none 55–61		¹ ODAKA	89 VNS	$\Gamma(X^0 \rightarrow e^+e^-)$. $B(X^0 \rightarrow \text{had.}) \gtrsim 0.2 \text{ MeV}$
>45	95	² DERRICK	86 HRS	$\Gamma(X^0 \rightarrow e^+e^-)=6 \text{ MeV}$
>46.6	95	³ ADEVA	85 MRKJ	$\Gamma(X^0 \rightarrow e^+e^-)=10 \text{ keV}$
>48	95	³ ADEVA	85 MRKJ	$\Gamma(X^0 \rightarrow e^+e^-)=4 \text{ MeV}$
		⁴ BERGER	85B PLUT	
none 39.8–45.5		⁵ ADEVA	84 MRKJ	$\Gamma(X^0 \rightarrow e^+e^-)=10 \text{ keV}$
>47.8	95	⁵ ADEVA	84 MRKJ	$\Gamma(X^0 \rightarrow e^+e^-)=4 \text{ MeV}$
none 39.8–45.2		⁵ BEHREND	84C CELL	
>47	95	⁵ BEHREND	84C CELL	$\Gamma(X^0 \rightarrow e^+e^-)=4 \text{ MeV}$

¹ ODAKA 89 looked for a narrow or wide scalar resonance in $e^+e^- \rightarrow \text{hadrons}$ at $E_{\text{cm}} = 55.0\text{--}60.8 \text{ GeV}$.

² DERRICK 86 found no deviation from the Standard Model Bhabha scattering at $E_{\text{cm}} = 29 \text{ GeV}$ and set limits on the possible scalar boson e^+e^- coupling. See their figure 4 for excluded region in the $\Gamma(X^0 \rightarrow e^+e^-)-m_{X^0}$ plane. Electronic chiral invariance requires a parity doublet of X^0 , in which case the limit applies for $\Gamma(X^0 \rightarrow e^+e^-) = 3 \text{ MeV}$.

³ ADEVA 85 first limit is from $2\gamma, \mu^+\mu^-$, hadrons assuming X^0 is a scalar. Second limit is from e^+e^- channel. $E_{\text{cm}} = 40\text{--}47 \text{ GeV}$. Supersedes ADEVA 84.

⁴ BERGER 85B looked for effect of spin-0 boson exchange in $e^+e^- \rightarrow e^+e^-$ and $\mu^+\mu^-$ at $E_{\text{cm}} = 34.7 \text{ GeV}$. See Fig. 5 for excluded region in the $m_{X^0} - \Gamma(X^0)$ plane.

⁵ ADEVA 84 and BEHREND 84C have $E_{\text{cm}} = 39.8\text{--}45.5 \text{ GeV}$. MARK-J searched X^0 in $e^+e^- \rightarrow \text{hadrons}, 2\gamma, \mu^+\mu^-, e^+e^-$ and CELLO in the same channels plus τ pair. No narrow or broad X^0 is found in the energy range. They also searched for the effect of X^0 with $m_{X^0} > E_{\text{cm}}$. The second limits are from Bhabha data and for spin-0 singlet. The same limits apply for $\Gamma(X^0 \rightarrow e^+e^-) = 2 \text{ MeV}$ if X^0 is a spin-0 doublet. The second limit of BEHREND 84C was read off from their figure 2. The original papers also list limits in other channels.

Search for X^0 Resonance in e^+e^- Collisions

The limit is for $\Gamma(X^0 \rightarrow e^+e^-) \cdot B(X^0 \rightarrow f)$, where f is the specified final state.

Spin 0 is assumed for X^0 .

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<10^3$	95	¹ ABE	93C VNS	$\Gamma(ee)$
$<(0.4\text{--}10)$	95	² ABE	93C VNS	$f = \gamma\gamma$
$<(0.3\text{--}5)$	95	^{3,4} ABE	93D TOPZ	$f = \gamma\gamma$
$<(2\text{--}12)$	95	^{3,4} ABE	93D TOPZ	$f = \text{hadrons}$
$<(4\text{--}200)$	95	^{4,5} ABE	93D TOPZ	$f = ee$
$<(0.1\text{--}6)$	95	^{4,5} ABE	93D TOPZ	$f = \mu\mu$
$<(0.5\text{--}8)$	90	⁶ STERNER	93 AMY	$f = \gamma\gamma$

- ¹ Limit is for $\Gamma(X^0 \rightarrow e^+e^-)$ $m_{X^0} = 56\text{--}63.5$ GeV for $\Gamma(X^0) = 0.5$ GeV.
² Limit is for $m_{X^0} = 56\text{--}61.5$ GeV and is valid for $\Gamma(X^0) \ll 100$ MeV. See their Fig. 5 for limits for $\Gamma = 1, 2$ GeV.
³ Limit is for $m_{X^0} = 57.2\text{--}60$ GeV.
⁴ Limit is valid for $\Gamma(X^0) \ll 100$ MeV. See paper for limits for $\Gamma = 1$ GeV and those for $J = 2$ resonances.
⁵ Limit is for $m_{X^0} = 56.6\text{--}60$ GeV.
⁶ STERNER 93 limit is for $m_{X^0} = 57\text{--}59.6$ GeV and is valid for $\Gamma(X^0) < 100$ MeV. See their Fig. 2 for limits for $\Gamma = 1, 3$ GeV.

Search for X^0 Resonance in $e p$ Collisions

VALUE	DOCUMENT ID	TECN	COMMENT
• • •	We do not use the following data for averages, fits, limits, etc. • • •		
	¹ CHEKANOV 02B	ZEUS	$X \rightarrow jj$
¹ CHEKANOV 02B search for photoproduction of X decaying into dijets in $e p$ collisions. See their Fig. 5 for the limit on the photoproduction cross section.			

Search for X^0 Resonance in $e^+e^- \rightarrow X^0\gamma$

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
• • •	We do not use the following data for averages, fits, limits, etc. • • •		
	¹ ABBIENDI 03D	OPAL	$X^0 \rightarrow \gamma\gamma$
	² ABREU 00Z	DLPH	X^0 decaying invisibly
	³ ADAM 96C	DLPH	X^0 decaying invisibly
¹ ABBIENDI 03D measure the $e^+e^- \rightarrow \gamma\gamma\gamma$ cross section at $\sqrt{s}=181\text{--}209$ GeV. The upper bound on the production cross section, $\sigma(e^+e^- \rightarrow X^0\gamma)$ times the branching ratio for $X^0 \rightarrow \gamma\gamma$, is less than 0.03 pb at 95%CL for X^0 masses between 20 and 180 GeV. See their Fig. 9b for the limits in the mass-cross section plane.			
² ABREU 00Z is from the single photon cross section at $\sqrt{s}=183, 189$ GeV. The production cross section upper limit is less than 0.3 pb for X^0 mass between 40 and 160 GeV. See their Fig. 4 for the limit in mass-cross section plane.			
³ ADAM 96C is from the single photon production cross at $\sqrt{s}=130, 136$ GeV. The upper bound is less than 3 pb for X^0 masses between 60 and 130 GeV. See their Fig. 5 for the exact bound on the cross section $\sigma(e^+e^- \rightarrow \gamma X^0)$.			

Search for X^0 Resonance in $Z \rightarrow f\bar{f}X^0$

The limit is for $B(Z \rightarrow f\bar{f}X^0) \cdot B(X^0 \rightarrow F)$ where f is a fermion and F is the specified final state. Spin 0 is assumed for X^0 .

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • •	We do not use the following data for averages, fits, limits, etc. • • •			
$< 3.7 \times 10^{-6}$	95	¹ ABREU 96T	DLPH	$f=e,\mu,\tau; F=\gamma\gamma$
		² ABREU 96T	DLPH	$f=\nu; F=\gamma\gamma$
		³ ABREU 96T	DLPH	$f=q; F=\gamma\gamma$
$< 6.8 \times 10^{-6}$	95	² ACTON 93E	OPAL	$f=e,\mu,\tau; F=\gamma\gamma$
$< 5.5 \times 10^{-6}$	95	² ACTON 93E	OPAL	$f=q; F=\gamma\gamma$
$< 3.1 \times 10^{-6}$	95	² ACTON 93E	OPAL	$f=\nu; F=\gamma\gamma$
$< 6.5 \times 10^{-6}$	95	² ACTON 93E	OPAL	$f=e,\mu; F=\ell\bar{\ell}, q\bar{q}, \nu\bar{\nu}$
$< 7.1 \times 10^{-6}$	95	² BUSKULIC 93F	ALEP	$f=e,\mu; F=\ell\bar{\ell}, q\bar{q}, \nu\bar{\nu}$
		⁴ ADRIANI 92F	L3	$f=q; F=\gamma\gamma$

- ¹ ABREU 96T obtain limit as a function of m_{X^0} . See their Fig. 6.
- ² Limit is for m_{X^0} around 60 GeV.
- ³ ABREU 96T obtain limit as a function of m_{X^0} . See their Fig. 15.
- ⁴ ADRIANI 92F give $\sigma_Z \cdot B(Z \rightarrow q\bar{q}X^0) \cdot B(X^0 \rightarrow \gamma\gamma) < (0.75-1.5)$ pb (95%CL) for $m_{X^0} = 10-70$ GeV. The limit is 1 pb at 60 GeV.

Search for X^0 Resonance in $W X^0$ final state

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
	¹ AALTONEN 13AA	CDF	$X^0 \rightarrow jj$
	² CHATRCHYAN 12BR	CMS	$X^0 \rightarrow jj$
	³ ABAZOV 11I	D0	$X^0 \rightarrow jj$
	⁴ ABE 97W	CDF	$X^0 \rightarrow b\bar{b}$
¹ AALTONEN 13AA search for X^0 production associated with W (or Z) in $p\bar{p}$ collisions at $E_{cm} = 1.96$ TeV. The upper limit on the cross section $\sigma(p\bar{p} \rightarrow W X^0)$ is 2.2 pb for $M_{X^0} = 145$ GeV.			
² CHATRCHYAN 12BR search for X^0 production associated with W in pp collisions at $E_{cm} = 7$ TeV. The upper limit on the cross section is 5.0 pb at 95% CL for $m_{X^0} = 150$ GeV.			
³ ABAZOV 11I search for X^0 production associated with W in $p\bar{p}$ collisions at $E_{cm} = 1.96$ TeV. The 95% CL upper limit on the cross section ranges from 2.57 to 1.28 pb for X^0 mass between 110 and 170 GeV.			
⁴ ABE 97W search for X^0 production associated with W in $p\bar{p}$ collisions at $E_{cm} = 1.8$ TeV. The 95%CL upper limit on the production cross section times the branching ratio for $X^0 \rightarrow b\bar{b}$ ranges from 14 to 19 pb for X^0 mass between 70 and 120 GeV. See their Fig. 3 for upper limits of the production cross section as a function of m_{X^0} .			

Search for X^0 Resonance in Quarkonium Decays

Limits are for branching ratios to modes shown. Spin 1 is assumed for X^0 .

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$< 3 \times 10^{-5} - 6 \times 10^{-3}$	90	¹ BALEST 95	CLE2	$\Upsilon(1S) \rightarrow X^0 \bar{X}^0 \gamma$, $m_{X^0} < 3.9$ GeV
¹ BALEST 95 three-body limit is for phase-space photon energy distribution and angular distribution same as for $\Upsilon \rightarrow gg\gamma$.				

REFERENCES FOR Searches for New Heavy Bosons (W' , Z' , leptoquarks, etc.)

AABOUD 18B	EPJ C78 24	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD 18F	PL B777 91	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD 18G	JHEP 1801 055	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
SIRUNYAN 18	PL B777 39	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN 18G	JHEP 1801 097	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AABOUD 17AK	PR D96 052004	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD 17AO	PL B774 494	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD 17AT	JHEP 1710 182	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD 17B	PL B765 32	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
KHACHATRYAN... 17AX	PL B773 563	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRYAN... 17H	JHEP 1702 048	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRYAN... 17J	JHEP 1703 077	V. Khachatryan <i>et al.</i>	(CMS Collab.)

KHACHATRY...	17T	PL B768 57	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	17U	PL B768 137	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	17W	PL B769 520	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	17Y	PL B770 257	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	17Z	PL B770 278	V. Khachatryan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	17A	JHEP 1703 162	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	17AK	PL B774 533	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	17AP	JHEP 1710 180	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	17H	JHEP 1707 121	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	17I	JHEP 1708 029	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	17Q	JHEP 1707 001	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	17R	EPJ C77 636	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	17T	PRL 119 111802	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	17V	JHEP 1709 053	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AABOUD	16	PL B759 229	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	16AA	EPJ C76 585	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	16AE	JHEP 1609 173	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	16P	EPJ C76 541	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	16U	PL B761 372	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	16V	PL B762 334	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AAD	16G	EPJ C76 5	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	16L	EPJ C76 210	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	16R	PL B755 285	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	16S	PL B754 302	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	16W	PL B758 249	G. Aad <i>et al.</i>	(ATLAS Collab.)
BARRANCO	16	JP G43 115004	J. Barranco <i>et al.</i>	
DEY	16	JHEP 1604 187	U.K. Dey, S. Mohanty	
KHACHATRY...	16AF	PR D93 032004	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	16AG	PR D93 032005	V. Khachatryan <i>et al.</i>	(CMS Collab.)
Also		PR D95 039906 (errat.)	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	16AO	JHEP 1602 122	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	16AP	JHEP 1602 145	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	16BD	EPJ C76 237	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	16BE	EPJ C76 317	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	16E	PR D93 012001	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	16K	PRL 116 071801	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	16L	PRL 117 031802	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	16O	PL B755 196	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KUMAR	16	PR D94 014022	G. Kumar	
AAD	15AM	JHEP 1507 157	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15AO	JHEP 1508 148	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15AT	EPJ C75 79	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15AU	EPJ C75 69	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15AV	EPJ C75 165	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15AZ	EPJ C75 209	G. Aad <i>et al.</i>	(ATLAS Collab.)
Also		EPJ C75 370 (errat.)	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15BB	EPJ C75 263	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15CD	PR D92 092001	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15CP	JHEP 1512 055	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15O	PRL 115 031801	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15R	PL B743 235	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15V	PR D91 052007	G. Aad <i>et al.</i>	(ATLAS Collab.)
AALTONEN	15C	PRL 115 061801	T. Aaltonen <i>et al.</i>	(CDF Collab.)
BESSAA	15	EPJ C75 97	A. Bessaa, S. Davidson	
KHACHATRY...	15AE	JHEP 1504 025	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	15AJ	JHEP 1507 042	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	15AV	JHEP 1509 201	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	15C	PL B740 83	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	15F	PRL 114 101801	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	15O	PL B748 255	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	15T	PR D91 092005	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	15V	PR D91 052009	V. Khachatryan <i>et al.</i>	(CMS Collab.)
SAHOO	15A	PR D91 094019	S. Sahoo, R. Mohanta	
AAD	14AI	JHEP 1409 037	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	14AT	PL B738 428	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	14S	PL B737 223	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	14V	PR D90 052005	G. Aad <i>et al.</i>	(ATLAS Collab.)
KHACHATRY...	14	JHEP 1408 173	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	14A	JHEP 1408 174	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	14O	EPJ C74 3149	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	14T	PL B739 229	V. Khachatryan <i>et al.</i>	(CMS Collab.)

MARTINEZ	14	PR D90 015028	R. Martinez, F. Ochoa	
PRIEELS	14	PR D90 112003	R. Prieels <i>et al.</i>	(LOUV, ETH, PSI+)
AAD	13AE	JHEP 1306 033	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	13AO	PR D87 112006	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	13AQ	PR D88 012004	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	13D	JHEP 1301 029	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	13G	JHEP 1301 116	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	13K	EPJ C73 2263	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	13S	PL B719 242	G. Aad <i>et al.</i>	(ATLAS Collab.)
AALTONEN	13A	PRL 110 121802	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	13AA	PR D88 092004	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	13R	PRL 111 031802	T. Aaltonen <i>et al.</i>	(CDF Collab.)
CHATRCHYAN	13A	JHEP 1301 013	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	13AF	PL B720 63	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	13AJ	PL B723 280	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	13AP	PR D87 072002	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	13AQ	PR D87 072005	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	13AS	PR D87 114015	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	13AU	PRL 110 141802	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	13BM	PRL 111 211804	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
	Also	PRL 112 119903 (errat.)	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	13E	PL B718 1229	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	13M	PRL 110 081801	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	13U	JHEP 1302 036	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
SAKAKI	13	PR D88 094012	Y. Sakaki <i>et al.</i>	
AAD	12AV	PRL 109 081801	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	12BB	PR D85 112012	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	12BV	JHEP 1209 041	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	12CC	JHEP 1211 138	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	12CK	PR D86 091103	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	12CR	EPJ C72 2241	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	12H	PL B709 158	G. Aad <i>et al.</i>	(ATLAS Collab.)
	Also	PL B711 442 (errat.)	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	12K	EPJ C72 2083	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	12M	EPJ C72 2056	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	12O	EPJ C72 2151	G. Aad <i>et al.</i>	(ATLAS Collab.)
AALTONEN	12AR	PR D86 112002	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	12N	PRL 108 211805	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	12R	PR D85 051101	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABRAMOWICZ	12A	PR D86 012005	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)
CHATRCHYAN	12AF	PRL 109 141801	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	12AG	PR D86 052013	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	12AI	JHEP 1208 110	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	12AQ	JHEP 1209 029	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
	Also	JHEP 1403 132 (errat.)	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	12AR	PL B717 351	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	12BG	PRL 109 261802	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	12BL	JHEP 1212 015	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	12BO	JHEP 1212 055	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	12BR	PRL 109 251801	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	12M	PL B714 158	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	12O	PL B716 82	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
KOSNIK	12	PR D86 055004	N. Kosnik	(LALO, STFN)
AAD	11D	PR D83 112006	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	11H	PRL 106 251801	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	11Z	EPJ C71 1809	G. Aad <i>et al.</i>	(ATLAS Collab.)
AALTONEN	11AD	PR D84 072003	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	11AE	PR D84 072004	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	11C	PR D83 031102	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	11I	PRL 106 121801	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AARON	11A	PL B701 20	F. D. Aaron <i>et al.</i>	(H1 Collab.)
AARON	11B	PL B704 388	F. D. Aaron <i>et al.</i>	(H1 Collab.)
AARON	11C	PL B705 52	F. D. Aaron <i>et al.</i>	(H1 Collab.)
ABAZOV	11A	PL B695 88	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	11H	PRL 107 011801	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	11I	PRL 107 011804	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	11L	PL B699 145	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	11V	PR D84 071104	V.M. Abazov <i>et al.</i>	(D0 Collab.)
BUENO	11	PR D84 032005	J.F. Bueno <i>et al.</i>	(TWIST Collab.)
	Also	PR D85 039908 (errat.)	J.F. Bueno <i>et al.</i>	(TWIST Collab.)
CHATRCHYAN	11N	PL B703 246	S. Chatrchyan <i>et al.</i>	(CMS Collab.)

CHATRCHYAN	11O	JHEP 1108 005	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	11Y	PL B704 123	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
DORSNER	11	JHEP 1111 002	I. Dorsner <i>et al.</i>	
KHACHATRY...	11D	PRL 106 201802	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	11E	PRL 106 201803	V. Khachatryan <i>et al.</i>	(CMS Collab.)
AALTONEN	10L	PL B691 183	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	10N	PRL 104 241801	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	10L	PL B693 95	V.M. Abazov <i>et al.</i>	(D0 Collab.)
DEL-AGUILA	10	JHEP 1009 033	F. del Aguila, J. de Blas, M. Perez-Victoria	(GRAN)
KHACHATRY...	10	PRL 105 211801	V. Khachatryan <i>et al.</i>	(CMS Collab.)
Also		PRL 106 029902	V. Khachatryan <i>et al.</i>	(CMS Collab.)
WAUTERS	10	PR C82 055502	F. Wauters <i>et al.</i>	(REZ, TAMU)
AALTONEN	09AC	PR D79 112002	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	09T	PRL 102 031801	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	09V	PRL 102 091805	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	09	PL B671 224	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	09AF	PL B681 224	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ERLER	09	JHEP 0908 017	J. Erler <i>et al.</i>	
AALTONEN	08D	PR D77 051102	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	08P	PR D77 091105	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	08Y	PRL 100 231801	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	08Z	PRL 101 071802	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	08AA	PL B668 98	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	08AD	PL B668 357	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	08AN	PRL 101 241802	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	08C	PRL 100 031804	V.M. Abazov <i>et al.</i>	(D0 Collab.)
MACDONALD	08	PR D78 032010	R.P. MacDonald <i>et al.</i>	(TWIST Collab.)
ZHANG	08	NP B802 247	Y. Zhang <i>et al.</i>	(PKGU, UMD)
AALTONEN	07H	PRL 99 171802	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	07E	PL B647 74	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	07J	PRL 99 061801	V.M. Abazov <i>et al.</i>	(D0 Collab.)
AKTAS	07A	EPJ C52 833	A. Aktas <i>et al.</i>	(H1 Collab.)
CHOUDHURY	07	PL B657 69	D. Choudhury <i>et al.</i>	
MELCONIAN	07	PL B649 370	D. Melconian <i>et al.</i>	(TRIUMF)
SCHAEEL	07A	EPJ C49 411	S. Schaeel <i>et al.</i>	(ALEPH Collab.)
SCHUMANN	07	PRL 99 191803	M. Schumann <i>et al.</i>	(HEID, ILLG, KARL+)
SMIRNOV	07	MPL A22 2353	A.D. Smirnov	
ABAZOV	06A	PL B636 183	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	06L	PL B640 230	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABDALLAH	06C	EPJ C45 589	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ABULENCIA	06L	PRL 96 211801	A. Abulencia <i>et al.</i>	(CDF Collab.)
ABULENCIA	06M	PRL 96 211802	A. Abulencia <i>et al.</i>	(CDF Collab.)
ABULENCIA	06T	PR D73 051102	A. Abulencia <i>et al.</i>	(CDF Collab.)
ABAZOV	05H	PR D71 071104	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABULENCIA	05A	PRL 95 252001	A. Abulencia <i>et al.</i>	(CDF Collab.)
ACOSTA	05I	PR D71 112001	D. Acosta <i>et al.</i>	(CDF Collab.)
ACOSTA	05P	PR D72 051107	D. Acosta <i>et al.</i>	(CDF Collab.)
ACOSTA	05R	PRL 95 131801	D. Acosta <i>et al.</i>	(CDF Collab.)
AKTAS	05B	PL B629 9	A. Aktas <i>et al.</i>	(H1 Collab.)
CHEKANOV	05	PL B610 212	S. Chekanov <i>et al.</i>	(HERA ZEUS Collab.)
CHEKANOV	05A	EPJ C44 463	S. Chekanov <i>et al.</i>	(ZEUS Collab.)
CYBURT	05	ASP 23 313	R.H. Cyburt <i>et al.</i>	
ABAZOV	04A	PRL 92 221801	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	04C	PR D69 111101	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABBIENDI	04G	EPJ C33 173	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	03D	EPJ C26 331	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	03R	EPJ C31 281	G. Abbiendi <i>et al.</i>	(OPAL)
ACOSTA	03B	PRL 90 081802	D. Acosta <i>et al.</i>	(CDF Collab.)
ADLOFF	03	PL B568 35	C. Adloff <i>et al.</i>	(H1 Collab.)
BARGER	03B	PR D67 075009	V. Barger, P. Langacker, H. Lee	
CHANG	03	PR D68 111101	M.-C. Chang <i>et al.</i>	(BELLE Collab.)
CHEKANOV	03B	PR D68 052004	S. Chekanov <i>et al.</i>	(ZEUS Collab.)
ABAZOV	02	PRL 88 191801	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABBIENDI	02B	PL B526 233	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
AFFOLDER	02C	PRL 88 071806	T. Affolder <i>et al.</i>	(CDF Collab.)
CHEKANOV	02	PR D65 092004	S. Chekanov <i>et al.</i>	(ZEUS Collab.)
CHEKANOV	02B	PL B531 9	S. Chekanov <i>et al.</i>	(ZEUS Collab.)
MUECK	02	PR D65 085037	A. Mueck, A. Pilafitsis, R. Rueckl	
ABAZOV	01B	PRL 87 061802	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	01D	PR D64 092004	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ADLOFF	01C	PL B523 234	C. Adloff <i>et al.</i>	(H1 Collab.)

AFFOLDER	01I	PRL 87 231803	T. Affolder <i>et al.</i>	(CDF Collab.)
BREITWEG	01	PR D63 052002	J. Breitweg <i>et al.</i>	(ZEUS Collab.)
CHEUNG	01B	PL B517 167	K. Cheung	
THOMAS	01	NP A694 559	E. Thomas <i>et al.</i>	
ABBIENDI	00M	EPJ C13 15	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBOTT	00C	PRL 84 2088	B. Abbott <i>et al.</i>	(D0 Collab.)
ABE	00	PRL 84 5716	F. Abe <i>et al.</i>	(CDF Collab.)
ABREU	00S	PL B485 45	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	00Z	EPJ C17 53	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	00P	PL B489 81	M. Acciarri <i>et al.</i>	(L3 Collab.)
ADLOFF	00	PL B479 358	C. Adloff <i>et al.</i>	(H1 Collab.)
AFFOLDER	00K	PRL 85 2056	T. Affolder <i>et al.</i>	(CDF Collab.)
BARATE	00I	EPJ C12 183	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARGER	00	PL B480 149	V. Barger, K. Cheung	
BREITWEG	00E	EPJ C16 253	J. Breitweg <i>et al.</i>	(ZEUS Collab.)
CHAY	00	PR D61 035002	J. Chay, K.Y. Lee, S. Nam	
CHO	00	MPL A15 311	G. Cho	
CORNET	00	PR D61 037701	F. Cornet, M. Relano, J. Rico	
DELGADO	00	JHEP 0001 030	A. Delgado, A. Pomarol, M. Quiros	
ERLER	00	PRL 84 212	J. Erler, P. Langacker	
GABRIELLI	00	PR D62 055009	E. Gabrielli	
RIZZO	00	PR D61 016007	T.G. Rizzo, J.D. Wells	
ROSNER	00	PR D61 016006	J.L. Rosner	
ZARNECKI	00	EPJ C17 695	A. Zarnecki	
ABBIENDI	99	EPJ C6 1	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBOTT	99J	PRL 83 2896	B. Abbott <i>et al.</i>	(D0 Collab.)
ABREU	99G	PL B446 62	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACKERSTAFF	99D	EPJ C8 3	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ADLOFF	99	EPJ C11 447	C. Adloff <i>et al.</i>	(H1 Collab.)
Also		EPJ C14 553 (errat.)	C. Adloff <i>et al.</i>	(H1 Collab.)
CASALBUONI	99	PL B460 135	R. Casalbuoni <i>et al.</i>	
CZAKON	99	PL B458 355	M. Czakon, J. Gluza, M. Zralek	
ERLER	99	PL B456 68	J. Erler, P. Langacker	
MARCIANO	99	PR D60 093006	W. Marciano	
MASIP	99	PR D60 096005	M. Masip, A. Pomarol	
NATH	99	PR D60 116004	P. Nath, M. Yamaguchi	
STRUMIA	99	PL B466 107	A. Strumia	
ABBOTT	98E	PRL 80 2051	B. Abbott <i>et al.</i>	(D0 Collab.)
ABBOTT	98J	PRL 81 38	B. Abbott <i>et al.</i>	(D0 Collab.)
ABE	98S	PRL 81 4806	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	98V	PRL 81 5742	F. Abe <i>et al.</i>	(CDF Collab.)
ACCIARRI	98J	PL B433 163	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACKERSTAFF	98V	EPJ C2 441	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
BARATE	98U	EPJ C4 571	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARENBOIM	98	EPJ C1 369	G. Barenboim	
CHO	98	EPJ C5 155	G. Cho, K. Hagiwara, S. Matsumoto	
CONRAD	98	RMP 70 1341	J.M. Conrad, M.H. Shaevitz, T. Bolton	
DONCHESKI	98	PR D58 097702	M.A. Doncheski, R.W. Robinett	
GROSS-PILCHER	98	hep-ex/9810015	C. Grosso-Pilcher, G. Landsberg, M. Paterno	
ABE	97F	PRL 78 2906	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	97G	PR D55 5263	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	97S	PRL 79 2192	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	97W	PRL 79 3819	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	97X	PRL 79 4327	F. Abe <i>et al.</i>	(CDF Collab.)
ACCIARRI	97Q	PL B412 201	M. Acciarri <i>et al.</i>	(L3 Collab.)
ARIMA	97	PR D55 19	T. Arima <i>et al.</i>	(VENUS Collab.)
BARENBOIM	97	PR D55 4213	G. Barenboim <i>et al.</i>	(VALE, IFIC)
DEANDREA	97	PL B409 277	A. Deandrea	(MARS)
DERRICK	97	ZPHY C73 613	M. Derrick <i>et al.</i>	(ZEUS Collab.)
GROSSMAN	97	PR D55 2768	Y. Grossman, Z. Ligeti, E. Nardi	(REHO, CIT)
JADACH	97	PL B408 281	S. Jadach, B.F.L. Ward, Z. Was	(CERN, INPK+)
STAHL	97	ZPHY C74 73	A. Stahl, H. Voss	(BONN)
ABACHI	96C	PRL 76 3271	S. Abachi <i>et al.</i>	(D0 Collab.)
ABREU	96T	ZPHY C72 179	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ADAM	96C	PL B380 471	W. Adam <i>et al.</i>	(DELPHI Collab.)
AID	96B	PL B369 173	S. Aid <i>et al.</i>	(H1 Collab.)
ALLET	96	PL B383 139	M. Allet <i>et al.</i>	(VILL, LEUV, LOUV, WISC)
ABACHI	95E	PL B358 405	S. Abachi <i>et al.</i>	(D0 Collab.)
ABE	95N	PRL 74 3538	F. Abe <i>et al.</i>	(CDF Collab.)
BALEST	95	PR D51 2053	R. Balest <i>et al.</i>	(CLEO Collab.)
KUZNETSOV	95	PRL 75 794	I.A. Kuznetsov <i>et al.</i>	(PNPI, KIAE, HARV+)

KUZNETSOV	95B	PAN 58 2113	A.V. Kuznetsov, N.V. Mikheev	(YARO)
		Translated from YAF 58 2228.		
MIZUKOSHI	95	NP B443 20	J.K. Mizukoshi, O.J.P. Eboli, M.C. Gonzalez-Garcia	
ABREU	94O	ZPHY C64 183	P. Abreu <i>et al.</i>	(DELPHI Collab.)
BHATTACH...	94	PL B336 100	G. Bhattacharyya, J. Ellis, K. Sridhar	(CERN)
Also		PL B338 522 (erratum)	G. Bhattacharyya, J. Ellis, K. Sridhar	(CERN)
BHATTACH...	94B	PL B338 522 (erratum)	G. Bhattacharyya, J. Ellis, K. Sridhar	(CERN)
DAVIDSON	94	ZPHY C61 613	S. Davidson, D. Bailey, B.A. Campbell	(CFPA+)
KUZNETSOV	94	PL B329 295	A.V. Kuznetsov, N.V. Mikheev	(YARO)
KUZNETSOV	94B	JETPL 60 315	I.A. Kuznetsov <i>et al.</i>	(PNPI, KIAE, HARV+)
		Translated from ZETFP 60 311.		
LEURER	94	PR D50 536	M. Leurer	(REHO)
LEURER	94B	PR D49 333	M. Leurer	(REHO)
Also		PRL 71 1324	M. Leurer	(REHO)
MAHANTA	94	PL B337 128	U. Mahanta	(MEHTA)
SEVERIJNS	94	PRL 73 611 (erratum)	N. Severijns <i>et al.</i>	(LOUV, WISC, LEUV+)
VILAIN	94B	PL B332 465	P. Vilain <i>et al.</i>	(CHARM II Collab.)
ABE	93C	PL B302 119	K. Abe <i>et al.</i>	(VENUS Collab.)
ABE	93D	PL B304 373	T. Abe <i>et al.</i>	(TOPAZ Collab.)
ABE	93G	PRL 71 2542	F. Abe <i>et al.</i>	(CDF Collab.)
ABREU	93J	PL B316 620	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACTON	93E	PL B311 391	P.D. Acton <i>et al.</i>	(OPAL Collab.)
ADRIANI	93M	PRPL 236 1	O. Adriani <i>et al.</i>	(L3 Collab.)
ALITTI	93	NP B400 3	J. Alitti <i>et al.</i>	(UA2 Collab.)
BHATTACH...	93	PR D47 3693	G. Bhattacharyya <i>et al.</i>	(CALC, JADA, ICTP+)
BUSKULIC	93F	PL B308 425	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
DERRICK	93	PL B306 173	M. Derrick <i>et al.</i>	(ZEUS Collab.)
RIZZO	93	PR D48 4470	T.G. Rizzo	(ANL)
SEVERIJNS	93	PRL 70 4047	N. Severijns <i>et al.</i>	(LOUV, WISC, LEUV+)
Also		PRL 73 611 (erratum)	N. Severijns <i>et al.</i>	(LOUV, WISC, LEUV+)
STERNER	93	PL B303 385	K.L. Sterner <i>et al.</i>	(AMY Collab.)
ABREU	92D	ZPHY C53 555	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ADRIANI	92F	PL B292 472	O. Adriani <i>et al.</i>	(L3 Collab.)
DECAMP	92	PRPL 216 253	D. Decamp <i>et al.</i>	(ALEPH Collab.)
IMAZATO	92	PRL 69 877	J. Imazato <i>et al.</i>	(KEK, INUS, TOKY+)
MISHRA	92	PRL 68 3499	S.R. Mishra <i>et al.</i>	(COLU, CHIC, FNAL+)
POLAK	92B	PR D46 3871	J. Polak, M. Zralek	(SILES)
ACTON	91	PL B268 122	D.P. Acton <i>et al.</i>	(OPAL Collab.)
ACTON	91B	PL B273 338	D.P. Acton <i>et al.</i>	(OPAL Collab.)
ADEVA	91D	PL B262 155	B. Adeva <i>et al.</i>	(L3 Collab.)
AQUINO	91	PL B261 280	M. Aquino, A. Fernandez, A. Garcia	(CINV, PUEB)
COLANGELO	91	PL B253 154	P. Colangelo, G. Nardulli	(BARI)
CUYPERS	91	PL B259 173	F. Cuypers, A.F. Falk, P.H. Frampton	(DURH, HARV+)
FARAGGI	91	MPL A6 61	A.E. Faraggi, D.V. Nanopoulos	(TAMU)
POLAK	91	NP B363 385	J. Polak, M. Zralek	(SILES)
RIZZO	91	PR D44 202	T.G. Rizzo	(WISC, ISU)
WALKER	91	APJ 376 51	T.P. Walker <i>et al.</i>	(HSCA, OSU, CHIC+)
ABE	90F	PL B246 297	K. Abe <i>et al.</i>	(VENUS Collab.)
ABE	90H	PR D41 1722	F. Abe <i>et al.</i>	(CDF Collab.)
AKRAWY	90J	PL B246 285	M.Z. Akrawy <i>et al.</i>	(OPAL Collab.)
GONZALEZ...	90D	PL B240 163	M.C. Gonzalez-Garcia, J.W.F. Valle	(VALE)
GRIFOLS	90	NP B331 244	J.A. Grifols, E. Masso	(BARC)
GRIFOLS	90D	PR D42 3293	J.A. Grifols, E. Masso, T.G. Rizzo	(BARC, CERN+)
KIM	90	PL B240 243	G.N. Kim <i>et al.</i>	(AMY Collab.)
LOPEZ	90	PL B241 392	J.L. Lopez, D.V. Nanopoulos	(TAMU)
BARBIERI	89B	PR D39 1229	R. Barbieri, R.N. Mohapatra	(PISA, UMD)
LANGACKER	89B	PR D40 1569	P. Langacker, S. Uma Sankar	(PENN)
ODAKA	89	JPSJ 58 3037	S. Odaka <i>et al.</i>	(VENUS Collab.)
ROBINETT	89	PR D39 834	R.W. Robinett	(PSU)
ALBAJAR	88B	PL B209 127	C. Albajar <i>et al.</i>	(UA1 Collab.)
BAGGER	88	PR D37 1188	J. Bagger, C. Schmidt, S. King	(HARV, BOST)
BALKE	88	PR D37 587	B. Balke <i>et al.</i>	(LBL, UCB, COLO, NWES+)
BERGSTROM	88	PL B212 386	L. Bergstrom	(STOH)
CUYPERS	88	PRL 60 1237	F. Cuypers, P.H. Frampton	(UNCCH)
DONCHESKI	88	PL B206 137	M.A. Doncheski, H. Grotch, R. Robinett	(PSU)
DONCHESKI	88B	PR D38 412	M.A. Doncheski, H. Grotch, R.W. Robinett	(PSU)
BARTEL	87B	ZPHY C36 15	W. Bartel <i>et al.</i>	(JADE Collab.)
BEHREND	86B	PL B178 452	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
DERRICK	86	PL 166B 463	M. Derrick <i>et al.</i>	(HRS Collab.)
Also		PR D34 3286	M. Derrick <i>et al.</i>	(HRS Collab.)

JODIDIO	86	PR D34 1967	A. Jodidio <i>et al.</i>	(LBL, NWES, TRIU)
Also		PR D37 237 (erratum)	A. Jodidio <i>et al.</i>	(LBL, NWES, TRIU)
MOHAPATRA	86	PR D34 909	R.N. Mohapatra	(UMD)
ADEVA	85	PL 152B 439	B. Adeva <i>et al.</i>	(Mark-J Collab.)
BERGER	85B	ZPHY C27 341	C. Berger <i>et al.</i>	(PLUTO Collab.)
STOKER	85	PRL 54 1887	D.P. Stoker <i>et al.</i>	(LBL, NWES, TRIU)
ADEVA	84	PRL 53 134	B. Adeva <i>et al.</i>	(Mark-J Collab.)
BEHREND	84C	PL 140B 130	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
BERGSMASMA	83	PL 122B 465	F. Bergsma <i>et al.</i>	(CHARM Collab.)
CARR	83	PRL 51 627	J. Carr <i>et al.</i>	(LBL, NWES, TRIU)
BEALL	82	PRL 48 848	G. Beall, M. Bander, A. Soni	(UCI, UCLA)
SHANKER	82	NP B204 375	O. Shanker	(TRIU)
