

$\Upsilon(2S)$ 

$$I^G(J^{PC}) = 0^-(1^{--})$$

### $\Upsilon(2S)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>10023.26±0.31 OUR AVERAGE</b>			
10023.5 ±0.5	<sup>1</sup> ARTAMONOV 00	MD1	$e^+e^- \rightarrow \text{hadrons}$
10023.1 ±0.4	BARBER 84	REDE	$e^+e^- \rightarrow \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
10023.6 ±0.5	<sup>2,3</sup> BARU	86B REDE	$e^+e^- \rightarrow \text{hadrons}$
<sup>1</sup> Reanalysis of BARU 86B using new electron mass (COHEN 87).			
<sup>2</sup> Reanalysis of ARTAMONOV 84.			
<sup>3</sup> Superseded by ARTAMONOV 00.			

### $m\Upsilon(3S) - m\Upsilon(2S)$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>331.50±0.02±0.13</b>	LEES	11C BABR	$e^+e^- \rightarrow \pi^+\pi^-X$

### $\Upsilon(2S)$ WIDTH

VALUE (keV)	DOCUMENT ID	COMMENT
<b>31.98±2.63 OUR EVALUATION</b>		See the Note on "Width Determinations of the $\Upsilon$ States"

### $\Upsilon(2S)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1$ $\Upsilon(1S)\pi^+\pi^-$	(17.85±0.26) %	
$\Gamma_2$ $\Upsilon(1S)\pi^0\pi^0$	(8.6 ± 0.4) %	
$\Gamma_3$ $\tau^+\tau^-$	(2.00±0.21) %	
$\Gamma_4$ $\mu^+\mu^-$	(1.93±0.17) %	S=2.2
$\Gamma_5$ $e^+e^-$	(1.91±0.16) %	
$\Gamma_6$ $\Upsilon(1S)\pi^0$	< 4 × 10 <sup>-5</sup>	CL=90%
$\Gamma_7$ $\Upsilon(1S)\eta$	(2.9 ± 0.4) × 10 <sup>-4</sup>	S=2.0
$\Gamma_8$ $J/\psi(1S)$ anything	< 6 × 10 <sup>-3</sup>	CL=90%
$\Gamma_9$ $J/\psi(1S)\eta_c$	< 5.4 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{10}$ $J/\psi(1S)\chi_{c0}$	< 3.4 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{11}$ $J/\psi(1S)\chi_{c1}$	< 1.2 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{12}$ $J/\psi(1S)\chi_{c2}$	< 2.0 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{13}$ $J/\psi(1S)\eta_c(2S)$	< 2.5 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{14}$ $J/\psi(1S)X(3940)$	< 2.0 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{15}$ $J/\psi(1S)X(4160)$	< 2.0 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{16}$ $\chi_{c1}$ anything	(2.2 ± 0.5) × 10 <sup>-4</sup>	

$\Gamma_{17}$	$\chi_{c1}(1P)^0 X_{tetra}$	$< 3.67 \times 10^{-5}$	CL=90%
$\Gamma_{18}$	$\chi_{c2}$ anything	$( 2.3 \pm 0.8 ) \times 10^{-4}$	
$\Gamma_{19}$	$\psi(2S)\eta_c$	$< 5.1 \times 10^{-6}$	CL=90%
$\Gamma_{20}$	$\psi(2S)\chi_{c0}$	$< 4.7 \times 10^{-6}$	CL=90%
$\Gamma_{21}$	$\psi(2S)\chi_{c1}$	$< 2.5 \times 10^{-6}$	CL=90%
$\Gamma_{22}$	$\psi(2S)\chi_{c2}$	$< 1.9 \times 10^{-6}$	CL=90%
$\Gamma_{23}$	$\psi(2S)\eta_c(2S)$	$< 3.3 \times 10^{-6}$	CL=90%
$\Gamma_{24}$	$\psi(2S)X(3940)$	$< 3.9 \times 10^{-6}$	CL=90%
$\Gamma_{25}$	$\psi(2S)X(4160)$	$< 3.9 \times 10^{-6}$	CL=90%
$\Gamma_{26}$	$\overline{2H}$ anything	$( 2.78_{-0.26}^{+0.30} ) \times 10^{-5}$	S=1.2
$\Gamma_{27}$	hadrons	$(94 \pm 11) \%$	
$\Gamma_{28}$	$ggg$	$(58.8 \pm 1.2) \%$	
$\Gamma_{29}$	$\gamma gg$	$( 1.87 \pm 0.28 ) \%$	
$\Gamma_{30}$	$\phi K^+ K^-$	$( 1.6 \pm 0.4 ) \times 10^{-6}$	
$\Gamma_{31}$	$\omega \pi^+ \pi^-$	$< 2.58 \times 10^{-6}$	CL=90%
$\Gamma_{32}$	$K^*(892)^0 K^- \pi^+ + c.c.$	$( 2.3 \pm 0.7 ) \times 10^{-6}$	
$\Gamma_{33}$	$\phi f_2'(1525)$	$< 1.33 \times 10^{-6}$	CL=90%
$\Gamma_{34}$	$\omega f_2(1270)$	$< 5.7 \times 10^{-7}$	CL=90%
$\Gamma_{35}$	$\rho(770) a_2(1320)$	$< 8.8 \times 10^{-7}$	CL=90%
$\Gamma_{36}$	$K^*(892)^0 \overline{K}_2^*(1430)^0 + c.c.$	$( 1.5 \pm 0.6 ) \times 10^{-6}$	
$\Gamma_{37}$	$K_1(1270)^\pm K^\mp$	$< 3.22 \times 10^{-6}$	CL=90%
$\Gamma_{38}$	$K_1(1400)^\pm K^\mp$	$< 8.3 \times 10^{-7}$	CL=90%
$\Gamma_{39}$	$b_1(1235)^\pm \pi^\mp$	$< 4.0 \times 10^{-7}$	CL=90%
$\Gamma_{40}$	$\rho \pi$	$< 1.16 \times 10^{-6}$	CL=90%
$\Gamma_{41}$	$\pi^+ \pi^- \pi^0$	$< 8.0 \times 10^{-7}$	CL=90%
$\Gamma_{42}$	$\omega \pi^0$	$< 1.63 \times 10^{-6}$	CL=90%
$\Gamma_{43}$	$\pi^+ \pi^- \pi^0 \pi^0$	$( 1.30 \pm 0.28 ) \times 10^{-5}$	
$\Gamma_{44}$	$K_S^0 K^+ \pi^- + c.c.$	$( 1.14 \pm 0.33 ) \times 10^{-6}$	
$\Gamma_{45}$	$K^*(892)^0 \overline{K}^0 + c.c.$	$< 4.22 \times 10^{-6}$	CL=90%
$\Gamma_{46}$	$K^*(892)^- K^+ + c.c.$	$< 1.45 \times 10^{-6}$	CL=90%
$\Gamma_{47}$	$f_1(1285)$ anything	$( 2.2 \pm 1.6 ) \times 10^{-3}$	
$\Gamma_{48}$	$f_1(1285) X_{tetra}$	$< 6.47 \times 10^{-5}$	CL=90%
$\Gamma_{49}$	Sum of 100 exclusive modes	$( 2.90 \pm 0.30 ) \times 10^{-3}$	

### Radiative decays

$\Gamma_{50}$	$\gamma \chi_{b1}(1P)$	$( 6.9 \pm 0.4 ) \%$	
$\Gamma_{51}$	$\gamma \chi_{b2}(1P)$	$( 7.15 \pm 0.35 ) \%$	
$\Gamma_{52}$	$\gamma \chi_{b0}(1P)$	$( 3.8 \pm 0.4 ) \%$	
$\Gamma_{53}$	$\gamma f_0(1710)$	$< 5.9 \times 10^{-4}$	CL=90%
$\Gamma_{54}$	$\gamma f_2'(1525)$	$< 5.3 \times 10^{-4}$	CL=90%
$\Gamma_{55}$	$\gamma f_2(1270)$	$< 2.41 \times 10^{-4}$	CL=90%
$\Gamma_{56}$	$\gamma f_J(2220)$		
$\Gamma_{57}$	$\gamma \eta_c(1S)$	$< 2.7 \times 10^{-5}$	CL=90%
$\Gamma_{58}$	$\gamma \chi_{c0}$	$< 1.0 \times 10^{-4}$	CL=90%



$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{27}\Gamma_5/\Gamma$

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.577±0.009 OUR AVERAGE</b>			
0.581±0.004±0.009	<sup>1</sup> ROSNER	06 CLEO	10.0 e <sup>+</sup> e <sup>-</sup> → hadrons
0.552±0.031±0.017	<sup>1</sup> BARU	96 MD1	e <sup>+</sup> e <sup>-</sup> → hadrons
0.54 ±0.04 ±0.02	<sup>1</sup> JAKUBOWSKI	88 CBAL	e <sup>+</sup> e <sup>-</sup> → hadrons
0.58 ±0.03 ±0.04	<sup>2</sup> GILES	84B CLEO	e <sup>+</sup> e <sup>-</sup> → hadrons
0.60 ±0.12 ±0.07	<sup>2</sup> ALBRECHT	82 DASP	e <sup>+</sup> e <sup>-</sup> → hadrons
0.54 ±0.07 <sup>+0.09</sup> <sub>-0.05</sub>	<sup>2</sup> NICZYPORUK	81C LENA	e <sup>+</sup> e <sup>-</sup> → hadrons
0.41 ±0.18	<sup>2</sup> BOCK	80 CNTR	e <sup>+</sup> e <sup>-</sup> → hadrons

<sup>1</sup> Radiative corrections evaluated following KURAEV 85.

<sup>2</sup> Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

**$\Upsilon(2S)$  PARTIAL WIDTHS**

$\Gamma(e^+e^-)$   $\Gamma_5$

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>
<b>0.612±0.011 OUR EVALUATION</b>	

**$\Upsilon(2S)$  BRANCHING RATIOS**

$\Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_1/\Gamma$

Abbreviation MM in the COMMENT field below stands for missing mass.

<u>VALUE (units 10<sup>-2</sup>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>17.85±0.26 OUR FIT</b>				
<b>17.92±0.26 OUR AVERAGE</b>				
16.8 ±1.1 ±1.3	906k	<sup>1</sup> LEES	11C BABR	e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> X
17.80±0.05±0.37	170k	<sup>2</sup> LEES	11L BABR	$\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$
18.02±0.02±0.61	851k	<sup>3</sup> BHARI	09 CLEO	e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> MM
17.22±0.17±0.75	11.8K	<sup>4</sup> AUBERT	08BP BABR	e <sup>+</sup> e <sup>-</sup> → γπ <sup>+</sup> π <sup>-</sup> ℓ <sup>+</sup> ℓ <sup>-</sup>
19.2 ±0.2 ±1.0	52.6k	<sup>5</sup> ALEXANDER	98 CLE2	π <sup>+</sup> π <sup>-</sup> ℓ <sup>+</sup> ℓ <sup>-</sup> , π <sup>+</sup> π <sup>-</sup> MM
18.1 ±0.5 ±1.0	11.6k	ALBRECHT	87 ARG	e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> MM
16.9 ±4.0		GELPHMAN	85 CBAL	e <sup>+</sup> e <sup>-</sup> → e <sup>+</sup> e <sup>-</sup> π <sup>+</sup> π <sup>-</sup>
19.1 ±1.2 ±0.6		BESSON	84 CLEO	π <sup>+</sup> π <sup>-</sup> MM
18.9 ±2.6		FONSECA	84 CUSB	e <sup>+</sup> e <sup>-</sup> → ℓ <sup>+</sup> ℓ <sup>-</sup> π <sup>+</sup> π <sup>-</sup>
21 ±7	7	NICZYPORUK	81B LENA	e <sup>+</sup> e <sup>-</sup> → ℓ <sup>+</sup> ℓ <sup>-</sup> π <sup>+</sup> π <sup>-</sup>

<sup>1</sup> LEES 11C reports  $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \Upsilon(2S)\text{anything})] = (1.78 \pm 0.02 \pm 0.11) \times 10^{-2}$  which we divide by our best value  $B(\Upsilon(3S) \rightarrow \Upsilon(2S)\text{anything}) = (10.6 \pm 0.8) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Using  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$ .

<sup>3</sup> A weighted average of the inclusive and exclusive results.

<sup>4</sup> Using  $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$ ,  $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$  and,  $\Gamma_{ee}(\Upsilon(2S)) = 0.612 \pm 0.011$  keV.

<sup>5</sup> Using  $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.52 \pm 0.17)\%$  and  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.07)\%$ .

$\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$   $\Gamma_2/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.6 ± 0.4 OUR AVERAGE</b>				
8.43 ± 0.16 ± 0.42	38k	<sup>1</sup> BHARI 09	CLEO	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
9.2 ± 0.6 ± 0.8	275	<sup>2</sup> ALEXANDER 98	CLE2	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
9.5 ± 1.9 ± 1.9	25	ALBRECHT 87	ARG	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
8.0 ± 1.5		GELPHMAN 85	CBAL	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
10.3 ± 2.3		FONSECA 84	CUSB	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$

<sup>1</sup> Authors assume  $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$ .

<sup>2</sup> Using  $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.52 \pm 0.17)\%$  and  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.07)\%$ .

 $\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$   $\Gamma_2/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.462 ± 0.037	<sup>1</sup> BHARI 09	CLEO	$e^+e^- \rightarrow \Upsilon(2S)$

<sup>1</sup> Not independent of other values reported by BHARI 09.

 $\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.00 ± 0.21 OUR AVERAGE</b>				
2.00 ± 0.12 ± 0.18	22k	<sup>1</sup> BESSON 07	CLEO	$e^+e^- \rightarrow \Upsilon(2S) \rightarrow \tau^+\tau^-$
1.7 ± 1.5 ± 0.6		HAAS 84B	CLEO	$e^+e^- \rightarrow \tau^+\tau^-$

<sup>1</sup> BESSON 07 reports  $[\Gamma(\Upsilon(2S) \rightarrow \tau^+\tau^-)/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \mu^+\mu^-)] = 1.04 \pm 0.04 \pm 0.05$  which we multiply by our best value  $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$   $\Gamma_4/\Gamma$ 

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.0193 ± 0.0017 OUR AVERAGE</b>					Error includes scale factor of 2.2. See the ideogram below.
0.0203 ± 0.0003 ± 0.0008		120k	ADAMS 05	CLEO	$e^+e^- \rightarrow \mu^+\mu^-$
0.0122 ± 0.0028 ± 0.0019			<sup>1</sup> KOBEL 92	CBAL	$e^+e^- \rightarrow \mu^+\mu^-$
0.0138 ± 0.0025 ± 0.0015			KAARSBERG 89	CSB2	$e^+e^- \rightarrow \mu^+\mu^-$
0.009 ± 0.006 ± 0.006			<sup>2</sup> ALBRECHT 85	ARG	$e^+e^- \rightarrow \mu^+\mu^-$
0.018 ± 0.008 ± 0.005			HAAS 84B	CLEO	$e^+e^- \rightarrow \mu^+\mu^-$


• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.038      90      NICZYPORUK 81C LENA  $e^+e^- \rightarrow \mu^+\mu^-$

<sup>1</sup> Taking into account interference between the resonance and continuum.

<sup>2</sup> Re-evaluated using  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 0.026$ .

WEIGHTED AVERAGE  
 $0.0193 \pm 0.0017$  (Error scaled by 2.2)



$\Gamma(\tau^+\tau^-)/\Gamma(\mu^+\mu^-)$

$\Gamma_3/\Gamma_4$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.04 \pm 0.04 \pm 0.05</math></b>	22k	BESSON 07	CLEO	$e^+e^- \rightarrow \Upsilon(2S)$

$\Gamma(\Upsilon(1S)\pi^0)/\Gamma_{total}$

$\Gamma_6/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4	90	<sup>1</sup> TAMPONI 13	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^0$
< 18	90	<sup>2</sup> HE 08A	CLEO	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
< 110	90	ALEXANDER 98	CLE2	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
< 800	90	LURZ 87	CBAL	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$

<sup>1</sup>TAMPONI 13 reports  $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0)/\Gamma_{total}] / [B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)] < 2.3 \times 10^{-4}$  which we multiply by our best value  $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-) = 17.85 \times 10^{-2}$ .

<sup>2</sup>Authors assume  $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$ .

$\Gamma(\Upsilon(1S)\pi^0)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$

$\Gamma_6/\Gamma_1$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 2.3</b>	90	TAMPONI 13	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^0$

$\Gamma(\Upsilon(1S)\eta)/\Gamma_{\text{total}}$

$\Gamma_7/\Gamma$

VALUE (units  $10^{-4}$ ) CL% EVTS DOCUMENT ID TECN COMMENT

**2.9 ± 0.4 OUR FIT** Error includes scale factor of 2.0.  
**2.9 ± 0.4 OUR AVERAGE** Error includes scale factor of 1.9. See the ideogram below.

2.39 ± 0.31 ± 0.14	112	<sup>1</sup> LEES	11L	BABR	$\Upsilon(2S) \rightarrow \ell^+ \ell^- \eta$
2.1 <sup>+0.7</sup> <sub>-0.6</sub> ± 0.3	14	<sup>2</sup> HE	08A	CLEO	$e^+ e^- \rightarrow \ell^+ \ell^- \eta$

• • • We use the following data for averages but not for fits. • • •

3.55 ± 0.32 ± 0.05	241	<sup>3</sup> TAMPONI	13	BELL	$e^+ e^- \rightarrow \Upsilon(1S)\eta$
--------------------	-----	----------------------	----	------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 9	90	<sup>1,4</sup> AUBERT	08BP	BABR	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- \pi^0 \ell^+ \ell^-$
< 28	90	ALEXANDER	98	CLE2	$e^+ e^- \rightarrow \ell^+ \ell^- \eta$
< 50	90	ALBRECHT	87	ARG	$e^+ e^- \rightarrow \pi^+ \pi^- \ell^+ \ell^-$ MM
< 70	90	LURZ	87	CBAL	$e^+ e^- \rightarrow \ell^+ \ell^- (\gamma\gamma, 3\pi^0)$
< 100	90	BESSON	84	CLEO	$e^+ e^- \rightarrow \pi^+ \pi^- \ell^+ \ell^-$ MM
< 20	90	FONSECA	84	CUSB	$e^+ e^- \rightarrow \ell^+ \ell^- (\gamma\gamma, \pi^+ \pi^- \pi^0)$


<sup>1</sup> Using  $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.38 \pm 0.11)\%$  and  $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$ .

<sup>2</sup> Authors assume  $B(\Upsilon(1S) \rightarrow e^+ e^-) + B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 4.96\%$ .

<sup>3</sup> TAMPONI 13 reports  $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\eta)/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+ \pi^-)] = (1.99 \pm 0.14 \pm 0.11) \times 10^{-3}$  which we multiply by our best value  $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+ \pi^-) = (17.85 \pm 0.26) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>4</sup> Using  $\Gamma_{ee}(\Upsilon(2S)) = 0.612 \pm 0.011$  keV.

WEIGHTED AVERAGE  
 2.9±0.4 (Error scaled by 1.9)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our 'best' values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

$\Gamma(\Upsilon(1S)\eta)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$   $\Gamma_7/\Gamma_1$ 

VALUE (units $10^{-3}$ )	CL% EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	----------	-------------	------	---------

**1.64±0.25 OUR FIT** Error includes scale factor of 2.0.**1.99±0.14±0.11** 241 TAMPONI 13 BELL  $e^+e^- \rightarrow \Upsilon(1S)\eta$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.35±0.17±0.08 <sup>1</sup> LEES 11L BABR  $\Upsilon(2S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\mu^+\mu^-$ < 5.2 90 <sup>2</sup> AUBERT 08BP BABR  $e^+e^- \rightarrow \gamma\pi^+\pi^-(\pi^0)\ell^+\ell^-$ <sup>1</sup> Not independent of other values reported by LEES 11L.<sup>2</sup> Not independent of other values reported by AUBERT 08BP. $\Gamma(\Upsilon(1S)\pi^0)/\Gamma(\Upsilon(1S)\eta)$   $\Gamma_6/\Gamma_7$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.13 90 TAMPONI 13 BELL  $e^+e^- \rightarrow \Upsilon(1S)\pi^0$  $\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_8/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.006 90 MASCHMANN 90 CBAL  $e^+e^- \rightarrow \text{hadrons}$  $\Gamma(J/\psi(1S)\eta_c)/\Gamma_{\text{total}}$   $\Gamma_9/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<5.4 × 10<sup>-6</sup> 90 YANG 14 BELL  $e^+e^- \rightarrow J/\psi X$  $\Gamma(J/\psi(1S)\chi_{c0})/\Gamma_{\text{total}}$   $\Gamma_{10}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<3.4 × 10<sup>-6</sup> 90 YANG 14 BELL  $e^+e^- \rightarrow J/\psi X$  $\Gamma(J/\psi(1S)\chi_{c1})/\Gamma_{\text{total}}$   $\Gamma_{11}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<1.2 × 10<sup>-6</sup> 90 YANG 14 BELL  $e^+e^- \rightarrow J/\psi X$  $\Gamma(J/\psi(1S)\chi_{c2})/\Gamma_{\text{total}}$   $\Gamma_{12}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<2.0 × 10<sup>-6</sup> 90 YANG 14 BELL  $e^+e^- \rightarrow J/\psi X$  $\Gamma(J/\psi(1S)\eta_c(2S))/\Gamma_{\text{total}}$   $\Gamma_{13}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<2.5 × 10<sup>-6</sup> 90 YANG 14 BELL  $e^+e^- \rightarrow J/\psi X$  $\Gamma(J/\psi(1S)X(3940))/\Gamma_{\text{total}}$   $\Gamma_{14}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<2.0 × 10<sup>-6</sup> 90 YANG 14 BELL  $e^+e^- \rightarrow J/\psi X$  $\Gamma(J/\psi(1S)X(4160))/\Gamma_{\text{total}}$   $\Gamma_{15}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<2.0 × 10<sup>-6</sup> 90 YANG 14 BELL  $e^+e^- \rightarrow J/\psi X$



$\Gamma(\chi_{c1} \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{16}/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>2.24 \pm 0.44 \pm 0.20</math></b>	376	JIA	17	BELL $\Upsilon(2S) \rightarrow \gamma J/\psi(1S)$

 $\Gamma(\chi_{c1}(1P)^0 X_{tetra})/\Gamma_{\text{total}}$   $\Gamma_{17}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 36.7 \times 10^{-6}</math></b>	90	<sup>1</sup> JIA	17A	BELL $e^+e^- \rightarrow \text{hadrons}$

<sup>1</sup> For a tetraquark state  $X_{tetra}$ , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of  $X_{tetra}$  mass and width range from  $4.4 \times 10^{-6}$  to  $36.7 \times 10^{-6}$ .

 $\Gamma(\chi_{c2} \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{18}/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>2.28 \pm 0.73 \pm 0.34</math></b>	JIA	17	BELL $\Upsilon(2S) \rightarrow \gamma J/\psi(1S)$

 $\Gamma(\psi(2S)\eta_c)/\Gamma_{\text{total}}$   $\Gamma_{19}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 5.1 \times 10^{-6}</math></b>	90	YANG	14	BELL $e^+e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)\chi_{c0})/\Gamma_{\text{total}}$   $\Gamma_{20}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 4.7 \times 10^{-6}</math></b>	90	YANG	14	BELL $e^+e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)\chi_{c1})/\Gamma_{\text{total}}$   $\Gamma_{21}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 2.5 \times 10^{-6}</math></b>	90	YANG	14	BELL $e^+e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)\chi_{c2})/\Gamma_{\text{total}}$   $\Gamma_{22}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 1.9 \times 10^{-6}</math></b>	90	YANG	14	BELL $e^+e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)\eta_c(2S))/\Gamma_{\text{total}}$   $\Gamma_{23}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 3.3 \times 10^{-6}</math></b>	90	YANG	14	BELL $e^+e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)X(3940))/\Gamma_{\text{total}}$   $\Gamma_{24}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 3.9 \times 10^{-6}</math></b>	90	YANG	14	BELL $e^+e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)X(4160))/\Gamma_{\text{total}}$   $\Gamma_{25}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 3.9 \times 10^{-6}</math></b>	90	YANG	14	BELL $e^+e^- \rightarrow \psi(2S)X$

$\Gamma(\overline{2H} \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{26}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>2.78^{+0.30}_{-0.26}</math></b>		<b>OUR AVERAGE</b>		Error includes scale factor of 1.2.

$2.64 \pm 0.11^{+0.26}_{-0.21}$		LEES	14G	BABR	$e^+ e^- \rightarrow \overline{2H} \chi$
$3.37 \pm 0.50 \pm 0.25$	58	ASNER	07	CLEO	$e^+ e^- \rightarrow \overline{2H} \chi$

 $\Gamma(g g g)/\Gamma_{\text{total}}$   $\Gamma_{28}/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b><math>58.8 \pm 1.2</math></b>	6M	<sup>1</sup> BESSON	06A	CLEO	$\Upsilon(2S) \rightarrow \text{hadrons}$

<sup>1</sup> Calculated using the value  $\Gamma(\gamma g g)/\Gamma(g g g) = (3.18 \pm 0.04 \pm 0.22 \pm 0.41)\%$  from BESSON 06A and PDG 08 values of  $B(\pi^+ \pi^- \Upsilon(1S)) = (18.1 \pm 0.4)\%$ ,  $B(\pi^0 \pi^0 \Upsilon(1S)) = (8.6 \pm 0.4)\%$ ,  $B(\mu^+ \mu^-) = (1.93 \pm 0.17)\%$ , and  $R_{\text{hadrons}} = 3.51$ . The statistical error is negligible and the systematic error is partially correlated with that of  $\Gamma(\gamma g g)/\Gamma_{\text{total}}$  measurement of BESSON 06A.

 $\Gamma(\gamma g g)/\Gamma(g g g)$   $\Gamma_{29}/\Gamma_{28}$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b><math>3.18 \pm 0.04 \pm 0.47</math></b>	6M	BESSON	06A	CLEO	$\Upsilon(2S) \rightarrow (\gamma +) \text{hadrons}$

 $\Gamma(\phi K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{30}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b><math>1.58 \pm 0.33 \pm 0.18</math></b>	58	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow 2(K^+ K^-)$

 $\Gamma(\omega \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{31}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<b><math>&lt;2.58</math></b>	90	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

 $\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{32}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b><math>2.32 \pm 0.40 \pm 0.54</math></b>	135	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

 $\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$   $\Gamma_{33}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<b><math>&lt;1.33</math></b>	90	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow 2(K^+ K^-)$

 $\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$   $\Gamma_{34}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<b><math>&lt;0.57</math></b>	90	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

 $\Gamma(\rho(770) a_2(1320))/\Gamma_{\text{total}}$   $\Gamma_{35}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<b><math>&lt;0.88</math></b>	90	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

$\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{36}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.53 \pm 0.52 \pm 0.19</math></b>	32	SHEN	12A	BELL $\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

 $\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$   $\Gamma_{37}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt; 3.22</math></b>	90	SHEN	12A	BELL $\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

 $\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$   $\Gamma_{38}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt; 0.83</math></b>	90	SHEN	12A	BELL $\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

 $\Gamma(b_1(1235)^\pm \pi^\mp)/\Gamma_{\text{total}}$   $\Gamma_{39}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt; 0.40</math></b>	90	SHEN	12A	BELL $\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

 $\Gamma(\rho\pi)/\Gamma_{\text{total}}$   $\Gamma_{40}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt; 1.16</math></b>	90	SHEN	13	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- \pi^0$

 $\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{41}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt; 0.80</math></b>	90	SHEN	13	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- \pi^0$

 $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{42}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt; 1.63</math></b>	90	SHEN	13	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0$

 $\Gamma(\pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{43}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>13.0 \pm 1.9 \pm 2.1</math></b>	$261 \pm 37$	SHEN	13	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0$

 $\Gamma(K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{44}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.14 \pm 0.30 \pm 0.13</math></b>	$40 \pm 10$		SHEN	13	BELL $\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 3.2$	90	<sup>1</sup> DOBBS	12A	$\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$
---------	----	--------------------	-----	--

<sup>1</sup> Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

 $\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{45}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt; 4.22</math></b>	90	SHEN	13	BELL $\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$

 $\Gamma(K^*(892)^- K^+ + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{46}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt; 1.45</math></b>	90	SHEN	13	BELL $\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$

$\Gamma(f_1(1285)\text{anything})/\Gamma_{\text{total}}$   $\Gamma_{47}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>2.20 \pm 1.50 \pm 0.63</math></b>	2.9k	JIA	17A	BELL $e^+e^- \rightarrow$ hadrons

$\Gamma(f_1(1285)X_{tetra})/\Gamma_{\text{total}}$   $\Gamma_{48}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt; 64.7 \times 10^{-6}</math></b>	90	<sup>1</sup> JIA	17A	BELL $e^+e^- \rightarrow$ hadrons

<sup>1</sup> For a tetraquark state  $X_{tetra}$ , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of  $X_{tetra}$  mass and width range from  $7.8 \times 10^{-6}$  to  $64.7 \times 10^{-6}$ .

$\Gamma(\text{Sum of 100 exclusive modes})/\Gamma_{\text{total}}$   $\Gamma_{49}/\Gamma$

VALUE (units $10^{-2}$ )	DOCUMENT ID	COMMENT
<b><math>0.29 \pm 0.03</math></b>	<sup>1,2</sup> DOBBS	12A $\Upsilon(2S) \rightarrow$ hadrons

<sup>1</sup> DOBBS 12A presents individual exclusive branching fractions or upper limits for 100 modes of four to ten pions, kaons, or protons.

<sup>2</sup> Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

$\Gamma(\gamma\chi_{b1}(1P))/\Gamma_{\text{total}}$   $\Gamma_{50}/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.069 \pm 0.004</math> OUR AVERAGE</b>				
$0.0693 \pm 0.0012 \pm 0.0041$	407k	ARTUSO	05	CLEO $e^+e^- \rightarrow \gamma X$
$0.069 \pm 0.005 \pm 0.009$		EDWARDS	99	CLE2 $\Upsilon(2S) \rightarrow \gamma\chi(1P)$
$0.091 \pm 0.018 \pm 0.022$		ALBRECHT	85E	ARG $e^+e^- \rightarrow \gamma\text{conv. } X$
$0.065 \pm 0.007 \pm 0.012$		NERNST	85	CBAL $e^+e^- \rightarrow \gamma X$
$0.080 \pm 0.017 \pm 0.016$		HAAS	84	CLEO $e^+e^- \rightarrow \gamma\text{conv. } X$
$0.059 \pm 0.014$		KLOPFEN...	83	CUSB $e^+e^- \rightarrow \gamma X$

$\Gamma(\gamma\chi_{b2}(1P))/\Gamma_{\text{total}}$   $\Gamma_{51}/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.0715 \pm 0.0035</math> OUR AVERAGE</b>				
$0.0724 \pm 0.0011 \pm 0.0040$	410k	ARTUSO	05	CLEO $e^+e^- \rightarrow \gamma X$
$0.074 \pm 0.005 \pm 0.008$		EDWARDS	99	CLE2 $\Upsilon(2S) \rightarrow \gamma\chi(1P)$
$0.098 \pm 0.021 \pm 0.024$		ALBRECHT	85E	ARG $e^+e^- \rightarrow \gamma\text{conv. } X$
$0.058 \pm 0.007 \pm 0.010$		NERNST	85	CBAL $e^+e^- \rightarrow \gamma X$
$0.102 \pm 0.018 \pm 0.021$		HAAS	84	CLEO $e^+e^- \rightarrow \gamma\text{conv. } X$
$0.061 \pm 0.014$		KLOPFEN...	83	CUSB $e^+e^- \rightarrow \gamma X$

$\Gamma(\gamma\chi_{b0}(1P))/\Gamma_{\text{total}}$   $\Gamma_{52}/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.038 \pm 0.004</math> OUR AVERAGE</b>				
$0.0375 \pm 0.0012 \pm 0.0047$	198k	ARTUSO	05	CLEO $e^+e^- \rightarrow \gamma X$
$0.034 \pm 0.005 \pm 0.006$		EDWARDS	99	CLE2 $\Upsilon(2S) \rightarrow \gamma\chi(1P)$
$0.064 \pm 0.014 \pm 0.016$		ALBRECHT	85E	ARG $e^+e^- \rightarrow \gamma\text{conv. } X$
$0.036 \pm 0.008 \pm 0.009$		NERNST	85	CBAL $e^+e^- \rightarrow \gamma X$
$0.044 \pm 0.023 \pm 0.009$		HAAS	84	CLEO $e^+e^- \rightarrow \gamma\text{conv. } X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.035 \pm 0.014$		KLOPFEN...	83	CUSB $e^+e^- \rightarrow \gamma X$
-------------------	--	------------	----	------------------------------------

$\Gamma(\gamma f_0(1710))/\Gamma_{\text{total}}$   $\Gamma_{53}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;59</b>	90	<sup>1</sup> ALBRECHT 89	ARG	$\Upsilon(2S) \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 5.9	90	<sup>2</sup> ALBRECHT 89	ARG	$\Upsilon(2S) \rightarrow \gamma \pi^+ \pi^-$
<sup>1</sup> Re-evaluated assuming $B(f_0(1710) \rightarrow K^+ K^-) = 0.19$ .				
<sup>2</sup> Includes unknown branching ratio of $f_0(1710) \rightarrow \pi^+ \pi^-$ .				

 $\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$   $\Gamma_{54}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;53</b>	90	<sup>1</sup> ALBRECHT 89	ARG	$\Upsilon(2S) \rightarrow \gamma K^+ K^-$
<sup>1</sup> Re-evaluated assuming $B(f'_2(1525) \rightarrow K \bar{K}) = 0.71$ .				

 $\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$   $\Gamma_{55}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;24.1</b>	90	<sup>1</sup> ALBRECHT 89	ARG	$\Upsilon(2S) \rightarrow \gamma \pi^+ \pi^-$
<sup>1</sup> Using $B(f_2(1270) \rightarrow \pi \pi) = 0.84$ .				

 $\Gamma(\gamma f_J(2220))/\Gamma_{\text{total}}$   $\Gamma_{56}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<6.8	90	<sup>1</sup> ALBRECHT 89	ARG	$\Upsilon(2S) \rightarrow \gamma K^+ K^-$
<sup>1</sup> Includes unknown branching ratio of $f_J(2220) \rightarrow K^+ K^-$ .				

 $\Gamma(\gamma \eta_c(1S))/\Gamma_{\text{total}}$   $\Gamma_{57}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;2.7 × 10<sup>-5</sup></b>	90	WANG 11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$

 $\Gamma(\gamma \chi_{c0})/\Gamma_{\text{total}}$   $\Gamma_{58}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.0 × 10<sup>-4</sup></b>	90	WANG 11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$

 $\Gamma(\gamma \chi_{c1})/\Gamma_{\text{total}}$   $\Gamma_{59}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;3.6 × 10<sup>-6</sup></b>	90	WANG 11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$

 $\Gamma(\gamma \chi_{c2})/\Gamma_{\text{total}}$   $\Gamma_{60}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.5 × 10<sup>-5</sup></b>	90	WANG 11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$

 $\Gamma(\gamma \chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi)/\Gamma_{\text{total}}$   $\Gamma_{61}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.8 × 10<sup>-6</sup></b>	90	WANG 11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$

$\Gamma(\gamma\chi_{c1}(3872) \rightarrow \pi^+\pi^-\pi^0 J/\psi)/\Gamma_{\text{total}}$   $\Gamma_{62}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.4 \times 10^{-6}$	90	WANG	11B BELL	$\Upsilon(2S) \rightarrow \gamma X$

 $\Gamma(\gamma X(3915) \rightarrow \omega J/\psi)/\Gamma_{\text{total}}$   $\Gamma_{63}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.8 \times 10^{-6}$	90	WANG	11B BELL	$\Upsilon(2S) \rightarrow \gamma X$

 $\Gamma(\gamma\chi_{c1}(4140) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$   $\Gamma_{64}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.2 \times 10^{-6}$	90	WANG	11B BELL	$\Upsilon(2S) \rightarrow \gamma X$

 $\Gamma(\gamma X(4350) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$   $\Gamma_{65}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.3 \times 10^{-6}$	90	WANG	11B BELL	$\Upsilon(2S) \rightarrow \gamma X$

 $\Gamma(\gamma\eta_b(1S))/\Gamma_{\text{total}}$   $\Gamma_{66}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$3.9 \pm 1.1^{+1.1}_{-0.9}$		$13 \pm 5k$	<sup>1</sup> AUBERT	09AQ BABR	$\Upsilon(2S) \rightarrow \gamma X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<21$	90	LEES	11J BABR	$\Upsilon(2S) \rightarrow X\gamma$
$< 8.4$	90	<sup>1</sup> BONVICINI	10 CLEO	$\Upsilon(2S) \rightarrow \gamma X$
$< 5.1$	90	<sup>2</sup> ARTUSO	05 CLEO	$e^+e^- \rightarrow \gamma X$

<sup>1</sup> Assuming  $\Gamma_{\eta_b(1S)} = 10$  MeV.

<sup>2</sup> Superseded by BONVICINI 10.

 $\Gamma(\gamma\eta_b(1S) \rightarrow \gamma \text{Sum of 26 exclusive modes})/\Gamma_{\text{total}}$   $\Gamma_{67}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.7 \times 10^{-6}$	90	SANDILYA	13 BELL	$\Upsilon(2S) \rightarrow \gamma \text{ hadrons}$

 $\Gamma(\gamma X_{b\bar{b}} \rightarrow \gamma \text{Sum of 26 exclusive modes})/\Gamma_{\text{total}}$   $\Gamma_{68}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$< 4.9$	90		SANDILYA	13 BELL	$\Upsilon(2S) \rightarrow \gamma \text{ hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$46.2^{+29.7}_{-14.2} \pm 10.6$		10	<sup>1</sup> DOBBS	12	$\Upsilon(2S) \rightarrow \gamma \text{ hadrons}$
---------------------------------	--	----	--------------------	----	---

<sup>1</sup> Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$   $\Gamma_{69}/\Gamma$   
( $1.5 \text{ GeV} < m_X < 5.0 \text{ GeV}$ )

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<1.95$	95	ROSNER	07A CLEO	$e^+e^- \rightarrow \gamma X$

$\Gamma(\gamma A^0 \rightarrow \gamma \text{hadrons})/\Gamma_{\text{total}}$   $\Gamma_{70}/\Gamma$   
 (0.3 GeV <  $m_{A^0}$  < 7 GeV)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 8 \times 10^{-5}$	90	<sup>1</sup> LEES	11H BABR	$\Upsilon(2S) \rightarrow \gamma \text{ hadrons}$

<sup>1</sup> For a narrow scalar or pseudoscalar  $A^0$ , excluding known resonances, with mass in the range 0.3–7 GeV. Measured 90% CL limits as a function of  $m_{A^0}$  range from  $1 \times 10^{-6}$  to  $8 \times 10^{-5}$ .

$\Gamma(\gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$   $\Gamma_{71}/\Gamma$

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$< 8.3$	90	<sup>1</sup> AUBERT	09Z BABR	$e^+ e^- \rightarrow \gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-$

<sup>1</sup> For a narrow scalar or pseudoscalar  $a_1^0$  with mass in the range 212–9300 MeV, excluding  $J/\psi$  and  $\psi(2S)$ . Measured 90% CL limits as a function of  $m_{a_1^0}$  range from  $0.26$ – $8.3 \times 10^{-6}$ .

———— LEPTON FAMILY NUMBER (LF) VIOLATING MODES ————

$\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$   $\Gamma_{72}/\Gamma$

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.2$	90	LEES	10B BABR	$e^+ e^- \rightarrow e^\pm \tau^\mp$

$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$   $\Gamma_{73}/\Gamma$

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.3$	90	LEES	10B BABR	$e^+ e^- \rightarrow \mu^\pm \tau^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 14.4$	95	LOVE	08A CLEO	$e^+ e^- \rightarrow \mu^\pm \tau^\mp$
----------	----	------	----------	--

$\Upsilon(2S)$  Cross-Particle Branching Ratios

$B(\Upsilon(2S) \rightarrow \pi^+ \pi^-) \times B(\Upsilon(3S) \rightarrow \Upsilon(2S) X)$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.78 \pm 0.02 \pm 0.11$	906k	LEES	11C BABR	$e^+ e^- \rightarrow \pi^+ \pi^- X$

$\Upsilon(2S)$  REFERENCES

JIA	17	PR D95 012001	S. Jia <i>et al.</i>	(BELLE Collab.)
JIA	17A	PR D96 112002	S. Jia <i>et al.</i>	(BELLE Collab.)
LEES	14G	PR D89 111102	J.P. Lees <i>et al.</i>	(BABAR Collab.)
YANG	14	PR D90 112008	S.D. Yang <i>et al.</i>	(BELLE Collab.)
SANDILYA	13	PRL 111 112001	S. Sandilya <i>et al.</i>	(BELLE Collab.)
SHEN	13	PR D88 011102	C.P. Shen <i>et al.</i>	(BELLE Collab.)
TAMPONI	13	PR D87 011104	U. Tamponi <i>et al.</i>	(BELLE Collab.)
DOBBS	12	PRL 109 082001	S. Dobbs <i>et al.</i>	
DOBBS	12A	PR D86 052003	S. Dobbs <i>et al.</i>	
SHEN	12A	PR D86 031102	C.P. Shen <i>et al.</i>	(BELLE Collab.)
LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11H	PRL 107 221803	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11L	PR D84 092003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
WANG	11B	PR D84 071107	X.L. Wang <i>et al.</i>	(BELLE Collab.)
BONVICINI	10	PR D81 031104	G. Bonvicini <i>et al.</i>	(CLEO Collab.)

LEES	10B	PRL 104 151802	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AUBERT	09AQ	PRL 103 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09Z	PRL 103 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)
BHARI	09	PR D79 011103	S.R. Bhari <i>et al.</i>	(CLEO Collab.)
AUBERT	08BP	PR D78 112002	B. Aubert <i>et al.</i>	(BABAR Collab.)
HE	08A	PRL 101 192001	Q. He <i>et al.</i>	(CLEO Collab.)
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
ASNER	07	PR D75 012009	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BESSON	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
BESSON	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
EDWARDS	99	PR D59 032003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
BARU	96	PRPL 267 71	S.E. Baru <i>et al.</i>	(NOVO)
KOBEL	92	ZPHY C53 193	M. Kobel <i>et al.</i>	(Crystal Ball Collab.)
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	89	ZPHY C42 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)
BUCHMUEL...	88	HE $e^+e^-$ Physics 412	W. Buchmueller, S. Cooper	(HANN, DESY, MIT)
Editors: A. Ali and P. Soeding, World Scientific, Singapore				
JAKUBOWSKI	88	ZPHY C40 49	Z. Jakubowski <i>et al.</i>	(Crystal Ball Collab.) IGJPC
ALBRECHT	87	ZPHY C35 283	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
LURZ	87	ZPHY C36 383	B. Lurz <i>et al.</i>	(Crystal Ball Collab.)
BARU	86B	ZPHY C32 622 (erratum)	S.E. Baru <i>et al.</i>	(NOVO)
ALBRECHT	85	ZPHY C28 45	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	85E	PL 160B 331	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
GELPHMAN	85	PR D32 2893	D. Gelpman <i>et al.</i>	(Crystal Ball Collab.)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
Translated from YAF 41 733.				
NERNST	85	PRL 54 2195	R. Nernst <i>et al.</i>	(Crystal Ball Collab.)
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)
BARBER	84	PL 135B 498	D.P. Barber <i>et al.</i>	(DESY, ARGUS Collab.+)
BESSON	84	PR D30 1433	D. Besson <i>et al.</i>	(CLEO Collab.)
FONSECA	84	NP B242 31	V. Fonseca <i>et al.</i>	(CUSB Collab.)
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)
HAAS	84	PRL 52 799	J. Haas <i>et al.</i>	(CLEO Collab.)
HAAS	84B	PR D30 1996	J. Haas <i>et al.</i>	(CLEO Collab.)
KLOPFEN...	83	PRL 51 160	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)
ALBRECHT	82	PL 116B 383	H. Albrecht <i>et al.</i>	(DESY, DORT, HEIDH+)
NICZYPORUK	81B	PL 100B 95	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
NICZYPORUK	81C	PL 99B 169	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
BOCK	80	ZPHY C6 125	P. Bock <i>et al.</i>	(HEIDP, MPIM, DESY, HAMB)