

$\Xi_c(2815)$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-) \text{ Status: } ***$$

Seen in both $\Xi_c \pi$ and $\Xi_c \pi \pi$ decays. The simplest assignment is that this belongs to the same SU(4) multiplet as the $\Lambda(1520)$ and the $\Lambda_c(2625)$, but the spin and parity have not been measured.

$\Xi_c(2815)$ MASSES

The masses are obtained from the mass-difference measurements that follow.

$\Xi_c(2815)^+$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

2816.67 ± 0.31 OUR FIT Error includes scale factor of 1.1.

• • • We do not use the following data for averages, fits, limits, etc. • • •

2817.0 ± 1.2 $\begin{smallmatrix} +0.7 \\ -0.8 \end{smallmatrix}$	73 ± 10	LESLIAK	08 BELL	$e^+ e^- \approx \gamma(4S)$
---	---------	---------	---------	------------------------------

$\Xi_c(2815)^0$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

2820.22 ± 0.32 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

2820.4 ± 1.4 $\begin{smallmatrix} +0.9 \\ -1.0 \end{smallmatrix}$	48 ± 8	LESLIAK	08 BELL	$e^+ e^- \approx \gamma(4S)$
---	--------	---------	---------	------------------------------

$\Xi_c(2815) - \Xi_c$ MASS DIFFERENCES

$m_{\Xi_c(2815)^+} - m_{\Xi_c^+}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

348.80 ± 0.10 OUR FIT

348.80 ± 0.08 ± 0.06 941 YELTON 16 BELL $e^+ e^-$, γ regions

• • • We do not use the following data for averages, fits, limits, etc. • • •

348.6 ± 0.6 ± 1.0	20	ALEXANDER	99B CLE2	$e^+ e^- \approx \gamma(4S)$
-------------------	----	-----------	----------	------------------------------

$m_{\Xi_c(2815)^0} - m_{\Xi_c^0}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

349.35 ± 0.11 OUR FIT

349.35 ± 0.08 ± 0.07 1258 YELTON 16 BELL $e^+ e^-$, γ regions

• • • We do not use the following data for averages, fits, limits, etc. • • •

347.2 ± 0.7 ± 2.0	9	ALEXANDER	99B CLE2	$e^+ e^- \approx \gamma(4S)$
-------------------	---	-----------	----------	------------------------------

$\Xi_c(2815)^+ - \Xi_c(2815)^0$ MASS DIFFERENCE

$m_{\Xi_c(2815)^+} - m_{\Xi_c(2815)^0}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

-3.55 ± 0.28 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

-3.47 ± 0.12 ± 0.48	YELTON	16 BELL	941 and 1258 evts
---------------------	--------	---------	-------------------

-3.4 ± 1.9 ± 0.9	LESLIAK	08 BELL	73 & 48 events
------------------	---------	---------	----------------

$\Xi_c(2815)$ WIDTHS **$\Xi_c(2815)^+$ WIDTH**

<u>VALUE (MeV)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.43 \pm 0.20 \pm 0.17$		941	YELTON	16 BELL	$e^+ e^-$, γ regions
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<3.5	90		ALEXANDER	99B CLE2	$e^+ e^- \approx \gamma(4S)$

 $\Xi_c(2815)^0$ WIDTH

<u>VALUE (MeV)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.54 \pm 0.18 \pm 0.17$		1258	YELTON	16 BELL	$e^+ e^-$, γ regions
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<6.5	90		ALEXANDER	99B CLE2	$e^+ e^- \approx \gamma(4S)$

 $\Xi_c(2815)$ DECAY MODES

The $\Xi_c \pi \pi$ modes are consistent with being entirely via $\Xi_c(2645) \pi$.

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \Xi_c^- \pi$	seen
$\Gamma_2 \quad \Xi_c^+ \pi^+ \pi^-$	seen
$\Gamma_3 \quad \Xi_c^0 \pi^+ \pi^-$	seen

 $\Xi_c(2815)$ REFERENCES

YELTON	16	PR D94 052011	J. Yelton <i>et al.</i>	(BELLE Collab.)
LESIK	08	PL B665 9	T. Lesiak <i>et al.</i>	(BELLE Collab.)
ALEXANDER	99B	PRL 83 3390	J.P. Alexander <i>et al.</i>	(CLEO Collab.)