

107. W' -Boson Searches

Revised October 2017 by B.A. Dobrescu (Fermilab) and S. Willocq (Univ. of Massachusetts).

The W' boson is a massive hypothetical particle of spin 1 and electric charge ± 1 , which is a color singlet and is predicted in various extensions of the Standard Model (SM).

107.1. W' couplings to quarks and leptons.

The Lagrangian terms describing couplings of a W'^+ boson to fermions are given by

$$\frac{W'^+}{\sqrt{2}} \left[\bar{u}_i \left(C_{qij}^R P_R + C_{qij}^L P_L \right) \gamma^\mu d_j + \bar{\nu}_i \left(C_{lij}^R P_R + C_{lij}^L P_L \right) \gamma^\mu e_j \right]. \quad (107.1)$$

Here u, d, ν and e are the SM fermions in the mass eigenstate basis, $i, j = 1, 2, 3$ label the fermion generation, and $P_{R,L} = (1 \pm \gamma_5)/2$. The coefficients C_{qij}^L , C_{qij}^R , C_{lij}^L , and C_{lij}^R are complex dimensionless parameters. If $C_{lij}^R \neq 0$, then the i th generation includes a right-handed neutrino. Using this notation, the SM W couplings are $C_q^L = g V_{\text{CKM}}$, $C_\ell^L = g \approx 0.63$ and $C_q^R = C_\ell^R = 0$.

Unitarity considerations imply that the W' boson is associated with a spontaneously-broken gauge symmetry. This is true even when it is a composite particle (*e.g.*, ρ^\pm -like bound states [1]) if its mass is much smaller than the compositeness scale, or a Kaluza-Klein mode in theories where the W boson propagates in extra dimensions [2]. The simplest extension of the electroweak gauge group that includes a W' boson is $SU(2)_1 \times SU(2)_2 \times U(1)$, but larger groups are encountered in some theories. A generic property of these gauge theories is that they also include a Z' boson [3]; the Z' -to- W' mass ratio is often a free parameter.

A tree-level mass mixing may be induced between the electrically-charged gauge bosons. Upon diagonalization of their mass matrix, the $W - Z$ mass ratio and the couplings of the observed W boson are shifted from the SM values. Their measurements imply that the mixing angle between the gauge eigenstates, θ_+ , must be smaller than about 10^{-2} . In certain theories the mixing is negligible (*e.g.* due to a new parity [4]), even when the W' mass is near the electroweak scale.

The W' coupling to WZ is fixed by Lorentz and gauge invariances, and to leading order in θ_+ is given by [5]

$$\frac{g \theta_+^i}{\cos \theta_W} \left[W_\mu'^+ \left(W_\nu^- Z^{\nu\mu} + Z_\nu W^{-\mu\nu} \right) + Z^\nu W^{-\mu} W_{\nu\mu}'^+ \right] + \text{H.c.}, \quad (107.2)$$

where $W^{\mu\nu} \equiv \partial^\mu W^\nu - \partial^\nu W^\mu$, etc. The θ_W dependence shown here corrects the one given in [6], which has been referred to as the Extended Gauge Model by the experimental collaborations. The W' coupling to Wh^0 , where h^0 is the SM Higgs boson, is

$$-\xi_h g_{W'} M_W W_\mu'^+ W^{\mu-} h^0 + \text{H.c.}, \quad (107.3)$$

where $g_{W'}$ is the gauge coupling of the W' boson, and the coefficient ξ_h satisfies $\xi_h \leq 1$ in simple Higgs sectors [5].

2 107. W' -boson searches

In models based on the “left-right symmetric” gauge group [7], $SU(2)_L \times SU(2)_R \times U(1)_{B-L}$, the SM fermions that couple to the W boson transform as doublets under $SU(2)_L$ while the other fermions transform as doublets under $SU(2)_R$. Consequently, the W' boson couples primarily to right-handed fermions; its coupling to left-handed fermions arises due to the θ_+ mixing, so that C_q^L is proportional to the CKM matrix and its elements are much smaller than the diagonal elements of C_q^R . Generically, C_q^R does not need to be proportional to V_{CKM} .

There are many other models based on the $SU(2)_1 \times SU(2)_2 \times U(1)$ gauge symmetry. In the “alternate left-right” model [8], all the couplings shown in Eq. (107.1) vanish, but there are some new fermions such that the W' boson couples to pairs involving a SM fermion and a new fermion. In the “unified SM” [9], the left-handed quarks are doublets under one $SU(2)$, and the left-handed leptons are doublets under a different $SU(2)$, leading to a mostly leptophobic W' boson: $C_{\ell ij}^L \ll C_{q ij}^L$ and $C_{\ell ij}^R = C_{q ij}^R = 0$. Fermions of different generations may also transform as doublets under different $SU(2)$ gauge groups [10]. In particular, the couplings to third generation quarks may be enhanced [11].

It is also possible that the W' couplings to SM fermions are highly suppressed. For example, if the quarks and leptons are singlets under one $SU(2)$ [12], then the couplings are proportional to the tiny mixing angle θ_+ . Similar suppressions may arise if some vectorlike fermions mix with the SM fermions [13].

Gauge groups that embed the electroweak symmetry, such as $SU(3)_W \times U(1)$ or $SU(4)_W \times U(1)$, also include one or more W' bosons [14].

107.2. Collider searches.

At LEP-II, W' bosons could have been produced in pairs via their photon and Z couplings. The production cross section is large enough to rule out $M_{W'} < \sqrt{s}/2 \approx 105$ GeV for most patterns of decay modes.

At hadron colliders, W' bosons can be detected through resonant pair production of fermions or electroweak bosons. Assuming that the W' width is much smaller than its mass, the contribution of the s -channel W' boson exchange to the total rate for $pp \rightarrow f\bar{f}'X$, where f and f' are fermions whose difference of electric charges is ± 1 , and X is any final state, may be approximated by the branching fraction $B(W' \rightarrow f\bar{f}')$ times the production cross section

$$\sigma(pp \rightarrow W'X) \simeq \frac{\pi}{48s} \sum_{i,j} \left[(C_{qij}^L)^2 + (C_{qij}^R)^2 \right] w_{ij} \left(M_{W'}^2/s, M_{W'} \right). \quad (107.4)$$

The functions w_{ij} include the information about proton structure, and are given to leading order in α_s by

$$w_{ij}(z, \mu) = \int_z^1 \frac{dx}{x} \left[u_i(x, \mu) \bar{d}_j \left(\frac{z}{x}, \mu \right) + \bar{u}_i(x, \mu) d_j \left(\frac{z}{x}, \mu \right) \right], \quad (107.5)$$




Figure 107.1: Upper limit on $\sigma(pp \rightarrow W'X) B(W' \rightarrow e\nu)$ from ATLAS [16], at 95% CL. The red line shows the theoretical prediction in the Sequential SM.

where $u_i(x, \mu)$ and $d_i(x, \mu)$ are the parton distributions inside the proton, at the factorization scale μ and parton momentum fraction x , for the up- and down-type quark of the i th generation, respectively. QCD corrections to W' production are sizable (they also include quark-gluon initial states), but preserve the above factorization of couplings at next-to-leading order [15].

The most commonly studied W' signal consists of a high-momentum electron or muon and large missing transverse momentum, with the transverse mass distribution forming a Jacobian peak with its endpoint at $M_{W'}$ (see Fig. 1a of [16]). Given that the branching fractions for $W' \rightarrow e\nu$ and $W' \rightarrow \mu\nu$ could be very different, the results in these channels should be presented separately. Searches in these channels often implicitly assume that the left-handed couplings vanish (no interference between W and W'), and that the right-handed neutrino is light compared to the W' boson and escapes the detector. An example of parameter values that satisfy these assumptions is $C_q^R = gV_{CKM}$, $C_\ell^R = g$, $C_q^L = C_\ell^L = 0$, which define a model that preserves lepton universality and is essentially equivalent to the Sequential SM used in many W' searches. However, if a W' boson were discovered and the final state fermions have left-handed helicity, then the effects of $W - W'$ interference could be observed [17], providing useful information about the W' couplings.

In the $e\nu$ channel, the ATLAS and CMS Collaborations set limits on the W' production cross section times branching fraction (and thus indirectly on the W' couplings) when $M_{W'}$ is in the 0.15 – 6 TeV range, based on 2–36 fb^{-1} at $\sqrt{s} = 13$ TeV [16,18], as shown in Fig. 107.1. ATLAS sets the strongest mass lower limit $M_{W'} > 5.2$ TeV in the Sequential SM (all limits in this mini-review are at the 95% CL). The coupling limits are much weaker for $M_{W'} < 150$ GeV, a range last explored with the Tevatron at $\sqrt{s} = 1.8$ TeV [19].

4 107. W' -boson searches

In the $\mu\nu$ channel, ATLAS and CMS set rate limits for $M_{W'}$ in the 0.15 – 6 TeV range from the same analyses as mentioned above, with the strongest lower mass limit of 4.5 TeV set by ATLAS [16] using 36.1 fb^{-1} of $\sqrt{s} = 13 \text{ TeV}$ data. When combined with the $e\nu$ channel assuming lepton universality, the upper limit on the $\sqrt{s} = 13 \text{ TeV}$ cross section times branching fraction to $\ell\nu$ varies between 0.2 and 4 fb for $M_{W'}$ between 1 and 5 TeV [16]. Only weak limits on $W' \rightarrow \mu\nu$ exist for $M_{W'} < 150 \text{ GeV}$ [20]. Note that masses of the order of the electroweak scale are interesting from a theory point of view, while lepton universality does not necessarily apply to a W' boson.

Dedicated searches for $W' \rightarrow \tau\nu$ have been performed by CMS at 8 TeV [21] and 13 TeV [22]. Limits are set on $\sigma \cdot B$ for $M_{W'}$ between 0.3 and 4 TeV for the former and between 1.0 and 5.8 TeV for the latter. A lower mass limit of 3.3 TeV is set in the Sequential SM.

The W' decay into a lepton and a right-handed neutrino, ν_R , may also be followed by the ν_R decay through a virtual W' boson into a lepton and two quark jets. The CMS [23,24] and ATLAS [25] searches in the $eejj$ and $\mu\mu jj$ channels have set limits on the cross section times branching fraction as a function of the ν_R mass or of $M_{W'}$. These searches are typically performed with same-charge lepton pairs that provide strong background reduction and are motivated by models with a left-right symmetry. However, it is also interesting to search in final states with opposite-charge lepton pairs, as done in the CMS analysis. A related W' search in the $\tau\tau jj$ channel with hadronic τ 's was also performed by CMS [26].

The $t\bar{b}$ channel is particularly important because a W' boson that couples only to right-handed fermions cannot decay to leptons when the right-handed neutrinos are heavier than the W' boson (additional motivations are provided by a W' boson with enhanced couplings to the third generation [11], and by a leptophobic W' boson). The usual signature consists of a leptonically-decaying W boson and two b -jets. Recent studies have also incorporated the fully hadronic decay channel for $M_{W'} \gg m_t$ with the use of jet substructure techniques to tag highly boosted top-jets. For a detailed discussion of this channel, see [27].

Searches for dijet resonances may be used to set limits on $W' \rightarrow q\bar{q}'$. CMS [28] and ATLAS [29] provide similar coverage in the $\sim 0.75 - 7.0 \text{ TeV}$ mass range with data collected at $\sqrt{s} = 8$ and 13 TeV, with the most stringent lower W' mass limit in the Sequential SM set to 3.6 TeV using 37 fb^{-1} of 13 TeV data. For lower masses, the best limits on W' couplings to quarks have been set by CDF [30] in the 300 – 500 GeV range, and by CMS [31] in the 500 – 750 GeV range. Limits for W' masses in the 50 – 300 GeV range can be derived from the dijet limits on Z' bosons set by CMS [32].

In some theories [4] the W' couplings to SM fermions are suppressed by discrete symmetries. W' production then occurs in pairs, through a photon or Z boson. The decay modes are model-dependent and often involve other new particles. The ensuing collider signals arise from cascade decays and typically include missing transverse momentum.

Searches for WZ resonances at the LHC have focused on the process $pp \rightarrow W' \rightarrow WZ$ with the production mainly from $u\bar{d} \rightarrow W'$ assuming SM-like couplings to quarks. ATLAS and CMS have set the strongest upper limits on the $W'WZ$ coupling for $M_{W'}$




Figure 107.2: Upper limits at 95% CL on W' production cross section times branching fraction into a W and a SM Higgs boson, from [41] (left) and [44] (right).

in the 0.2 – 5.0 TeV range with a combination of fully leptonic, semi-leptonic and fully hadronic channels (see also [27]) at both 8 and 13 TeV [33]-[37]. The strongest lower limit on the mass is set by CMS at 13 TeV with 35.5 fb^{-1} in the $WZ \rightarrow (jj)(\nu\bar{\nu})$ final state, where the parentheses represent a resonance; the limit is $M_{W'} > 3.2 \text{ TeV}$ in the context of the Heavy Vector Triplet (HVT) weakly-coupled scenario A [38].

A fermiophobic W' boson that couples to WZ may be produced at hadron colliders in association with a Z boson, or via WZ fusion. This would give rise to $(WZ)Z$ and $(WZ)jj$ final states [39].

W' bosons have also been searched for recently in final states with a W boson and a SM Higgs boson in the channels $W \rightarrow \ell\nu$ or $W \rightarrow qq$ and $h^0 \rightarrow b\bar{b}$ or $h^0 \rightarrow WW$ by ATLAS [40,41] and CMS [36,42,43,44] at $\sqrt{s} = 8$ and 13 TeV. Cross section limits are set for W' masses in the range between 0.4 and 4.5 TeV. The ATLAS and CMS 13 TeV analyses both set the most stringent lower limit on the mass: $M_{W'} > 2.4 \text{ TeV}$ for the HVT weakly-coupled scenario A, as shown in Fig. 107.2.

107.3. Low-energy constraints.

The properties of W' bosons are also constrained by measurements of processes at energies much below $M_{W'}$. The bounds on $W - W'$ mixing [45] are mostly due to the change in W properties compared to the SM. Limits on deviations in the ZWW couplings provide a leading constraint for fermiophobic W' bosons [13].

Constraints arising from low-energy effects of W' exchange are strongly model-dependent. If the W' couplings to quarks are not suppressed, then box diagrams involving a W and a W' boson contribute to neutral meson-mixing. In the case of W' couplings to right-handed quarks as in the left-right symmetric model, the limit from $K_L - K_S$ mixing is severe: $M_{W'} > 2.9 \text{ TeV}$ for $C_q^R = gV_{\text{CKM}}$ [46]. However, if no correlation between the W' and W couplings is assumed, then the limit on $M_{W'}$ may be significantly relaxed [47].

W' exchange also contributes at tree level to various low-energy processes. In particular, it would impact the measurement of the Fermi constant G_F in muon decay,

6 107. W' -boson searches

which in turn would change the predictions of many other electroweak processes. A recent test of parity violation in polarized muon decay [48] has set limits of about 600 GeV on $M_{W'}$, assuming W' couplings to right-handed leptons as in left-right symmetric models and a light ν_R . There are also W' contributions to the neutron electric dipole moment, β decays, and other processes [45].

If right-handed neutrinos have Majorana masses, then there are tree-level contributions to neutrinoless double-beta decay, and a limit on $M_{W'}$ versus the ν_R mass may be derived [49]. For ν_R masses below a few GeV, the W' boson contributes to leptonic and semileptonic B meson decays, so that limits may be placed on various combinations of W' parameters [47]. For ν_R masses below ~ 30 MeV, the most stringent constraints on $M_{W'}$ are due to the limits on ν_R emission from supernovae.

References:

1. M. Bando, T. Kugo, and K. Yamawaki, Phys. Rept. **164**, 217 (1988).
2. H.C. Cheng *et al.*, Phys. Rev. D **64**, 065007 (2001).
3. See the Section on “ Z' -boson searches” in this *Review*.
4. H.C. Cheng and I. Low, JHEP **0309**, 051 (2003).
5. B.A. Dobrescu and Z. Liu, JHEP **1510**, 118 (2015).
6. G. Altarelli, B. Mele and M. Ruiz-Altaba, Z. Phys. C **45**, 109 (1989) [Z. Phys. C **47**, 676 (1990)].
7. R.N. Mohapatra and J.C. Pati, Phys. Rev. D **11**, 566 (1975); G. Senjanovic and R.N. Mohapatra, Phys. Rev. D **12**, 1502 (1975).
8. K.S. Babu, X.G. He, and E. Ma, Phys. Rev. D **36**, 878 (1987).
9. H. Georgi, E.E. Jenkins, and E.H. Simmons, Nucl. Phys. B **331**, 541 (1990).
10. See, *e.g.*, X. Li and E. Ma, J. Phys. G **19**, 1265 (1993).
11. D.J. Muller and S. Nandi, Phys. Lett. B **383**, 345 (1996); E. Malkawi, T. Tait, and C.P. Yuan, Phys. Lett. B **385**, 304 (1996).
12. A. Donini *et al.*, Nucl. Phys. B **507**, 51 (1997).
13. R.S. Chivukula *et al.*, Phys. Rev. D **74**, 075011 (2006); H.J. He, T. Tait, and C.P. Yuan, Phys. Rev. D **62**, 011702 (2000).
14. F. Pisano and V. Pleitez, Phys. Rev. D **46**, 410 (1992); **51**, 3865 (1995).
15. Z. Sullivan, Phys. Rev. D **66**, 075011 (2002).
16. ATLAS Collab., arXiv:1706.04786.
17. T.G. Rizzo, JHEP **0705**, 037 (2007); E. Boos *et al.*, Phys. Lett. B **655**, 245 (2007).
18. CMS Collab., Phys. Lett. B **770**, 278 (2017).
19. CDF Collab., F. Abe *et al.*, Phys. Rev. Lett. **74**, 2900 (1995); D0 Collab., S. Abachi *et al.*, Phys. Lett. B **358**, 405 (1995).
20. CDF Collab., F. Abe *et al.*, Phys. Rev. Lett. **67**, 2609 (1991).
21. CMS Collab., Phys. Lett. B **755**, 196 (2016).
22. CMS Collab., PAS EXO-16-006, Mar. 2016.
23. CMS Collab., Eur. Phys. J. C **74**, 3149 (2014).
24. CMS Collab., PAS EXO-16-045, May 2017.
25. ATLAS Collab., JHEP **1507**, 162 (2015).
26. CMS Collab., JHEP **1703**, 077 (2017).

27. K.M. Black *et al.*, “Dynamical electroweak symmetry breaking” in this *Review*.
28. CMS Collab., Phys. Lett. B **769**, 520 (2017); Phys. Rev. D **91**, 052009 (2015).
29. ATLAS Collab., arXiv:1703.09127; Phys. Rev. D **91**, 052007 (2015); CONF-2016-030, June 2016.
30. CDF Collab., F. Abe *et al.*, Phys. Rev. D **55**, R5263 (1997); T. Aaltonen *et al.*, Phys. Rev. D **79**, 112002 (2009).
31. CMS Collab., Phys. Rev. Lett. **117**, 031802 (2016).
32. CMS Collab., arXiv:1705.10532; PAS-EXO-17-001, May 2017.
33. ATLAS Collab., Phys. Lett. B **755**, 285 (2016).
34. ATLAS Collab., arXiv:1708.04445; arXiv:1708.09638; CONF-2017-051, Jul. 2017.
35. CMS Collab., JHEP **1408**, 173 (2014); JHEP **1408**, 174 (2014); Phys. Lett. B **740**, 83 (2015).
36. CMS Collab., arXiv:1705.09171.
37. CMS Collab., arXiv:1708.05379; PAS EXO-15-002, Dec. 2015; PAS B2G-16-020, Aug. 2016; PAS B2G-16-022, Jan. 2017; PAS B2G-17-005, Jul. 2017.
38. D. Pappadopulo *et al.*, JHEP **1409**, 060 (2014).
39. H.J. He *et al.*, Phys. Rev. D **78**, 031701 (2008).
40. ATLAS Collab., Eur. Phys. J. C **75**, 263 (2015).
41. ATLAS Collab., arXiv:1707.06958.
42. CMS Collab., JHEP **1602**, 145 (2016).
43. CMS Collab., Phys. Lett. B **768**, 137 (2017).
44. CMS Collab., arXiv:1707.01303.
45. See the particle listings for W' in this *Review*.
46. Y. Zhang *et al.*, Phys. Rev. D **76**, 091301 (2007); S. Bertolini, A. Maiezza, and F. Nesti, Phys. Rev. D **89**, 095028 (2014).
47. P. Langacker and S.U. Sankar, Phys. Rev. D **40**, 1569 (1989).
48. TWIST Collab., J.F. Bueno *et al.*, Phys. Rev. D **84**, 032005 (2011).
49. See Fig. 5 of G. Prezeau, M. Ramsey-Musolf, and P. Vogel, Phys. Rev. D **68**, 034016 (2003).