

CHARMED BARYONS ($C = +1$)

$$\Lambda_c^+ = udc, \quad \Sigma_c^{++} = uuc, \quad \Sigma_c^+ = udc, \quad \Sigma_c^0 = ddc,$$

$$\Xi_c^+ = usc, \quad \Xi_c^0 = dsc, \quad \Omega_c^0 = ssc$$

Λ_c^+

$$I(J^P) = 0(\frac{1}{2}^+)$$

Mass $m = 2286.46 \pm 0.14$ MeV

Mean life $\tau = (200 \pm 6) \times 10^{-15}$ s ($S = 1.6$)

$c\tau = 59.9$ μm

Decay asymmetry parameters

$$\Lambda\pi^+ \quad \alpha = -0.91 \pm 0.15$$

$$\Sigma^+\pi^0 \quad \alpha = -0.45 \pm 0.32$$

$$\Lambda\ell^+\nu_\ell \quad \alpha = -0.86 \pm 0.04$$

$$(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha}) \text{ in } \Lambda_c^+ \rightarrow \Lambda\pi^+, \bar{\Lambda}_c^- \rightarrow \bar{\Lambda}\pi^- = -0.07 \pm 0.31$$

$$(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha}) \text{ in } \Lambda_c^+ \rightarrow \Lambda e^+\nu_e, \bar{\Lambda}_c^- \rightarrow \bar{\Lambda}e^-\bar{\nu}_e = 0.00 \pm 0.04$$

Branching fractions marked with a footnote, e.g. [a], have been corrected for decay modes not observed in the experiments. For example, the sub-mode fraction $\Lambda_c^+ \rightarrow p\bar{K}^*(892)^0$ seen in $\Lambda_c^+ \rightarrow pK^-\pi^+$ has been multiplied up to include $\bar{K}^*(892)^0 \rightarrow \bar{K}^0\pi^0$ decays.

Λ_c^+ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
Hadronic modes with a p or n: $S = -1$ final states			
pK_S^0	(1.58 \pm 0.08) %	S=1.1	873
$pK^-\pi^+$	(6.23 \pm 0.33) %	S=1.4	823
$p\bar{K}^*(892)^0$	[a] (1.94 \pm 0.27) %		685
$\Delta(1232)^{++}K^-$	(1.07 \pm 0.25) %		710
$\Lambda(1520)\pi^+$	[a] (2.2 \pm 0.5) %		627
$pK^-\pi^+$ nonresonant	(3.4 \pm 0.4) %		823
$pK_S^0\pi^0$	(1.96 \pm 0.13) %	S=1.1	823
$nK_S^0\pi^+$	(1.82 \pm 0.25) %		821
$p\bar{K}^0\eta$	(1.6 \pm 0.4) %		568
$pK_S^0\pi^+\pi^-$	(1.59 \pm 0.12) %	S=1.2	754
$pK^-\pi^+\pi^0$	(4.42 \pm 0.31) %	S=1.5	759
$pK^*(892)^-\pi^+$	[a] (1.4 \pm 0.5) %		580

$\rho(K^- \pi^+)_{\text{nonresonant}} \pi^0$	(4.5 \pm 0.8) %	759
$\Delta(1232) \bar{K}^*(892)$	seen	419
$\rho K^- 2\pi^+ \pi^-$	(1.4 \pm 0.9) $\times 10^{-3}$	671
$\rho K^- \pi^+ 2\pi^0$	(10 \pm 5) $\times 10^{-3}$	678

Hadronic modes with a p : $S = 0$ final states

$p\pi^0$	< 2.7 $\times 10^{-4}$	CL=90%	945
$p\eta$	(1.24 \pm 0.30) $\times 10^{-3}$		856
$p\pi^+ \pi^-$	(4.2 \pm 0.4) $\times 10^{-3}$		927
$\rho f_0(980)$	[a] (3.4 \pm 2.3) $\times 10^{-3}$		614
$\rho 2\pi^+ 2\pi^-$	(2.2 \pm 1.4) $\times 10^{-3}$		852
$\rho K^+ K^-$	(10 \pm 4) $\times 10^{-4}$		616
$p\phi$	[a] (1.06 \pm 0.14) $\times 10^{-3}$		590
$\rho K^+ K^- \text{ non-}\phi$	(5.2 \pm 1.2) $\times 10^{-4}$		616
$p\phi\pi^0$	(10 \pm 4) $\times 10^{-5}$		460
$\rho K^+ K^- \pi^0 \text{ nonresonant}$	< 6.3 $\times 10^{-5}$	CL=90%	494

Hadronic modes with a hyperon: $S = -1$ final states

$\Lambda\pi^+$	(1.29 \pm 0.07) %	S=1.2	864
$\Lambda\pi^+ \pi^0$	(7.0 \pm 0.4) %	S=1.1	844
$\Lambda\rho^+$	< 6 %	CL=95%	636
$\Lambda\pi^- 2\pi^+$	(3.61 \pm 0.29) %	S=1.5	807
$\Sigma(1385)^+ \pi^+ \pi^-, \Sigma^{*+} \rightarrow$ $\Lambda\pi^+$	(1.0 \pm 0.5) %		688
$\Sigma(1385)^- 2\pi^+, \Sigma^{*-} \rightarrow$ $\Lambda\pi^-$	(7.6 \pm 1.4) $\times 10^{-3}$		688
$\Lambda\pi^+ \rho^0$	(1.4 \pm 0.6) %		524
$\Sigma(1385)^+ \rho^0, \Sigma^{*+} \rightarrow \Lambda\pi^+$	(5 \pm 4) $\times 10^{-3}$		363
$\Lambda\pi^- 2\pi^+ \text{ nonresonant}$	< 1.1 %	CL=90%	807
$\Lambda\pi^- \pi^0 2\pi^+ \text{ total}$	(2.2 \pm 0.8) %		757
$\Lambda\pi^+ \eta$	[a] (2.2 \pm 0.5) %		691
$\Sigma(1385)^+ \eta$	[a] (1.06 \pm 0.32) %		570
$\Lambda\pi^+ \omega$	[a] (1.5 \pm 0.5) %		517
$\Lambda\pi^- \pi^0 2\pi^+, \text{ no } \eta \text{ or } \omega$	< 8 $\times 10^{-3}$	CL=90%	757
$\Lambda K^+ \bar{K}^0$	(5.6 \pm 1.1) $\times 10^{-3}$	S=1.9	443
$\Xi(1690)^0 K^+, \Xi^{*0} \rightarrow \Lambda \bar{K}^0$	(1.6 \pm 0.5) $\times 10^{-3}$		286
$\Sigma^0 \pi^+$	(1.28 \pm 0.07) %	S=1.1	825
$\Sigma^+ \pi^0$	(1.24 \pm 0.10) %		827
$\Sigma^+ \eta$	(6.9 \pm 2.3) $\times 10^{-3}$		713
$\Sigma^+ \pi^+ \pi^-$	(4.42 \pm 0.28) %	S=1.2	804
$\Sigma^+ \rho^0$	< 1.7 %	CL=95%	575
$\Sigma^- 2\pi^+$	(1.86 \pm 0.18) %		799
$\Sigma^0 \pi^+ \pi^0$	(2.2 \pm 0.8) %		803
$\Sigma^0 \pi^- 2\pi^+$	(1.10 \pm 0.30) %		763
$\Sigma^+ \pi^+ \pi^- \pi^0$	—		767

$\Sigma^+ \omega$	[a]	$(1.69 \pm 0.21) \%$		569
$\Sigma^- \pi^0 2\pi^+$		$(2.1 \pm 0.4) \%$		762
$\Sigma^+ K^+ K^-$		$(3.4 \pm 0.4) \times 10^{-3}$	S=1.1	349
$\Sigma^+ \phi$	[a]	$(3.8 \pm 0.6) \times 10^{-3}$	S=1.1	295
$\Xi(1690)^0 K^+, \Xi^{*0} \rightarrow$ $\Sigma^+ K^-$		$(10.0 \pm 2.5) \times 10^{-4}$		286
$\Sigma^+ K^+ K^-$ nonresonant		$< 8 \times 10^{-4}$	CL=90%	349
$\Xi^0 K^+$		$(4.9 \pm 1.2) \times 10^{-3}$		653
$\Xi^- K^+ \pi^+$		$(6.2 \pm 0.6) \times 10^{-3}$	S=1.1	565
$\Xi(1530)^0 K^+, \Xi^0 \rightarrow$ $\Xi^- \pi^+$		$(3.3 \pm 1.2) \times 10^{-3}$		473

Hadronic modes with a hyperon: S = 0 final states

ΛK^+		$(6.0 \pm 1.2) \times 10^{-4}$		781
$\Lambda K^+ \pi^+ \pi^-$		$< 5 \times 10^{-4}$	CL=90%	637
$\Sigma^0 K^+$		$(5.1 \pm 0.8) \times 10^{-4}$		735
$\Sigma^0 K^+ \pi^+ \pi^-$		$< 2.6 \times 10^{-4}$	CL=90%	574
$\Sigma^+ K^+ \pi^-$		$(2.1 \pm 0.6) \times 10^{-3}$		670
$\Sigma^+ K^*(892)^0$	[a]	$(3.4 \pm 1.0) \times 10^{-3}$		469
$\Sigma^- K^+ \pi^+$		$< 1.2 \times 10^{-3}$	CL=90%	664

Doubly Cabibbo-suppressed modes

$p K^+ \pi^-$		$(1.46 \pm 0.23) \times 10^{-4}$		823
---------------	--	----------------------------------	--	-----

Semileptonic modes

$\Lambda e^+ \nu_e$		$(3.6 \pm 0.4) \%$		871
$\Lambda \mu^+ \nu_\mu$		$(3.5 \pm 0.5) \%$		867

Inclusive modes

e^+ anything		$(4.5 \pm 1.7) \%$		—
$p e^+$ anything		$(1.8 \pm 0.9) \%$		—
p anything		$(50 \pm 16) \%$		—
p anything (no Λ)		$(12 \pm 19) \%$		—
n anything		$(50 \pm 16) \%$		—
n anything (no Λ)		$(29 \pm 17) \%$		—
Λ anything		$(35 \pm 11) \%$	S=1.4	—
Σ^\pm anything	[b]	$(10 \pm 5) \%$		—
3prongs		$(24 \pm 8) \%$		—

**$\Delta C = 1$ weak neutral current (C1) modes, or
Lepton Family number (LF), or Lepton number (L), or
Baryon number (B) violating modes**

$p e^+ e^-$	C1	$< 5.5 \times 10^{-6}$	CL=90%	951
$p \mu^+ \mu^-$	C1	$< 4.4 \times 10^{-5}$	CL=90%	937
$p e^+ \mu^-$	LF	$< 9.9 \times 10^{-6}$	CL=90%	947
$p e^- \mu^+$	LF	$< 1.9 \times 10^{-5}$	CL=90%	947

$\bar{p}2e^+$	L,B	< 2.7	$\times 10^{-6}$	CL=90%	951
$\bar{p}2\mu^+$	L,B	< 9.4	$\times 10^{-6}$	CL=90%	937
$\bar{p}e^+\mu^+$	L,B	< 1.6	$\times 10^{-5}$	CL=90%	947
$\Sigma^-\mu^+\mu^+$	L	< 7.0	$\times 10^{-4}$	CL=90%	812

 $\Lambda_c(2595)^+$

$$I(J^P) = 0(\frac{1}{2}^-)$$

The spin-parity follows from the fact that $\Sigma_c(2455)\pi$ decays, with little available phase space, are dominant. This assumes that $J^P = 1/2^+$ for the $\Sigma_c(2455)$.

$$\begin{aligned} \text{Mass } m &= 2592.25 \pm 0.28 \text{ MeV} \\ m - m_{\Lambda_c^+} &= 305.79 \pm 0.24 \text{ MeV} \\ \text{Full width } \Gamma &= 2.6 \pm 0.6 \text{ MeV} \end{aligned}$$

$\Lambda_c^+\pi\pi$ and its submode $\Sigma_c(2455)\pi$ — the latter just barely — are the only strong decays allowed to an excited Λ_c^+ having this mass; and the submode seems to dominate.

$\Lambda_c(2595)^+$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_c^+\pi^+\pi^-$	[c] —	117
$\Sigma_c(2455)^{++}\pi^-$	$24 \pm 7\%$	†
$\Sigma_c(2455)^0\pi^+$	$24 \pm 7\%$	†
$\Lambda_c^+\pi^+\pi^-$ 3-body	$18 \pm 10\%$	117
$\Lambda_c^+\pi^0$	[d] not seen	258
$\Lambda_c^+\gamma$	not seen	288

 $\Lambda_c(2625)^+$

$$I(J^P) = 0(\frac{3}{2}^-)$$

J^P has not been measured; $\frac{3}{2}^-$ is the quark-model prediction.

$$\begin{aligned} \text{Mass } m &= 2628.11 \pm 0.19 \text{ MeV} \quad (S = 1.1) \\ m - m_{\Lambda_c^+} &= 341.65 \pm 0.13 \text{ MeV} \quad (S = 1.1) \\ \text{Full width } \Gamma &< 0.97 \text{ MeV, CL} = 90\% \end{aligned}$$

$\Lambda_c^+\pi\pi$ and its submode $\Sigma_c(2455)\pi$ are the only strong decays allowed to an excited Λ_c^+ having this mass.

$\Lambda_c(2625)^+$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$\Lambda_c^+\pi^+\pi^-$	$\approx 67\%$		184
$\Sigma_c(2455)^{++}\pi^-$	< 5	90%	102

$\Sigma_c(2455)^0 \pi^+$	<5	90%	102
$\Lambda_c^+ \pi^+ \pi^-$ 3-body	large		184
$\Lambda_c^+ \pi^0$	[d] not seen		293
$\Lambda_c^+ \gamma$	not seen		319

$\Lambda_c(2860)^+$

$$I(J^P) = 0(\frac{3}{2}^+)$$

Mass $m = 2856.1^{+2.3}_{-6.0}$ MeV

Full width $\Gamma = 68^{+12}_{-22}$ MeV

$\Lambda_c(2860)^+$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^0 p$	seen	259

$\Lambda_c(2880)^+$

$$I(J^P) = 0(\frac{5}{2}^+)$$

Mass $m = 2881.63 \pm 0.24$ MeV

$m - m_{\Lambda_c^+} = 595.17 \pm 0.28$ MeV

Full width $\Gamma = 5.6^{+0.8}_{-0.6}$ MeV

$\Lambda_c(2880)^+$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_c^+ \pi^+ \pi^-$	seen	471
$\Sigma_c(2455)^{0,++} \pi^\pm$	seen	376
$\Sigma_c(2520)^{0,++} \pi^\pm$	seen	317
$p D^0$	seen	316

$\Lambda_c(2940)^+$

$$I(J^P) = 0(\frac{3}{2}^-)$$

$J^P = 3/2^-$ is favored, but is not certain

Mass $m = 2939.6^{+1.3}_{-1.5}$ MeV

Full width $\Gamma = 20^{+6}_{-5}$ MeV

$\Lambda_c(2940)^+$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$p D^0$	seen	420
$\Sigma_c(2455)^{0,++} \pi^\pm$	seen	—

$\Sigma_c(2455)$

$$I(J^P) = 1(\frac{1}{2}^+)$$

$$\begin{aligned}
\Sigma_c(2455)^{++} \text{ mass } m &= 2453.97 \pm 0.14 \text{ MeV} \\
\Sigma_c(2455)^+ \text{ mass } m &= 2452.9 \pm 0.4 \text{ MeV} \\
\Sigma_c(2455)^0 \text{ mass } m &= 2453.75 \pm 0.14 \text{ MeV} \\
m_{\Sigma_c^{++}} - m_{\Lambda_c^+} &= 167.510 \pm 0.017 \text{ MeV} \\
m_{\Sigma_c^+} - m_{\Lambda_c^+} &= 166.4 \pm 0.4 \text{ MeV} \\
m_{\Sigma_c^0} - m_{\Lambda_c^+} &= 167.290 \pm 0.017 \text{ MeV} \\
m_{\Sigma_c^{++}} - m_{\Sigma_c^0} &= 0.220 \pm 0.013 \text{ MeV} \\
m_{\Sigma_c^+} - m_{\Sigma_c^0} &= -0.9 \pm 0.4 \text{ MeV} \\
\Sigma_c(2455)^{++} \text{ full width } \Gamma &= 1.89_{-0.18}^{+0.09} \text{ MeV} \quad (S = 1.1) \\
\Sigma_c(2455)^+ \text{ full width } \Gamma &< 4.6 \text{ MeV, CL} = 90\% \\
\Sigma_c(2455)^0 \text{ full width } \Gamma &= 1.83_{-0.19}^{+0.11} \text{ MeV} \quad (S = 1.2)
\end{aligned}$$

$\Lambda_c^+ \pi$ is the only strong decay allowed to a Σ_c having this mass.

$\Sigma_c(2455)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_c^+ \pi$	$\approx 100\%$	94

 $\Sigma_c(2520)$

$$I(J^P) = 1(\frac{3}{2}^+)$$

J^P has not been measured; $\frac{3}{2}^+$ is the quark-model prediction.

$$\begin{aligned}
\Sigma_c(2520)^{++} \text{ mass } m &= 2518.41_{-0.19}^{+0.21} \text{ MeV} \quad (S = 1.1) \\
\Sigma_c(2520)^+ \text{ mass } m &= 2517.5 \pm 2.3 \text{ MeV} \\
\Sigma_c(2520)^0 \text{ mass } m &= 2518.48 \pm 0.20 \text{ MeV} \quad (S = 1.1) \\
m_{\Sigma_c(2520)^{++}} - m_{\Lambda_c^+} &= 231.95_{-0.12}^{+0.17} \text{ MeV} \quad (S = 1.3) \\
m_{\Sigma_c(2520)^+} - m_{\Lambda_c^+} &= 231.0 \pm 2.3 \text{ MeV} \\
m_{\Sigma_c(2520)^0} - m_{\Lambda_c^+} &= 232.02_{-0.14}^{+0.15} \text{ MeV} \quad (S = 1.3) \\
m_{\Sigma_c(2520)^{++}} - m_{\Sigma_c(2520)^0} &= 0.01 \pm 0.15 \text{ MeV} \\
\Sigma_c(2520)^{++} \text{ full width } \Gamma &= 14.78_{-0.40}^{+0.30} \text{ MeV} \\
\Sigma_c(2520)^+ \text{ full width } \Gamma &< 17 \text{ MeV, CL} = 90\% \\
\Sigma_c(2520)^0 \text{ full width } \Gamma &= 15.3_{-0.5}^{+0.4} \text{ MeV}
\end{aligned}$$

$\Lambda_c^+ \pi$ is the only strong decay allowed to a Σ_c having this mass.

$\Sigma_c(2520)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_c^+ \pi$	$\approx 100\%$	179

$\Sigma_c(2800)$

$$I(J^P) = 1(?^?)$$

$$\begin{aligned} \Sigma_c(2800)^{++} \text{ mass } m &= 2801_{-6}^{+4} \text{ MeV} \\ \Sigma_c(2800)^+ \text{ mass } m &= 2792_{-5}^{+14} \text{ MeV} \\ \Sigma_c(2800)^0 \text{ mass } m &= 2806_{-7}^{+5} \text{ MeV} \quad (S = 1.3) \\ m_{\Sigma_c(2800)^{++}} - m_{\Lambda_c^+} &= 514_{-6}^{+4} \text{ MeV} \\ m_{\Sigma_c(2800)^+} - m_{\Lambda_c^+} &= 505_{-5}^{+14} \text{ MeV} \\ m_{\Sigma_c(2800)^0} - m_{\Lambda_c^+} &= 519_{-7}^{+5} \text{ MeV} \quad (S = 1.3) \\ \Sigma_c(2800)^{++} \text{ full width } \Gamma &= 75_{-17}^{+22} \text{ MeV} \\ \Sigma_c(2800)^+ \text{ full width } \Gamma &= 62_{-40}^{+60} \text{ MeV} \\ \Sigma_c(2800)^0 \text{ full width } \Gamma &= 72_{-15}^{+22} \text{ MeV} \end{aligned}$$

$\Sigma_c(2800)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_c^+ \pi$	seen	443

Ξ_c^+

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

$$\begin{aligned} \text{Mass } m &= 2467.87 \pm 0.30 \text{ MeV} \quad (S = 1.1) \\ \text{Mean life } \tau &= (442 \pm 26) \times 10^{-15} \text{ s} \quad (S = 1.3) \\ c\tau &= 132 \text{ } \mu\text{m} \end{aligned}$$

Branching fractions marked with a footnote, e.g. [a], have been corrected for decay modes not observed in the experiments. For example, the sub-mode fraction $\Xi_c^+ \rightarrow \Sigma^+ \bar{K}^*(892)^0$ seen in $\Xi_c^+ \rightarrow \Sigma^+ K^- \pi^+$ has been multiplied up to include $\bar{K}^*(892)^0 \rightarrow \bar{K}^0 \pi^0$ decays.

Ξ_c^+ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	P (MeV/c)
-----------------------	--------------------------------	------------------	-------------

**No absolute branching fractions have been measured.
The following are branching ratios relative to $\Xi^- 2\pi^+$.**

Cabibbo-favored ($S = -2$) decays — relative to $\Xi^- 2\pi^+$

$p 2K_S^0$	0.087 ± 0.021		767
$\Lambda \bar{K}^0 \pi^+$	—		852
$\Sigma(1385)^+ \bar{K}^0$	[a] 1.0 ± 0.5		746
$\Lambda K^- 2\pi^+$	0.323 ± 0.033		787
$\Lambda \bar{K}^*(892)^0 \pi^+$	[a] <0.16	90%	608
$\Sigma(1385)^+ K^- \pi^+$	[a] <0.23	90%	678

$\Sigma^+ K^- \pi^+$		0.94 ± 0.10		811
$\Sigma^+ \bar{K}^*(892)^0$	[a]	0.81 ± 0.15		658
$\Sigma^0 K^- 2\pi^+$		0.27 ± 0.12		735
$\Xi^0 \pi^+$		0.55 ± 0.16		877
$\Xi^- 2\pi^+$		DEFINED AS 1		851
$\Xi(1530)^0 \pi^+$	[a]	<0.10	90%	750
$\Xi^0 \pi^+ \pi^0$		2.3 ± 0.7		856
$\Xi^0 \pi^- 2\pi^+$		1.7 ± 0.5		818
$\Xi^0 e^+ \nu_e$		2.3 $\begin{smallmatrix} +0.7 \\ -0.8 \end{smallmatrix}$		884
$\Omega^- K^+ \pi^+$		0.07 ± 0.04		399

Cabibbo-suppressed decays — relative to $\Xi^- 2\pi^+$

$\rho K^- \pi^+$		0.21 ± 0.04		944
$\rho \bar{K}^*(892)^0$	[a]	0.116 ± 0.030		828
$\Sigma^+ \pi^+ \pi^-$		0.48 ± 0.20		922
$\Sigma^- 2\pi^+$		0.18 ± 0.09		918
$\Sigma^+ K^+ K^-$		0.15 ± 0.06		579
$\Sigma^+ \phi$	[a]	<0.11	90%	549
$\Xi(1690)^0 K^+, \Xi^0 \rightarrow \Sigma^+ K^-$		<0.05	90%	501



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

$$\text{Mass } m = 2470.87^{+0.28}_{-0.31} \text{ MeV}$$

$$m_{\Xi_c^0} - m_{\Xi_c^+} = 3.00 \pm 0.24 \text{ MeV}$$

$$\text{Mean life } \tau = (112^{+13}_{-10}) \times 10^{-15} \text{ s}$$

$$c\tau = 33.6 \text{ } \mu\text{m}$$

Decay asymmetry parameters

$$\Xi^- \pi^+ \quad \alpha = -0.6 \pm 0.4$$

Branching fractions marked with a footnote, e.g. [a], have been corrected for decay modes not observed in the experiments. For example, the sub-mode fraction $\Xi_c^0 \rightarrow p K^- \bar{K}^*(892)^0$ seen in $\Xi_c^0 \rightarrow p K^- K^- \pi^+$ has been multiplied up to include $\bar{K}^*(892)^0 \rightarrow \bar{K}^0 \pi^0$ decays.

Ξ_c^0 DECAY MODES Fraction (Γ_i/Γ) p (MeV/c)


**No absolute branching fractions have been measured.
The following are branching *ratios* relative to $\Xi^- \pi^+$.**

Cabibbo-favored (S = -2) decays — relative to $\Xi^- \pi^+$

$p K^- K^- \pi^+$	0.34 ± 0.04	676
$p K^- \bar{K}^*(892)^0$	[a] 0.21 ± 0.05	413
$p K^- K^- \pi^+$ (no \bar{K}^{*0})	0.21 ± 0.04	676
ΛK_S^0	0.210 ± 0.028	906
$\Lambda K^- \pi^+$	1.07 ± 0.14	856
$\Lambda \bar{K}^0 \pi^+ \pi^-$	seen	787
$\Lambda K^- \pi^+ \pi^+ \pi^-$	seen	703
$\Xi^- \pi^+$	DEFINED AS 1	875
$\Xi^- \pi^+ \pi^+ \pi^-$	3.3 ± 1.4	816
$\Omega^- K^+$	0.297 ± 0.024	522
$\Xi^- e^+ \nu_e$	3.1 ± 1.1	882
$\Xi^- \ell^+$ anything	1.0 ± 0.5	—

Cabibbo-suppressed decays — relative to $\Xi^- \pi^+$

$\Xi^- K^+$	0.028 ± 0.006	790
$\Lambda K^+ K^-$ (no ϕ)	0.029 ± 0.007	648
$\Lambda \phi$	[a] 0.034 ± 0.007	621



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

Mass $m = 2577.4 \pm 1.2$ MeV (S = 2.9)

$$m_{\Xi_c^{'+}} - m_{\Xi_c^+} = 109.5 \pm 1.2 \text{ MeV} \quad (S = 3.7)$$

$$m_{\Xi_c^{'+}} - m_{\Xi_c^0} = -1.4 \pm 1.3 \text{ MeV} \quad (S = 2.5)$$

The $\Xi_c^{'+} - \Xi_c^+$ mass difference is too small for any strong decay to occur.

$\Xi_c^{'+}$ DECAY MODES Fraction (Γ_i/Γ) p (MeV/c)

$\Xi_c^+ \gamma$	seen	107
------------------	------	-----

$\Xi_c^{\prime 0}$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

$$\text{Mass } m = 2578.8 \pm 0.5 \text{ MeV} \quad (S = 1.2)$$

$$m_{\Xi_c^{\prime 0}} - m_{\Xi_c^0} = 108.0 \pm 0.4 \text{ MeV} \quad (S = 1.2)$$

The $\Xi_c^{\prime 0} - \Xi_c^0$ mass difference is too small for any strong decay to occur.

$\Xi_c^{\prime 0}$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c^{\prime 0} \gamma$	seen	106

 $\Xi_c(2645)$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^+)$$

J^P has not been measured; $\frac{3}{2}^+$ is the quark-model prediction.

$$\Xi_c(2645)^+ \text{ mass } m = 2645.53 \pm 0.31 \text{ MeV}$$

$$\Xi_c(2645)^0 \text{ mass } m = 2646.32 \pm 0.31 \text{ MeV} \quad (S = 1.1)$$

$$m_{\Xi_c(2645)^+} - m_{\Xi_c^0} = 174.66 \pm 0.09 \text{ MeV}$$

$$m_{\Xi_c(2645)^0} - m_{\Xi_c^+} = 178.44 \pm 0.11 \text{ MeV} \quad (S = 1.1)$$

$$m_{\Xi_c(2645)^+} - m_{\Xi_c(2645)^0} = -0.79 \pm 0.27 \text{ MeV}$$

$$\Xi_c(2645)^+ \text{ full width } \Gamma = 2.14 \pm 0.19 \text{ MeV} \quad (S = 1.1)$$

$$\Xi_c(2645)^0 \text{ full width } \Gamma = 2.35 \pm 0.22 \text{ MeV}$$

$\Xi_c \pi$ is the only strong decay allowed to a Ξ_c resonance having this mass.

$\Xi_c(2645)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c^0 \pi^+$	seen	102
$\Xi_c^+ \pi^-$	seen	106

 $\Xi_c(2790)$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^-)$$

J^P has not been measured; $\frac{1}{2}^-$ is the quark-model prediction.

$$\Xi_c(2790)^+ \text{ mass } = 2792.0 \pm 0.5 \text{ MeV} \quad (S = 1.2)$$

$$\Xi_c(2790)^0 \text{ mass } = 2792.8 \pm 1.2 \text{ MeV} \quad (S = 2.9)$$

$$m_{\Xi_c(2790)^+} - m_{\Xi_c^0} = 321.1 \pm 0.4 \text{ MeV} \quad (S = 1.2)$$

$$m_{\Xi_c(2790)^0} - m_{\Xi_c^+} = 324.9 \pm 1.2 \text{ MeV} \quad (S = 3.7)$$

$$m_{\Xi_c(2790)^+} - m_{\Xi_c^0} = 213.10 \pm 0.26 \text{ MeV} \quad (S = 1.2)$$

$$\begin{aligned}
 m_{\Xi_c(2790)^0} - m_{\Xi_c'} &= 215.4 \pm 0.8 \text{ MeV} \quad (S = 3.7) \\
 m_{\Xi_c(2790)^+} - m_{\Xi_c(2790)^0} &= -0.9 \pm 1.3 \text{ MeV} \quad (S = 2.5) \\
 \Xi_c(2790)^+ \text{ width} &= 8.9 \pm 1.0 \text{ MeV} \\
 \Xi_c(2790)^0 \text{ width} &= 10.0 \pm 1.1 \text{ MeV}
 \end{aligned}$$

$\Xi_c(2790)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c \pi$	seen	—
$\Xi_c' \pi$	seen	160

$\Xi_c(2815)$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$$

J^P has not been measured; $\frac{3}{2}^-$ is the quark-model prediction.

$$\begin{aligned}
 \Xi_c(2815)^+ \text{ mass } m &= 2816.67 \pm 0.31 \text{ MeV} \quad (S = 1.1) \\
 \Xi_c(2815)^0 \text{ mass } m &= 2820.22 \pm 0.32 \text{ MeV} \\
 m_{\Xi_c(2815)^+} - m_{\Xi_c^+} &= 348.80 \pm 0.10 \text{ MeV} \\
 m_{\Xi_c(2815)^0} - m_{\Xi_c^0} &= 349.35 \pm 0.11 \text{ MeV} \\
 m_{\Xi_c(2815)^+} - m_{\Xi_c(2815)^0} &= -3.55 \pm 0.28 \text{ MeV} \\
 \Xi_c(2815)^+ \text{ full width } \Gamma &= 2.43 \pm 0.26 \text{ MeV} \\
 \Xi_c(2815)^0 \text{ full width } \Gamma &= 2.54 \pm 0.25 \text{ MeV}
 \end{aligned}$$

The $\Xi_c \pi \pi$ modes are consistent with being entirely via $\Xi_c(2645)\pi$.

$\Xi_c(2815)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c \pi$	seen	—
$\Xi_c^+ \pi^+ \pi^-$	seen	196
$\Xi_c^0 \pi^+ \pi^-$	seen	191

$\Xi_c(2970)$

$$I(J^P) = \frac{1}{2}(??)$$

$$\begin{aligned}
 \Xi_c(2970)^+ m &= 2969.4 \pm 0.8 \text{ MeV} \quad (S = 1.1) \\
 \Xi_c(2970)^0 m &= 2967.8 \pm 0.8 \text{ MeV} \quad (S = 1.1) \\
 m_{\Xi_c(2970)^+} - m_{\Xi_c^0} &= 498.5 \pm 0.8 \text{ MeV} \quad (S = 1.1) \\
 m_{\Xi_c(2970)^0} - m_{\Xi_c^+} &= 499.9^{+0.8}_{-0.7} \text{ MeV} \quad (S = 1.1) \\
 m_{\Xi_c(2970)^+} - m_{\Xi_c(2970)^0} &= 1.6 \pm 1.1 \text{ MeV} \quad (S = 1.1) \\
 \Xi_c(2970)^+ \text{ width } \Gamma &= 20.9^{+2.4}_{-3.5} \text{ MeV} \quad (S = 1.2) \\
 \Xi_c(2970)^0 \text{ width } \Gamma &= 28.1^{+3.4}_{-4.0} \text{ MeV} \quad (S = 1.5)
 \end{aligned}$$

$\Xi_c(2970)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_c^+ \bar{K} \pi$	seen	231
$\Sigma_c(2455) \bar{K}$	seen	133
$\Lambda_c^+ \bar{K}$	not seen	414
$\Xi_c 2\pi$	seen	385
$\Xi_c(2645) \pi$	seen	277

$\Xi_c(3055)$

$$I(J^P) = ?(??)$$

Mass $m = 3055.9 \pm 0.4$ MeV

Full width $\Gamma = 7.8 \pm 1.9$ MeV

$\Xi_c(3055)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Sigma^{++} K^-$	seen	—
ΛD^+	seen	316

$\Xi_c(3080)$

$$I(J^P) = \frac{1}{2}(??)$$

$\Xi_c(3080)^+ m = 3077.2 \pm 0.4$ MeV

$\Xi_c(3080)^0 m = 3079.9 \pm 1.4$ MeV (S = 1.3)

$\Xi_c(3080)^+ \text{ width } \Gamma = 3.6 \pm 1.1$ MeV (S = 1.5)

$\Xi_c(3080)^0 \text{ width } \Gamma = 5.6 \pm 2.2$ MeV

$\Xi_c(3080)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_c^+ \bar{K} \pi$	seen	415
$\Sigma_c(2455) \bar{K}$	seen	342
$\Sigma_c(2455)^{++} K^-$	seen	342
$\Sigma_c(2520)^{++} K^-$	seen	239
$\Sigma_c(2455) \bar{K} + \Sigma_c(2520) \bar{K}$	seen	—
$\Lambda_c^+ \bar{K}$	not seen	536
$\Lambda_c^+ \bar{K} \pi^+ \pi^-$	not seen	144
ΛD^+	seen	362

Ω_c^0

$$I(J^P) = 0(\frac{1}{2}^+)$$

J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

Mass $m = 2695.2 \pm 1.7$ MeV (S = 1.3)

Mean life $\tau = (69 \pm 12) \times 10^{-15}$ s

$c\tau = 21$ μm

Ω_c^0 DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
--------------------------	--------------------------------	------------------	-------------

**No absolute branching fractions have been measured.
The following are branching *ratios* relative to $\Omega^- \pi^+$.**

Cabibbo-favored ($S = -3$) decays — relative to $\Omega^- \pi^+$

$\Omega^- \pi^+$	DEFINED AS 1		821
$\Omega^- \pi^+ \pi^0$	1.80 ± 0.33		797
$\Omega^- \rho^+$	> 1.3	90%	532
$\Omega^- \pi^- 2\pi^+$	0.31 ± 0.05		753
$\Omega^- e^+ \nu_e$	2.4 ± 1.2		829
$\Xi^0 \bar{K}^0$	1.64 ± 0.29		950
$\Xi^0 K^- \pi^+$	1.20 ± 0.18		901
$\Xi^0 \bar{K}^{*0}, \bar{K}^{*0} \rightarrow K^- \pi^+$	0.68 ± 0.16		764
$\Xi^- \bar{K}^0 \pi^+$	2.12 ± 0.28		895
$\Xi^- K^- 2\pi^+$	0.63 ± 0.09		830
$\Xi(1530)^0 K^- \pi^+, \Xi^{*0} \rightarrow$	0.21 ± 0.06		757
$\Xi^- \bar{K}^{*0} \pi^+$	0.34 ± 0.11		653
$\Sigma^+ K^- K^- \pi^+$	< 0.32	90%	689
$\Lambda \bar{K}^0 \bar{K}^0$	1.72 ± 0.35		837

$\Omega_c(2770)^0$

$$I(J^P) = 0(\frac{3}{2}^+)$$

J^P has not been measured; $\frac{3}{2}^+$ is the quark-model prediction.

$$\text{Mass } m = 2765.9 \pm 2.0 \text{ MeV } (S = 1.2)$$

$$m_{\Omega_c(2770)^0} - m_{\Omega_c^0} = 70.7^{+0.8}_{-0.9} \text{ MeV}$$

The $\Omega_c(2770)^0 - \Omega_c^0$ mass difference is too small for any strong decay to occur.

$\Omega_c(2770)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Omega_c^0 \gamma$	presumably 100%	70

$\Omega_c(3000)^0$

$$I(J^P) = ?(??)$$

$$\text{Mass } m = 3000.4 \pm 0.4 \text{ MeV}$$

$$\text{Full width } \Gamma = 4.5 \pm 0.7 \text{ MeV}$$

$\Omega_c(3000)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c^+ K^-$	seen	181

 $\Omega_c(3050)^0$

$$I(J^P) = ?(??)$$

Mass $m = 3050.2 \pm 0.33$ MeVFull width $\Gamma < 1.2$ MeV, CL = 95%

$\Omega_c(3050)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c^+ K^-$	seen	278

 $\Omega_c(3065)^0$

$$I(J^P) = ?(??)$$

Mass $m = 3065.6 \pm 0.4$ MeVFull width $\Gamma = 3.5 \pm 0.4$ MeV

$\Omega_c(3065)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c^+ K^-$	seen	303

 $\Omega_c(3090)^0$

$$I(J^P) = ?(??)$$

Mass $m = 3090.2 \pm 0.7$ MeVFull width $\Gamma = 8.7 \pm 1.3$ MeV

$\Omega_c(3090)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c^+ K^-$	seen	339

 $\Omega_c(3120)^0$

$$I(J^P) = ?(??)$$

Mass $m = 3119.1 \pm 1.0$ MeVFull width $\Gamma < 2.6$ MeV, CL = 95%

$\Omega_c(3120)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c^+ K^-$	seen	379

NOTES

- [a] This branching fraction includes all the decay modes of the final-state resonance.
- [b] The value is for the sum of the charge states or particle/antiparticle states indicated.
- [c] See AALTONEN 11H, Fig. 8, for the calculated ratio of $\Lambda_c^+ \pi^0 \pi^0$ and $\Lambda_c^+ \pi^+ \pi^-$ partial widths as a function of the $\Lambda_c(2595)^+ - \Lambda_c^+$ mass difference. At our value of the mass difference, the ratio is about 4.
- [d] A test that the isospin is indeed 0, so that the particle is indeed a Λ_c^+ .