

Z_c(3900)

I^G(J^{PC}) = 1⁺(1⁺⁻)

was X(3900)

Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on non- $q\bar{q}$ states.

Charged Z_c(3900) seen as a peak in the invariant mass distribution of the J/ ψ π^\pm system by BES III (ABLIKIM 13T) in $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ at c.m. energy of 4.26 GeV and by radiative return from

e^+e^- collisions at \sqrt{s} from 9.46 to 10.86 GeV at Belle (LIU 13B).

Partial wave analysis of ABLIKIM 17J determines $J^P = 1^+$ with more than 7σ significance. Neutral Z_c(3900) seen in the J/ $\psi\pi^0$ invariant mass distribution in $e^+e^- \rightarrow \pi^0\pi^0 J/\psi$ at c.m. energies of 4.23, 4.26, and 4.36 GeV by BES III (ABLIKIM 15U) and at 4.17 GeV by XIAO 13A. Peaks in $(D\bar{D}^*)^{0,\pm}$ reported by BES III (ABLIKIM 14A, ABLIKIM 15AB) are assumed to be related.

Z_c(3900) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
3887.2±2.3 OUR AVERAGE		Error includes scale factor of 1.6. See the ideogram below.			
3895.0±5.2 ^{+ 4.0} _{- 2.7}	502	1 ABAZOV	18B D0		1.96 TeV $p\bar{p} \rightarrow J/\psi\pi^+\pi^- X$
3885.7 ^{+4.3} _{-5.7} ± 8.4		2 ABLIKIM	15AB BES3	0	$e^+e^- \rightarrow \pi^0(D\bar{D}^*)^0$
3881.7±1.6± 1.6	1.2k	2 ABLIKIM	15AC BES3	±	$e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$
3894.8±2.3± 3.2	356	2 ABLIKIM	15U BES3	0	$e^+e^- \rightarrow \pi^0\pi^0 J/\psi$
3883.9±1.5± 4.2	1.2k	2 ABLIKIM	14A BES3	±	$e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$
3899.0±3.6± 4.9	307	2 ABLIKIM	13T BES3	±	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
3894.5±6.6± 4.5	159	2 LIU	13B BELL	±	$e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$
3886 ± 4 ± 2	81	2,3 XIAO	13A	±	4.17 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$
3904 ± 9 ± 5	25	2,3 XIAO	13A	0	4.17 $e^+e^- \rightarrow \pi^0\pi^0 J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •


3881.2±4.2±52.7 6k ⁴ ABLIKIM 17J BES3 ± $e^+e^- \rightarrow \pi^+\pi^- J/\psi$

¹ The signal of the Z_c(3900) is correlated with a parent J/ $\psi\pi^+\pi^-$ system in the invariant mass range 4.2–4.7 GeV.

² Neglecting interference between the Z_c(3900) and non-resonant continuum.

³ For $M^2(\pi^+\pi^-) < 0.65$ GeV². Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

⁴ Pole mass obtained from a fit to a Flatte-like formula.



Z_c(3900) WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
28.2± 2.6 OUR AVERAGE					
51.8± 4.6± 36.0	6 k	1 ABLIKIM	17J BES3	±	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$
35 ±11 ±12	±15	2 ABLIKIM	15AB BES3	0	$e^+ e^- \rightarrow \pi^0 (D\bar{D}^*)^0$
26.6± 2.0± 2.1	1248	2 ABLIKIM	15AC BES3	±	$e^+ e^- \rightarrow \pi^\pm (D\bar{D}^*)^\mp$
29.6± 8.2± 8.2	356	2 ABLIKIM	15U BES3	0	$e^+ e^- \rightarrow \pi^0 \pi^0 J/\psi$
24.8± 3.3± 11.0	1212	2 ABLIKIM	14A BES3	±	$e^+ e^- \rightarrow \pi^\pm (D\bar{D}^*)^\mp$
46 ±10 ±20	307	2 ABLIKIM	13T BES3	±	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$
63 ±24 ±26	159	2 LIU	13B BELL	±	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$
37 ± 4 ± 8	81	2,3 XIAO	13A	±	$4.17 e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$

¹ Pole width obtained from a fit to a Flatte-like formula.

² Neglecting interference between the Z_c(3900) and non-resonant continuum.

³ For $M^2(\pi^+ \pi^-) < 0.65 \text{ GeV}^2$. Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

Z_c(3900) DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 J/\psi \pi$	seen
$\Gamma_2 h_c \pi^\pm$	not seen
$\Gamma_3 \eta_c \pi^+ \pi^-$	not seen

Γ_4	$(D\bar{D}^*)^\pm$	seen
Γ_5	$D^0 D^{*-} + \text{c.c.}$	seen
Γ_6	$D^- D^{*0} + \text{c.c.}$	seen
Γ_7	$\omega\pi^\pm$	not seen
Γ_8	$J/\psi\eta$	not seen
Γ_9	$D^+ D^{*-} + \text{c.c.}$	seen
Γ_{10}	$D^0 \bar{D}^{*0} + \text{c.c.}$	seen

 $Z_c(3900)$ BRANCHING RATIOS **$\Gamma(J/\psi\pi)/\Gamma_{\text{total}}$** **$\Gamma_1/\Gamma$**

VALUE	CL%	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
seen	356		ABLIKIM	15U	BES3	0
seen	307		ABLIKIM	13T	BES3	\pm
seen	25	¹ XIAO		13A		$e^+ e^- \rightarrow \pi^0 \pi^0 J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen 90 ² ADOLPH 15D COMP \pm $\gamma N \rightarrow J/\psi\pi^\pm N$ ¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.² ADOLPH 15D measure $B(Z_c(3900)^\pm \rightarrow J/\psi\pi^\pm) \sigma(\gamma N \rightarrow Z_c(3900)^\pm N) / \sigma(\gamma N \rightarrow J/\psi N) < 3.7 \times 10^{-3}$ at 90% CL. **$\Gamma(h_c\pi^\pm)/\Gamma_{\text{total}}$** **$\Gamma_2/\Gamma$**

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
not seen	ABLIKIM	13X	BES3	\pm $e^+ e^- \rightarrow h_c\pi^+\pi^-$

 $\Gamma(\eta_c\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_3/Γ**

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
not seen	¹ VINOKUROVA 15	BELL	0	$B^+ \rightarrow K^+ \eta_c\pi^+\pi^-$

¹ VINOKUROVA 15 reports $B(B^+ \rightarrow K^+ Z_c(3900)^0) \times B(X \rightarrow \eta_c\pi^+\pi^-) < 4.7 \times 10^{-5}$ at 90% CL. **$\Gamma((D\bar{D}^*)^\pm)/\Gamma(J/\psi\pi)$** **$\Gamma_4/\Gamma_1$**

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
6.2±1.1±2.7	¹ ABLIKIM	14A	BES3	\pm $e^+ e^- \rightarrow \pi^\pm (D\bar{D}^*)^\mp$

¹ Assuming the same origin of the $(D\bar{D}^*)^\pm$ and $\pi^\pm J/\psi$ decay modes. **$\Gamma(D^0 D^{*-} + \text{c.c.})/\Gamma_{\text{total}}$** **$\Gamma_5/\Gamma$**

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
seen	ABLIKIM	15AC	BES3	\pm $e^+ e^- \rightarrow \pi^+ D^0 D^{*-} + \text{c.c.}$
seen	ABLIKIM	14A	BES3	\pm $e^+ e^- \rightarrow \pi^+ D^0 D^{*-} + \text{c.c.}$

 $\Gamma(D^- D^{*0} + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_6/Γ**

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
seen	ABLIKIM	15AC	BES3	\pm $e^+ e^- \rightarrow \pi^+ D^- D^{*0} + \text{c.c.}$
seen	ABLIKIM	14A	BES3	\pm $e^+ e^- \rightarrow \pi^+ D^- D^{*0} + \text{c.c.}$

$\Gamma(\omega\pi^\pm)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_7/Γ
not seen	ABLIKIM	15R	BES3	\pm	$e^+ e^- \rightarrow \omega\pi^+\pi^-$

 $\Gamma(J/\psi\eta)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_8/Γ
not seen	ABLIKIM	15Q	BES3	0	$4.0\text{--}4.6 e^+ e^- \rightarrow J/\psi\eta\pi^0$

 $\Gamma(J/\psi\eta)/\Gamma(J/\psi\pi)$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_8/Γ_1
<0.15	90	ABLIKIM	15Q	BES3	0	$4.226 e^+ e^- \rightarrow J/\psi\eta\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.65	90	ABLIKIM	15Q	BES3	0	$4.257 e^+ e^- \rightarrow J/\psi\eta\pi^0$

 $\Gamma(D^+D^{*-} + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_9/Γ
seen	ABLIKIM	15AB	BES3	0	$e^+ e^- \rightarrow \pi^0(D\bar{D}^*)^0$

 $\Gamma(D^0\bar{D}^{*0} + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_{10}/Γ
seen	ABLIKIM	15AB	BES3	0	$e^+ e^- \rightarrow \pi^0(D\bar{D}^*)^0$

 $\Gamma(D^+D^{*-} + \text{c.c.})/\Gamma(D^0\bar{D}^{*0} + \text{c.c.})$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_9/Γ_{10}
0.96±0.18±0.12	ABLIKIM	15AB	BES3	0	$e^+ e^- \rightarrow \pi^0(D\bar{D}^*)^0$

 $Z_c(3900)$ REFERENCES

ABAZOV	18B	PR D98 052010	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABLIKIM	17J	PRL 119 072001	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	15AB	PRL 115 222002	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	15AC	PR D92 092006	M. Ablikim <i>et al.</i>	(BES III Collab.) JP
ABLIKIM	15Q	PR D92 012008	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	15R	PR D92 032009	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	15U	PRL 115 112003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ADOLPH	15D	PL B742 330	C. Adolph <i>et al.</i>	(COMPASS Collab.)
VINOKUROVA	15	JHEP 1506 132	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
Also		JHEP 1702 088 (errat.)	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
ABLIKIM	14A	PRL 112 022001	M. Ablikim <i>et al.</i>	(BES III Collab.) JP
ABLIKIM	13T	PRL 110 252001	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13X	PRL 111 242001	M. Ablikim <i>et al.</i>	(BES III Collab.)
LIU	13B	PRL 110 252002	Z.Q. Liu <i>et al.</i>	(BELLE Collab.)
XIAO	13A	PL B727 366	T. Xiao <i>et al.</i>	(NWES)