



$I(J^P) = 0(\frac{1}{2}^+)$ Status: ****

The parity of the Λ_c^+ is defined to be positive (as are the parities of the proton, neutron, and Λ). The quark content is $u d c$. Results of an analysis of $p K^- \pi^+$ decays (JEZABEK 92) are consistent with $J = 1/2$. Nobody doubts that the spin is indeed 1/2.

We have omitted some results that have been superseded by later experiments. The omitted results may be found in earlier editions.

Λ_c^+ MASS

Our value in 2004, 2284.9 ± 0.6 MeV, was the average of the measurements now filed below as “not used.” The BABAR measurement is so much better that we use it alone. Note that it is about 2.6 (old) standard deviations above the 2004 value.

The fit also includes $\Sigma_c - \Lambda_c^+$ and $\Lambda_c^{*+} - \Lambda_c^+$ mass-difference measurements, but this doesn’t affect the Λ_c^+ mass. The new (in 2006) Λ_c^+ mass simply pushes all those other masses higher.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2286.46 ± 0.14 OUR FIT				
2286.46 ± 0.14	4891	¹ AUBERT,B	05S BABR	$\Lambda K_S^0 K^+$ and $\Sigma^0 K_S^0 K^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2284.7 ± 0.6 ± 0.7	1134	AVERY	91 CLEO	Six modes
2281.7 ± 2.7 ± 2.6	29	ALVAREZ	90B NA14	$p K^- \pi^+$
2285.8 ± 0.6 ± 1.2	101	BARLAG	89 NA32	$p K^- \pi^+$
2284.7 ± 2.3 ± 0.5	5	AGUILAR-...	88B LEBC	$p K^- \pi^+$
2283.1 ± 1.7 ± 2.0	628	ALBRECHT	88C ARG	$p K^- \pi^+, p \bar{K}^0, \Lambda 3\pi$
2286.2 ± 1.7 ± 0.7	97	ANJOS	88B E691	$p K^- \pi^+$
2281 ± 3	2	JONES	87 HBC	$p K^- \pi^+$
2283 ± 3	3	BOSETTI	82 HBC	$p K^- \pi^+$
2290 ± 3	1	CALICCHIO	80 HYBR	$p K^- \pi^+$


¹AUBERT,B 05S uses low-Q $\Lambda K_S^0 K^+$ and $\Sigma^0 K_S^0 K^+$ decays to minimize systematic errors. The error above includes systematic as well as statistical errors. Many cross checks and adjustments to properties of the BABAR detector, as well as the large number of clean events, make this by far the best measurement of the Λ_c^+ mass.

Λ_c^+ MEAN LIFE

Measurements with an error $\geq 100 \times 10^{-15}$ s or with fewer than 20 events have been omitted from the Listings.

VALUE (10^{-15} s)	EVTS	DOCUMENT ID	TECN	COMMENT
200 ± 6 OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.				
204.6 $\pm 3.4 \pm 2.5$	8034	LINK	02C FOCS	$p K^- \pi^+$
198.1 $\pm 7.0 \pm 5.6$	1630	KUSHNIR...	01 SELX	$\Lambda_c^+ \rightarrow p K^- \pi^+$

$179.6 \pm 6.9 \pm 4.4$	4749	MAHMOOD	01	CLE2	$e^+ e^- \approx \gamma(4S)$
$215 \pm 16 \pm 8$	1340	FRABETTI	93D	E687	$\gamma Be, \Lambda_c^+ \rightarrow p K^- \pi^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$180 \pm 30 \pm 30$	29	ALVAREZ	90	NA14	$\gamma, \Lambda_c^+ \rightarrow p K^- \pi^+$
$200 \pm 30 \pm 30$	90	FRABETTI	90	E687	$\gamma Be, \Lambda_c^+ \rightarrow p K^- \pi^+$
196^{+23}_{-20}	101	BARLAG	89	NA32	$p K^- \pi^+ + c.c.$
$220 \pm 30 \pm 20$	97	ANJOS	88B	E691	$p K^- \pi^+ + c.c.$



Λ_c^+ DECAY MODES

Branching fractions marked with a footnote, e.g. [a], have been corrected for decay modes not observed in the experiments. For example, the sub-mode fraction $\Lambda_c^+ \rightarrow p \bar{K}^*(892)^0$ seen in $\Lambda_c^+ \rightarrow p K^- \pi^+$ has been multiplied up to include $\bar{K}^*(892)^0 \rightarrow \bar{K}^0 \pi^0$ decays.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Hadronic modes with a p or n: $S = -1$ final states		
$\Gamma_1 \quad p K_S^0$	(1.59 ± 0.08) %	$S=1.1$
$\Gamma_2 \quad p K^- \pi^+$	(6.28 ± 0.32) %	$S=1.4$
$\Gamma_3 \quad p \bar{K}^*(892)^0$	[a] (1.96 ± 0.27) %	
$\Gamma_4 \quad \Delta(1232)^{++} K^-$	(1.08 ± 0.25) %	
$\Gamma_5 \quad \Lambda(1520) \pi^+$	[a] (2.2 ± 0.5) %	

Γ_6	$pK^-\pi^+$ nonresonant	(3.5 \pm 0.4) %	
Γ_7	$pK_S^0\pi^0$	(1.97 \pm 0.13) %	S=1.1
Γ_8	$nK_S^0\pi^+$	(1.82 \pm 0.25) %	
Γ_9	$p\bar{K}^0\eta$	(1.6 \pm 0.4) %	
Γ_{10}	$pK_S^0\pi^+\pi^-$	(1.60 \pm 0.12) %	S=1.1
Γ_{11}	$pK^-\pi^+\pi^0$	(4.46 \pm 0.30) %	S=1.5
Γ_{12}	$pK^*(892)^-\pi^+$	[a] (1.4 \pm 0.5) %	
Γ_{13}	$p(K^-\pi^+)_{\text{nonresonant}}\pi^0$	(4.6 \pm 0.8) %	
Γ_{14}	$\Delta(1232)\bar{K}^*(892)$	seen	
Γ_{15}	$pK^-2\pi^+\pi^-$	(1.4 \pm 0.9) $\times 10^{-3}$	
Γ_{16}	$pK^-\pi^+2\pi^0$	(1.0 \pm 0.5) %	

Hadronic modes with a p : $S = 0$ final states

Γ_{17}	$p\pi^0$	< 2.7 $\times 10^{-4}$	CL=90%
Γ_{18}	$p\eta$	(1.24 \pm 0.30) $\times 10^{-3}$	
Γ_{19}	$p\omega(782)^0$	(9 \pm 4) $\times 10^{-4}$	
Γ_{20}	$p\pi^+\pi^-$	(4.61 \pm 0.28) $\times 10^{-3}$	
Γ_{21}	$p f_0(980)$	[a] (3.5 \pm 2.3) $\times 10^{-3}$	
Γ_{22}	$p2\pi^+2\pi^-$	(2.3 \pm 1.4) $\times 10^{-3}$	
Γ_{23}	pK^+K^-	(1.06 \pm 0.06) $\times 10^{-3}$	
Γ_{24}	$p\phi$	[a] (1.06 \pm 0.14) $\times 10^{-3}$	
Γ_{25}	pK^+K^- non- ϕ	(5.3 \pm 1.2) $\times 10^{-4}$	
Γ_{26}	$p\phi\pi^0$	(10 \pm 4) $\times 10^{-5}$	
Γ_{27}	$pK^+K^-\pi^0$ nonresonant	< 6.3 $\times 10^{-5}$	CL=90%

Hadronic modes with a hyperon: $S = -1$ final states

Γ_{28}	$\Lambda\pi^+$	(1.30 \pm 0.07) %	S=1.1
Γ_{29}	$\Lambda\pi^+\pi^0$	(7.1 \pm 0.4) %	S=1.1
Γ_{30}	$\Lambda\rho^+$	< 6 %	CL=95%
Γ_{31}	$\Lambda\pi^-2\pi^+$	(3.64 \pm 0.29) %	S=1.4
Γ_{32}	$\Sigma(1385)^+\pi^+\pi^-$, $\Sigma^{*+} \rightarrow \Lambda\pi^+$	(1.0 \pm 0.5) %	
Γ_{33}	$\Sigma(1385)^-2\pi^+$, $\Sigma^{*-} \rightarrow \Lambda\pi^-$	(7.6 \pm 1.4) $\times 10^{-3}$	
Γ_{34}	$\Lambda\pi^+\rho^0$	(1.5 \pm 0.6) %	
Γ_{35}	$\Sigma(1385)^+\rho^0$, $\Sigma^{*+} \rightarrow \Lambda\pi^+$	(5 \pm 4) $\times 10^{-3}$	
Γ_{36}	$\Lambda\pi^-2\pi^+$ nonresonant	< 1.1 %	CL=90%
Γ_{37}	$\Lambda\pi^-\pi^02\pi^+$ total	(2.3 \pm 0.8) %	
Γ_{38}	$\Lambda\pi^+\eta$	[a] (2.2 \pm 0.5) %	
Γ_{39}	$\Sigma(1385)^+\eta$	[a] (1.07 \pm 0.32) %	
Γ_{40}	$\Lambda\pi^+\omega$	[a] (1.5 \pm 0.5) %	
Γ_{41}	$\Lambda\pi^-\pi^02\pi^+$, no η or ω	< 8 $\times 10^{-3}$	CL=90%
Γ_{42}	$\Lambda K^+\bar{K}^0$	(5.7 \pm 1.1) $\times 10^{-3}$	S=1.9
Γ_{43}	$\Xi(1690)^0K^+$, $\Xi^{*0} \rightarrow \Lambda\bar{K}^0$	(1.6 \pm 0.5) $\times 10^{-3}$	
Γ_{44}	$\Sigma^0\pi^+$	(1.29 \pm 0.07) %	S=1.1
Γ_{45}	$\Sigma^+\pi^0$	(1.25 \pm 0.10) %	

Γ_{46}	$\Sigma^+ \eta$	$(6.9 \pm 2.3) \times 10^{-3}$	
Γ_{47}	$\Sigma^+ \pi^+ \pi^-$	$(4.50 \pm 0.25) \%$	S=1.3
Γ_{48}	$\Sigma^+ \rho^0$	$< 1.7 \%$	CL=95%
Γ_{49}	$\Sigma^- 2\pi^+$	$(1.87 \pm 0.18) \%$	
Γ_{50}	$\Sigma^0 \pi^+ \pi^0$	$(3.5 \pm 0.4) \%$	
Γ_{51}	$\Sigma^+ \pi^0 \pi^0$	$(1.55 \pm 0.15) \%$	
Γ_{52}	$\Sigma^0 \pi^- 2\pi^+$	$(1.11 \pm 0.30) \%$	
Γ_{53}	$\Sigma^+ \pi^+ \pi^- \pi^0$	—	
Γ_{54}	$\Sigma^+ \omega$	[a] $(1.70 \pm 0.21) \%$	
Γ_{55}	$\Sigma^- \pi^0 2\pi^+$	$(2.1 \pm 0.4) \%$	
Γ_{56}	$\Sigma^+ K^+ K^-$	$(3.5 \pm 0.4) \times 10^{-3}$	S=1.1
Γ_{57}	$\Sigma^+ \phi$	[a] $(3.9 \pm 0.6) \times 10^{-3}$	S=1.1
Γ_{58}	$\Xi(1690)^0 K^+, \Xi^{*0} \rightarrow \Sigma^+ K^-$	$(1.02 \pm 0.25) \times 10^{-3}$	
Γ_{59}	$\Sigma^+ K^+ K^-$ nonresonant	$< 8 \times 10^{-4}$	CL=90%
Γ_{60}	$\Xi^0 K^+$	$(5.5 \pm 0.7) \times 10^{-3}$	
Γ_{61}	$\Xi^- K^+ \pi^+$	$(6.2 \pm 0.6) \times 10^{-3}$	S=1.1
Γ_{62}	$\Xi(1530)^0 K^+$	$(4.3 \pm 0.9) \times 10^{-3}$	S=1.1

Hadronic modes with a hyperon: $S = 0$ final states

Γ_{63}	ΛK^+	$(6.1 \pm 1.2) \times 10^{-4}$	
Γ_{64}	$\Lambda K^+ \pi^+ \pi^-$	$< 5 \times 10^{-4}$	CL=90%
Γ_{65}	$\Sigma^0 K^+$	$(5.2 \pm 0.8) \times 10^{-4}$	
Γ_{66}	$\Sigma^0 K^+ \pi^+ \pi^-$	$< 2.6 \times 10^{-4}$	CL=90%
Γ_{67}	$\Sigma^+ K^+ \pi^-$	$(2.1 \pm 0.6) \times 10^{-3}$	
Γ_{68}	$\Sigma^+ K^*(892)^0$	[a] $(3.5 \pm 1.0) \times 10^{-3}$	
Γ_{69}	$\Sigma^- K^+ \pi^+$	$< 1.2 \times 10^{-3}$	CL=90%

Doubly Cabibbo-suppressed modes

Γ_{70}	$p K^+ \pi^-$	$(1.11 \pm 0.18) \times 10^{-4}$
---------------	---------------	------------------------------------

Semileptonic modes

Γ_{71}	$\Lambda e^+ \nu_e$	$(3.6 \pm 0.4) \%$
Γ_{72}	$\Lambda \mu^+ \nu_\mu$	$(3.5 \pm 0.5) \%$

Inclusive modes

Γ_{73}	$e^+ \text{ anything}$	$(3.95 \pm 0.35) \%$
Γ_{74}	$p \text{ anything}$	$(50 \pm 16) \%$
Γ_{75}	$n \text{ anything}$	$(50 \pm 16) \%$
Γ_{76}	$\Lambda \text{ anything}$	$(38.2 \pm 2.9) \%$
Γ_{77}	3prongs	$(24 \pm 8) \%$

**$\Delta C = 1$ weak neutral current (*C1*) modes, or
Lepton Family number (*LF*), or Lepton number (*L*), or
Baryon number (*B*) violating modes**

Γ_{78}	$p e^+ e^-$	<i>C1</i>	< 5.5	$\times 10^{-6}$	CL=90%
Γ_{79}	$p \mu^+ \mu^-$ non-resonant	<i>C1</i>	< 7.7	$\times 10^{-8}$	CL=90%
Γ_{80}	$p e^+ \mu^-$	<i>LF</i>	< 9.9	$\times 10^{-6}$	CL=90%
Γ_{81}	$p e^- \mu^+$	<i>LF</i>	< 1.9	$\times 10^{-5}$	CL=90%
Γ_{82}	$\bar{p} 2e^+$	<i>L,B</i>	< 2.7	$\times 10^{-6}$	CL=90%
Γ_{83}	$\bar{p} 2\mu^+$	<i>L,B</i>	< 9.4	$\times 10^{-6}$	CL=90%
Γ_{84}	$\bar{p} e^+ \mu^+$	<i>L,B</i>	< 1.6	$\times 10^{-5}$	CL=90%
Γ_{85}	$\Sigma^- \mu^+ \mu^+$	<i>L</i>	< 7.0	$\times 10^{-4}$	CL=90%

[a] This branching fraction includes all the decay modes of the final-state resonance.

CONSTRAINED FIT INFORMATION

An overall fit to 41 branching ratios uses 62 measurements and one constraint to determine 21 parameters. The overall fit has a $\chi^2 = 47.4$ for 42 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_{47}	36								
x_{49}	4	8							
x_{52}	7	14	1						
x_{54}	14	29	3	5					
x_{56}	17	45	4	6	13				
x_{57}	13	37	3	5	11	16			
x_{60}	6	13	1	2	5	6	5		
x_{61}	19	34	4	7	11	16	13	6	
x_{62}	4	10	1	2	3	4	4	2	4
	x_{45}	x_{47}	x_{49}	x_{52}	x_{54}	x_{56}	x_{57}	x_{60}	x_{61}

x_2	54									
x_7	46	55								
x_{10}	44	64	39							
x_{11}	51	61	40	60						
x_{28}	54	66	44	42	43					
x_{29}	45	61	41	38	36	65				
x_{31}	51	37	28	41	60	45	36			
x_{42}	16	22	14	14	14	26	19	12		
x_{44}	51	55	38	37	40	74	58	44	20	
x_{45}	38	39	30	25	29	34	33	23	10	29
x_{47}	51	88	50	60	61	59	55	39	20	50
x_{49}	5	9	5	6	6	6	6	3	2	5
x_{52}	13	14	9	12	15	13	11	20	4	12
x_{54}	19	30	18	23	26	19	18	18	6	16
x_{56}	23	41	23	28	28	27	25	18	9	23
x_{57}	19	32	19	22	23	22	20	14	7	18
x_{60}	8	15	8	10	9	10	9	6	3	8
x_{61}	29	39	25	25	25	51	35	24	14	38
x_{62}	6	11	6	7	7	7	7	4	2	6
	x_1	x_2	x_7	x_{10}	x_{11}	x_{28}	x_{29}	x_{31}	x_{42}	x_{44}

Λ_c^+ BRANCHING RATIOS

A few really obsolete results have been omitted.

— Hadronic modes with a p : $S = -1$ final states —

$\Gamma(pK_S^0)/\Gamma_{\text{total}}$	Γ_1/Γ			
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.59 ± 0.08 OUR FIT	Error includes scale factor of 1.1.			
$1.52 \pm 0.08 \pm 0.03$	1243	ABLIKIM	16	BES3 $e^+ e^- \rightarrow \Lambda_c \bar{\Lambda}_c$, 4.599 GeV

$\Gamma(pK_S^0)/\Gamma(pK^-\pi^+)$ Γ_1/Γ_2

Measurements given as a \bar{K}^0 ratio have been divided by 2 to convert to a K_S^0 ratio.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.254 ± 0.012 OUR FIT	Error includes scale factor of 1.4.			
0.234 ± 0.020 OUR AVERAGE				
0.23 ± 0.01 ± 0.02	1025	ALAM	98	CLE2 $e^+ e^- \approx \gamma(4S)$
0.22 ± 0.04 ± 0.03	133	AVERY	91	CLEO $e^+ e^-$ 10.5 GeV
0.28 ± 0.09 ± 0.07	45	ANJOS	90	E691 γ Be 70–260 GeV
0.31 ± 0.08 ± 0.02	73	ALBRECHT	88C	ARG $e^+ e^-$ 10 GeV

$\Gamma(pK^-\pi^+)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
6.28±0.32 OUR FIT	Error includes scale factor of 1.4.			
6.3 ±0.5 OUR AVERAGE	Error includes scale factor of 2.0.			
5.84±0.27±0.23	6.3k	ABLIKIM	16	BES3 $e^+e^- \rightarrow \Lambda_c \bar{\Lambda}_c$, 4.599 GeV
6.84±0.24 ^{+0.21} _{-0.27}	1.4k	¹ ZUPANC	14	BELL $e^+e^- \rightarrow D^{(*)}-\bar{p}\pi^+$ recoil
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5.0 ±1.3		² PDG	02	See footnote

¹ This ZUPANC 14 value is the FIRST-EVER model-independent measurement of a Λ_c^+ branching fraction.

² See the note by P. Burchat, " Λ_c^+ Branching Fractions," in any edition of the Review from 2002 through 2014 for how this value was obtained. It is now obsolete.

 $\Gamma(p\bar{K}^*(892)^0)/\Gamma(pK^-\pi^+)$ Γ_3/Γ_2

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.31±0.04 OUR AVERAGE				
0.29±0.04±0.03		¹ AITALA	00	E791 $\pi^- N$, 500 GeV
0.35 ^{+0.06} _{-0.07} ±0.03	39	BOZEK	93	NA32 $\pi^- Cu$ 230 GeV
0.42±0.24	12	BASILE	81B CNTR	$p p \rightarrow \Lambda_c^+ e^- X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.35±0.11		BARLAG	90D NA32	See BOZEK 93

¹ AITALA 00 makes a coherent 5-dimensional amplitude analysis of 946 ± 38 $\Lambda_c^+ \rightarrow p K^- \pi^+$ decays.

 $\Gamma(\Delta(1232)^{++}K^-)/\Gamma(pK^-\pi^+)$ Γ_4/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.17±0.04 OUR AVERAGE	Error includes scale factor of 1.1.			
0.18±0.03±0.03		¹ AITALA	00	E791 $\pi^- N$, 500 GeV
0.12 ^{+0.04} _{-0.05} ±0.05	14	BOZEK	93	NA32 $\pi^- Cu$ 230 GeV
0.40±0.17	17	BASILE	81B CNTR	$p p \rightarrow \Lambda_c^+ e^- X$

¹ AITALA 00 makes a coherent 5-dimensional amplitude analysis of 946 ± 38 $\Lambda_c^+ \rightarrow p K^- \pi^+$ decays.

 $\Gamma(\Lambda(1520)\pi^+)/\Gamma(pK^-\pi^+)$ Γ_5/Γ_2

Unseen decay modes of the $\Lambda(1520)$ are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.35±0.08 OUR AVERAGE				
0.34±0.08±0.05		¹ AITALA	00	E791 $\pi^- N$, 500 GeV
0.40 ^{+0.18} _{-0.13} ±0.09	12	BOZEK	93	NA32 $\pi^- Cu$ 230 GeV

¹ AITALA 00 makes a coherent 5-dimensional amplitude analysis of 946 ± 38 $\Lambda_c^+ \rightarrow p K^- \pi^+$ decays.

$\Gamma(pK^-\pi^+\text{nonresonant})/\Gamma(pK^-\pi^+)$ Γ_6/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.55±0.06 OUR AVERAGE				
0.55±0.06±0.04		¹ AITALA	00	E791 $\pi^- N$, 500 GeV
$0.56^{+0.07}_{-0.09} \pm 0.05$	71	BOZEK	93	NA32 $\pi^- Cu$ 230 GeV

¹ AITALA 00 makes a coherent 5-dimensional amplitude analysis of 946 ± 38 $\Lambda_c^+ \rightarrow pK^-\pi^+$ decays.

 $\Gamma(pK_S^0\pi^0)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.97±0.13 OUR FIT Error includes scale factor of 1.1.				
1.87±0.13±0.05	558	ABLIKIM	16	BES3 $e^+e^- \rightarrow \Lambda_c\bar{\Lambda}_c$, 4.599 GeV

 $\Gamma(pK_S^0\pi^0)/\Gamma(pK^-\pi^+)$ Γ_7/Γ_2

Measurements given as a \bar{K}^0 ratio have been divided by 2 to convert to a K_S^0 ratio.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.314±0.018 OUR FIT				
0.33 ± 0.03 ± 0.04	774	ALAM	98	CLE2 $e^+e^- \approx \gamma(4S)$

 $\Gamma(nK_S^0\pi^+)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.82±0.23±0.11	83	ABLIKIM	17H	BES3 e^+e^- at 4.6 GeV

 $\Gamma(p\bar{K}^0\eta)/\Gamma(pK^-\pi^+)$ Γ_9/Γ_2

Unseen decay modes of the η are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.25±0.04±0.04	57	AMMAR	95	CLE2 $e^+e^- \approx \gamma(4S)$

 $\Gamma(pK_S^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.60±0.12 OUR FIT Error includes scale factor of 1.1.				
1.53±0.11±0.09	485	ABLIKIM	16	BES3 $e^+e^- \rightarrow \Lambda_c\bar{\Lambda}_c$, 4.599 GeV

 $\Gamma(pK_S^0\pi^+\pi^-)/\Gamma(pK^-\pi^+)$ Γ_{10}/Γ_2

Measurements given as a \bar{K}^0 ratio have been divided by 2 to convert to a K_S^0 ratio.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.255±0.015 OUR FIT Error includes scale factor of 1.1.				
0.257±0.031 OUR AVERAGE				
0.26 ± 0.02 ± 0.03	985	ALAM	98	CLE2 $e^+e^- \approx \gamma(4S)$
0.22 ± 0.06 ± 0.02	83	AVERY	91	CLEO e^+e^- 10.5 GeV
0.49 ± 0.18 ± 0.04	12	BARLAG	90D	NA32 π^- 230 GeV

 $\Gamma(pK^-\pi^+\pi^0)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
4.46±0.30 OUR FIT Error includes scale factor of 1.5.				
4.53±0.23±0.30	1849	ABLIKIM	16	BES3 $e^+e^- \rightarrow \Lambda_c\bar{\Lambda}_c$, 4.599 GeV

$\Gamma(pK^-\pi^+\pi^0)/\Gamma(pK^-\pi^+)$ Γ_{11}/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.71 ± 0.04 OUR FIT				Error includes scale factor of 2.4.
0.685 ± 0.019 OUR AVERAGE				
0.685 ± 0.007 ± 0.018	242k	PAL	17	BELL $e^+e^- \approx \gamma(4S), \gamma(5S)$
0.67 ± 0.04 ± 0.11	2.6k	ALAM	98	CLE2 $e^+e^- \approx \gamma(4S)$

 $\Gamma(pK^*(892)^-\pi^+)/\Gamma(pK_S^0\pi^+\pi^-)$ Γ_{12}/Γ_{10} Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.88 ± 0.28	17	ALEEV	94	BIS2 nN 20–70 GeV

 $\Gamma(p(K^-\pi^+)_{\text{nonresonant}}\pi^0)/\Gamma(pK^-\pi^+)$ Γ_{13}/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.73 ± 0.12 ± 0.05	67	BOZEK	93	NA32 π^- Cu 230 GeV

 $\Gamma(\Delta(1232)\bar{K}^*(892))/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	35	AMENDOLIA	87	SPEC γ Ge-Si

 $\Gamma(pK^-2\pi^+\pi^-)/\Gamma(pK^-\pi^+)$ Γ_{15}/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.022 ± 0.015		BARLAG	90D	NA32 π^- 230 GeV

 $\Gamma(pK^-\pi^+2\pi^0)/\Gamma(pK^-\pi^+)$ Γ_{16}/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.16 ± 0.07 ± 0.03	15	BOZEK	93	NA32 π^- Cu 230 GeV

Hadronic modes with a p : $S = 0$ final states

 $\Gamma(p\pi^0)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.7 × 10⁻⁴	90	ABLIKIM	17Q	BES3 e^+e^- at 4.6 GeV

 $\Gamma(p\eta)/\Gamma_{\text{total}}$ Γ_{18}/Γ Unseen decay modes of the η are included.

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
1.24 ± 0.28 ± 0.10	52	ABLIKIM	17Q	BES3 $\eta \rightarrow 2\gamma, \pi^+\pi^0\pi^-$

 $\Gamma(p\omega(782)^0)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
9.4 ± 3.2 ± 2.2	13	AAIJ	18N	LHCb Seen in $\Lambda_c^+ \rightarrow p\mu^+\mu^-$

 $\Gamma(p\pi^+\pi^-)/\Gamma(pK^-\pi^+)$ Γ_{20}/Γ_2

VALUE (units 10 ⁻²)	EVTS	DOCUMENT ID	TECN	COMMENT
7.35 ± 0.24 OUR AVERAGE				Error includes scale factor of 1.3.

7.44 ± 0.08 ± 0.18	20k	AAIJ	18V	LHCb $\Lambda_b^0 \rightarrow \Lambda_c^+\mu^-\chi$
6.70 ± 0.48 ± 0.25	495	ABLIKIM	16U	BES3 e^+e^- at 4.599 GeV
6.9 ± 3.6	5	BARLAG	90D	NA32 π^- 230 GeV

$\Gamma(p f_0(980))/\Gamma(p K^- \pi^+)$ Unseen decay modes of the $f_0(980)$ are included. Γ_{21}/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
0.055±0.036	BARLAG	90D	π^- 230 GeV

 $\Gamma(p 2\pi^+ 2\pi^-)/\Gamma(p K^- \pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.036±0.023	BARLAG	90D	π^- 230 GeV

 $\Gamma(p K^+ K^-)/\Gamma(p K^- \pi^+)$

VALUE (units 10^{-2})	EVTS
1.70±0.04 OUR AVERAGE	

		DOCUMENT ID	TECN	COMMENT
1.70±0.03±0.03	3.4k	AAIJ	18V	LHCb $\Lambda_b^0 \rightarrow \Lambda_c^+ \mu^- X$
1.4 ± 0.2 ± 0.2	676	ABE	02C	BELL $e^+ e^- \approx \gamma(4S)$
3.9 ± 0.9 ± 0.7	214	ALEXANDER	96C	CLE2 $e^+ e^- \approx \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
9.6 ± 2.9 ± 1.0	30	FRABETTI	E687	γ Be, \bar{E}_γ 220 GeV
4.8 ± 2.7		BARLAG	90D	π^- 230 GeV

 $\Gamma(p \phi)/\Gamma(p K^- \pi^+)$ Unseen decay modes of the ϕ are included. Γ_{24}/Γ_2

VALUE (units 10^{-2})	EVTS
1.70±0.21 OUR AVERAGE	

		DOCUMENT ID	TECN	COMMENT
1.81±0.33±0.13	44	ABLIKIM	16U	BES3 $e^+ e^-$ at 4.599 GeV
1.5 ± 0.2 ± 0.2	345	ABE	02C	BELL $e^+ e^- \approx \gamma(4S)$
2.4 ± 0.6 ± 0.3	54	ALEXANDER	96C	CLE2 $e^+ e^- \approx \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.0 ± 2.7		BARLAG	90D	π^- 230 GeV

 $\Gamma(p K^+ K^- \text{non-}\phi)/\Gamma(p K^- \pi^+)$ Γ_{25}/Γ_2

VALUE (units 10^{-3})	EVTS
8.4 ± 1.8 OUR AVERAGE	

		DOCUMENT ID	TECN	COMMENT
9.36±2.22±0.71	38	ABLIKIM	16U	BES3 $e^+ e^-$ at 4.599 GeV
7 ± 2 ± 2	344	ABE	02C	BELL $e^+ e^- \approx \gamma(4S)$

 $\Gamma(p \phi \pi^0)/\Gamma(p K^- \pi^+)$ Γ_{26}/Γ_2

VALUE (units 10^{-3})	EVTS
1.538±0.641^{+0.077}_{-0.100}	

PAL	DOCUMENT ID	TECN	COMMENT
	17	BELL	$e^+ e^- \approx \gamma(4S), \gamma(5S)$

 $\Gamma(p K^+ K^- \pi^0 \text{nonresonant})/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE	CL%
<6.3 × 10⁻⁵	90

PAL	DOCUMENT ID	TECN	COMMENT
	17	BELL	$e^+ e^- \approx \gamma(4S), \gamma(5S)$

———— Hadronic modes with a hyperon: $S = -1$ final states —— $\Gamma(\Lambda \pi^+)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (%)	EVTS
1.30±0.07 OUR FIT	Error includes scale factor of 1.1.

1.24±0.07±0.03	706	ABLIKIM	16	BES3 $e^+ e^- \rightarrow \Lambda_c \bar{\Lambda}_c$, 4.599 GeV
-----------------------	-----	---------	----	--

$\Gamma(\Lambda\pi^+)/\Gamma(pK^-\pi^+)$					Γ_{28}/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.207±0.009 OUR FIT	Error includes scale factor of 1.2.				
0.204±0.019 OUR AVERAGE					
0.217±0.013±0.020	750	LINK	05F	FOCS	γ nucleus, $\bar{E}_\gamma \approx 180$ GeV
0.18 ± 0.03 ± 0.04		ALBRECHT	92	ARG	$e^+e^- \approx 10.4$ GeV
0.18 ± 0.03 ± 0.03	87	AVERY	91	CLEO	$e^+e^- 10.5$ GeV
$\Gamma(\Lambda\pi^+\pi^0)/\Gamma_{\text{total}}$					Γ_{29}/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
7.1 ± 0.4 OUR FIT	Error includes scale factor of 1.1.				
7.01±0.37±0.19	1497	ABLIKIM	16	BES3	$e^+e^- \rightarrow \Lambda_c\bar{\Lambda}_c$, 4.599 GeV
$\Gamma(\Lambda\pi^+\pi^0)/\Gamma(pK^-\pi^+)$					Γ_{29}/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
1.12±0.05 OUR FIT	Error includes scale factor of 1.1.				
0.73±0.09±0.16	464	AVERY	94	CLE2	$e^+e^- \approx \gamma(3S), \gamma(4S)$
$\Gamma(\Lambda\rho^+)/\Gamma(pK^-\pi^+)$					Γ_{30}/Γ_2
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.95	95	AVERY	94	CLE2	$e^+e^- \approx \gamma(3S), \gamma(4S)$
$\Gamma(\Lambda\pi^-2\pi^+)/\Gamma_{\text{total}}$					Γ_{31}/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
3.64±0.29 OUR FIT	Error includes scale factor of 1.4.				
3.81±0.24±0.18	609	ABLIKIM	16	BES3	$e^+e^- \rightarrow \Lambda_c\bar{\Lambda}_c$, 4.599 GeV
$\Gamma(\Lambda\pi^-2\pi^+)/\Gamma(pK^-\pi^+)$					Γ_{31}/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.58 ± 0.04 OUR FIT	Error includes scale factor of 1.9.				
0.522±0.032 OUR AVERAGE					
0.508±0.024±0.024	1356	LINK	05F	FOCS	γ nucleus, $\bar{E}_\gamma \approx 180$ GeV
0.65 ± 0.11 ± 0.12	289	AVERY	91	CLEO	$e^+e^- 10.5$ GeV
0.82 ± 0.29 ± 0.27	44	ANJOS	90	E691	γ Be 70–260 GeV
0.94 ± 0.41 ± 0.13	10	BARLAG	90D	NA32	$\pi^- 230$ GeV
0.61 ± 0.16 ± 0.04	105	ALBRECHT	88C	ARG	$e^+e^- 10$ GeV
$\Gamma(\Sigma(1385)^+\pi^+\pi^-, \Sigma^{*+} \rightarrow \Lambda\pi^+)/\Gamma(\Lambda\pi^-2\pi^+)$					Γ_{32}/Γ_{31}
VALUE	DOCUMENT ID	TECN	COMMENT		
0.28±0.10±0.08	LINK	05F	FOCS	γ nucleus, $\bar{E}_\gamma \approx 180$ GeV	
$\Gamma(\Sigma(1385)^-2\pi^+, \Sigma^{*-} \rightarrow \Lambda\pi^-)/\Gamma(\Lambda\pi^-2\pi^+)$					Γ_{33}/Γ_{31}
VALUE	DOCUMENT ID	TECN	COMMENT		
0.21±0.03±0.02	LINK	05F	FOCS	γ nucleus, $\bar{E}_\gamma \approx 180$ GeV	
$\Gamma(\Lambda\pi^+\rho^0)/\Gamma(\Lambda\pi^-2\pi^+)$					Γ_{34}/Γ_{31}
VALUE	DOCUMENT ID	TECN	COMMENT		
0.40±0.12±0.12	LINK	05F	FOCS	γ nucleus, $\bar{E}_\gamma \approx 180$ GeV	

$\Gamma(\Sigma(1385)^+\rho^0, \Sigma^{*+} \rightarrow \Lambda\pi^+)/\Gamma(\Lambda\pi^-2\pi^+)$ Γ_{35}/Γ_{31}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.14±0.09±0.07	LINK	05F FOCS	γ nucleus, $\bar{E}_\gamma \approx 180$ GeV

 $\Gamma(\Lambda\pi^-2\pi^+ \text{ nonresonant})/\Gamma(\Lambda\pi^-2\pi^+)$ Γ_{36}/Γ_{31}

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.3	90	LINK	05F FOCS	γ nucleus, $\bar{E}_\gamma \approx 180$ GeV

 $\Gamma(\Lambda\pi^-\pi^02\pi^+ \text{ total})/\Gamma(pK^-\pi^+)$ Γ_{37}/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.36±0.09±0.09	50	¹ CRONIN-HEN..03	CLE3	$e^+e^- \approx \gamma(4S)$

¹ CRONIN-HENNESSY 03 finds this channel to be dominantly $\Lambda\eta\pi^+$ and $\Lambda\omega\pi^+$; see below.

 $\Gamma(\Lambda\pi^+\eta)/\Gamma(pK^-\pi^+)$ Γ_{38}/Γ_2

Unseen decay modes of the η are included.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.36±0.07 OUR AVERAGE				
0.41±0.17±0.10	11	CRONIN-HEN..03	CLE3	$e^+e^- \approx \gamma(4S)$
0.35±0.05±0.06	116	AMMAR	95	CLE2 $e^+e^- \approx \gamma(4S)$

 $\Gamma(\Sigma(1385)^+\eta)/\Gamma(pK^-\pi^+)$ Γ_{39}/Γ_2

Unseen decay modes of the $\Sigma(1385)^+$ and η are included.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.17±0.04±0.03	54	AMMAR	95	CLE2 $e^+e^- \approx \gamma(4S)$

 $\Gamma(\Lambda\pi^+\omega)/\Gamma(pK^-\pi^+)$ Γ_{40}/Γ_2

Unseen decay modes of the ω are included.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.24±0.06±0.06	32	CRONIN-HEN..03	CLE3	$e^+e^- \approx \gamma(4S)$

 $\Gamma(\Lambda\pi^-\pi^02\pi^+, \text{ no } \eta \text{ or } \omega)/\Gamma(pK^-\pi^+)$ Γ_{41}/Γ_2

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.13	90	CRONIN-HEN..03	CLE3	$e^+e^- \approx \gamma(4S)$

 $\Gamma(\Lambda K^+\bar{K}^0)/\Gamma(pK^-\pi^+)$ Γ_{42}/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.090±0.017 OUR FIT				Error includes scale factor of 1.9.

0.131±0.020 OUR AVERAGE

0.142±0.018±0.022	251	LINK	05F FOCS	γ nucleus, $\bar{E}_\gamma \approx 180$ GeV
0.12 ± 0.02 ± 0.02	59	AMMAR	95	CLE2 $e^+e^- \approx \gamma(4S)$

 $\Gamma(\Xi(1690)^0 K^+, \Xi^*0 \rightarrow \Lambda\bar{K}^0)/\Gamma(\Lambda K^+\bar{K}^0)$ Γ_{43}/Γ_{42}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.28±0.07 OUR AVERAGE				
0.32±0.10±0.04	84±24	LINK	05F FOCS	γ nucleus, $\bar{E}_\gamma \approx 180$ GeV
0.26±0.08±0.03	93	ABE	02C BELL	$e^+e^- \approx \gamma(4S)$

$\Gamma(\Lambda K^+ \bar{K}^0)/\Gamma(\Lambda \pi^+)$					Γ_{42}/Γ_{28}
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.44 ± 0.08 OUR FIT		Error includes scale factor of 2.0.			
0.395 ± 0.026 ± 0.036	460 ± 30	AUBERT	07U	BABR $e^+ e^- \approx \gamma(4S)$	
$\Gamma(\Sigma^0 \pi^+)/\Gamma_{\text{total}}$					Γ_{44}/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
1.29 ± 0.07 OUR FIT		Error includes scale factor of 1.1.			
1.27 ± 0.08 ± 0.03	522	ABLIKIM	16	BES3 $e^+ e^- \rightarrow \Lambda_c \bar{\Lambda}_c$, 4.599 GeV	
$\Gamma(\Sigma^0 \pi^+)/\Gamma(p K^- \pi^+)$					Γ_{44}/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.206 ± 0.010 OUR FIT		Error includes scale factor of 1.2.			
0.20 ± 0.04 OUR AVERAGE					
0.21 ± 0.02 ± 0.04	196	AVERY	94	CLE2 $e^+ e^- \approx \gamma(3S), \gamma(4S)$	
0.17 ± 0.06 ± 0.04		ALBRECHT	92	ARG $e^+ e^- \approx 10.4$ GeV	
$\Gamma(\Sigma^0 \pi^+)/\Gamma(\Lambda \pi^+)$					Γ_{44}/Γ_{28}
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.99 ± 0.04 OUR FIT					
0.98 ± 0.05 OUR AVERAGE					
0.977 ± 0.015 ± 0.051	33k	AUBERT	07U	BABR $e^+ e^- \approx \gamma(4S)$	
1.09 ± 0.11 ± 0.19	750	LINK	05F	FOCS γ nucleus, $\bar{E}_\gamma \approx 180$ GeV	
$\Gamma(\Sigma^+ \pi^0)/\Gamma_{\text{total}}$					Γ_{45}/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
1.25 ± 0.10 OUR FIT					
1.18 ± 0.10 ± 0.03	309	ABLIKIM	16	BES3 $e^+ e^- \rightarrow \Lambda_c \bar{\Lambda}_c$, 4.599 GeV	
$\Gamma(\Sigma^+ \pi^0)/\Gamma(p K^- \pi^+)$					Γ_{45}/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.199 ± 0.015 OUR FIT					
0.20 ± 0.03 ± 0.03	93	KUBOTA	93	CLE2 $e^+ e^- \approx \gamma(4S)$	
$\Gamma(\Sigma^+ \eta)/\Gamma(p K^- \pi^+)$					Γ_{46}/Γ_2
Unseen decay modes of the η are included.					
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.11 ± 0.03 ± 0.02	26	AMMAR	95	CLE2 $e^+ e^- \approx \gamma(4S)$	
$\Gamma(\Sigma^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$					Γ_{47}/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
4.50 ± 0.25 OUR FIT		Error includes scale factor of 1.3.			
4.25 ± 0.24 ± 0.20	1156	ABLIKIM	16	BES3 $e^+ e^- \rightarrow \Lambda_c \bar{\Lambda}_c$, 4.599 GeV	

$\Gamma(\Sigma^+ \pi^+ \pi^-)/\Gamma(p K^- \pi^+)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.716±0.019 OUR FIT**0.720±0.024 OUR AVERAGE**

0.719±0.003±0.024	2.7M	BERGER	18	BELL $e^+ e^- \approx \gamma(4S)$
0.74 ± 0.07 ± 0.09	487	KUBOTA	93	CLE2 $e^+ e^- \approx \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.72 ± 0.14	47 ± 9	VAZQUEZ-JA...08	SELX	Σ^- nucleus, 600 GeV
0.54 +0.18 -0.15	11	BARLAG	92	NA32 π^- Cu 230 GeV

 Γ_{47}/Γ_2  $\Gamma(\Sigma^+ \rho^0)/\Gamma(p K^- \pi^+)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.27	95	KUBOTA	93	CLE2 $e^+ e^- \approx \gamma(4S)$

 Γ_{48}/Γ_2  $\Gamma(\Sigma^- 2\pi^+)/\Gamma_{\text{total}}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.87±0.18 OUR FIT				
1.81±0.17±0.09	161	ABLIKIM	17Y	BES3 $e^+ e^-$ at 4.6 GeV

 $\Gamma(\Sigma^- 2\pi^+)/\Gamma(p K^- \pi^+)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.297±0.030 OUR FIT				Error includes scale factor of 1.1.
0.314±0.067	30 ± 6	VAZQUEZ-JA...08	SELX	Σ^- nucleus, 600 GeV

 Γ_{49}/Γ_2  $\Gamma(\Sigma^- 2\pi^+)/\Gamma(\Sigma^+ \pi^+ \pi^-)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.42±0.04 OUR FIT				Error includes scale factor of 1.1.
0.53±0.15±0.07	56	FRABETTI	94E	E687 γ Be, \bar{E}_γ 220 GeV

 Γ_{49}/Γ_{47}  $\Gamma(\Sigma^0 \pi^+ \pi^0)/\Gamma(p K^- \pi^+)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.56 ± 0.05 OUR AVERAGE				Error includes scale factor of 1.5.
0.575±0.005±0.036	2.7M	BERGER	18	BELL $e^+ e^- \approx \gamma(4S)$
0.36 ± 0.09 ± 0.10	117	AVERY	94	CLE2 $e^+ e^- \approx \gamma(3S), \gamma(4S)$

 Γ_{50}/Γ_2  $\Gamma(\Sigma^+ \pi^0 \pi^0)/\Gamma(p K^- \pi^+)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.247±0.006±0.019	925k	BERGER	18	BELL $e^+ e^- \approx \gamma(4S)$

 Γ_{51}/Γ_2  $\Gamma(\Sigma^0 \pi^- 2\pi^+)/\Gamma(p K^- \pi^+)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.18±0.05 OUR FIT				
0.21±0.05±0.05	90	AVERY	94	CLE2 $e^+ e^- \approx \gamma(3S), \gamma(4S)$

 Γ_{52}/Γ_2  $\Gamma(\Sigma^0 \pi^- 2\pi^+)/\Gamma(\Lambda \pi^- 2\pi^+)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.31±0.08 OUR FIT				
0.26±0.06±0.09	480	LINK	05F	FOCS γ nucleus, $\bar{E}_\gamma \approx 180$ GeV

 Γ_{52}/Γ_{31} 

$\Gamma(\Sigma^+ \omega)/\Gamma_{\text{total}}$ Γ_{54}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.70±0.21 OUR FIT				
1.56±0.20±0.07	157	ABLIKIM	16	BES3 $e^+ e^- \rightarrow \Lambda_c \bar{\Lambda}_c$, 4.599 GeV

 $\Gamma(\Sigma^+ \omega)/\Gamma(p K^- \pi^+)$ Γ_{54}/Γ_2 Unseen decay modes of the ω are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.271±0.031 OUR FIT				
0.54 ±0.13 ±0.06	107	KUBOTA	93	CLE2 $e^+ e^- \approx \gamma(4S)$

 $\Gamma(\Sigma^- \pi^0 2\pi^+)/\Gamma_{\text{total}}$ Γ_{55}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
2.11±0.33±0.14	88	ABLIKIM	17Y	BES3 $e^+ e^-$ at 4.6 GeV

 $\Gamma(\Sigma^+ K^+ K^-)/\Gamma(p K^- \pi^+)$ Γ_{56}/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.056±0.006 OUR FIT				
0.070±0.011±0.011	59	AVERY	93	CLE2 $e^+ e^- \approx 10.5$ GeV

 $\Gamma(\Sigma^+ K^+ K^-)/\Gamma(\Sigma^+ \pi^+ \pi^-)$ Γ_{56}/Γ_{47}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.078±0.008 OUR FIT				
0.074±0.009 OUR AVERAGE				
0.076±0.007±0.009	246	ABE	02C	BELL $e^+ e^- \approx \gamma(4S)$
0.071±0.011±0.011	103	LINK	02G	FOCS γ nucleus, ≈ 180 GeV

 $\Gamma(\Sigma^+ \phi)/\Gamma(p K^- \pi^+)$ Γ_{57}/Γ_2 Unseen decay modes of the ϕ are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.062±0.009 OUR FIT				
0.069±0.023±0.016	26	AVERY	93	CLE2 $e^+ e^- \approx 10.5$ GeV

 $\Gamma(\Sigma^+ \phi)/\Gamma(\Sigma^+ \pi^+ \pi^-)$ Γ_{57}/Γ_{47} Unseen decay modes of the ϕ are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.087±0.012 OUR FIT				
0.086±0.012 OUR AVERAGE				
0.085±0.012±0.012	129	ABE	02C	BELL $e^+ e^- \approx \gamma(4S)$
0.087±0.016±0.006	57	LINK	02G	FOCS γ nucleus, ≈ 180 GeV

 $\Gamma(\Xi(1690)^0 K^+, \Xi^{*0} \rightarrow \Sigma^+ K^-)/\Gamma(\Sigma^+ \pi^+ \pi^-)$ Γ_{58}/Γ_{47}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.023±0.005 OUR AVERAGE				
0.023±0.005±0.005	75	ABE	02C	BELL $e^+ e^- \approx \gamma(4S)$
0.022±0.006±0.006	34	LINK	02G	FOCS γ nucleus, ≈ 180 GeV

 $\Gamma(\Sigma^+ K^+ K^- \text{ nonresonant})/\Gamma(\Sigma^+ \pi^+ \pi^-)$ Γ_{59}/Γ_{47}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.018	90	ABE	02C	BELL $e^+ e^- \approx \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.028	90	LINK	02G	FOCS γ nucleus, ≈ 180 GeV

$\Gamma(\Xi^0 K^+)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS
5.5 \pm 0.7 OUR FIT	
5.90 \pm 0.86 \pm 0.39	68

Γ_{60}/Γ



$\Gamma(\Xi^0 K^+)/\Gamma(p K^- \pi^+)$

VALUE	EVTS
0.088 \pm 0.012 OUR FIT	
0.078 \pm 0.013 \pm 0.013	56

Γ_{60}/Γ_2

$\Gamma(\Xi^- K^+ \pi^+)/\Gamma(p K^- \pi^+)$


VALUE	EVTS
0.099 \pm 0.009 OUR FIT	
0.098 \pm 0.021 OUR AVERAGE	

Γ_{61}/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.14 \pm 0.03 \pm 0.02	34	ALBRECHT	95B	ARG $e^+ e^- \approx 10.4$ GeV
0.079 \pm 0.013 \pm 0.014	60	AVERY	93	CLE2 $e^+ e^- \approx 10.5$ GeV
0.15 \pm 0.04 \pm 0.03	30	AVERY	91	CLEO $e^+ e^- 10.5$ GeV

WEIGHTED AVERAGE

0.098 \pm 0.021 (Error scaled by 1.3)



$\Gamma(\Xi^- K^+ \pi^+)/\Gamma(\Lambda \pi^+)$

VALUE	EVTS
0.48 \pm 0.04 OUR FIT	
0.480 \pm 0.016 \pm 0.039	2665 ± 84

Γ_{61}/Γ_{28}

$\Gamma(\Xi(1530)^0 K^+)/\Gamma_{\text{total}}$	Γ_{62}/Γ			
<u>VALUE (units 10^{-3})</u> <u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>				
4.3 ± 0.9 OUR FIT Error includes scale factor of 1.1.				
5.02±0.99±0.31	60	ABLIKIM	18Y	BES3 $e^+ e^-$ at 4.6 GeV
$\Gamma(\Xi(1530)^0 K^+)/\Gamma(pK^-\pi^+)$	Γ_{62}/Γ_2			
<u>VALUE</u> <u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>				
0.068±0.014 OUR FIT Error includes scale factor of 1.1.				
0.053±0.016±0.010	24	AVERY	93	CLE2 $e^+ e^- \approx 10.5$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.05 ± 0.02 ± 0.01	11	ALBRECHT	95B	ARG $e^+ e^- \approx 10.4$ GeV

———— Hadronic modes with a hyperon: $S = 0$ final states ——

$\Gamma(\Lambda K^+)/\Gamma(\Lambda\pi^+)$	Γ_{63}/Γ_{28}			
<u>VALUE</u> <u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>				
0.047±0.009 OUR AVERAGE Error includes scale factor of 1.8.				
0.044±0.004±0.003	1162 ± 101	AUBERT	07U	BABR $e^+ e^- \approx \gamma(4S)$
0.074±0.010±0.012	265	ABE	02C	BELL $e^+ e^- \approx \gamma(4S)$

$\Gamma(\Lambda K^+\pi^+\pi^-)/\Gamma(\Lambda\pi^+)$	Γ_{64}/Γ_{28}			
<u>VALUE</u> <u>CL%</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>				
<4.1 × 10⁻²	90	AUBERT	07U	BABR $e^+ e^- \approx \gamma(4S)$

$\Gamma(\Sigma^0 K^+)/\Gamma(\Sigma^0\pi^+)$	Γ_{65}/Γ_{44}			
<u>VALUE</u> <u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>				
0.040±0.006 OUR AVERAGE				
0.038±0.005±0.003	366 ± 52	AUBERT	07U	BABR $e^+ e^- \approx \gamma(4S)$
0.056±0.014±0.008	75	ABE	02C	BELL $e^+ e^- \approx \gamma(4S)$

$\Gamma(\Sigma^0 K^+\pi^+\pi^-)/\Gamma(\Sigma^0\pi^+)$	Γ_{66}/Γ_{44}			
<u>VALUE</u> <u>CL%</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>				
<2.0 × 10⁻²	90	AUBERT	07U	BABR $e^+ e^- \approx \gamma(4S)$

$\Gamma(\Sigma^+ K^+\pi^-)/\Gamma(\Sigma^+\pi^+\pi^-)$	Γ_{67}/Γ_{47}			
<u>VALUE</u> <u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>				
0.047±0.011±0.008				
0.047 ± 0.011 ± 0.008	105	ABE	02C	BELL $e^+ e^- \approx \gamma(4S)$

$\Gamma(\Sigma^+ K^*(892)^0)/\Gamma(\Sigma^+\pi^+\pi^-)$	Γ_{68}/Γ_{47}			
Unseen decay modes of the $K^*(892)^0$ are included.				
0.078±0.018±0.013	49	LINK	02G	FOCS γ nucleus, ≈ 180 GeV

$\Gamma(\Sigma^- K^+\pi^+)/\Gamma(\Sigma^+ K^*(892)^0)$	Γ_{69}/Γ_{68}			
<u>VALUE</u> <u>CL%</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>				
<0.35				
<0.35	90	LINK	02G	FOCS γ nucleus, ≈ 180 GeV

Doubly Cabibbo-suppressed modes

$\Gamma(pK^+\pi^-)/\Gamma(pK^-\pi^+)$					Γ_{70}/Γ_2
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.77 ± 0.27 OUR AVERAGE	Error includes scale factor of 1.9.				
$1.65 \pm 0.15 \pm 0.05$	392	AAIJ	18V	LHCb	$\Lambda_b^0 \rightarrow \Lambda_c^+ \mu^- X$
$2.35 \pm 0.27 \pm 0.21$	3379	YANG	16	BELL	At or near γ_s

Semileptonic modes

$\Gamma(\Lambda e^+ \nu_e)/\Gamma_{\text{total}}$					Γ_{71}/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
$3.63 \pm 0.38 \pm 0.20$	104	ABLIKIM	15Y	BES3	$567 \text{ pb}^{-1}, 4.599 \text{ GeV}$

$\Gamma(\Lambda e^+ \nu_e)/\Gamma(e^+ \text{anything})$					Γ_{71}/Γ_{73}
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
$91.9 \pm 12.5 \pm 5.4$	214	ABLIKIM	18AF	BES3	$e^+ e^- 4.6 \text{ GeV}$

$\Gamma(\Lambda e^+ \nu_e)/\Gamma(pK^-\pi^+)$					Γ_{71}/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.43 ± 0.08	^{1,2} BERGFELD	94	CLE2	$e^+ e^- \approx \gamma(4S)$
0.38 ± 0.14	^{2,3} ALBRECHT	91G	ARG	$e^+ e^- \approx 10.4 \text{ GeV}$

¹ BERGFELD 94 measures $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e) = (4.87 \pm 0.28 \pm 0.69) \text{ pb}$.

² To extract $\Gamma(\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e)/\Gamma(\Lambda_c^+ \rightarrow pK^-\pi^+)$, we use $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow pK^-\pi^+) = (11.2 \pm 1.3) \text{ pb}$, which is the weighted average of measurements from ARGUS (ALBRECHT 96E) and CLEO (AVERY 91).

³ ALBRECHT 91G measures $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e) = (4.20 \pm 1.28 \pm 0.71) \text{ pb}$.

$\Gamma(\Lambda \mu^+ \nu_\mu)/\Gamma_{\text{total}}$					Γ_{72}/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
$3.49 \pm 0.46 \pm 0.27$	79	ABLIKIM	17D	BES3	$e^+ e^- \text{ at } 4.6 \text{ GeV}$

$\Gamma(\Lambda \mu^+ \nu_\mu)/\Gamma(pK^-\pi^+)$					Γ_{72}/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.40 ± 0.09	^{1,2} BERGFELD	94	CLE2	$e^+ e^- \approx \gamma(4S)$
0.35 ± 0.20	^{2,3} ALBRECHT	91G	ARG	$e^+ e^- \approx 10.4 \text{ GeV}$

¹ BERGFELD 94 measures $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow \Lambda \mu^+ \nu_\mu) = (4.43 \pm 0.51 \pm 0.64) \text{ pb}$.

² To extract $\Gamma(\Lambda_c^+ \rightarrow \Lambda \mu^+ \nu_\mu)/\Gamma(\Lambda_c^+ \rightarrow pK^-\pi^+)$, we use $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow pK^-\pi^+) = (11.2 \pm 1.3) \text{ pb}$, which is the weighted average of measurements from ARGUS (ALBRECHT 96E) and CLEO (AVERY 91).

³ ALBRECHT 91G measures $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow \Lambda \mu^+ \nu_\mu) = (3.91 \pm 2.02 \pm 0.90) \text{ pb}$.

$\Gamma(\Lambda\mu^+\nu_\mu)/\Gamma(\Lambda e^+\nu_e)$ Γ_{72}/Γ_{71}

VALUE	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$0.96 \pm 0.16 \pm 0.04$	¹ ABLIKIM	17D BES3	e^+e^- at 4.6 GeV
¹ This is the ratio of the ABLIKIM 17D $\Lambda\mu^+\nu_e$ branching fraction and the ABLIKIM 15Y $\Lambda e^+\nu_e$ branching fraction (see above), and so is not an independent measurement.			

Inclusive modes

$\Gamma(e^+ \text{anything})/\Gamma_{\text{total}}$ Γ_{73}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$3.95 \pm 0.34 \pm 0.09$	214	ABLIKIM	18AF BES3	e^+e^- 4.6 GeV

$\Gamma(p \text{ anything})/\Gamma_{\text{total}}$ Γ_{74}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$0.50 \pm 0.08 \pm 0.14$	¹ CRAWFORD 92	CLEO	e^+e^- 10.5 GeV

¹ This CRAWFORD 92 value includes protons from Λ decay. The value is model dependent, but account is taken of this in the systematic error.

$\Gamma(n \text{ anything})/\Gamma_{\text{total}}$ Γ_{75}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$0.50 \pm 0.08 \pm 0.14$	¹ CRAWFORD 92	CLEO	e^+e^- 10.5 GeV

¹ This CRAWFORD 92 value includes neutrons from Λ decay. The value is model dependent, but account is taken of this in the systematic error.

$\Gamma(\Lambda \text{ anything})/\Gamma_{\text{total}}$ Γ_{76}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$38.2^{+2.8}_{-2.2} \pm 0.9$	700	ABLIKIM	18E BES3	e^+e^- at 4.6 GeV

$\Gamma(3\text{prongs})/\Gamma_{\text{total}}$ Γ_{77}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$0.24 \pm 0.07 \pm 0.04$	KAYIS-TOPAK.03	CHRS	ν_μ emulsion, $\bar{E}=27$ GeV

Rare or forbidden modes

$\Gamma(pe^+e^-)/\Gamma_{\text{total}}$ Γ_{78}/Γ

A test for the $\Delta C=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<5.5 \times 10^{-6}$	90	4.0 ± 7.1	LEES	11G BABR	$e^+e^- \approx \gamma(4S)$

$\Gamma(p\mu^+\mu^- \text{ non-resonant})/\Gamma_{\text{total}}$ Γ_{79}/Γ

A test for the $\Delta C=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.7 \times 10^{-8}$	90	AAIJ	18N LHCb	Ratio to $p\phi$, $\phi \rightarrow \mu^+\mu^-$

$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

$<4.4 \times 10^{-5}$	90	LEES	11G BABR	$e^+e^- \approx \gamma(4S)$
$<3.4 \times 10^{-4}$	90	KODAMA	95 E653	π^- emulsion 600 GeV

$\Gamma(p e^+ \mu^-)/\Gamma_{\text{total}}$

A test of lepton family-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<9.9 \times 10^{-6}$	90	-0.7 ± 3.0	LEES	11G BABR	$e^+ e^- \approx \gamma(4S)$

 Γ_{80}/Γ $\Gamma(p e^- \mu^+)/\Gamma_{\text{total}}$

A test of lepton family-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<19 \times 10^{-6}$	90	6.2 ± 4.9	LEES	11G BABR	$e^+ e^- \approx \gamma(4S)$

 Γ_{81}/Γ $\Gamma(\bar{p}2e^+)/\Gamma_{\text{total}}$

A test of lepton- and baryon-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.7 \times 10^{-6}$	90	-1.5 ± 4.5	LEES	11G BABR	$e^+ e^- \approx \gamma(4S)$

 Γ_{82}/Γ $\Gamma(\bar{p}2\mu^+)/\Gamma_{\text{total}}$

A test of lepton- and baryon-number conservation and of lepton family-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<9.4 \times 10^{-6}$	90	0.0 ± 2.2	LEES	11G BABR	$e^+ e^- \approx \gamma(4S)$

 Γ_{83}/Γ $\Gamma(\bar{p}e^+\mu^+)/\Gamma_{\text{total}}$

A test of lepton- and baryon-number conservation and of lepton family-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<16 \times 10^{-6}$	90	10.1 ± 6.8	LEES	11G BABR	$e^+ e^- \approx \gamma(4S)$

 Γ_{84}/Γ $\Gamma(\Sigma^-\mu^+\mu^+)/\Gamma_{\text{total}}$

A test of lepton-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<7.0 \times 10^{-4}$	90	0	KODAMA	95 E653	π^- emulsion 600 GeV

 Γ_{85}/Γ $\Lambda_c^+ \text{ DECAY PARAMETERS}$

See the note on "Baryon Decay Parameters" in the neutron Listings.

 $\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Lambda \pi^+$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.91 ± 0.15 OUR AVERAGE				
$-0.78 \pm 0.16 \pm 0.19$		LINK	06A FOCS	$\gamma A, \bar{E}_\gamma \approx 180$ GeV
$-0.94 \pm 0.21 \pm 0.12$	414	¹ BISHAI	95 CLE2	$e^+ e^- \approx \gamma(4S)$
-0.96 ± 0.42		ALBRECHT	92 ARG	$e^+ e^- \approx 10.4$ GeV
-1.1 ± 0.4	86	AVERY	90B CLEO	$e^+ e^- \approx 10.6$ GeV

¹ BISHAI 95 actually gives $\alpha = -0.94^{+0.21+0.12}_{-0.06-0.06}$, chopping the errors at the physical limit -1.0 . However, for $\alpha \approx -1.0$, some experiments should get unphysical values ($\alpha < -1.0$), and for averaging with other measurements such values (or errors that extend below -1.0) should *not* be chopped.

 $\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Sigma^+ \pi^0$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.45 \pm 0.31 \pm 0.06$	89	BISHAI	95 CLE2	$e^+ e^- \approx \gamma(4S)$

α FOR $\Lambda_c^+ \rightarrow \Lambda \ell^+ \nu_\ell$

The experiments don't cover the complete (or same incomplete) $M(\Lambda \ell^+)$ range, but we average them together anyway.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.86±0.04 OUR AVERAGE				
-0.86±0.03±0.02	3201	¹ HINSON	05	CLEO $e^+ e^- \approx \gamma(4S)$
-0.91±0.42±0.25		² ALBRECHT	94B	ARG $e^+ e^- \approx 10$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.82 ^{+0.09} _{-0.06} ^{+0.06} _{-0.03}	700	³ CRAWFORD	95	CLE2 See HINSON 05
-0.89 ^{+0.17} _{-0.11} ^{+0.09} _{-0.05}	350	⁴ BERGFELD	94	CLE2 See CRAWFORD 95

¹ HINSON 05 measures the form-factor ratio $R \equiv f_2/f_1$ for $\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e$ events to be $-0.31 \pm 0.05 \pm 0.04$ and the pole mass to be $2.21 \pm 0.08 \pm 0.14$ GeV/c², and from these calculates α , averaged over q^2 , where $\langle q^2 \rangle = 0.67$ (GeV/c)².

² ALBRECHT 94B uses Λe^+ and $\Lambda \mu^+$ events in the mass range $1.85 < M(\Lambda \ell^+) < 2.20$ GeV.

³ CRAWFORD 95 measures the form-factor ratio $R \equiv f_2/f_1$ for $\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e$ events to be $-0.25 \pm 0.14 \pm 0.08$ and from this calculates α , averaged over q^2 , to be the above.

⁴ BERGFELD 94 uses Λe^+ events.

$\Lambda_c^+, \bar{\Lambda}_c^-$ CP-VIOLATING DECAY ASYMMETRIES

$(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha})$ in $\Lambda_c^+ \rightarrow \Lambda \pi^+, \bar{\Lambda}_c^- \rightarrow \bar{\Lambda} \pi^-$

This is zero if CP is conserved.

VALUE	DOCUMENT ID	TECN	COMMENT
-0.07±0.19±0.24	LINK	06A	FOCS $\gamma A, \bar{E}_\gamma \approx 180$ GeV

$(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha})$ in $\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e, \bar{\Lambda}_c^- \rightarrow \bar{\Lambda} e^- \bar{\nu}_e$

This is zero if CP is conserved.

VALUE	DOCUMENT ID	TECN	COMMENT
0.00±0.03±0.02	HINSON	05	CLEO $e^+ e^- \approx \gamma(4S)$

$A_{CP}(\Lambda X)$ in $\Lambda_c \rightarrow \Lambda X, \bar{\Lambda}_c \rightarrow \bar{\Lambda} X$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
2.1^{+7.0}_{-6.6}±1.6	700	ABLIKIM	18E	BES3 $e^+ e^-$ at 4.6 GeV

$$\Delta A_{CP} = A_{CP}(\Lambda_c^+ \rightarrow p K^+ K^-) - A_{CP}(\Lambda_c^+ \rightarrow p \pi^+ \pi^-)$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
0.30±0.91±0.61	¹ AAIJ	18R	LHCb $p p$ 7, 8 TeV

¹ AAIJ 18R applies phase-space-dependent weights to the $\Lambda_c^+ \rightarrow p \pi^+ \pi^-$ sample to align its kinematics with the $\Lambda_c^+ \rightarrow p K^+ K^-$ sample.

Λ_c^+ REFERENCES

We have omitted some papers that have been superseded by later experiments. The omitted papers may be found in our 1992 edition (Physical Review **D45**, 1 June, Part II) or in earlier editions.

AAIJ	18N	PR D97 091101	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	18R	JHEP 1803 182	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	18V	JHEP 1803 043	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	18AF	PRL 121 251801	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	18E	PRL 121 062003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	18Y	PL B783 200	M. Ablikim <i>et al.</i>	(BES III Collab.)
BERGER	18	PR D98 112006	M. Berger <i>et al.</i>	(BELLE Collab.)
ABLIKIM	17D	PL B767 42	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	17H	PRL 118 112001	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	17Q	PR D95 111102	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	17Y	PL B772 388	M. Ablikim <i>et al.</i>	(BES III Collab.)
PAL	17	PR D96 051102	B. Pal <i>et al.</i>	(BELLE Collab.)
ABLIKIM	16	PRL 116 052001	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	16U	PRL 117 232002	M. Ablikim <i>et al.</i>	(BES III Collab.)
YANG	16	PRL 117 011801	S.B. Yang <i>et al.</i>	(BELLE Collab.)
ABLIKIM	15Y	PRL 115 221805	M. Ablikim <i>et al.</i>	(BES III Collab.)
ZUPANC	14	PRL 113 042002	A. Zupanc <i>et al.</i>	(BELLE Collab.)
LEES	11G	PR D84 072006	J.P. Lees <i>et al.</i>	(BABAR Collab.)
VAZQUEZ-JA... 08		PL B666 299	E. Vazquez-Jauregui <i>et al.</i>	(SELEX Collab.)
AUBERT	07U	PR D75 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
LINK	06A	PL B634 165	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
AUBERT,B	05S	PR D72 052006	B. Aubert <i>et al.</i>	(BABAR Collab.)
HINSON	05	PRL 94 191801	J.W. Hinson <i>et al.</i>	(CLEO Collab.)
LINK	05F	PL B624 22	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
CRONIN-HEN... 03		PR D67 012001	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
KAYIS-TOPAKU..03		PL B555 156	A. Kayis-Topaksu <i>et al.</i>	(CERN CHORUS Collab.)
ABE	02C	PL B524 33	K. Abe <i>et al.</i>	(KEK BELLE Collab.)
LINK	02C	PRL 88 161801	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	02G	PL B540 25	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
PDG	02	PR D66 010001	K. Hagiwara <i>et al.</i>	(PDG Collab.)
KUSHNIR...	01	PRL 86 5243	A. Kushnirenko <i>et al.</i>	(FNAL SELEX Collab.)
MAHMOOD	01	PRL 86 2232	A.H. Mahmood <i>et al.</i>	(CLEO Collab.)
AITALA	00	PL B471 449	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
ALAM	98	PR D57 4467	M.S. Alam <i>et al.</i>	(CLEO Collab.)
ALBRECHT	96E	PRPL 276 223	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	96C	PR D53 1013	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
ALBRECHT	95B	PL B342 397	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
AMMAR	95	PRL 74 3534	R. Ammar <i>et al.</i>	(CLEO Collab.)
BISHAI	95	PL B350 256	M. Bishai <i>et al.</i>	(CLEO Collab.)
CRAWFORD	95	PRL 75 624	G. Crawford <i>et al.</i>	(CLEO Collab.)
KODAMA	95	PL B345 85	K. Kodama <i>et al.</i>	(FNAL E653 Collab.)
ALBRECHT	94B	PL B326 320	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEEV	94	PAN 57 1370	A.N. Aleev <i>et al.</i>	(Serpukhov BIS-2 Collab.)
		Translated from YF 57 1443.		
AVERY	94	PL B325 257	P. Avery <i>et al.</i>	(CLEO Collab.)
BERGFELD	94	PL B323 219	T. Bergfeld <i>et al.</i>	(CLEO Collab.)
FRAEBETTI	94E	PL B328 193	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
AVERY	93	PRL 71 2391	P. Avery <i>et al.</i>	(CLEO Collab.)
BOZEK	93	PL B312 247	A. Bozek <i>et al.</i>	(CERN NA32 Collab.)
FRAEBETTI	93D	PRL 70 1755	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRAEBETTI	93H	PL B314 477	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
KUBOTA	93	PRL 71 3255	Y. Kubota <i>et al.</i>	(CLEO Collab.)
ALBRECHT	92	PL B274 239	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BARLAG	92	PL B283 465	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
CRAWFORD	92	PR D45 752	G. Crawford <i>et al.</i>	(CLEO Collab.)
JEZABEK	92	PL B286 175	M. Jezabek, K. Rybicki, R. Rylko	(CRAC)
ALBRECHT	91G	PL B269 234	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
AVERY	91	PR D43 3599	P. Avery <i>et al.</i>	(CLEO Collab.)
ALVAREZ	90	ZPHY C47 539	M.P. Alvarez <i>et al.</i>	(CERN NA14/2 Collab.)
ALVAREZ	90B	PL B246 256	M.P. Alvarez <i>et al.</i>	(CERN NA14/2 Collab.)
ANJOS	90	PR D41 801	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
AVERY	90B	PR D 65 2842	P. Avery <i>et al.</i>	(CLEO Collab.)
BARLAG	90D	ZPHY C48 29	S. Barlag <i>et al.</i>	(ACCMOR Collab.)

FRAEBETTI	90	PL B251 639	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
BARLAG	89	PL B218 374	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
AGUILAR-...	88B	ZPHY C40 321	M. Aguilar-Benitez <i>et al.</i>	(LEBC-EHS Collab.)
Also		PL B189 254	M. Aguilar-Benitez <i>et al.</i>	(LEBC-EHS Collab.)
Also		PL B199 462	M. Aguilar-Benitez <i>et al.</i>	(LEBC-EHS Collab.)
Also		SJNP 48 833	M. Begalli <i>et al.</i>	(LEBC-EHS Collab.)
		Translated from YAF 48 1310.		
ALBRECHT	88C	PL B207 109	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ANJOS	88B	PRL 60 1379	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
AMENDOLIA	87	ZPHY C36 513	S.R. Amendolia <i>et al.</i>	(CERN NA1 Collab.)
JONES	87	ZPHY C36 593	G.T. Jones <i>et al.</i>	(CERN WA21 Collab.)
BOSETTI	82	PL 109B 234	P.C. Bosetti <i>et al.</i>	(AACH3, BONN, CERN+)
BASILE	81B	NC 62A 14	M. Basile <i>et al.</i>	(CERN, BGNA, PGIA, FRAS)
CALICCHIO	80	PL 93B 521	M. Calicchio <i>et al.</i>	(BARI, BIRM, BRUX+)

OTHER RELATED PAPERS

MIGLIOZZI	99	PL B462 217	P. Migliozzi <i>et al.</i>
DUNIETZ	98	PR D58 094010	I. Dunietz