

# b' (4<sup>th</sup> Generation) Quark, Searches for

## b'(-1/3)-quark/hadron mass limits in p-p̄ and pp collisions

| VALUE (GeV)   | CL% | DOCUMENT ID     | TECN      | COMMENT   |
|---|-----|-----------------|-----------|---|
| >1130   | 95  | 1 SIRUNYAN      | 19AQ CMS  | $B(b' \rightarrow Z b) = 1$                               |
| >1230   | 95  | 2 SIRUNYAN      | 19BWCMS   | $B(b' \rightarrow W t) = 1$                               |
| >1350   | 95  | 3 AABOUD        | 18AW ATLS | $B(b' \rightarrow W t) = 1$                               |
| >1000   | 95  | 4 AABOUD        | 18CE ATLS | $\geq 2\ell + \cancel{E}_T + \geq 1b_j$                   |
| > 950   | 95  | 5 AABOUD        | 18CL ATLS | $W t, Z b, h b$ modes                                     |
| >1010   | 95  | 6,7 AABOUD      | 18CP ATLS | 2,3 $\ell$ , singlet model                                |
| >1140   | 95  | 5,8 AABOUD      | 18CP ATLS | 2,3 $\ell$ , doublet model                                |
| >1220   | 95  | 9,10 AABOUD     | 18CR ATLS | singlet b'. ATLAS Combination                             |
| >1370   | 95  | 9,11 AABOUD     | 18CR ATLS | b' in a weak isospin doublet (t', b'). ATLAS combination. |
| > 910   | 95  | 12 SIRUNYAN     | 18BM CMS  | $W t, Z b, h b$ modes                                     |
| > 845   | 95  | 13 SIRUNYAN     | 18Q CMS   | $B(b' \rightarrow W u) = 1$                               |
| > 730   | 95  | 14 SIRUNYAN     | 17AU CMS  |   |
| > 880   | 95  | 15 KHACHATRY... | 16AN CMS  | $B(b' \rightarrow W t) = 1$                               |
| > 620   | 95  | 16 AAD          | 15BY ATLS | $W t, Z b, h b$ modes                                     |
| > 730   | 95  | 17 AAD          | 15BY ATLS | $B(b' \rightarrow W t) = 1$                               |
| > 810   | 95  | 18 AAD          | 15Z ATLS  |   |
| > 755   | 95  | 19 AAD          | 14AZ ATLS | $B(b' \rightarrow W t) = 1$                               |
| > 675   | 95  | 20 CHATRCHYAN   | 13I CMS   | $B(b' \rightarrow W t) = 1$                               |
| > 190   | 95  | 21 ABAZOV       | 08X D0    | $c\tau = 200\text{mm}$                                    |
| > 190   | 95  | 22 ACOSTA       | 03 CDF    | quasi-stable b'   |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |     |                 |           |   |
| <350, 580–635, >700   | 95  | 23 AAD          | 15AR ATLS | $B(b' \rightarrow H b) = 1$                               |
| > 690   | 95  | 24 AAD          | 15CN ATLS | $B(b' \rightarrow W q) = 1$ (q=u)                         |
| > 480   | 95  | 25 AAD          | 12AT ATLS | $B(b' \rightarrow W t) = 1$                               |
| > 400   | 95  | 26 AAD          | 12AU ATLS | $B(b' \rightarrow Z b) = 1$                               |
| > 350   | 95  | 27 AAD          | 12BC ATLS | $B(b' \rightarrow W q) = 1$<br>(q=u,c)                    |
| > 450   | 95  | 28 AAD          | 12BE ATLS | $B(b' \rightarrow W t) = 1$                               |
| > 685   | 95  | 29 CHATRCHYAN   | 12BH CMS  | $m_{t'} = m_{b'}$   |
| > 611   | 95  | 30 CHATRCHYAN   | 12X CMS   | $B(b' \rightarrow W t) = 1$                               |
| > 372   | 95  | 31 AALTONEN     | 11J CDF   | $b' \rightarrow W t$                                      |
| > 361   | 95  | 32 CHATRCHYAN   | 11L CMS   | Repl. by CHATRCHYAN 12X                                   |
| > 338   | 95  | 33 AALTONEN     | 10H CDF   | $b' \rightarrow W t$                                      |
| > 380–430   | 95  | 34 FLACCO       | 10 RVUE   | $m_{b'} > m_{t'}$   |
| > 268   | 95  | 35,36 AALTONEN  | 07C CDF   | $B(b' \rightarrow Z b) = 1$                               |
| > 199   | 95  | 37 AFFOLDER     | 00 CDF    | NC: $b' \rightarrow Z b$                                  |
| > 148   | 95  | 38 ABE          | 98N CDF   | NC: $b' \rightarrow Z b + \text{vertex}$                  |
| > 96  | 95  | 39 ABACHI       | 97D D0    | NC: $b' \rightarrow b\gamma$                              |

|       |    |    |             |     |      |  |
|-------|----|----|-------------|-----|------|--|
| > 128 | 95 | 40 | ABACHI      | 95F | D0   | $\ell\ell + \text{jets}, \ell + \text{jets}$ |
| > 75  | 95 | 41 | MUKHOPAD... | 93  | RVUE | NC: $b' \rightarrow b\ell\ell$               |
| > 85  | 95 | 42 | ABE         | 92  | CDF  | CC: $\ell\ell$                               |
| > 72  | 95 | 43 | ABE         | 90B | CDF  | CC: $e + \mu$                                |
| > 54  | 95 | 44 | AKESSON     | 90  | UA2  | CC: $e + \text{jets} + \cancel{E}_T$         |
| > 43  | 95 | 45 | ALBAJAR     | 90B | UA1  | CC: $\mu + \text{jets}$                      |
| > 34  | 95 | 46 | ALBAJAR     | 88  | UA1  | CC: $e \text{ or } \mu + \text{jets}$        |

<sup>1</sup> SIRUNYAN 19AQ based on  $35.9 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13 \text{ TeV}$ . Pair production of vector-like  $b'$  is searched for with one  $b'$  decaying into  $Zb$  and the other  $b'$  decaying into  $Wt, Zb, hb$ . Events with an opposite-sign lepton pair consistent with coming from  $Z$  and jets are used. Mass limits are obtained for a variety of branching ratios of  $b'$ .

<sup>2</sup> SIRUNYAN 19BW based on  $35.9 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13 \text{ TeV}$ . The limit is for the pair-produced vector-like  $b'$  using all-hadronic final state. The analysis is made for the  $Zb, Wt, hb$  modes and mass limits are obtained for a variety of branching ratios.

<sup>3</sup> AABOUD 18AW based on  $36.1 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13 \text{ TeV}$ . The limit is for the pair-produced vector-like  $b'$  using lepton-plus-jets final state. The search is also sensitive to the decays into  $Zb$  and  $Hb$  final states.

<sup>4</sup> AABOUD 18CE based on  $36.1 \text{ fb}^{-1}$  of proton-proton data taken at  $\sqrt{s} = 13 \text{ TeV}$ . Events including a same-sign lepton pair are used. The limit is for a singlet model, assuming the branching ratios of  $b'$  into  $Zb, Wt$  and  $Hb$  as predicted by the model.

<sup>5</sup> AABOUD 18CL, AABOUD 18CP based on  $36.1 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13 \text{ TeV}$ . The limit is for the pair-produced vector-like  $b'$  using all-hadronic final state. The analysis is particularly powerful for the  $b' \rightarrow hb$  mode. Assuming the pure decay only in this mode sets a limit  $m_{b'} > 1010 \text{ GeV}$ .

<sup>6</sup> AABOUD 18CP based on  $36.1 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13 \text{ TeV}$ . Pair and single production of vector-like  $b'$  are searched for with at least one  $b'$  decaying into  $Zb$ . In the case of  $B(b' \rightarrow Zb) = 1$ , the limit is  $m_{b'} > 1220 \text{ GeV}$ .

<sup>7</sup> The limit is for the singlet model, assuming that the branching ratios into  $Wt, Zb, hb$  add up to one.

<sup>8</sup> The limit is for the doublet model, assuming that the branching ratios into  $Wt, Zb, hb$  add up to one.

<sup>9</sup> AABOUD 18CR based on  $36.1 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13 \text{ TeV}$ . A combination of searches for the pair-produced vector-like  $b'$  in various decay channels ( $b' \rightarrow Wt, Zb, hb$ ). Also a model-independent limit is obtained as  $m_{b'} > 1.03 \text{ TeV}$ , assuming that the branching ratios into  $Zb, Wt$ , and  $hb$  add up to one.

<sup>10</sup> The limit is for the singlet  $b'$ .

<sup>11</sup> The limit is for  $b'$  in a weak isospin doublet ( $t', b'$ ) and  $|V_{t'b}| \ll |V_{tb}|$ . For a  $b'$  in a doublet with a charge  $-4/3$  vector-like quark, the limit  $m_{b'} > 1.14 \text{ TeV}$  is obtained.

<sup>12</sup> SIRUNYAN 18BM based on  $35.9 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13 \text{ TeV}$ . The limit is for the pair-produced vector-like  $b'$ . Three channels (single lepton, same-charge 2 leptons, or at least 3 leptons) are considered for various branching fraction combinations. Assuming  $B(tW) = 1$ , the limit is  $1240 \text{ GeV}$  and for  $B(bZ) = 1$  it is  $960 \text{ GeV}$ .

<sup>13</sup> SIRUNYAN 18Q based on  $19.7 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 8 \text{ TeV}$ . The limit is for the pair-produced vector-like  $b'$  that couple only to light quarks. Upper cross section limits on the single production of a  $b'$  and constraints for other decay channels ( $Zq$  and  $Hq$ ) are also given in the paper.

<sup>14</sup> SIRUNYAN 17AU based on  $2.3\text{--}2.6 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13 \text{ TeV}$ . Limit on pair-produced singlet vector-like  $b'$  using one lepton and several jets. The mass bound is given for a  $b'$  transforming as a singlet under the electroweak symmetry group, assumed

- to decay through  $W$ ,  $Z$  or Higgs boson (which decays to jets) and to a third generation quark.
- 15 KHACHATRYAN 16AN based on  $19.7 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 8 \text{ TeV}$ . Limit on pair-produced vector-like  $b'$  using 1, 2, and  $>2$  leptons as well as fully hadronic final states. Other limits depending on the branching fractions to  $tW$ ,  $bZ$ , and  $bH$  are given in Table IX.
  - 16 AAD 15BY based on  $20.3 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 8 \text{ TeV}$ . Limit on pair-produced vector-like  $b'$  assuming the branching fractions to  $W$ ,  $Z$ , and  $h$  modes of the singlet model. Used events containing  $\geq 2\ell + \cancel{E}_T + \geq 2j$  ( $\geq 1 b$ ) and including a same-sign lepton pair.
  - 17 AAD 15BY based on  $20.3 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 8 \text{ TeV}$ . Limit on pair-produced chiral  $b'$ -quark. Used events containing  $\geq 2\ell + \cancel{E}_T + \geq 2j$  ( $\geq 1 b$ ) and including a same-sign lepton pair.
  - 18 AAD 15Z based on  $20.3 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 8 \text{ TeV}$ . Used events with  $\ell + \cancel{E}_T + \geq 6j$  ( $\geq 1 b$ ) and at least one pair of jets from weak boson decay, primarily designed to select the signature  $b'\bar{b}' \rightarrow WWt\bar{t} \rightarrow WWWWb\bar{b}$ . This is a limit on pair-produced vector-like  $b'$ . The lower mass limit is 640 GeV for a vector-like singlet  $b'$ .
  - 19 Based on  $20.3 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 8 \text{ TeV}$ . No significant excess over SM expectation is found in the search for pair production or single production of  $b'$  in the events with dilepton from a high  $p_T Z$  and additional jets ( $\geq 1 b$ -tag). If instead of  $B(b' \rightarrow Wt) = 1$  an electroweak singlet with  $B(b' \rightarrow Wt) \sim 0.45$  is assumed, the limit reduces to 685 GeV.
  - 20 Based on  $5.0 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 7 \text{ TeV}$ . CHATRCHYAN 13l looked for events with one isolated electron or muon, large  $\cancel{E}_T$ , and at least four jets with large transverse momenta, where one jet is likely to originate from the decay of a bottom quark.
  - 21 Result is based on  $1.1 \text{ fb}^{-1}$  of data. No signal is found for the search of long-lived particles which decay into final states with two electrons or photons, and upper bound on the cross section times branching fraction is obtained for  $2 < c\tau < 7000 \text{ fm}$ ; see Fig. 3. 95% CL excluded region of  $b'$  lifetime and mass is shown in Fig. 4.
  - 22 ACOSTA 03 looked for long-lived fourth generation quarks in the data sample of  $90 \text{ pb}^{-1}$  of  $\sqrt{s}=1.8 \text{ TeV } p\bar{p}$  collisions by using the muon-like penetration and anomalously high ionization energy loss signature. The corresponding lower mass bound for the charge  $(2/3)e$  quark ( $t'$ ) is 220 GeV. The  $t'$  bound is higher than the  $b'$  bound because  $t'$  is more likely to produce charged hadrons than  $b'$ . The 95% CL upper bounds for the production cross sections are given in their Fig. 3.
  - 23 AAD 15AR based on  $20.3 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 8 \text{ TeV}$ . Used lepton-plus-jets final state. See Fig. 24 for mass limits in the plane of  $B(b' \rightarrow Wt)$  vs.  $B(b' \rightarrow Hb)$  from  $b'\bar{b}' \rightarrow Hb + X$  searches.
  - 24 AAD 15CN based on  $20.3 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 8 \text{ TeV}$ . Limit on pair-production of chiral  $b'$ -quark. Used events with  $\ell + \cancel{E}_T + \geq 4j$  (non- $b$ -tagged). Limits on a heavy vector-like quark, which decays into  $Wq$ ,  $Zq$ ,  $hq$ , are presented in the plane  $B(Q \rightarrow Wq)$  vs.  $B(Q \rightarrow hq)$  in Fig. 12.
  - 25 Based on  $1.04 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 7 \text{ TeV}$ . No signal is found for the search of heavy quark pair production that decay into  $W$  and a  $t$  quark in the events with a high  $p_T$  isolated lepton, large  $\cancel{E}_T$ , and at least 6 jets in which one, two or more dijets are from  $W$ .
  - 26 Based on  $2.0 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 7 \text{ TeV}$ . No  $b' \rightarrow Zb$  invariant mass peak is found in the search of heavy quark pair production that decay into  $Z$  and a  $b$  quark in events with  $Z \rightarrow e^+e^-$  and at least one  $b$ -jet. The lower mass limit is 358 GeV for a vector-like singlet  $b'$  mixing solely with the third SM generation.
  - 27 Based on  $1.04 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 7 \text{ TeV}$ . No signal is found for the search of heavy quark pair production that decay into  $W$  and a quark in the events with dileptons, large  $\cancel{E}_T$ , and  $\geq 2$  jets.

- <sup>28</sup> Based on  $1.04 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 7 \text{ TeV}$ . AAD 12BE looked for events with two isolated like-sign leptons and at least 2 jets, large  $\cancel{E}_T$  and  $H_T > 350 \text{ GeV}$ .
- <sup>29</sup> Based on  $5 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 7 \text{ TeV}$ . CHATRCHYAN 12BH searched for QCD and EW production of single and pair of degenerate 4'th generation quarks that decay to  $bW$  or  $tW$ . Absence of signal in events with one lepton, same-sign dileptons or tripleptons gives the bound. With a mass difference of  $25 \text{ GeV}/c^2$  between  $m_{t'}$  and  $m_{b'}$ , the corresponding limit shifts by about  $\pm 20 \text{ GeV}/c^2$ .
- <sup>30</sup> Based on  $4.9 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 7 \text{ TeV}$ . CHATRCHYAN 12X looked for events with tripleptons or same-sign dileptons and at least one  $b$  jet.
- <sup>31</sup> Based on  $4.8 \text{ fb}^{-1}$  of data in  $p\bar{p}$  collisions at  $1.96 \text{ TeV}$ . AALTONEN 11J looked for events with  $\ell + \cancel{E}_T + \geq 5j$  ( $\geq 1$   $b$  or  $c$ ). No signal is observed and the bound  $\sigma(b'\bar{b}') < 30 \text{ fb}$  for  $m_{b'} > 375 \text{ GeV}$  is found for  $B(b' \rightarrow Wt) = 1$ .
- <sup>32</sup> Based on  $34 \text{ pb}^{-1}$  of data in  $pp$  collisions at  $7 \text{ TeV}$ . CHATRCHYAN 11L looked for multi-jet events with tripleptons or same-sign dileptons. No excess above the SM background excludes  $m_{b'}$  between 255 and 361 GeV at 95% CL for  $B(b' \rightarrow Wt) = 1$ .
- <sup>33</sup> Based on  $2.7 \text{ fb}^{-1}$  of data in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96 \text{ TeV}$ . AALTONEN 10H looked for pair production of heavy quarks which decay into  $tW^-$  or  $tW^+$ , in events with same sign dileptons ( $e$  or  $\mu$ ), several jets and large missing  $E_T$ . The result is obtained for  $b'$  which decays into  $tW^-$ . For the charge  $5/3$  quark ( $T_{5/3}$ ) which decays into  $tW^+$ ,  $m_{T_{5/3}} > 365 \text{ GeV}$  (95% CL) is found when it has the charge  $-1/3$  partner  $B$  of the same mass.
- <sup>34</sup> FLACCO 10 result is obtained from AALTONEN 10H result of  $m_{b'} > 338 \text{ GeV}$ , by relaxing the condition  $B(b' \rightarrow Wt) = 100\%$  when  $m_{b'} > m_{t'}$ .
- <sup>35</sup> Result is based on  $1.06 \text{ fb}^{-1}$  of data. No excess from the SM  $Z$ +jet events is found when  $Z$  decays into  $ee$  or  $\mu\mu$ . The  $m_{b'}$  bound is found by comparing the resulting upper bound on  $\sigma(b'\bar{b}') [1-(1-B(b' \rightarrow Zb))^2]$  and the LO estimate of the  $b'$  pair production cross section shown in Fig. 38 of the article.
- <sup>36</sup> HUANG 08 reexamined the  $b'$  mass lower bound of 268 GeV obtained in AALTONEN 07C that assumes  $B(b' \rightarrow Zb) = 1$ , which does not hold for  $m_{b'} > 255 \text{ GeV}$ . The lower mass bound is given in the plane of  $\sin^2(\theta_{tb'})$  and  $m_{b'}$ .
- <sup>37</sup> AFFOLDER 00 looked for  $b'$  that decays into  $b+Z$ . The signal searched for is  $bbZZ$  events where one  $Z$  decays into  $e^+e^-$  or  $\mu^+\mu^-$  and the other  $Z$  decays hadronically. The bound assumes  $B(b' \rightarrow Zb) = 100\%$ . Between 100 GeV and 199 GeV, the 95%CL upper bound on  $\sigma(b' \rightarrow \bar{b}') \times B^2(b' \rightarrow Zb)$  is also given (see their Fig. 2).
- <sup>38</sup> ABE 98N looked for  $Z \rightarrow e^+e^-$  decays with displaced vertices. Quoted limit assumes  $B(b' \rightarrow Zb) = 1$  and  $c\tau_{b'} = 1 \text{ cm}$ . The limit is lower than  $m_Z + m_b$  ( $\sim 96 \text{ GeV}$ ) if  $c\tau > 22 \text{ cm}$  or  $c\tau < 0.009 \text{ cm}$ . See their Fig. 4.
- <sup>39</sup> ABACHI 97D searched for  $b'$  that decays mainly via FCNC. They obtained 95%CL upper bounds on  $B(b'\bar{b}' \rightarrow \gamma + 3 \text{ jets})$  and  $B(b'\bar{b}' \rightarrow 2\gamma + 2 \text{ jets})$ , which can be interpreted as the lower mass bound  $m_{b'} > m_Z + m_b$ .
- <sup>40</sup> ABACHI 95F bound on the top-quark also applies to  $b'$  and  $t'$  quarks that decay predominantly into  $W$ . See FROGGATT 97.
- <sup>41</sup> MUKHOPADHYAYA 93 analyze CDF dilepton data of ABE 92G in terms of a new quark decaying via flavor-changing neutral current. The above limit assumes  $B(b' \rightarrow b\ell^+\ell^-) = 1\%$ . For an exotic quark decaying only via virtual  $Z$  [ $B(b\ell^+\ell^-) = 3\%$ ], the limit is 85 GeV.
- <sup>42</sup> ABE 92 dilepton analysis limit of  $>85 \text{ GeV}$  at CL=95% also applies to  $b'$  quarks, as discussed in ABE 90B.
- <sup>43</sup> ABE 90B exclude the region 28–72 GeV.

- <sup>44</sup> AKESSON 90 searched for events having an electron with  $p_T > 12$  GeV, missing momentum  $> 15$  GeV, and a jet with  $E_T > 10$  GeV,  $|\eta| < 2.2$ , and excluded  $m_{b'}$  between 30 and 69 GeV.
- <sup>45</sup> For the reduction of the limit due to non-charged-current decay modes, see Fig. 19 of ALBAJAR 90B.
- <sup>46</sup> ALBAJAR 88 study events at  $E_{cm} = 546$  and 630 GeV with a muon or isolated electron, accompanied by one or more jets and find agreement with Monte Carlo predictions for the production of charm and bottom, without the need for a new quark. The lower mass limit is obtained by using a conservative estimate for the  $b'\bar{b}'$  production cross section and by assuming that it cannot be produced in  $W$  decays. The value quoted here is revised using the full  $O(\alpha_s^3)$  cross section of ALTARELLI 88.

### $b'(-1/3)$ mass limits from single production in $p\bar{p}$ and $pp$ collisions

| VALUE (GeV)     | CL% | DOCUMENT ID                  | TECN      | COMMENT   |
|-----------------|-----|------------------------------|-----------|---|
| >1500           | 95  | <sup>1</sup> AAD             | 16AH ATLS | $gb \rightarrow b' \rightarrow tW, B(b' \rightarrow tW)=1$  |
| >1390           | 95  | <sup>2</sup> KHACHATRY...16I | CMS       | $gb \rightarrow b' \rightarrow tW, B(b' \rightarrow tW)=1$  |
| >1430           | 95  | <sup>3</sup> KHACHATRY...16I | CMS       | $gb \rightarrow b' \rightarrow tW, B(b' \rightarrow tW)=1$  |
| <b>&gt;1530</b> | 95  | <sup>4</sup> KHACHATRY...16I | CMS       | $gb \rightarrow b' \rightarrow tW, B(b' \rightarrow tW)=1$  |
| <b>&gt; 693</b> | 95  | <sup>5</sup> ABAZOV          | 11F D0    | $qu \rightarrow q'b' \rightarrow q'(Wu)$<br>$\tilde{\kappa}_{ub'}=1, B(b' \rightarrow Wu)=1$      |
| <b>&gt; 430</b> | 95  | <sup>5</sup> ABAZOV          | 11F D0    | $qd \rightarrow qb' \rightarrow q(Zd)$<br>$\tilde{\kappa}_{db'}=\sqrt{2}, B(b' \rightarrow Zd)=1$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>6</sup> SIRUNYAN 19AI CMS  $bZ/tW \rightarrow b' \rightarrow tW$

<sup>1</sup> AAD 16AH based on  $20.3 \text{ fb}^{-1}$  of data in  $pp$  collisions at 8 TeV. No significant excess over SM expectation is found in the search for a vector-like  $b'$  in the single-lepton and dilepton channels ( $\ell$  or  $\ell\ell$ ) + 1,2,3  $j$  ( $\geq 1b$ ). The model assumes that the  $b'$  has the excited quark couplings.

<sup>2</sup> Based on  $19.7 \text{ fb}^{-1}$  of data in  $pp$  collisions at 8 TeV. Limit on left-handed  $b'$  assuming 100% decay to  $tW$  and using all-hadronic, lepton + jets, and dilepton final states.

<sup>3</sup> Based on  $19.7 \text{ fb}^{-1}$  of data in  $pp$  collisions at 8 TeV. Limit on right-handed  $b'$  assuming 100% decay to  $tW$  and using all-hadronic, lepton + jets, and dilepton final states.

<sup>4</sup> Based on  $19.7 \text{ fb}^{-1}$  of data in  $pp$  collisions at 8 TeV. Limit on vector-like  $b'$  assuming 100% decay to  $tW$  and using all-hadronic, lepton+jets, and dilepton final states.

<sup>5</sup> Based on  $5.4 \text{ fb}^{-1}$  of data in  $p\bar{p}$  collisions at 1.96 TeV. ABAZOV 11F looked for single production of  $b'$  via the  $W$  or  $Z$  coupling to the first generation up or down quarks, respectively. Model independent cross section limits for the single production processes  $p\bar{p} \rightarrow b'q \rightarrow Wuq$ , and  $p\bar{p} \rightarrow b'q \rightarrow Zdq$  are given in Figs. 3 and 4, respectively, and the mass limits are obtained for the model of ATRE 09 with degenerate bi-doublets of vector-like quarks.

<sup>6</sup> SIRUNYAN 19AI based on  $35.9 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13$  TeV. Exclusion limits are set on the product of the production cross section and branching fraction for the  $b'(-1/3) + b$  and  $b'(-1/3) + t$  modes as a function of the vector-like quark mass in Figs. 7 and 8 and in Tab. 2 for relative vector-like quark widths between 1 and 30% for left- and right-handed vector-like quark couplings. No significant deviation from the SM prediction is observed.

**MASS LIMITS for  $b'$  (4<sup>th</sup> Generation) Quark or Hadron in  $e^+e^-$  Collisions**Search for hadrons containing a fourth-generation  $-1/3$  quark denoted  $b'$ .The last column specifies the assumption for the decay mode ( $CC$  denotes the conventional charged-current decay) and the event signature which is looked for.

| VALUE (GeV)   | CL% | DOCUMENT ID             | TECN     | COMMENT   |
|---|-----|-------------------------|----------|---|
| <b>&gt;46.0</b>   | 95  | <sup>1</sup> DECAMP     | 90F ALEP | any decay   |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |     |                         |          |   |
| none 96–103   | 95  | <sup>2</sup> ABDALLAH   | 07 DLPH  | $b' \rightarrow bZ, cW$                                     |
|   |     | <sup>3</sup> ADRIANI    | 93G L3   | Quarkonium  |
| >44.7   | 95  | ADRIANI                 | 93M L3   | $\Gamma(Z)$   |
| >45   | 95  | ABREU                   | 91F DLPH | $\Gamma(Z)$   |
| none 19.4–28.2  | 95  | ABE                     | 90D VNS  | Any decay; event shape                                      |
| >45.0   | 95  | ABREU                   | 90D DLPH | $B(CC) = 1$ ; event shape                                   |
| >44.5   | 95  | <sup>4</sup> ABREU      | 90D DLPH | $b' \rightarrow cH^-, H^- \rightarrow \bar{c}s, \tau^- \nu$ |
| >40.5   | 95  | <sup>5</sup> ABREU      | 90D DLPH | $\Gamma(Z \rightarrow \text{hadrons})$                      |
| >28.3   | 95  | ADACHI                  | 90 TOPZ  | $B(\text{FCNC})=100\%$ ; isol. $\gamma$ or 4 jets           |
| >41.4   | 95  | <sup>6</sup> AKRAWY     | 90B OPAL | Any decay; acoplanarity                                     |
| >45.2   | 95  | <sup>6</sup> AKRAWY     | 90B OPAL | $B(CC) = 1$ ; acoplanarity                                  |
| >46   | 95  | <sup>7</sup> AKRAWY     | 90J OPAL | $b' \rightarrow \gamma + \text{any}$                        |
| >27.5   | 95  | <sup>8</sup> ABE        | 89E VNS  | $B(CC) = 1$ ; $\mu, e$                                      |
| none 11.4–27.3  | 95  | <sup>9</sup> ABE        | 89G VNS  | $B(b' \rightarrow b\gamma) > 10\%$ ; isolated $\gamma$      |
| >44.7   | 95  | <sup>10</sup> ABRAMS    | 89C MRK2 | $B(CC) = 100\%$ ; isol. track                               |
| >42.7   | 95  | <sup>10</sup> ABRAMS    | 89C MRK2 | $B(bg) = 100\%$ ; event shape                               |
| >42.0   | 95  | <sup>10</sup> ABRAMS    | 89C MRK2 | Any decay; event shape                                      |
| >28.4   | 95  | <sup>11,12</sup> ADACHI | 89C TOPZ | $B(CC) = 1$ ; $\mu$   |
| >28.8   | 95  | <sup>13</sup> ENO       | 89 AMY   | $B(CC) \gtrsim 90\%$ ; $\mu, e$                             |
| >27.2   | 95  | <sup>13,14</sup> ENO    | 89 AMY   | any decay; event shape                                      |
| >29.0   | 95  | <sup>13</sup> ENO       | 89 AMY   | $B(b' \rightarrow bg) \gtrsim 85\%$ ; event shape           |
| >24.4   | 95  | <sup>15</sup> IGARASHI  | 88 AMY   | $\mu, e$  |
| >23.8   | 95  | <sup>16</sup> SAGAWA    | 88 AMY   | event shape   |
| >22.7   | 95  | <sup>17</sup> ADEVA     | 86 MRKJ  | $\mu$   |
| >21   |     | <sup>18</sup> ALTHOFF   | 84C TASS | $R$ , event shape   |
| >19   |     | <sup>19</sup> ALTHOFF   | 84I TASS | Aplanarity  |

<sup>1</sup> DECAMP 90F looked for isolated charged particles, for isolated photons, and for four-jet final states. The modes  $b' \rightarrow bg$  for  $B(b' \rightarrow bg) > 65\%$   $b' \rightarrow b\gamma$  for  $B(b' \rightarrow b\gamma) > 5\%$  are excluded. Charged Higgs decay were not discussed.

<sup>2</sup> ABDALLAH 07 searched for  $b'$  pair production at  $E_{\text{cm}}=196\text{--}209$  GeV, with  $420 \text{ pb}^{-1}$ . No signal leads to the 95% CL upper limits on  $B(b' \rightarrow bZ)$  and  $B(b' \rightarrow cW)$  for  $m_{b'} = 96$  to  $103$  GeV.

<sup>3</sup> ADRIANI 93G search for vector quarkonium states near  $Z$  and give limit on quarkonium- $Z$  mixing parameter  $\delta m^2 < (10\text{--}30) \text{ GeV}^2$  (95%CL) for the mass  $88\text{--}94.5$  GeV. Using Richardson potential, a  $1S (b'\bar{b}')$  state is excluded for the mass range  $87.7\text{--}94.7$  GeV. This range depends on the potential choice.

<sup>4</sup> ABREU 90D assumed  $m_{H^-} < m_{b'} - 3$  GeV.

- <sup>5</sup> Superseded by ABREU 91F.
- <sup>6</sup> AKRAWY 90B search was restricted to data near the  $Z$  peak at  $E_{\text{cm}} = 91.26$  GeV at LEP. The excluded region is between 23.6 and 41.4 GeV if no  $H^+$  decays exist. For charged Higgs decays the excluded regions are between  $(m_{H^+} + 1.5 \text{ GeV})$  and 45.5 GeV.
- <sup>7</sup> AKRAWY 90J search for isolated photons in hadronic  $Z$  decay and derive  $B(Z \rightarrow b'\bar{b}') \cdot B(b' \rightarrow \gamma X) / B(Z \rightarrow \text{hadrons}) < 2.2 \times 10^{-3}$ . Mass limit assumes  $B(b' \rightarrow \gamma X) > 10\%$ .
- <sup>8</sup> ABE 89E search at  $E_{\text{cm}} = 56\text{--}57$  GeV at TRISTAN for multihadron events with a spherical shape (using thrust and acoplanarity) or containing isolated leptons.
- <sup>9</sup> ABE 89G search was at  $E_{\text{cm}} = 55\text{--}60.8$  GeV at TRISTAN.
- <sup>10</sup> If the photonic decay mode is large ( $B(b' \rightarrow b\gamma) > 25\%$ ), the ABRAMS 89C limit is 45.4 GeV. The limit for Higgs decay ( $b' \rightarrow cH^-, H^- \rightarrow \bar{c}s$ ) is 45.2 GeV.
- <sup>11</sup> ADACHI 89C search was at  $E_{\text{cm}} = 56.5\text{--}60.8$  GeV at TRISTAN using multi-hadron events accompanying muons.
- <sup>12</sup> ADACHI 89C also gives limits for any mixture of  $CC$  and  $bg$  decays.
- <sup>13</sup> ENO 89 search at  $E_{\text{cm}} = 50\text{--}60.8$  at TRISTAN.
- <sup>14</sup> ENO 89 considers arbitrary mixture of the charged current,  $bg$ , and  $b\gamma$  decays.
- <sup>15</sup> IGARASHI 88 searches for leptons in low-thrust events and gives  $\Delta R(b') < 0.26$  (95% CL) assuming charged current decay, which translates to  $m_{b'} > 24.4$  GeV.
- <sup>16</sup> SAGAWA 88 set limit  $\sigma(\text{top}) < 6.1$  pb at CL=95% for top-flavored hadron production from event shape analyses at  $E_{\text{cm}} = 52$  GeV. By using the quark parton model cross-section formula near threshold, the above limit leads to lower mass bounds of 23.8 GeV for charge  $-1/3$  quarks.
- <sup>17</sup> ADEVA 86 give 95%CL upper bound on an excess of the normalized cross section,  $\Delta R$ , as a function of the minimum c.m. energy (see their figure 3). Production of a pair of  $1/3$  charge quarks is excluded up to  $E_{\text{cm}} = 45.4$  GeV.
- <sup>18</sup> ALTHOFF 84C narrow state search sets limit  $\Gamma(e^+e^-)B(\text{hadrons}) < 2.4$  keV CL = 95% and heavy charge  $1/3$  quark pair production  $m > 21$  GeV, CL = 95%.
- <sup>19</sup> ALTHOFF 84i exclude heavy quark pair production for  $7 < m < 19$  GeV ( $1/3$  charge) using aplanarity distributions (CL = 95%).

---

## REFERENCES FOR Searches for (Fourth Generation) $b'$ Quark

|              |      |                |                              |                 |
|--------------|------|----------------|------------------------------|-----------------|
| SIRUNYAN     | 19AI | EPJ C79 90     | A.M. Sirunyan <i>et al.</i>  | (CMS Collab.)   |
| SIRUNYAN     | 19AQ | EPJ C79 364    | A.M. Sirunyan <i>et al.</i>  | (CMS Collab.)   |
| SIRUNYAN     | 19BW | PR D100 072001 | A.M. Sirunyan <i>et al.</i>  | (CMS Collab.)   |
| AABOUD       | 18AW | JHEP 1808 048  | M. Aaboud <i>et al.</i>      | (ATLAS Collab.) |
| AABOUD       | 18CE | JHEP 1812 039  | M. Aaboud <i>et al.</i>      | (ATLAS Collab.) |
| AABOUD       | 18CL | PR D98 092005  | M. Aaboud <i>et al.</i>      | (ATLAS Collab.) |
| AABOUD       | 18CP | PR D98 112010  | M. Aaboud <i>et al.</i>      | (ATLAS Collab.) |
| AABOUD       | 18CR | PRL 121 211801 | M. Aaboud <i>et al.</i>      | (ATLAS Collab.) |
| SIRUNYAN     | 18BM | JHEP 1808 177  | A.M. Sirunyan <i>et al.</i>  | (CMS Collab.)   |
| SIRUNYAN     | 18Q  | PR D97 072008  | A.M. Sirunyan <i>et al.</i>  | (CMS Collab.)   |
| SIRUNYAN     | 17AU | JHEP 1711 085  | A.M. Sirunyan <i>et al.</i>  | (CMS Collab.)   |
| AAD          | 16AH | JHEP 1602 110  | G. Aad <i>et al.</i>         | (ATLAS Collab.) |
| KHACHATRY... | 16AN | PR D93 112009  | V. Khachatryan <i>et al.</i> | (CMS Collab.)   |
| KHACHATRY... | 16I  | JHEP 1601 166  | V. Khachatryan <i>et al.</i> | (CMS Collab.)   |
| AAD          | 15AR | JHEP 1508 105  | G. Aad <i>et al.</i>         | (ATLAS Collab.) |
| AAD          | 15BY | JHEP 1510 150  | G. Aad <i>et al.</i>         | (ATLAS Collab.) |
| AAD          | 15CN | PR D92 112007  | G. Aad <i>et al.</i>         | (ATLAS Collab.) |
| AAD          | 15Z  | PR D91 112011  | G. Aad <i>et al.</i>         | (ATLAS Collab.) |
| AAD          | 14AZ | JHEP 1411 104  | G. Aad <i>et al.</i>         | (ATLAS Collab.) |
| CHATRCHYAN   | 13I  | JHEP 1301 154  | S. Chatrchyan <i>et al.</i>  | (CMS Collab.)   |
| AAD          | 12AT | PRL 109 032001 | G. Aad <i>et al.</i>         | (ATLAS Collab.) |
| AAD          | 12AU | PRL 109 071801 | G. Aad <i>et al.</i>         | (ATLAS Collab.) |
| AAD          | 12BC | PR D86 012007  | G. Aad <i>et al.</i>         | (ATLAS Collab.) |
| AAD          | 12BE | JHEP 1204 069  | G. Aad <i>et al.</i>         | (ATLAS Collab.) |

|             |      |                |   |                   |
|-------------|------|----------------|---|-------------------|
| CHATRCHYAN  | 12BH | PR D86 112003  | S. Chatrchyan <i>et al.</i>             | (CMS Collab.)     |
| CHATRCHYAN  | 12X  | JHEP 1205 123  | S. Chatrchyan <i>et al.</i>             | (CMS Collab.)     |
| AALTONEN    | 11J  | PRL 106 141803 | T. Aaltonen <i>et al.</i>               | (CDF Collab.)     |
| ABAZOV      | 11F  | PRL 106 081801 | V.M. Abazov <i>et al.</i>               | (D0 Collab.)      |
| CHATRCHYAN  | 11L  | PL B701 204    | S. Chatrchyan <i>et al.</i>             | (CMS Collab.)     |
| AALTONEN    | 10H  | PRL 104 091801 | T. Aaltonen <i>et al.</i>               | (CDF Collab.)     |
| FLACCO      | 10   | PRL 105 111801 | C.J. Flacco <i>et al.</i>               | (UCI, HAIF)       |
| ATRE        | 09   | PR D79 054018  | A. Atre <i>et al.</i>                   |                   |
| ABAZOV      | 08X  | PRL 101 111802 | V.M. Abazov <i>et al.</i>               | (D0 Collab.)      |
| HUANG       | 08   | PR D77 037302  | P.Q. Hung, M. Sher                      | (UVA, WILL)       |
| AALTONEN    | 07C  | PR D76 072006  | T. Aaltonen <i>et al.</i>               | (CDF Collab.)     |
| ABDALLAH    | 07   | EPJ C50 507    | J. Abdallah <i>et al.</i>               | (DELPHI Collab.)  |
| ACOSTA      | 03   | PRL 90 131801  | D. Acosta <i>et al.</i>                 | (CDF Collab.)     |
| AFFOLDER    | 00   | PRL 84 835     | A. Affolder <i>et al.</i>               | (CDF Collab.)     |
| ABE         | 98N  | PR D58 051102  | F. Abe <i>et al.</i>                    | (CDF Collab.)     |
| ABACHI      | 97D  | PRL 78 3818    | S. Abachi <i>et al.</i>                 | (D0 Collab.)      |
| FROGGATT    | 97   | ZPHY C73 333   | C.D. Froggatt, D.J. Smith, H.B. Nielsen | (GLAS+)           |
| ABACHI      | 95F  | PR D52 4877    | S. Abachi <i>et al.</i>                 | (D0 Collab.)      |
| ADRIANI     | 93G  | PL B313 326    | O. Adriani <i>et al.</i>                | (L3 Collab.)      |
| ADRIANI     | 93M  | PRPL 236 1     | O. Adriani <i>et al.</i>                | (L3 Collab.)      |
| MUKHOPAD... | 93   | PR D48 2105    | B. Mukhopadhyaya, D.P. Roy              | (TATA)            |
| ABE         | 92   | PRL 68 447     | F. Abe <i>et al.</i>                    | (CDF Collab.)     |
| Also        |      | PR D45 3921    | F. Abe <i>et al.</i>                    | (CDF Collab.)     |
| ABE         | 92G  | PR D45 3921    | F. Abe <i>et al.</i>                    | (CDF Collab.)     |
| ABREU       | 91F  | NP B367 511    | P. Abreu <i>et al.</i>                  | (DELPHI Collab.)  |
| ABE         | 90B  | PRL 64 147     | F. Abe <i>et al.</i>                    | (CDF Collab.)     |
| ABE         | 90D  | PL B234 382    | K. Abe <i>et al.</i>                    | (VENUS Collab.)   |
| ABREU       | 90D  | PL B242 536    | P. Abreu <i>et al.</i>                  | (DELPHI Collab.)  |
| ADACHI      | 90   | PL B234 197    | I. Adachi <i>et al.</i>                 | (TOPAZ Collab.)   |
| AKESSON     | 90   | ZPHY C46 179   | T. Akesson <i>et al.</i>                | (UA2 Collab.)     |
| AKRAWY      | 90B  | PL B236 364    | M.Z. Akrawy <i>et al.</i>               | (OPAL Collab.)    |
| AKRAWY      | 90J  | PL B246 285    | M.Z. Akrawy <i>et al.</i>               | (OPAL Collab.)    |
| ALBAJAR     | 90B  | ZPHY C48 1     | C. Albajar <i>et al.</i>                | (UA1 Collab.)     |
| DECAMP      | 90F  | PL B236 511    | D. Decamp <i>et al.</i>                 | (ALEPH Collab.)   |
| ABE         | 89E  | PR D39 3524    | K. Abe <i>et al.</i>                    | (VENUS Collab.)   |
| ABE         | 89G  | PRL 63 1776    | K. Abe <i>et al.</i>                    | (VENUS Collab.)   |
| ABRAMS      | 89C  | PRL 63 2447    | G.S. Abrams <i>et al.</i>               | (Mark II Collab.) |
| ADACHI      | 89C  | PL B229 427    | I. Adachi <i>et al.</i>                 | (TOPAZ Collab.)   |
| ENO         | 89   | PRL 63 1910    | S. Eno <i>et al.</i>                    | (AMY Collab.)     |
| ALBAJAR     | 88   | ZPHY C37 505   | C. Albajar <i>et al.</i>                | (UA1 Collab.)     |
| ALTARELLI   | 88   | NP B308 724    | G. Altarelli <i>et al.</i>              | (CERN, ROMA, ETH) |
| IGARASHI    | 88   | PRL 60 2359    | S. Igarashi <i>et al.</i>               | (AMY Collab.)     |
| SAGAWA      | 88   | PRL 60 93      | H. Sagawa <i>et al.</i>                 | (AMY Collab.)     |
| ADEVA       | 86   | PR D34 681     | B. Adeva <i>et al.</i>                  | (Mark-J Collab.)  |
| ALTHOFF     | 84C  | PL 138B 441    | M. Althoff <i>et al.</i>                | (TASSO Collab.)   |
| ALTHOFF     | 84I  | ZPHY C22 307   | M. Althoff <i>et al.</i>                | (TASSO Collab.)   |